Demonstrations

H.M.

Dekking

Groningen

1. Some Cases of Vascular Anomalies.

A man with markedly distended conjunctival vessels in a sharply demarcated area. No changes during four years' observation.

A child with naevus flammeus and buphthalmos.

A new form of phacomatosis? A girl, aged 14 years, with dilated vessels in the left upper eyelid and arteriovenous aneurysm in the retina of the same eye. The latter is remarkable on account of the spontaneous regression: in the course of about one year the markedly dilated artery and vein were restored to their normal width, so that the connection is hardly to be seen. In the meantime other arterio-venous aneurysms have developed. The rest of the retina is entirely normal. The visual acuity is 6/6.

Vascular anomalies in the retina in dystrophia adiposogenitalis. Throughout the fundus there are vessels which show a white sheath over a short distance. In the periphery there is a large crescent-shaped white spot, looking like a retinal exudate. The retinal vessels running over and near this spot show aneurysms, as is the case in Coats' disease. However, there is no further sign of this affection: the appearance of the vascular sheaths and the white spot has not changed for a year.

2. Familial Macular Degeneration in Two Generations. In some members of this family the affection manifests itself as a whitish-yellow spot in the macula. In two others this spot is connected with the papilla by a white cord. In another member the affection set in as a typical Kuhnt-Julius disciform macular degeneration. All examinations, including those for toxoplasmosis, were negative.

D e k k i n g , Demonstrations

81

Kaiser-Fleischer's Ring in Wilson's Disease.

Ulcus Rodens Corneae (?). Microscopy of the enucleated eye did not yield any further information.

Congenital Ptosis in Four Generations.

Discussion.

Rochat: The pathological course and the aspect of the repeatedly recurring conjunctival ulcer slightly suggest self-mutilation by some chemical substance as the cause. In this connection, Leber's conjunctivitis pelrificans will be remembered, which, after having been regarded as a disease for some years, finally appeared to be the result of automutilation.

Flieringa: (a) The recurrent ulcerous conjunctivitis in the patient presented suggested an artefact to me, in agreement with Rochat. I remember that Weve demonstrated the patient a few years ago and that he had found a great number of fungus threads. Has Dekking also searched for them in this patient?

(b) Has the patient with dilated pericorneal vessels been examined gonio-scopically? In cases of aneurysm of the conjunctival vessels, dilated vessels were often observed in the angulus iridis.

Dekking replied

to Rochat: I do not believe that the fourth patient is a case of automutilation. In the first place, we have observed the patient in the clinic for a long time and very carefully. In the second place, she does not give the impression of being hysterical. Further, the other eye was quite normal;

to Flieringa: With regard to the same patient: Fungi were not found in spite of repeated bacteriological examinations. Gonioscopy was not carried out in patient Ia). In your case there were aneurysms, in our case only distended vessels.

Clinical Contribution to the Knowledge of the Course of the Nerve Fibres in the Retina.

By A. C. COPPER (Zutphen).

After a short historical survey of the opinions on the course of the retinal nerve fibres, the much disputed problem of the depth localization of the fibres is dealt with. Do the peripheral fibres show a preference for the deep or for the superficial parts of the nerve fibre layer, or for the peripheral or the central part of the papilla? Experimental and histological investigations give varying results.