Two Optical Systems of Cylinder Glasses

M.C.

Colenbrander

Leyden

To demonstrate what can be done with cylinder glasses, I have mounted two non-focal systems of cylinder glasses in a tube. One system consists of two cylinders of $-f-6$ at $1/6$ -r $1/6$ metre = $33 \frac{1}{8}$ cm. distance, and the other of one cylinder of $-f-2$ and one cylinder of $-f-6$ at $1/8 - 1/6$ metre = also $33 \frac{1}{8}$ cm. apart.

Colenbrander, Two Optical Systems of Cylinder Glasses 85

With these two combinations distant objects are seen absolutely sharply.

One system produces reversion in one direction. Rotation of the tube produces rotation of the image at twice the angular speed. The rotating cylinders can be used to observe what occurs in a rotating centrifuge.

The other system produces a magnification or a diminution in one direction, consequently a distortion of the image, whose direction changes on rotation of the tube.

Discussion.

Roelofs reported that there is a French system in which a narrowed image of an object is obtained by means of a cylindrical object glass. The image is projected at its natural width by a cylinder. He had learned that Bouwers uses curved mirrors for the same purpose, which gives less distortion. This system is known as "Delrama".

"Naphazoline", an Effective Curative in the Treatment of Irido-Cyclitis and Glaucoma.

By R. W. VON NORDHEIM (Utrecht).

(In collaboration with F. A. NELEMANS, Utrecht.)

Naphazoline, in a watery solution and applied locally, has a decongestive influence on mucous membranes. As such it is used by rhinologists, laryngologists, otologists and also by ophthalmologists, to reduce the swelling of mucous membranes.

Some time ago one of us proved that a solution of naphazoline dropped into the conjunctival cul-de-sac penetrates into the bulbus oculi. Intraocularly, naphazoline causes a moderate dilatation of the pupil; an intensifying action was observed with atropine as regards the dilatation of the pupil.

Having proved that naphazoline

after administration to the conjunctival cul-de-sac

could also be detected intraocularly

we have not only used this medicament in case of swelling and irritation of the conjunctiva for which it is recommended

but also in processes localized intraocularly.