The Surgical Treatment of Post-traumatic Intercalary Staphyloma

R. W. J. N.

Hoppenbrouwers

Utrecht

Author's address: Dr. R. W. J. N. Hoppenbrouwers, Maliesingel 7, Utrecht (Holland)

In a patient of twenty-one a staphyloma along the corneal limbus, covering more than a quadrant, developed following a blunt trauma. As a result of irregular astigmatism visus amounted to 1/60. The lens was clear but over the area of the staphyloma the zonular fibres were torn. Since the staphyloma was progressive it was necessary to operate.

The following possibilities were taken into consideration:

Fig. 1. Posttraumatic staphyloma.

Freeing and cutting away the staphyloma, followed by stitching of the edges.

Superimposition of the area with Tenon's capsule.

Covering the area with fascia lata.

Making a corneo-scleral transplantation. The first method was selected.

300 Hoppenbrouwers, The Surgical Treatment

$\phi \leq \Lambda$

o

Fig. 2.

Fig. 3.

Fig. 2 and 3. Procedure in the case of complicating rupture of the cornea.
of Post-traumatic Intercalary Staphyloma

301

While preparing the conjunctiva it turned out that besides the staphyloma there was a sharp gash penetrating the cornea but just covered by the conjunctiva. Since it was now no longer possible to suture the edges it was necessary to take rectangular stitches, two parallel sides being applied through healthy cornea and sclera (see figures). The advantage of this method was that the stitches were tied outside the conjunctiva, so that they could later be entirely removed, while the area on which they were supported was in completely healthy tissue.

It is important to lay emphasis on the fact that if an eye is afflicted by a blunt trauma it is essential to be aware that somewhere else in the cornea and sclera a sharp gash may be found.

This also proved true in the tests of Cusiodes as communicated by him at the meeting of the German Ophthalmological Society in Berlin in 1960.