Probiotics in the Treatment and Prevention of Atopic Dermatitis

R. Fölster-Holst

Universitätsklinikum Schleswig-Holstein, Campus Kiel, Kiel, Germany

Atopic dermatitis (AD) is the most common chronic inflammatory skin disease in infancy, which has increased steadily in the industrialized countries over the last 3 decades. Clinically the disease is characterized by dry skin, intractable pruritus and current relapses or chronic course which is associated with reduced quality of life for patients and their families. The therapy is mainly symptomatic and includes moisturizing the skin and topical anti-inflammatory treatment such as corticosteroids and calcineurin inhibitors. Additionally patients have to avoid irritants and allergens.

The main cause of AD is still not clearly understood. Apparently, it is the result of a complex interaction between genetic and environmental factors leading to disturbed epidermal differentiation, impaired epidermal barrier function and dysbalance of the immune system. The latter is characterized by a disruption of the Th1/Th2 cytokine balance towards an activation of Th2 cells, which is followed by induction of IgE and the activation and recruitment of eosinophils. It has been suggested that the absence of the exposure to microbes in the early childhood predisposes an infant to develop atopic diseases (hygiene hypothesis) [1]. Furthermore, patients with AD have significantly increased colonization by superantigen-secreting *Staphylococcus aureus*, which is associated with immunosuppressive activity of regulatory T cells [2].

The environmental factors which may influence atopic diseases also include probiotics such as lactobacilli or bifidobacteria. They are defined as specific microbial cultures which confer health benefits by prophylactic or therapeutic effects. These effects include the stabilization of the intestinal barrier, stimulation of intestinal IgA production, and modulation of specific and nonspecific immune responses to environmental factors like allergens [3–6].

In the end probiotic bacteria lead to a reduction in Th2 cytokines and increased production of IL-10 and TGF- β by regulatory T cells [5, 6–9]. However, there are conflicting results, as recently reported by authors from Denmark [10]. They showed that certain probiotic strains can modify antigen-presenting cells to cause reduced activity of regulatory T cells. So far the results regarding the immunomodulatory effects of probiotics appear variable, which may be due to using different probiotic strains and different methods of stimulating cytokine production.

Based on these findings probiotics have been considered to improve and prevent AD. The Finnish group headed by Isolauri (1997) was the first to show significant reduction in SCORAD (Scoring Atopic Dermatitis) in the probiotic group compared to a placebo group [11]. Four years later the same study group reported a positive effect

Table 1. Probiotics in the prevention of AD (randomized controlled studies)

First author, year	Patients	Probiotic	Duration of application	Age at follow-up years	Effect of probiotics
Kalliomäki [12–14] 2001, 2003, 2007	132	<i>Lactobacillus rhamnosus</i> GG	Prenatal 4 weeks, postnatal 6 months	2 4 7	Significantly reduced rate of AD in the verum group
Abrahamsson [18] 2007	188	<i>Lactobacillus reuteri</i>	Prenatal 2 weeks, postnatal 1 year	1	Significantly reduced rate of AD in the verum group (only IgE-associated AD)
Kukkonen [19] 2007	925	Mix of various probiotics + prebiotics	Prenatal 2–4 weeks, postnatal 6 months	2	Significantly reduced rate of AD in the verum group
Taylor [20] 2007	178	<i>Lactobacillus acidophilus</i>	Only postnatal (6 months)	1	No difference between verum and placebo
Wickens [21] 2008	474	<i>Lactobacillus rhamnosus</i> or <i>Bifidobacterium animalis</i> subsp. lactis	Prenatal 4 weeks, postnatal 2 years	2	Significantly reduced rate of AD in the verum group (only for <i>L. rhamnosus</i>)
Kopp [22] 2008	94	<i>Lactobacillus rhamnosus</i> GG	Prenatal 4–6 weeks, postnatal 6 months	2	No difference between verum and placebo
Kim [23] 2009	112	Mix of probiotics	Prenatal 4–6 weeks, postnatal 6 months	1	Significantly reduced rate of AD in the verum group
Niers [24] 2009	102	Mix of probiotics	Prenatal 4–6 weeks, postnatal 12 months	2	Significantly reduced rate of AD in the verum group
Soh [25] 2009	253	<i>Lactobacillus rhamnosus</i> + <i>Bifidobacterium longum</i>	Only postnatal (6 months)	1	No difference between verum and placebo

in the prevention of AD. The results showed a 50% reduction in the incidence of AD in the probiotic group compared to a placebo group at a 2-year follow-up [12]. In their follow-up study prevention of AD by probiotics extended beyond infancy [13, 14].

Up to now there are only a few randomized controlled studies using probiotics for the prevention and treatment of AD [15–17], summarized in tables 1 and 2.

Two meta-analyses of trials studying the effects of probiotics on AD risk were recently carried out by Osborn and Sinn [35] as well as Lee et al. [36]. The emphasis of Osborn and Sinn [35] was the effect of probiotics in the prevention of food hypersensitivity and allergic diseases, while the meta-analysis of Lee et al. [36] focused on the prevention and treatment of AD. Lee et al. concluded that current evidence is more convincing for the efficacy of probiotics in the prevention than in the treatment of pediatric AD. Osborn and Sinn stated that there is insufficient evidence to recommend probiotics

for the prevention of allergic disease and food hypersensitivity.

Both meta-analyses considered the research conducted by Rautava et al. [37] as well as Kalliomäki et al. [12, 13] as independent studies, even though both surveys originated from one and the same study.

Taken together there are conflicting results regarding the effects of probiotics in the treatment and prevention of AD. Probably, this is related to differences in study design (characterization of patients: number, age, severity of AD, sensitization to food and inhalative allergens, additional respiratory atopy), dosage of the probiotics, application of 1 probiotic strain or composition of different strains, duration of application, length of follow-up and time slot of administration. Therefore, at present probiotics cannot be generally recommended for the treatment and the prevention of AD. However, based on epidemiological, experimental and clinical results, it appears worthwhile to perform further randomized controlled

Table 2. Probiotics in the treatment of AD (randomized controlled trials)

First author year	Patients/age	Probiotic	Duration of application weeks	Effect of probiotics
Majamaa [11] 1997	27/2.5–15.7 months	<i>Lactobacillus rhamnosus</i> GG	4	Significant reduction in SCORAD in the verum group
Kirjavainen [26], 2003	35/5.5 months	<i>Lactobacillus rhamnosus</i> GG	7.5	Significant reduction in SCORAD in the verum group
Rosenfeldt [27] 2003	43/5.2 years	<i>Lactobacillus rhamnosus</i> + <i>Lactobacillus reuteri</i>	6	Significant reduction in SCORAD in the verum group (only IgE-associated AD)
Weston [28] 2005	53/10.9 months	<i>Lactobacillus fermentum</i>	8	Significant reduction in SCORAD in the verum group
Viljanen [29] 2005	230/6.4 months	<i>Lactobacillus rhamnosus</i> or mix of various probiotics	4	Significant reduction in SCORAD in the verum group (only IgE-associated AD)
Sistek [30] 2006	59/4.1 years	<i>Lactobacillus rhamnosus</i> + <i>Bifidobacterium lactis</i>	12	Significant reduction in SCORAD in the verum group (only food-sensitized patients)
Brouwer [31] 2006	50/5.2 months	<i>Lactobacillus rhamnosus</i> or <i>Lactobacillus rhamnosus</i> GG	12	No effect of probiotics
Passeron [32] 2006	39/5.9 years	Symbiotics or prebiotics (placebo)	12	Significant reduction in SCORAD in both groups
Fölster-Holst [33], 2006	47/18.8 months	<i>Lactobacillus rhamnosus</i> GG	8	No effect of <i>Lactobacillus rhamnosus</i> GG
Grüber [34] 2007	102/7.4 months	<i>Lactobacillus rhamnosus</i> GG	8	No effect of <i>Lactobacillus rhamnosus</i> GG

multicenter studies with standardized parameters in a representative number of patients. Regarding the heterogeneity of AD it is important to characterize the patients very accurately in order to define subgroups which could be susceptible to using probiotics as therapeutical and preventive measurements.

Disclosure Statement

R.F.-H. does not have any relationship to disclose.

References

- 1 Strachan DP: Family size, infection and atopy: the first decade of the 'hygiene hypothesis'. *Thorax* 2000;55(suppl 1):S2–S10.
- 2 Ou LS, Goleva E, Hall C, Leung DY: T regulatory cells in atopic dermatitis and subversion of their activity by superantigens. *J Allergy Clin Immunol* 2004;113:756–763.
- 3 Isolauri E, Kalliomäki M, Laitinen K, Salminen S: Modulation of the murine gut barrier and microbiota: a novel target in allergic disease. *Curr Pharm Des* 2008;14:1368–1375.
- 4 Rosenfeldt V, Benfeldt E, Valerius NH, Paerregaard A, Michaelsen KF: Effect of probiotics on gastrointestinal symptoms and small intestinal permeability in children with atopic dermatitis. *J Pediatr* 2004;145:612–616.
- 5 Pochard P, Gosset P, Grangette C, Andre C, Tonnel AB, Pestel J, Mercenier A: Lactic acid bacteria inhibit TH2 cytokine production by mononuclear cells from allergic patients. *J Allergy Clin Immunol* 2002;110:617–623.
- 6 Ghadimi D, Fölster-Holst R, de Vrese M, Winkler P, Heller KJ, Schrezenmeier J: Effects of probiotic bacteria and their genomic DNA on TH1/TH2-cytokine production by peripheral blood mononuclear cells (PBMCs) of healthy and allergic subjects. *Immunobiology* 2008;213:677–692.
- 7 Hacini-Rachinel F, Gheit H, Le Duque JB, Dif F, Nancey S, Kaiserlian D: Oral probiotic control skin inflammation by acting on both effector and regulatory T cells. *PLoS One* 2009;4:e4903.

- 8 Mastrangeli G, Corinti S, Butteroni C, Af-ferni C, Bonura A, Boirivant M, Colombo P, Di Felice G: Effects of live and inactivated VSL#3 probiotic preparations in the modulation of in vitro and in vivo allergen-induced Th2 responses. *Int Arch Allergy Immunol* 2009;150:133–143.
- 9 Smits HH, Engering A, van der Kleij D, de Jong EC, Schipper K, van Capel TM, Zaat BA, Yazdanbakhsh M, Wierenga EA, van Kooyk Y, Kapsenberg ML: Selective probiotic bacteria induce IL-10-producing regulatory T cells in vitro by modulating dendritic cell function through dendritic cell-specific intercellular adhesion molecule 3-grabbing nonintegrin. *J Allergy Clin Immunol* 2005;115:1260–1267.
- 10 Schmidt EG, Claesson MH, Jensen SS, Ravn P, Kristensen NN: Antigen-presenting cells exposed to *Lactobacillus acidophilus* NCFM, *Bifidobacterium bifidum* BI-98, and BI-504 reduce regulatory T cell activity. *Inflamm Bowel Dis* 2010;16:390–400.
- 11 Majamaa H, Isolauri E: Probiotics: a novel approach in the management of food allergy. *J Allergy Clin Immunol* 1997;99:179–185.
- 12 Kalliomäki M, Salminen S, Arvilommi H, Kero P, Koskinen P, Isolauri E: Probiotics in primary prevention of atopic disease: a randomized placebo-controlled trial. *Lancet* 2001;357:1076–1079.
- 13 Kalliomäki M, Salminen S, Poussa T, Arvilommi H, Isolauri E: Probiotics and prevention of atopic disease: 4-year follow-up of a randomised placebo-controlled trial. *Lancet* 2003;361:1869–1871.
- 14 Kalliomäki M, Salminen S, Poussa T, Isolauri E: Probiotics during the first 7 years of life: a cumulative risk reduction of eczema in a randomized, placebo-controlled trial. *J Allergy Clin Immunol* 2007;119:1019–1021.
- 15 Van der Aa LB, Heymans HS, van Aalderen WM, Sprinkelman AB: Probiotics and prebiotics in atopic dermatitis: review of the theoretical background and clinical evidence. *Pediatr Allergy Immunol* 2009, Epub ahead of print.
- 16 Betsi GI, Papadavid E, Falagas ME: Probiotics for the treatment or prevention of atopic dermatitis: a review of the evidence from randomized controlled trials. *Am J Clin Dermatol* 2008;9:93–103.
- 17 Fölster-Holst R, Offick B, Proksch E, Schrezenmeier J: Probiotics in treatment and/or prevention of allergies; in Michail S, Sherman PM (eds): *Probiotics in Pediatric Medicine*. Totowa, Humana Press, Springer, 2009, 243–267.
- 18 Abrahamsson TR, Jakobsson T, Böttcher MF, Fredrikson M, Jenmalm MC, Björkstén B, Oldaeus G: Probiotics in prevention of IgE-associated eczema: a double-blind, randomized, placebo-controlled trial. *J Allergy Clin Immunol* 2007;119:1174–1180.
- 19 Kukkunen K, Savilahti E, Haahtela T, Juntunen-Backman K, Korpela R, Poussa T, Ture T, Kuitunen M: Probiotics and prebiotic galacto-oligosaccharides in the prevention of allergic diseases: a randomized, double-blind, placebo-controlled trial. *J Allergy Clin Immunol* 2007;119:192–198.
- 20 Taylor AL, Dunstan JA, Prescott SL: Probiotic supplementation for the first 6 months of life fails to reduce the risk of atopic dermatitis and increases the risk of allergen sensitization in high-risk children: a randomized controlled trial. *J Allergy Clin Immunol* 2007;119:184–191.
- 21 Wickens K, Black PN, Stanley TV, Mitchell E, Fitzharris P, Tannock GW, Purdie G, Crane J, Probiotic Study Group: A differential effect of 2 probiotics in the prevention of eczema and atopy: a double-blind, randomized, placebo-controlled trial. *J Allergy Clin Immunol* 2008;122:788–794.
- 22 Kopp MV, Hennemuth I, Heinzmann A, Urbanek R: Randomized, double-blind, placebo-controlled trial of probiotics for primary prevention: no clinical effects of *Lactobacillus* GG supplementation. *Pediatrics* 2008;121:e850–e856.
- 23 Kim JY, Kwon JH, Ahn SH, Lee SI, Han YS, Choi YO, Lee SY, Ahn KM, Ji GE: Effect of probiotic mix (*Bifidobacterium bifidum*, *Bifidobacterium lactis*, *Lactobacillus acidophilus*) in the primary prevention of eczema: a double-blind, randomized, placebo-controlled trial. *Pediatr Allergy Immunol* 2009, Epub ahead of print.
- 24 Niers L, Martín R, Rijkers G, Sengers F, Timmerman H, van Uden N, Smidt H, Kimpfen J, Hoekstra M: The effects of selected probiotic strains on the development of eczema (the PandA study). *Allergy* 2009;64:1349–1358.
- 25 Soh SE, Aw M, Gerez I, Chong YS, Rauff M, Ng YP, Wong HB, Pai N, Lee BW, Shek LP: Probiotic supplementation in the first 6 months of life in at risk Asian infants – effects on eczema and atopic sensitization at the age of 1 year. *Clin Exp Allergy* 2009;39:571–578.
- 26 Kirjavainen PV, Salminen SJ, Isolauri E: Probiotic bacteria in the management of atopic disease: underscoring the importance of viability. *J Pediatr Gastroenterol Nutr* 2003;36:223–227.
- 27 Rosenfeldt V, Benfeldt E, Nielsen SD, Michaelsen KF, Jeppesen DL, Valerius NH, Paerregaard A: Effect of probiotic *Lactobacillus* strains in children with atopic dermatitis. *J Allergy Clin Immunol* 2003;111:389–395.
- 28 Weston S, Halbert A, Richmond P, Prescott SL: Effects of probiotics on atopic dermatitis: a randomised controlled trial. *Arch Dis Child* 2005;90:892–897.
- 29 Viljanen M, Savilahti E, Haahtela T, Juntunen-Backman K, Korpela R, Poussa T, Ture T, Kuitunen M: Probiotics in the treatment of atopic eczema/dermatitis syndrome in infants: a double-blind placebo-controlled trial. *Allergy* 2005;60:494–500.
- 30 Sistek D, Kelly R, Wickens K, Stanley T, Fitzharris P, Crane J: Is the effect of probiotics on atopic dermatitis confined to food sensitized children? *Clin Exp Allergy* 2006;36:629–633.
- 31 Brouwer ML, Wolt-Plompen SAA, Dubois AEJ, van der Heide S, Jansen DJ, Hoijer MA, Kauffman HF, Duiverman EJ: No effects of probiotics on atopic dermatitis in infancy: a randomized placebo-controlled trial. *Clin Exp Allergy* 2006;36:899–906.
- 32 Passeron T, Lacour JP, Fontas E, Ortonne JP: Prebiotics and synbiotics: two promising approaches for the treatment of atopic dermatitis in children above 2 years. *Allergy* 2006;61:431–437.
- 33 Fölster-Holst R, Müller F, Schnopp N, Abeck D, Kreiselmaier I, Lenz T, von Rügen U, Schrezenmeier J, Christophers E, Weichen-thal M: Prospective, randomized controlled trial on *Lactobacillus rhamnosus* in infants with moderate to severe atopic dermatitis. *Br J Dermatol* 2006;155:1256–1261.
- 34 Grüber C, Wendt M, Sulser C, Lau S, Kulig M, Wahn U, Werfel T, Niggemann B: Randomized, placebo-controlled trial of *Lactobacillus rhamnosus* GG as treatment of atopic dermatitis in infancy. *Allergy* 2007;62:1270–1276.
- 35 Osborn DA, Sinn JK: Probiotics in infants for prevention of allergic disease and food hypersensitivity. *Cochrane Database Syst Rev* 2007 Oct 17;CD006475.
- 36 Lee J, Seto D, Bielory L: Meta-analysis of clinical trials of probiotics for prevention and treatment of pediatric atopic dermatitis. *J Allergy Clin Immunol* 2008;121:116–121.e11.
- 37 Rautava S, Isolauri E: The development of gut immune responses and gut microbiota: effects of probiotics in prevention and treatment of allergic disease. *Curr Issues Intest Microbiol* 2002;3:15–22.