Human GnRH Deficiency: A Unique Disease Model to Unravel the Ontogeny of GnRH Neurons

Ravikumar Balasubramanian Andrew Dwyer Stephanie B. Seminara
Nelly Pitteloud Ursula B. Kaiser William F. Crowley, Jr.

Harvard Reproductive Endocrine Sciences Center of Excellence, Reproductive Endocrine Unit of the Department of Medicine of the Massachusetts General Hospital and Endocrine Division of the Brigham and Women's Hospital, Boston, Mass., USA

Key Words

Gonadotropin-releasing hormone deficiency ·
Gonadotropin-releasing hormone neuronal ontogeny ·
Puberty · Kallmann syndrome · Hypogonadotropic
hypogonadism

Abstract

Evolutionary survival of a species is largely a function of its reproductive fitness. In mammals, a sparsely populated and widely dispersed network of hypothalamic neurons, the gonadotropin-releasing hormone (GnRH) neurons, serve as the pilot light of reproduction via coordinated secretion of GnRH. Since its first description, human GnRH deficiency has been recognized both clinically and genetically as a heterogeneous disease. A spectrum of different reproductive phenotypes comprised of congenital GnRH deficiency with anosmia (Kallmann syndrome), congenital GnRH deficiency with normal olfaction (normosmic idiopathic hypogonadotropic hypogonadism), and adult-onset hypogonadotropic hypogonadism has been described. In the last two decades, several genes and pathways which govern GnRH ontogeny have been discovered by studying humans with GnRH deficiency. More importantly, detailed study of these patients has highlighted the emerging theme of oligogenicity and

genotypic synergism, and also expanded the phenotypic diversity with the documentation of reversal of GnRH deficiency later in adulthood in some patients. The underlying genetic defect has also helped understand the associated non-reproductive phenotypes seen in some of these patients. These insights now provide practicing clinicians with targeted genetic diagnostic strategies and also impact on clinical management.

Copyright © 2010 S. Karger AG, Basel

Introduction

A unique neural network within the hypothalamus consisting of approximately 1,500 neurons termed the parvocellular system initiates and maintains reproductive function in mammals [1, 2]. It accomplishes this task by coordinating the synthesis and pulsatile secretion of a single neuroendocrine decapeptide, gonadotropin-releasing hormone (GnRH), from this neural network. In humans, this GnRH neuronal network is fully active resulting in adult levels of activity of the hypothalamic-pituitary-gonadal axis during the late stages of gestation and the early neonatal period. It then becomes quiescent during childhood until its full reactivation at puberty

[3, 4]. These GnRH neurons reside predominantly in the medial preoptic area, project their axonal processes into the median eminence of the hypothalamus, and secrete GnRH into the hypophyseal portal circulation in a coordinated and pulsatile manner [5–7]. Once at its cognate pituitary receptor on the gonadotropes, GnRH stimulates both the synthesis and secretion of the two pituitary gonadotropic hormones: luteinizing hormone (LH) and follicle-stimulating hormone that, in turn, stimulate steroidogenesis and govern the maturation and function of the germ cells in the gonads in both sexes.

Remarkably, despite the critical evolutionary role of the reproductive system in survival of the species, human reproductive activity is invested in a single gene encoding GnRH and this unique network of hypothalamic neurons. Considering the ever-changing environmental demands that have threatened survival of mammals over the millennia, it is likely that a multi-tiered, overlapping network of genetic and cellular pathways has evolved to regulate the secretion of GnRH. The existence of such a complex yet fascinating biological system ‘upstream’ of GnRH secretion has been hinted at by the study of humans presenting with GnRH deficiency. These insights have been enabled by combining careful clinical investigations in these patients with the new powerful tools of developmental and molecular biology, genetics, animal models, and structural biology. Integration of these various tools and investigative approaches has allowed several groups around the world to begin to dissect genetic components of this unique neuroendocrine network and begin to define a functional map of the ontogeny of the GnRH neurons. This review attempts to put into perspective the evolving genetic architecture and molecular pathways governing the ontogeny of GnRH neurons as elucidated by the study of humans with GnRH deficiency, a truly prismatic disorder.

GnRH Neuronal Development

The first major advance that provided insight into the complexities of the developmental biology of the GnRH neurons came from investigations in rodents where GnRH neurons were demonstrated to originate outside the central nervous system (CNS) in the medial olfactory placode during embryological development [8–10]. From such humble origins where these neurons first specify their biological fate, committed GnRH neurons migrate from the olfactory placode along the olfactory epithelium to the olfactory bulb and then, via a structural scaffolding

created by the olfactory bulb and tract, ultimately migrate to take up their final residence over a widely dispersed network within the medial preoptic area of the hypothalamus [11, 12] (fig. 1a, b). These precise anatomic locations presumably reflect their need to determine and hence monitor a wide variety of sensory systems and other neuronal systems that provide data to the reproductive system required for survival (fig. 1b). These include body weight and nutritional status; circadian and light/dark cycles; stress and adrenal function; olfaction which helps locate predators as well as potential mating options; and sight and hearing of other environmental variables that might threaten the species. The GnRH neurons integrate these complex signals and determine whether the timing and environment are optimal to support full capability of the reproductive system by both sex steroid secretion and fertility or whether the optimal reproductive status is to silence and suppress both. The fine balance of timing of these two different settings on the GnRH scale then determines the ability of the individual and species to adapt and survive various environmental threats.

Once in place, these GnRH neurons extend axonal projections to the median eminence where they coordinate their secretion of GnRH in a pulsatile pattern of secretion that is absolutely required by the gonadotrope to evoke physiologic gonadotropin secretion [13] and gonadal function (fig. 1b). These GnRH migration studies are in line with the previously documented observation that when normal fetal GnRH neurons were transplanted into the third ventricles of GnRH-deficient adult female hypogonadal mice (hpg mice), transplanted GnRH neurons proved capable of projecting GnRH-containing processes to the median eminence in the host [14], initiating pulsatile LH secretion [15], and restoring limited reproductive competence [14] to those GnRH-deficient animals. The precise factors guiding this peripatetic journey of the GnRH neurons are unknown but experimental evidence supports involvement of a wide variety of genetic and biochemical cues including axonal guidance molecules, cell adhesion molecules, and various transcription and growth factors [16].

Human GnRH Deficiency: A Clinically and Genetically Heterogeneous Disease Model

Reactivation of the GnRH pulse generator during adolescence heralds the onset of puberty (fig. 2a). In boys, pubertal onset is marked by testicular enlargement, which is followed by penile and pubic hair growth. In

Fig. 1. GnRH migration. **a** Migratory route of GnRH neurons in the embryonic mouse brain. GnRH neurons (black dots) originate in the medial wall of the olfactory placode, migrate along the olfactory axons and enter the brain perforating the cribriform plate. They then migrate to reach their final destination in the preoptic area (poa) of the hypothalamus. GnRH neurons on embryonic day 11 are seen in the vomeronasal organ (vno) and medial wall of the olfactory placode. In the 16-day-old fetal mouse brain, most of the GnRH neurons have reached the hypothalamus. gt = Ganglion terminale; ob = olfactory bulb. Reproduced with permission from Schwanzel-Fukuda and Pfaff [8]. **b** Cartoon of GnRH migration in humans. GnRH neurons originate in the embryonic olfactory placode and migrate along olfactory axons, penetrate the cribriform plate and migrate across the olfactory bulb, aided by various chemoattractive and repulsion factors and eventually reach the preoptic area of the hypothalamus, where they synchronize pulsatile GnRH secretion. Kisspeptin (KISS1) and the tachykinin (TAC) neurons serve as upstream modulators of GnRH secretion. Upon GnRH stimulation, the pituitary (PIT) secretes LH and follicle-stimulating hormone (FSH), which in turn regulate gonadal steroidogenesis and gametogenesis.

girls, the first sign of puberty is the onset of breast budding (thelarche), followed by pubic hair growth and menarche. Normal ages for puberty are 8–13 years in girls and 9–14 years in boys. Puberty is ‘delayed’ when the secondary sexual characteristics fail to develop beyond 2 standard deviations from the mean. This delay in the activation of the GnRH pulse generator resulting in delayed puberty is the mildest form of human GnRH deficiency (fig. 2b). On the contrary, premature initiation of the

GnRH pulse generator results in central precocious puberty (fig. 2c).

In 1856, Maestre de San Juan [17] first documented the pathological association of hypogonadism and absence of the olfactory system. A more detailed description of the syndrome was reported in 1944 by Kallmann [18] using patients in three affected pedigrees with hypogonadism and anosmia. His observations were the first to hint at a broader spectrum of clinical defects and identify the fa-

Fig. 2. GnRH pulsatile secretion in normal puberty and disorders of pubertal development. **a** Pulsatile secretion of GnRH is fully active during the early neonatal period ('mini-puberty of infancy'), followed by quiescence during childhood, and reactivates in adolescence signalling normal puberty. **b** Delayed activation of the GnRH pulse generator results in delayed puberty. **c** Premature activation of the GnRH pulse generator in early childhood results in precocious puberty.

milial nature in the clinical syndrome that was seen in both sexes and accompanied by multiple congenital anomalies. In 1954, de Morsier [19] first noted the link between hypogonadism and neuroanatomical defects including agenesis of the olfactory bulb and tract and other midline neuroanatomical defects and coined the term 'olfacto-genital dysplasia'. Since then, hypogonadotropic hypogonadism accompanied by anosmia has been classically described as Kallmann syndrome (KS) (fig. 3a). Eventually, the discovery of GnRH in 1971 aided the linking of the etiology of KS to hypogonadotropic hypogonadism [20, 21]. Although the variability of the pituitary gonadotropin response to a single bolus of GnRH initially failed to definitively confirm hypothalamic GnRH deficiency [22], subsequent normalization of the pituitary-gonadal axis in response to a physiologically defined regimen of exogenous GnRH administration in most of these patients confirmed the hypothalamic defect in these patients [23]. However, a pituitary defect in GnRH action was subsequently demonstrated by the discovery of mutations in GnRH receptor (GNRHR) in some patients with hypogonadotropic hypogonadism [24]. Hypogonadotropic hypogonadism is also known to occur in the presence of normal olfaction and is classified as normosmic idiopathic hypogonadotropic hypogonadism (nIHH) (fig. 3b). In this regard, it is noteworthy that even

in Kallmann's [18] initial report, 3 of the original 12 hypogonadal individuals had no obvious defect in olfaction.

Mutations in a number of different genes have now been linked with human GnRH deficiency, but only approximately 40% of the genetic heritability has been uncovered, with a number of genes yet to be discovered (table 1). However, the growing appreciation that both KS and nIHH can occur within a given family (i.e. with the same genetic milieu) demonstrated that both of these variants of GnRH deficiency are part of a continuous spectrum of clinical and genetic defects rather than single unique phenotypes. Those with anosmia presumably signal a joint involvement of the olfactory system and GnRH neurons in the pathophysiology of this disorder, whereas those with a normal sense of smell indicate processes occurring either before or after migration of GnRH neurons. In addition, several patients with KS and nIHH present with varying degrees of pubertal delay, with some presenting with complete absence of puberty suggesting complete GnRH deficiency (fig. 4a) and others with partial pubertal development indicating partial GnRH deficiency (fig. 4b). Several other additional phenotypes including isolated anosmia and delayed puberty also form part of this clinical spectrum [25–28]. Moreover, the presence of phenotypically normal individuals who carry a similar genetic burden as other affected family members has uncovered the critical role played by oligogenicity and genotypic synergism in the observed clinical phenotype [29, 30].

Human GnRH deficiency had traditionally been thought to be a monogenic disorder with complete and permanent failure of sexual maturation due to silencing of the GnRH pulse generator. However, it has become increasingly clear that human GnRH deficiency is not merely a congenital arrest of GnRH neuronal development as was implied by the initial studies of *KALI*-deleted patients (see below). The identification of adult-onset idiopathic hypogonadotropic hypogonadism (AHH) in men confirmed normal sexual development until the cessation of GnRH pulse generator function that occurs well into adulthood [28] (fig. 4c). Similarly, spontaneous reversals of well-established GnRH deficiency following long-term therapy with sex steroids has added yet another clinical variant suggesting a more complex pathophysiology than was initially thought [31] (fig. 4d).

Thus, it is becoming increasingly clear that GnRH deficiency in humans represents a unique family of oligogenic disorders that present with a continuum of reproductive and non-reproductive phenotypes. These phenotypes belie both developmental and functional

Fig. 3. KS and nIHH. **a** KS results from a developmental defect in olfactory bulb (OB) morphogenesis and GnRH neuronal migration resulting in failure of GnRH neurons to arrive at the preoptic area (POA) in the hypothalamus and failure of GnRH secretion. The GnRH neurons are arrested in their extracerebral route. Mutations in *KAL1*, *PROK2*, *PROKR2*, *FGFR1*, *FGF8*, *NELF* and *CHD7* have been associated with KS. **b** nIHH results from either defective pulsatile secretion of GnRH or defective GnRH action at the pituitary. Mutations in *GNRH1*, *KISS1R*, *GNRHR*, *PROK2*, *PROKR2*, *FGFR1*, *TAC3*, *TACR3*, *CHD7* and *NROB1* genes have been associated with nIHH.

pathophysiological defects and in some cases can reverse completely in adulthood. Detailed clinical investigations of human GnRH deficiency aided by diverse basic research tools have proven to provide a prism into the fascinating complexities of GnRH deficiency and the ontog-

eny of this neuronal network. It is possible that the rare genetic variants that underlie human GnRH deficiency and possibly central precocious puberty may eventually explain the heritability of delayed puberty and early puberty seen within the normal population (fig. 5).

Fig. 4. Phenotypic heterogeneity in human GnRH deficiency. GnRH deficiency can either be complete (a) or partial (b). c Some patients develop GnRH deficiency in adulthood following normal GnRH activation in the neonatal period (AHH). d Occasionally, following complete or partial GnRH deficiency, GnRH pulse generator activates in adulthood (reversal of GnRH deficiency).

Fig. 5. Genetics of puberty. The precise genetic basis of the extreme tails of normal puberty is unclear. The genetic variants causing GnRH deficiency and precocious puberty may possibly explain the heritability of early and delayed puberty in the normal population.

Table 1. Genetics of human GnRH deficiency

Gene (chromosomal locus)	OMIM No.	Reproductive phenotypes	Associated phenotypes
<i>KAL1</i> (Xp22.3)	308700	KS	unilateral renal agenesis, bimanual synkinesia, high-arched palate
<i>KISS1R</i> (19p13.3)	604161	nIHH	none
<i>FGFR1</i> (8p11.2–p11.1)	136350	KS	cleft lip/cleft palate
<i>FGF8</i> (10q.24)	600483	nIHH AHH	skeletal anomalies (hand and foot) external ear hypoplasia
<i>PROK2</i> (3p21.1)	607002	KS	? circadian/sleep dysregulation
<i>PROKR2</i> (20p.13)	607123	nIHH AHH	
<i>GNRH1</i> (8p21–p11.2)	152760	nIHH	none reported
<i>GNRHR</i> (4q21.2)	138850	nIHH	none reported
<i>TAC3</i> (12q13–q21)	162332	nIHH	microphallus, cryptorchidism
<i>TACR3</i> (4q25)	162330		reversal of GnRH deficiency
<i>NELF</i> (9q34.3)	608137	KS	none reported
<i>NROB1</i> (Xp.21.3–p21.2)	300473	nIHH	part of X-linked adrenal hypoplasia congenita
<i>LEP</i> (7q31.3)	164160	nIHH	severe obesity
<i>LEPR</i> (1p31)	601007		
<i>CHD7</i> (8q12.1)	608892	KS nIHH	part of CHARGE syndrome

CHARGE syndrome = eye Coloboma, Heart anomalies, choanal Atresia, growth and developmental Retardation, Genitourinary anomalies and Ear abnormalities.

Fig. 6. *KAL1* mutations in human GnRH deficiency. **a** Gene deletions (black lines) in *KAL1* gene causing KS. **b** Schematic of exons of *KAL1* gene. **c** Schematic of the anosmin-1 protein displaying known point mutations (in hexagons) causing KS. Anosmin-1, a multidomain protein, consists of a N-terminal signal peptide fol-

lowed by a cysteine-rich region (CYS), a WAP-like four-disulfide core motif, four tandem FnIII repeats and a C-terminal histidine-rich region (H) (insertions, deletions, duplications and intronic changes not shown).

Discovery of *KAL1*

The first human gene responsible for KS was localized to the distal portion of the X chromosome by investigators studying a pedigree with 5 male patients with X-linked ichthyosis, hypogonadotropic hypogonadism and anosmia [32]. This finding was extended in a subsequent study of a male infant born with KS associated with ichthyosis, chondroplasia punctata and choanal atresia and chromosomal analysis showed deletion of the Xp22.3 region of the X chromosome inherited from his unaffected mother [33]. The child subsequently died and an autopsy showed complete absence of olfactory bulbs and tracts and absence of cribriform plate perforations [33]. A second male child was then conceived by the same mother and amniocentesis revealed an identical defective X chromosome. Neuroanatomical studies of the aborted fetus at 19 weeks of gestation revealed that the GnRH neurons had developed, migrated from the olfactory placode towards the CNS, yet were arrested in their migration at the cribriform plate of the ethmoid bone, presumably since there were no olfactory axons to guide their transport further [34].

Eventually, the first KS gene, *KAL1*, was identified by positional cloning of the distal portion of the X chromosome (Xp22.3) simultaneously by two independent groups [35, 36]. The *KAL1* gene (fig. 6b) is comprised of 14 exons, is yet to be discovered in the mouse, and encodes a 680-amino acid secreted extracellular-matrix glycoprotein, anosmin-1 (fig. 6c). Anosmin-1 is a secreted multi-domain protein consisting of an N-terminal signal peptide followed by a cysteine-rich region (cys-box), a whey acidic protein (WAP)-like four disulfide core motif, four tandem fibronectin type III (FnIII) repeats and a C-terminal histidine-rich region [35, 36].

Anosmin-1 is required for the formation of the olfactory guidance platform for GnRH neuronal migration, but its precise biological activity in achieving this function is unclear. Similarly, a wave of its expression in the kidney precedes the appearance of glomeruli during renal development. This may well explain why some KS patients with mutations in *KAL1* have congenital defects of the kidney including complete renal agenesis. Human anosmin-1 shares significant homology with neural cell adhesion molecules that have fibronectin repeats [35, 36].

Although the search for the rodent homologue of anosmin-1 has been elusive, known *KAL1* orthologs in other species are highly conserved, especially at the WAP and FnIII domains [37]. The eight cysteine residues in the WAP domain of anosmin-1 which form four disulfide bonds are also highly conserved amongst other serine protease inhibitors that are members of this WAP protein family [38]. Since anosmin-1 is a secreted protein, it is thought to act locally at the cell surface via its interaction with the cell membrane that depends largely on the binding of one of the 4 FnIII domains to heparan sulfate proteoglycans [38, 39]. Expression of anosmin-1 is regionally and temporally restricted during human organogenesis [40]. Migration of immortalized rodent GnRH neurons is stimulated by human anosmin-1 [16]. Anosmin-1 immunoreactivity in human embryos has been shown to be present in the nasal placode as early as embryonic week 4.5 and its expression is seen in the presumptive olfactory bulb by the 5th embryonic week even prior to the olfactory nerve synaptogenesis with the olfactory bulb [40]. However, the absence of anosmin-1 in the olfactory epithelium or the extracerebral route of the olfactory axonal elongation suggests that the role of anosmin-1 may primarily rest in the intracerebral olfactory axonal elongation and subsequent GnRH neuronal migration along this course. This suggestion is further supported by the arrest of GnRH neuronal migration at the cribriform plate in the KS fetus lacking *KAL1* [34]. Taken together, these findings suggest that the biological role of anosmin-1 in the presumptive olfactory bulb may be critical to attracting olfactory axons towards the forebrain. In the absence of anosmin-1, this trajectory is lost and GnRH neurons thus fail to reach the olfactory bulb initially and subsequently the hypothalamus [41]. Apart from its role in olfactory axonal attraction, it is likely anosmin-1 may independently play a role in olfactory bulb morphogenesis per se and/or neurite outgrowth beyond the olfactory bulb through the lateral olfactory tract [42].

Despite the critical developmental role played by anosmin-1, deletions in *KAL1* (fig. 6a) and mutations in *KAL1* (fig. 6c) only account for up to 10–14% of familial KS [43] and 8–11% of sporadic KS [44] patients. A wide spectrum of mutations spanning the entire coding sequence of the *KAL1* gene have been reported in X-linked KS subjects [45]. Many of the point mutations cluster around the FnIII domain [46], confirming the key molecular role played by this domain in protein activity [39, 47]. In human subjects with *KAL1* mutations, several nonreproductive features, including unilateral renal agenesis, synkinesia or mirror movements, hearing loss,

midline facial defects, and bony abnormalities are variably seen [44, 48]. In keeping with the clinical signatures, the wider developmental roles of anosmin-1 are strongly supported by the detection of the protein in the extracellular matrices of various nonreproductive tissues such as bronchial tubes, mesonephric tubules, ureteric bud, digestive tract, large blood vessels, and inner ear [40].

KISS1R (GPR54)

The relative rarity of human GnRH deficiency, its multiple inheritance patterns, its associated reduced fertility, and consequent small pedigree sizes have rendered traditional linkage analysis less effective in aiding novel gene discovery in this phenotype. However, in 2003, using linkage analysis in inbred families, two groups independently identified *GPR54*, a G protein-coupled receptor, and its cognate ligand, kisspeptin, to be upstream gatekeepers of GnRH neurons [49, 50]. Coupled with complementary mouse genetics and in vitro confirmation of loss of biological activity of the mutant receptor protein, *GPR54* was implicated as a key regulator of puberty [50] and a number of mutations in this receptor have now been described in nIHH patients (fig. 7a, b). *GPR54*, which has recently been redesignated as *KISS1R*, is a member of the rhodopsin family of G protein-coupled receptors with sequence homologies to the members of the galanin receptor family [51]. Kisspeptins, an overlapping family of RF-amide peptides, serve as endogenous ligands for *KISS1R* and are potent stimulators of gonadotropin release in all mammalian species studied to date [51, 52]. Kisspeptin is coded by the *KISS1* gene and its proteolytic processing results in different-length kisspeptins, all of those which retain the amidated carboxy-terminal decapeptide are potent *KISS1R* agonists [51, 53]. The longest kisspeptin, kisspeptin-1, is a 54-amino acid peptide and was initially termed metastatin due to its ability to inhibit tumor metastasis [53]. So far, *KISS1* mutations causing human GnRH deficiency have not been reported.

Review of human and mouse *KISS1R* investigations have provided fascinating insights into the role of *KISS1/KISS1R* biology in human reproduction. Both *KISS1*^{−/−} and *KISS1R*^{−/−} mice are phenocopies of nIHH and interestingly show normal GnRH content in the hypothalamus, providing the first indication that mutations in *KISS1R* do not affect GnRH neuronal migration or GnRH synthesis but rather GnRH release [50, 54]. Several key

Fig. 7. *KISS1R* and human GnRH deficiency. **a** Schematic of the predicted *KISS1R* protein displaying known mutations causing nHH in humans. **b** Dose-response curves for the ligand-stimulated production of inositol phosphate in *KISS1R* mutant constructs, corrected for protein content. **i** Curve for the L148S mutation (three independent experiments, each performed in triplicate). **ii** Curve for the R331X mutation (two independent experiments, each performed in triplicate). **iii** Curve for the X399R polyA stop mutation (two independent experiments, each performed in triplicate). The percentages on the y-axis represent the percentages of the maximal stimulation for each GPR54 construct. **iv** The relative quantification of the wild-type and mutant GPR54 allele expression in lymphoblastoid cell lines as measured by quantitative RT-PCR. Reproduced with permission from Seminara et al. [50].

observations have emerged from the study of human GnRH-deficient subjects with *KISS1R* mutations. Neuroendocrine profiling of probands with *KISS1R* mutations has generally shown dampened but present low-amplitude LH pulses [50, 55] suggesting some degree of endogenous GnRH secretion that is synchronized but reduced in pulse amplitude. An African-American proband with a compound heterozygous mutation in *KISS1R* also showed a striking leftward shift in his LH dose-response relationship to exogenous GnRH. This observation suggests some degree of endogenous pituitary priming by intact but dampened GnRH pulsatility [50]. In one published report of a male with *KISS1R* mutation who presented with cryptorchidism and micropenis in infancy, neuroendocrine evaluation at 2 months of age revealed undetectable gonadotropins also suggesting a role for the *KISS1/KISS1R* system in the 'mini-puberty' of infancy [56]. Some patients with *KISS1R* mutations are able to undergo folliculogenesis, spermatogenesis and successful pregnancies following therapy with exogenous GnRH, suggesting some degree of intact pituitary gonadotropin function with no significant primary gonadal defects or defects in placentation in these individuals [57].

Although a variety of diverse phenotypes is often seen in human GnRH-deficient subjects, nonreproductive phenotypes have yet to be described in subjects with *KISS1R* mutations. However, both *KISS1* and *KISS1R* expression are not restricted to the hypothalamic-pituitary-gonadal axis and both the ligand and receptor expression are reported elsewhere in the body such as the adrenal, placenta, and pancreatic islet beta cells [53]. Moreover, both *KISS1* and *KISS1R* were implicated as metastasis suppressors in cells from breast, melanoma, and pancreatic cancers [53]. The *KISS1/KISS1R* pathway has also been implicated in hypothalamic control of energy balance [58], glucose-induced insulin secretion in the pancreas [59] and has been postulated to be a potential integrator of the gonadal hormonal and circadian signals and relay to the GnRH neurons [60].

Although mutations in the *KISS1/KISS1R* pathway do not seem to be a significant contributor to human GnRH deficiency (<5%), their relative rarity probably suggests an evolutionarily critical role in reproduction and species propagation that might well have undergone negative selection. The recently reported documentation of a ligand-independent gain-of-function mutation in GPR54 in a girl with idiopathic central precocious puberty demonstrates the crucial role of this system in the maturation of the reproductive axis [61]. Furthermore, constant infu-

sion of the terminal decapeptide of kisspeptin to rhesus monkeys is capable of inducing a homologous desensitization of the *KISS1R* receptor despite retained GnRH responsiveness at the level of the gonadotrope [62]. Thus, the discovery of this previously unsuspected pathway in human reproduction that has now been discovered to represent a key control point of GnRH at sexual maturation in all mammals studied to date has ignited a burst of basic and clinical research into the *KISS1/KISS1R* axis in mammalian physiology. This system offers a further approach to therapeutically modulate human GnRH function and potentially translate the biology of this pathway into wider clinical applications across diverse reproductive phenotypes.

Fibroblast Growth Factor Signalling and GnRH Ontogeny

The family of 23 fibroblast growth factors (FGFs) and their 4 cognate receptors (FGFRs) have multiple roles in central nervous and skeletal system development and include brain patterning, branching morphogenesis and limb development [63]. In 2 individuals with a contiguous gene syndrome with a dominant form of KS, Dodé et al. [64] first identified an overlapping interstitial deletion at chromosome 8p11–p12 using fluorescent in situ hybridization and reported loss-of-function mutations in *FGFR1* as the underlying cause of KS. This finding not only convincingly demonstrated *FGFR1* to be *KAL2* but also explained the long-standing paradox of patients with both GnRH deficiency and hereditary spherocytosis due to ankyrin deficiency. Both these genes are located in a region quite close to the *GnRH* gene itself which had not been identified as a cause of GnRH deficiency until recently. Again, it was GnRH-deficient patients with mutations in this gene that heralded the recognition of the relatively crucial role of the FGFs in mammalian reproduction.

There are 23 mammalian FGFs that act by binding and activating one of the 4 FGFR family of tyrosine kinase receptors in a heparan sulfate proteoglycan-dependent manner [63]. Four *FGFR* genes (*FGFR1–FGFR4*) encode their respective receptors. Each of these 4 FGFRs consists of three extracellular immunoglobulin domains (D1–D3), a single-pass transmembrane domain and a cytoplasmic tyrosine kinase domain [65] (fig. 8a). An 'acidic box' consisting of a serine-rich sequence present in the D1–D2 linker and in conjunction with the D1 domain is thought to mediate autoinhibition [63]. Within the first

Fig. 8. FGF pathway in human GnRH deficiency. **a** Schematic of *FGFR1* protein displaying mutations causing KS (blue) and nIHH (black). The extracellular domain of *FGFR1* contains a signal peptide (SP), three extracellular immunoglobulin-like (D1, D2, and D3) domains, followed by the TM, and an intracellular domain comprising two tyrosine kinase subdomains (PTK). The acidic box (AB) and the docking protein FRS2 domain in the extracellular domain and intracellular domain, respectively, are specific features of FGF receptors. **b** Genomic structure and differential splicing of the human *FGF8* gene. **i** Structure of the *FGF8* gene. Boxes denote exons; lines denote introns. **ii** Schematic of the four *FGF8* isoforms identified in humans with mutations identified to date are indicated by arrows and numbered according to the *FGF8f* and *FGF8b* protein isoforms (* homozygous mutations). Reproduced from Falardeau et al. [71].

half of D2 there is a site that binds the FGFR coreceptor, heparan sulfate proteoglycan. The D2 and D3 domains are responsible for ligand binding and specificity. Alternative splicing of the *FGFR* genes produces 7 principal isoforms and each isoform exhibits differential ligand binding specificities [66]. After binding of their cognate ligand(s) and heparan sulfate proteoglycan(s), the FGFRs dimerize, resulting in autophosphorylation and stimulation of the protein tyrosine kinase activity. This activation step is followed by the stimulation of an extensive intracellular signalling cascade involving various signal proteins, coreceptor proteins and various sec-

ond messengers that are described in detail elsewhere [63]. This *FGFR1* signalling activity is tightly regulated and in quiescent cells, the acidic box domain of these receptors is thought to bind to the positively charged heparan sulfate binding site and result in a 'closed' inactive state that is in equilibrium with the 'open' active one. Upon ligand availability, the equilibrium shifts towards the 'open' state.

A large number of individual point mutations and deletions of the *FGFR1* gene have now been reported in subjects with both KS [26, 42] and nIHH [67] (fig. 8a). These mutations span all the functional domains of the

FGFR1 receptor and, in contrast to *KALI*, account for up to 10% of each KS and nIHH patient series [67, 68]. Structural and biochemical studies using recombinant *FGFR1* proteins to study the structure-activity relationships of these mutations have been particularly informative. Mutations in the D2 and D3 loops of the receptor were shown to cause misfolding and impaired cell-surface expression of the receptor and/or ligand-receptor interaction and binding [27, 67]. Mutations in the cytoplasmic tyrosine kinase domain appear to cause structural perturbations and reduce catalytic activity of this domain [67]. At least 11 different FGFs can activate *FGFR1* [69]. However, the specific ligand implicated in GnRH neuron ontogeny was unknown until a proband with KS and cleft palate with a L342S mutation in the *FGFR1c* gene demonstrated that this mutation results in a selective and dramatic loss of binding affinity of the FGF8b isoform to the *FGFR1* receptor. Integrating this observation with the described overlapping patterns of FGF8 and *FGFR1* expression in the brain and defective olfactory bulb neurogenesis and loss of fate specification of GnRH neurons seen in FGF8 hypomorphic mice [70], a candidate gene approach revealed six *FGF8* gene mutations in GnRH-deficient unrelated probands [71] (fig. 8b). Two of the probands also carried additional *FGFR1* mutations confirming oligogenicity in the FGF8 signalling pathway in human GnRH deficiency. The structural and functional impact of these *FGF8* mutations were confirmed by a crystal structure analysis of the FGF8b/*FGFR2c*/heparin model and reduced *FGF8* function in vitro [71].

These findings of *FGFR1* and *FGF8* mutations in both KS and nIHH add a new dimension to the complexity of GnRH ontogeny. Impaired FGF signalling can clearly result in loss of fate specification of GnRH neurons and/or GnRH migratory defects with impaired olfactory epithelium/bulb neurogenesis resulting in the KS phenotype and through alternative mechanisms affect GnRH neuronal survival beyond olfactory neurogenesis resulting in a nIHH phenotype. The effect of FGF signalling on olfactory bulb neurogenesis may, in fact, relate to the association between anosmin and *FGFR1* proteins. Both anosmin-1 and *FGFR1* bind to heparan sulfate proteoglycans. Anosmin-1 has been shown in vitro to enhance the mitogenic effect of the *FGFR1c* isoform on lymphoid cells in the presence of FGF2 [72]. These above associations have led to the postulation that X-linked *KALI* mutations may be secondary to a FGF signalling defect on GnRH neuronal development [73]. FGF signalling has previously been implicated in GnRH neurite outgrowth

and GnRH axonal guidance to the median eminence and this observation suggests a chemoattractant role for FGF8 [74]. The importance of the FGF signalling pathway in GnRH neuron ontogeny is further supported by the murine models with targeted ablation of *FGFR1* in the telencephalon [75] and *FGF8* hypomorphic mice [70], both of which lack olfactory bulbs. In addition, mice expressing a dominant-negative form of *FGFR1* in their GnRH neurons and *FGF8* hypomorphic mice exhibit reduced or absent GnRH neurons in the hypothalamus [76]. Since complete reversal of GnRH deficiency has been documented in subjects carrying mutations in *FGF8* and *FGFR1*, there may well be a role for FGF signalling in GnRH neuronal survival and apoptosis pathways [26, 31, 71, 77].

Prokineticin 2/Prokineticin Receptor 2 and GnRH Ontogeny

Targeted gene deletions in animals often unearth unsuspected phenotypes. The prokineticin system is a prime example of this phenomenon as it was initially studied because of its importance in gastrointestinal function. Prokineticins (PROK1 and 2) are the mammalian protein orthologs of proteins discovered in the black mamba snake's venomous secretions [78]. PROKR1 and PROKR2 are their respective receptors. These two receptors differ little in their ligand preference since they are >85% homologous in their DNA sequence. In fact, they were originally thought to be two different isoforms of the same receptor (i.e. GPR73a and GPR73b) until it was appreciated that they were encoded by two different genes. However, unanticipated hypogonadotropism occurred in murine deletions of the PROK2 pathway. While *PROK1*^{-/-} [79] and *PROKR1*^{-/-} [80] animals lack any CNS phenotypes, *PROK2*^{-/-} [79, 81] and *PROKR2*^{-/-} [80] mice show olfactory bulb anomalies (fig. 9b), decreased GnRH neuronal migration to the hypothalamus, and hypogonadotropic hypogonadism responsive to exogenous gonadotropins. Although the olfactory bulb morphogenesis is completely disrupted in *PROKR2*^{-/-} mice, in 50% of *PROK2*^{-/-} mice, some olfactory bulb morphogenesis is preserved, suggesting that although PROK2 plays an important role in neuronal migration, several other key molecules/gene products are likely to be involved in the orchestration of GnRH neuronal migration and olfactory bulb development. Likewise, the evolving role of oligogenicity in GnRH deficiency suggests that although individual gene prod-

Fig. 9. Prokineticin 2 pathway and human GnRH deficiency. **a** Schematic of the predicted PROKR2 protein displaying mutations causing KS (yellow boxes) and nIHH (white boxes). **b** Morphological examination of wild-type *PROK2*^{+/+} and *PROK2*^{-/-} mice showing hypoplasia of olfactory bulbs (red circle). Reproduced with permission from Ng et al. [79]. **c** In vitro activity of a selection of *PROKR2* mutants in an intracellular Ca²⁺ assay. Reproduced with permission from Cole et al. [90].

ucts play a critical role in determining the reproductive phenotype, interaction and synergy between various genes make a cumulative contribution to the eventual clinical phenotype.

The ligands, PROK1 (85 amino acids) and PROK2 (81 amino acids), are multifunctional secreted proteins with only 44% shared homology [78]. However, both ligands share 10 highly conserved cysteines that presumably pair in 5 bridges and have an identical basic N-terminal amino acid sequence, AVITGA, that is essential for their biological activity [82]. PROKR1 and PROKR2 are G-protein-coupled receptors whose major biological differences lie more in their differential pattern of expression than in their ability to distinguish between ligands. Previously, the 'prokineticin system' drew attention because of its broadly diverse biologic functions including its ability to stimulate gastrointestinal motility [83], angiogenesis [84], hematopoiesis [85] and its effect on pain modulation [86]. In contrast to the more widely expressed PROKR1,

the ligand PROK2, and its cognate receptor, PROKR2, have a unique expression profile within the CNS including olfactory system, arcuate nucleus, suprachiasmatic nuclei (SCN) and median eminence [79, 80]. Yet another fascinating feature of PROK2 is its known function as a chemoattractant for neural progenitor cells that ultimately populate the olfactory bulb and assist in its dynamic function during life [79]. Within the CNS, therefore, the combined anatomic expression pattern of PROK2/PROKR2 matches with systems that coordinate smell, reproduction, temperature, and circadian rhythms – all critical factors for successful evolution of species. Sensing animals of the opposite sex ready for breeding, determining internal reproductive readiness, and facilitating sensing of light/dark changes are all functions essential for reproductive 'fitness'.

Because of the strong expression of *PROK2*/*PROKR2* in the SCN and of *PROKR2* in the paraventricular nucleus, a key SCN output target for the endocrine system, this

signalling system is a key candidate for a potential link between the reproductive and circadian systems. In addition, the induction of *PROK2*'s expression by the clock genes, *BMAL1* and *CLOCK* [79], support the notion that this ligand/receptor pair are strong candidates to serve as a potential second messenger system from the SCN's master clock to the more peripheral and diverse circadian endocrine rhythms such as the reproductive system. Furthermore, both *PROK2*^{-/-} and *PROKR2*^{-/-} animals exhibit disruptions of some of their circadian rhythms including abnormal thermogenesis, increased nocturnal physical activity, impaired circadian cortisol, and abnormal glucose regulation [87–89].

In keeping with these observations in *PROK2*^{-/-} and *PROKR2*^{-/-} mice, humans with loss-of-function mutations in both *PROK2* and *PROKR2* have been documented to have both KS and nIHH [81, 90, 91] (fig. 9a, c). Similar to the subjects with *FGF8/FGFR1* mutations, considerable phenotypic variability is evident within family members and some asymptomatic individuals carry identical mutations to other affected individuals within sibships. This variable expressivity and incomplete penetrance that are now emerging in several genes causing GnRH deficiency strongly suggest the existence of a digenic/oligogenic phenomenon as previously documented [90, 91]. Subjects with mutations in the *PROK2/PROKR2* system also have been documented to undergo reversal of their hypogonadotropism following treatment with sex steroids. This observation suggests that some degree of GnRH neuronal plasticity in adulthood, possibly modulated by sex steroids, is responsible for their improvement [90]. Any connection between the integrity of the circadian system in the SCN and the GnRH system is yet to be determined in the humans. However, several nonreproductive phenotypes are seen with prokineticin mutations including sleep disorders and obesity [90, 91]. However, detailed circadian assessment in patients with mutations in this system will be required to ascertain if prokineticin is a major output molecule linking the circadian clock and reproduction.

GNRH1* and *GNRHR

The human *GNRH1* gene is located at 8p21–8p11.2, consists of 4 exons and encodes the preprohormone that is ultimately processed to produce GnRH decapeptide [92]. Mutation in *GNRH1* is an obvious candidate as an etiology for human GnRH deficiency and in keeping with this, in the hypogonadal (hpg) mouse model, a homozygous deletion of the *GNRH1* gene resulted in hypogonad-

otropic hypogonadism [14]. Although human *GNRH1* mutations have been elusive for many years, two independent groups have recently described homozygous frameshift *GNRH1* mutations in patients with GnRH deficiency [93, 94]. In both studies, consistent with the critical role played by GnRH, subjects with homozygous mutations in *GNRH1* showed severe hypogonadism with affected males having micropallus. In addition to homozygous frameshift mutations, heterozygous rare sequence variants in *GNRH1* have also been described in one of these studies [93]. As seen with other genes implicated in GnRH neuronal ontogeny, it is likely that oligogenicity and genotypic synergy with known/unknown genes may operate to produce the phenotype. No non-reproductive features were reported in these patients.

The amino acid sequence of GnRHR was first deduced for the mouse receptor cloned from the pituitary α T3 gonadotrope cell line [95]. The human *GNRHR* gene maps to chromosome 4q13.2–13.3 and is comprised of three exons that encode a 328-amino acid protein, with >85% homology within mammalian species, with near identity in the transmembrane domains. The GnRHR contains seven transmembrane (TM) domains, six of which alternate extra- and intracellular loops with an extracellular amino terminus. However, GnRHR is unique among the rhodopsin family of GPCRs in its lack of an intracellular carboxy terminus [96]. The extracellular domains and superficial regions of the TMs are involved in binding of GnRH, and the TMs are believed to be involved in receptor configuration and conformational change associated with signal propagation (receptor activation) [97]. These changes are thought to propagate into conformational changes in the intracellular domains involved in interacting with G proteins and other proteins for intracellular signal transduction [97]. GnRH binds to the GnRHR in a hairpin structure with the amino- and carboxy-terminal domains contributing mainly to receptor binding and activation [98].

Since the original reports of *GNRHR* mutations causing nIHH [99], a variety of inactivating mutations have been described. Most are missense mutations and a significant number of compound heterozygous changes are seen. The most common mutations occur in the first extracellular and third intracellular loops, although they span across the receptor. The first extracellular loop mutations reduce ligand affinity and the third intracellular loop mutations reduce signal transduction [100]. Recently, a cell-permeant small molecule that was a GnRH antagonist was shown to rescue most of the naturally occurring mutants by increasing their expression, presumably

by stabilizing their intracellular processing and transport [101]. These small molecular ‘chaperones’ offer future therapeutic options for patients with *GNRHR* and other GPCR mutations.

GNRHR mutations can account for up to 40% of familial cases of nIHH and perhaps up to 17% of sporadic cases of nIHH [102]. Patients with *GNRHR* mutations present with a wide spectrum of reproductive symptoms ranging from severe hypogonadotropism including microphallus and undescended testes in males at birth to failure of pubertal development in adolescence [100] as well as infertility in adults [24, 99, 103]. However, partial defects are also seen with significant variations in phenotypes despite similar genetic functional defects [102]. This variability of clinical phenotypes presumably reflects several issues including the severity of intrinsic disruption of the *GNRHR* processing and/or function, the dosing of genes involved (heterozygous, biallelic/homozygous or compound heterozygous mutations), the coexistence of mutations in other GnRH deficiency causing genes (oligogenicity) [29], and as yet unapparent epigenetic and environmental factors.

TAC3 and TACR3

Using a similar strategy of studying consanguineous families with nIHH that led to the key discovery of the *KISS1R* mutations in human GnRH deficiency, Topaloglu et al. [104] recently reported loss-of-function mutations in both *TACR3*, the gene encoding a G protein-coupled receptor, neurokinin B (NKB) receptor and *TAC3*, the gene encoding NKB, the ligand for *TACR3*. This human genetic investigation highlights a hitherto unrecognized role of the *TAC3/TACR3* pathway in regulation of the GnRH pulse generator. Neurokinin B belongs to a phylogenetically conserved family of proteins which also includes substance P, neurokinin A and hemokinin-1 [105]. *TACR3* is predominantly expressed in the CNS including the hypothalamus [106] with high and preferential binding to NKB [107].

Male patients with *TACR3* mutations characteristically have micropenis and fail to enter puberty during adolescence. These reproductive phenotypes strongly suggest an important role of the *TAC3/TACR3* pathway in both the ‘mini-puberty’ of infancy and gonadotropin activation at puberty. None of the patients reported in the first human study had KS, suggesting a primary role of this *TAC3/TACR3* pathway in functional integrity of the GnRH pulse generator. The receptor *TACR3* is expressed

on the GnRH axons and the KNDy (kisspeptin, neurokinin and dynorphin) neuronal population and neurons coexpressing *KISS1* and *TAC3* have been described in the arcuate nucleus and postulated to be the primary pathway mediating the sex-steroid feedback to the GnRH neurons [108, 109]. However, murine deletion of the murine ortholog of *TACR3* has not been associated with reproductive abnormalities as seen in humans [110]. Recently, Gianetti et al. have reported several *TAC3/TACR3* mutations in a large outbred cohort of patients with nIHH [111]. In this detailed report, micropenis was observed in a majority of male subjects. In addition, evidence of neuroendocrine recovery of the hypothalamo-pituitary-gonadal axis was observed in a significant number of male and female subjects during adulthood. These observations strongly suggest that the *TAC3/TACR3* signaling pathway is critical during the neonatal period and puberty, but is relatively less-critical in adulthood. No non-reproductive features have been reported in patients with *TACR3* mutations, but *TAC3* mutation have been associated with learning disabilities [104]. In addition, the potential oligogenic interactions of the *TAC3/TACR3* pathway with the other genes related to GnRH ontogeny will need to be determined. The hypothalamic interplay between *KISS1* and *NKB* is already well recognized [109, 112, 113] and this interaction is likely to be a subject of intense study in the coming years.

NELF

The human nasal embryonic LHRH factor gene, *NELF*, emerged as a strong candidate gene for KS following its strong association with axonal guidance of olfactory and GnRH neurons in mice [114]. Although a unique rare sequence variant in *NELF* has been reported in human GnRH deficiency, the biology of *NELF* in humans is still unclear [115]. However, the emerging theme of oligogenicity in human GnRH deficiency implies that *NELF* may be a critical modifier gene that synergizes with other pathogenic genes to result in the observed human phenotype [29].

Genetics of Other Multisystem Disorders Associated with KS/nIHH

Various multisystem disorders with overlapping features of KS/nIHH have been reported (table 1). Mutations of the *DAX1* (*NROB1*) gene (dosage-sensitive sex

reversal-adrenal hypoplasia congenita critical region on the X chromosome gene 1), encoding a novel orphan nuclear receptor, have been identified in patients with X-linked adrenal hypoplasia congenita and nIHH [116]. Mutations in both leptin (*LEP*) and leptin receptor (*LEPR*) are associated with obesity and hypogonadotropic hypogonadism [117]. Heterozygous mutations in *CHD7* (chromodomain helicase DNA-binding protein), a gene that encodes a large chromatin-modelling CHD7 protein, cause the CHARGE syndrome (eye Coloboma, Heart anomalies, choanal Atresia, growth and developmental Retardation, Genitourinary anomalies and Ear abnormalities) [118]. Some patients with CHARGE syndrome also share overlapping features of KS [119]. Recently, mutations in *CHD7* have been identified in patients with both KS and nIHH [120], and in particular, KS/nIHH subjects who show presence of some of the CHARGE syndrome features such as deafness, dysmorphic ears and/or hypoplasia of the semicircular canals in addition to their hypogonadism, are more likely to harbor mutations in *CHD7* [121].

Future Directions

In the last decade, the study of humans with GnRH deficiency has been prismatic in revealing the fascinating ontogeny of GnRH neurons. However, several key

molecules/pathways that govern GnRH neuronal migration remain undiscovered and since pathogenic genetic mutations only account for approximately 40% of such patients, a substantial number of genes underlying GnRH neuron ontogeny are yet to be discovered. In addition, the emerging evidence of oligogenicity, modifier genes and potential gene-environment interactions emphasize the pressing need for an integrated investigatory approach encompassing detailed human phenotyping, use of next generation gene sequencing to aid novel gene discovery and the judicious use of bioinformatics. Although genotyping of GnRH deficiency is currently done predominantly in a research setting in most countries, the accumulating genetic knowledge will eventually allow the incorporation of genetic testing and genetic counseling into standard clinical care of these patients. The precise genetic basis of the onset of human puberty has been cited as one of the critical unanswered questions in the 21st century [122]. The prismatic human model of GnRH deficiency presents a unique biologic opportunity to unravel the molecular basis of the arousal of the GnRH pulse generator to signal puberty.

References

- 1 Wray S, Hoffman G: A developmental study of the quantitative distribution of LHRH neurons within the central nervous system of postnatal male and female rats. *J Comp Neurol* 1986;252:522–531.
- 2 Merchenthaler I, et al: Gonadotropin-releasing hormone (GnRH) neurons and pathways in the rat brain. *Cell Tissue Res* 1984;237:15–29.
- 3 Grumbach MM: A window of opportunity: the diagnosis of gonadotropin deficiency in the male infant. *J Clin Endocrinol Metab* 2005;90:3122–3127.
- 4 Conn PM, Crowley WF Jr: Gonadotropin-releasing hormone and its analogs. *Annu Rev Med* 1994;45:391–405.
- 5 Antunes JL, et al: Luteinizing hormone-releasing hormone in human pituitary blood. *J Neurosurg* 1978;49:382–386.
- 6 Terasawa E: Luteinizing hormone-releasing hormone (LHRH) neurons: mechanism of pulsatile LHRH release. *Vitam Horm* 2001; 63:91–129.
- 7 Lehman MN, et al: Immunocytochemical localization of luteinizing hormone-releasing hormone (LHRH) pathways in the sheep brain during anestrus and the mid-luteal phase of the estrous cycle. *J Comp Neurol* 1986;244:19–35.
- 8 Schwanzel-Fukuda M, Pfaff DW: Origin of luteinizing hormone-releasing hormone neurons. *Nature* 1989;338:161–164.
- 9 Wray S, Grant P, Gainer H: Evidence that cells expressing luteinizing hormone-releasing hormone mRNA in the mouse are derived from progenitor cells in the olfactory placode. *Proc Natl Acad Sci USA* 1989;86: 8132–8136.
- 10 Wray S, Nieburgs A, Elkabes S: Spatiotemporal cell expression of luteinizing hormone-releasing hormone in the prenatal mouse: evidence for an embryonic origin in the olfactory placode. *Brain Res Dev Brain Res* 1989;46:309–318.
- 11 Fueshko S, Wray S: LHRH cells migrate on peripherin fibers in embryonic olfactory explant cultures: an in vitro model for neurophilic neuronal migration. *Dev Biol* 1994; 166:331–348.
- 12 Schwanzel-Fukuda M: Origin and migration of luteinizing hormone-releasing hormone neurons in mammals. *Microsc Res Tech* 1999;44:2–10.
- 13 Belchetz PE, et al: Hypophysial responses to continuous and intermittent delivery of hypothalamic gonadotropin-releasing hormone. *Science* 1978;202:631–633.
- 14 Gibson MJ, et al: Mating and pregnancy can occur in genetically hypogonadal mice with preoptic area brain grafts. *Science* 1984;225: 949–951.
- 15 Gibson MJ, Miller GM, Silverman AJ: Pulsatile luteinizing hormone secretion in normal female mice and in hypogonadal female mice with preoptic area implants. *Endocrinology* 1991;128:965–971.

- 16 Cariboni A, et al: The product of X-linked Kallmann's syndrome gene (KAL1) affects the migratory activity of gonadotropin-releasing hormone (GnRH)-producing neurons. *Hum Mol Genet* 2004;13:2781–2791.
- 17 Maestre de San Juan A: Teratologia: falta total de los nervios olfactorios con anosmia en un individuo en quien existia una atrofia congénita de los testículos y miembro viril. *Siglo Medico* 1856;3:211–221.
- 18 Kallmann F, Schoenfeld W, Barrera S: The genetic aspects of primary eunuchoidism. *Am J Ment Defic* 1944;48:203–236.
- 19 de Morsier G: Etudes sur les dysraphies crânio-encephaliques. *Schweiz Arch Neurol Psychiatr* 1954;74:309–361.
- 20 Schally AV, et al: Isolation and properties of the FSH and LH-releasing hormone. *Biochem Biophys Res Commun* 1971;43:393–399.
- 21 Amoss M, et al: Purification, amino acid composition and N-terminus of the hypothalamic luteinizing hormone releasing factor (LRF) of ovine origin. *Biochem Biophys Res Commun* 1971;44:205–210.
- 22 Naftolin F, Harris GW, Bobrow M: Effect of purified luteinizing hormone releasing factor on normal and hypogonadotropic anomic men. *Nature* 1971;232:496–497.
- 23 Hoffman AR, Crowley WF Jr: Induction of puberty in men by long-term pulsatile administration of low-dose gonadotropin-releasing hormone. *N Engl J Med* 1982;307:1237–1241.
- 24 de Roux N, et al: Loss of function mutations of the GnRH receptor: a new cause of hypogonadotropic hypogonadism. *J Pediatr Endocrinol Metab* 1999;12(suppl 1):267–275.
- 25 Waldstreicher J, et al: The genetic and clinical heterogeneity of gonadotropin-releasing hormone deficiency in the human. *J Clin Endocrinol Metab* 1996;81:4388–4395.
- 26 Pitteloud N, et al: Reversible Kallmann syndrome, delayed puberty, and isolated anosmia occurring in a single family with a mutation in the fibroblast growth factor receptor 1 gene. *J Clin Endocrinol Metab* 2005;90:1317–1322.
- 27 Pitteloud N, et al: Mutations in fibroblast growth factor receptor 1 cause Kallmann syndrome with a wide spectrum of reproductive phenotypes. *Mol Cell Endocrinol* 2006;254–255:60–69.
- 28 Nachtigall LB, et al: Adult-onset idiopathic hypogonadotropic hypogonadism – a treatable form of male infertility. *N Engl J Med* 1997;336:410–415.
- 29 Pitteloud N, et al: Digenic mutations account for variable phenotypes in idiopathic hypogonadotropic hypogonadism. *J Clin Invest* 2007;117:457–463.
- 30 Dodé C, Hardelin JP: Kallmann syndrome. *Eur J Hum Genet* 2009;17:139–146.
- 31 Raivio T, et al: Reversal of idiopathic hypogonadotropic hypogonadism. *N Engl J Med* 2007;357:863–873.
- 32 Ballabio A, et al: X-linked ichthyosis, due to steroid sulphatase deficiency, associated with Kallmann syndrome (hypogonadotropic hypogonadism and anosmia): linkage relationships with Xg and cloned DNA sequences from the distal short arm of the X chromosome. *Hum Genet* 1986;72:237–240.
- 33 Bick D, et al: Male infant with ichthyosis, Kallmann syndrome, chondrodysplasia punctata, and an Xp chromosome deletion. *Am J Med Genet* 1989;33:100–107.
- 34 Schwanzel-Fukuda M, Bick D, Pfaff DW: Luteinizing hormone-releasing hormone (LHRH)-expressing cells do not migrate normally in an inherited hypogonadal (Kallmann) syndrome. *Brain Res Mol Brain Res* 1989;6:311–326.
- 35 Franco B, et al: A gene deleted in Kallmann's syndrome shares homology with neural cell adhesion and axonal path-finding molecules. *Nature* 1991;353:529–536.
- 36 Legouis R, et al: The candidate gene for the X-linked Kallmann syndrome encodes a protein related to adhesion molecules. *Cell* 1991;67:423–435.
- 37 Hardelin JP: Kallmann syndrome: towards molecular pathogenesis. *Mol Cell Endocrinol* 2001;179:75–81.
- 38 Hu Y, et al: Cross-talk of anosmin-1, the protein implicated in X-linked Kallmann's syndrome, with heparan sulphate and urokinase-type plasminogen activator. *Biochem J* 2004;384:495–505.
- 39 Bulow HE, et al: Heparan sulfate proteoglycan-dependent induction of axon branching and axon misrouting by the Kallmann syndrome gene kal-1. *Proc Natl Acad Sci USA* 2002;99:6346–6351.
- 40 Hardelin JP, et al: Anosmin-1 is a regionally restricted component of basement membranes and interstitial matrices during organogenesis: implications for the developmental anomalies of X chromosome-linked Kallmann syndrome. *Dev Dyn* 1999;215:26–44.
- 41 Cadman SM, et al: Molecular pathogenesis of Kallmann's syndrome. *Horm Res* 2007;67:231–242.
- 42 Kim SH, et al: Diversity in fibroblast growth factor receptor 1 regulation: learning from the investigation of Kallmann syndrome. *J Neuroendocrinol* 2008;20:141–163.
- 43 Oliveira LM, et al: The importance of autosomal genes in Kallmann syndrome: genotype-phenotype correlations and neuroendocrine characteristics. *J Clin Endocrinol Metab* 2001;86:1532–1538.
- 44 Quinton R, et al: Idiopathic gonadotrophin deficiency: genetic questions addressed through phenotypic characterization. *Clin Endocrinol (Oxf)* 2001;55:163–174.
- 45 Hardelin JP, Dodé C: The complex genetics of Kallmann syndrome: KAL1, FGFR1, FGF8, PROKR2, PROK2, et al. *Sex Dev* 2008;2:181–193.
- 46 Tsai PS, Gill JC: Mechanisms of disease: insights into X-linked and autosomal-dominant Kallmann syndrome. *Nat Clin Pract Endocrinol Metab* 2006;2:160–171.
- 47 Andrenacci D, et al: Functional dissection of the Drosophila Kallmann's syndrome protein DmKal-1. *BMC Genet* 2006;7:47.
- 48 Hardelin JP, et al: Heterogeneity in the mutations responsible for X chromosome-linked Kallmann syndrome. *Hum Mol Genet* 1993;2:373–377.
- 49 de Roux N, et al: Hypogonadotropic hypogonadism due to loss of function of the KiSS1-derived peptide receptor GPR54. *Proc Natl Acad Sci USA* 2003;100:10972–10976.
- 50 Seminara SB, et al: The GPR54 gene as a regulator of puberty. *N Engl J Med* 2003;349:1614–1627.
- 51 Muir AI, et al: AXOR12, a novel human G protein-coupled receptor, activated by the peptide KiSS-1. *J Biol Chem* 2001;276:28969–28975.
- 52 Colledge WH: Kisspeptins and GnRH neuronal signalling. *Trends Endocrinol Metab* 2009;20:115–121.
- 53 Ohtaki T, et al: Metastasis suppressor gene KiSS-1 encodes peptide ligand of a G-protein-coupled receptor. *Nature* 2001;411:613–617.
- 54 Chan YM, Broder-Fingert S, Seminara SB: Reproductive functions of kisspeptin and Gpr54 across the life cycle of mice and men. *Peptides* 2009;30:42–48.
- 55 Tenenbaum-Rakover Y, et al: Neuroendocrine phenotype analysis in five patients with isolated hypogonadotropic hypogonadism due to a L102P inactivating mutation of GPR54. *J Clin Endocrinol Metab* 2007;92:1137–1144.
- 56 Semple RK, et al: Two novel missense mutations in g protein-coupled receptor 54 in a patient with hypogonadotropic hypogonadism. *J Clin Endocrinol Metab* 2005;90:1849–1855.
- 57 Pallais JC, et al: Neuroendocrine, gonadal, placental, and obstetric phenotypes in patients with IHH and mutations in the G-protein coupled receptor, GPR54. *Mol Cell Endocrinol* 2006;254–255:70–77.
- 58 Castellano JM, et al: KiSS-1/kisspeptins and the metabolic control of reproduction: physiologic roles and putative pathophysiological implications. *Peptides* 2009;30:139–145.
- 59 Hauge-Evans AC, et al: A role for kisspeptin in islet function. *Diabetologia* 2006;49:2131–2135.
- 60 Robertson JL, et al: Circadian regulation of Kiss1 neurons: implications for timing the preovulatory gonadotropin-releasing hormone/luteinizing hormone surge. *Endocrinology* 2009;150:3664–3671.
- 61 Teles MG, et al: A GPR54-activating mutation in a patient with central precocious puberty. *N Engl J Med* 2008;358:709–715.

- 62 Plant TM, Ramaswamy S: Kisspeptin and the regulation of the hypothalamic-pituitary-gonadal axis in the rhesus monkey (*Macaca mulatta*). *Peptides* 2009;30:67–75.
- 63 Beenken A, Mohammadi M: The FGF family: biology, pathophysiology and therapy. *Nat Rev Drug Discov* 2009;8:235–253.
- 64 Dodé C, et al: Loss-of-function mutations in FGFR1 cause autosomal dominant Kallmann syndrome. *Nat Genet* 2003;33:463–465.
- 65 Mohammadi M, Olsen SK, Ibrahimi OA: Structural basis for fibroblast growth factor receptor activation. *Cytokine Growth Factor Rev* 2005;16:107–137.
- 66 Yeh BK, et al: Structural basis by which alternative splicing confers specificity in fibroblast growth factor receptors. *Proc Natl Acad Sci USA* 2003;100:2266–2271.
- 67 Pitteloud N, et al: Mutations in fibroblast growth factor receptor 1 cause both Kallmann syndrome and normosmic idiopathic hypogonadotropic hypogonadism. *Proc Natl Acad Sci USA* 2006;103:6281–6286.
- 68 Trarbach EB, et al: Novel fibroblast growth factor receptor 1 mutations in patients with congenital hypogonadotropic hypogonadism with and without anosmia. *J Clin Endocrinol Metab* 2006;91:4006–4012.
- 69 Zhang X, et al: Receptor specificity of the fibroblast growth factor family: the complete mammalian FGF family. *J Biol Chem* 2006;281:15694–15700.
- 70 Meyers EN, Lewandoski M, Martin GR: An Fgf8 mutant allelic series generated by Cre- and Flp-mediated recombination. *Nat Genet* 1998;18:136–141.
- 71 Falardeau J, et al: Decreased FGF8 signaling causes deficiency of gonadotropin-releasing hormone in humans and mice. *J Clin Invest* 2008;118:2822–2831.
- 72 Gonzalez-Martinez D, et al: Anosmin-1 modulates fibroblast growth factor receptor 1 signaling in human gonadotropin-releasing hormone olfactory neuroblasts through a heparan sulfate-dependent mechanism. *J Neurosci* 2004;24:10384–10392.
- 73 Dodé C, Hardelin JP: Kallmann syndrome: fibroblast growth factor signaling insufficiency? *J Mol Med* 2004;82:725–734.
- 74 Gibson MJ, Ingraham L, Dobrjansky A: Soluble factors guide gonadotropin-releasing hormone axonal targeting to the median eminence. *Endocrinology* 2000;141:3065–3071.
- 75 Hebert JM, et al: FGF signaling through FGFR1 is required for olfactory bulb morphogenesis. *Development* 2003;130:1101–1111.
- 76 Tsai PS, et al: Targeted expression of a dominant-negative fibroblast growth factor (FGF) receptor in gonadotropin-releasing hormone (GnRH) neurons reduces FGF responsiveness and the size of GnRH neuronal population. *Mol Endocrinol* 2005;19:225–236.
- 77 Storm EE, Rubenstein JL, Martin GR: Dosage of Fgf8 determines whether cell survival is positively or negatively regulated in the developing forebrain. *Proc Natl Acad Sci USA* 2003;100:1757–1762.
- 78 Li M, et al: Identification of two prokineticin cDNAs: recombinant proteins potently contract gastrointestinal smooth muscle. *Mol Pharmacol* 2001;59:692–698.
- 79 Ng KL, et al: Dependence of olfactory bulb neurogenesis on prokineticin 2 signaling. *Science* 2005;308:1923–1927.
- 80 Matsumoto S, et al: Abnormal development of the olfactory bulb and reproductive system in mice lacking prokineticin receptor PKR2. *Proc Natl Acad Sci USA* 2006;103:4140–4145.
- 81 Pitteloud N, et al: Loss-of-function mutation in the prokineticin 2 gene causes Kallmann syndrome and normosmic idiopathic hypogonadotropic hypogonadism. *Proc Natl Acad Sci USA* 2007;104:17447–17452.
- 82 Bullock CM, Li JD, Zhou QY: Structural determinants required for the bioactivities of prokineticins and identification of prokineticin receptor antagonists. *Mol Pharmacol* 2004;65:582–588.
- 83 Schweitz H, et al: MIT(1), a black mamba toxin with a new and highly potent activity on intestinal contraction. *FEBS Lett* 1999;461:183–188.
- 84 LeCouter J, Lin R, Ferrara N: The role of EG-VEGF in the regulation of angiogenesis in endocrine glands. *Cold Spring Harb Symp Quant Biol* 2002;67:217–221.
- 85 LeCouter J, et al: Bv8 and endocrine gland-derived vascular endothelial growth factor stimulate hematopoiesis and hematopoietic cell mobilization. *Proc Natl Acad Sci USA* 2004;101:16813–16818.
- 86 Negri L, et al: Nociceptive sensitization by the secretory protein Bv8. *Br J Pharmacol* 2002;137:1147–1154.
- 87 Hu WP, et al: Altered circadian and homeostatic sleep regulation in prokineticin 2-deficient mice. *Sleep* 2007;30:247–256.
- 88 Cheng MY, et al: Prokineticin 2 transmits the behavioural circadian rhythm of the suprachiasmatic nucleus. *Nature* 2002;417:405–410.
- 89 Prosser HM, et al: Prokineticin receptor 2 (Prokr2) is essential for the regulation of circadian behavior by the suprachiasmatic nuclei. *Proc Natl Acad Sci USA* 2007;104:648–653.
- 90 Cole LW, et al: Mutations in prokineticin 2 and prokineticin receptor 2 genes in human gonadotrophin-releasing hormone deficiency: molecular genetics and clinical spectrum. *J Clin Endocrinol Metab* 2008;93:3551–3559.
- 91 Dodé C, et al: Kallmann syndrome: mutations in the genes encoding prokineticin-2 and prokineticin receptor-2. *PLoS Genet* 2006;2:e175.
- 92 Cheng CK, Leung PC: Molecular biology of gonadotropin-releasing hormone (GnRH)-I, GnRH-II, and their receptors in humans. *Endocr Rev* 2005;26:283–306.
- 93 Chan YM, et al: GNRH1 mutations in patients with idiopathic hypogonadotropic hypogonadism. *Proc Natl Acad Sci USA* 2009;106:11703–11708.
- 94 Bouligand J, et al: Isolated familial hypogonadotropic hypogonadism and a GNRH1 mutation. *N Engl J Med* 2009;360:2742–2748.
- 95 Tsutsumi M, et al: Cloning and functional expression of a mouse gonadotropin-releasing hormone receptor. *Mol Endocrinol* 1992;6:1163–1169.
- 96 Stojilkovic SS, Reinhart J, Catt KJ: Gonadotropin-releasing hormone receptors: structure and signal transduction pathways. *Endocr Rev* 1994;15:462–499.
- 97 Millar RP, et al: Gonadotropin-releasing hormone receptors. *Endocr Rev* 2004;25:235–275.
- 98 Karten MJ, Rivier JE: Gonadotropin-releasing hormone analog design. Structure-function studies toward the development of agonists and antagonists: rationale and perspective. *Endocr Rev* 1986;7:44–66.
- 99 de Roux N, et al: A family with hypogonadotropic hypogonadism and mutations in the gonadotropin-releasing hormone receptor. *N Engl J Med* 1997;337:1597–1602.
- 100 Karges B, Karges W, de Roux N: Clinical and molecular genetics of the human GnRH receptor. *Hum Reprod Update* 2003;9:523–530.
- 101 Leanos-Miranda A, Janovick JA, Conn PM: Receptor-misrouting: an unexpectedly prevalent and rescuable etiology in gonadotropin-releasing hormone receptor-mediated hypogonadotropic hypogonadism. *J Clin Endocrinol Metab* 2002;87:4825–4828.
- 102 Beranova M, et al: Prevalence, phenotypic spectrum, and modes of inheritance of gonadotropin-releasing hormone receptor mutations in idiopathic hypogonadotropic hypogonadism. *J Clin Endocrinol Metab* 2001;86:1580–1588.
- 103 Pitteloud N, et al: The fertile eunuch variant of idiopathic hypogonadotropic hypogonadism: spontaneous reversal associated with a homozygous mutation in the gonadotropin-releasing hormone receptor. *J Clin Endocrinol Metab* 2001;86:2470–2475.
- 104 Topaloglu AK, et al: TAC3 and TACR3 mutations in familial hypogonadotropic hypogonadism reveal a key role for neurokinin B in the central control of reproduction. *Nat Genet* 2009;41:354–358.
- 105 Pennefather JN, et al: Tachykinins and tachykinin receptors: a growing family. *Life Sci* 2004;74:1445–1463.
- 106 Mileusnic D, et al: Neurokinin-3 receptor distribution in rat and human brain: an immunohistochemical study. *Neuroscience* 1999;89:1269–1290.

- 107 Satake H, Kawada T: Overview of the primary structure, tissue-distribution, and functions of tachykinins and their receptors. *Curr Drug Targets* 2006;7:963–974.
- 108 Rance NE, Bruce TR: Neurokinin B gene expression is increased in the arcuate nucleus of ovariectomized rats. *Neuroendocrinology* 1994;60:337–345.
- 109 Rance NE: Menopause and the human hypothalamus: evidence for the role of kisspeptin/neurokinin B neurons in the regulation of estrogen negative feedback. *Peptides* 2009;30:111–122.
- 110 Siuciak JA, et al: Disruption of the neurokinin-3 receptor (NK3) in mice leads to cognitive deficits. *Psychopharmacology (Berl)* 2007;194:185–195.
- 111 Gianetti E, et al: *TAC3/TACR3* mutations reveal preferential activation of gonadotropin-releasing hormone release by neurokinin B in neonatal life followed by reversal in adulthood. *J Clin Endocrinol Metab* 2010;95:2857–2867.
- 112 Wakabayashi Y, et al: Neurokinin B and dynorphin A in kisspeptin neurons of the arcuate nucleus participate in generation of periodic oscillation of neural activity driving pulsatile gonadotropin-releasing hormone secretion in the goat. *J Neurosci* 2010;30:3124–3132.
- 113 Navarro VM, et al: Regulation of gonadotropin-releasing hormone secretion by kisspeptin/dynorphin/neurokinin B neurons in the arcuate nucleus of the mouse. *J Neurosci* 2009;29:11859–11866.
- 114 Kramer PR, Wray S: Novel gene expressed in nasal region influences outgrowth of olfactory axons and migration of luteinizing hormone-releasing hormone (LHRH) neurons. *Genes Dev* 2000;14:1824–1834.
- 115 Miura K, Acierno JS Jr, Seminara SB: Characterization of the human nasal embryonic LHRH factor gene, NELF, and a mutation screening among 65 patients with idiopathic hypogonadotropic hypogonadism (IHH). *J Hum Genet* 2004;49:265–268.
- 116 Yu RN, et al: The role of DAX-1 in reproduction. *Trends Endocrinol Metab* 1998;9:169–175.
- 117 Farooqi S, O’Rahilly S: Genetics of obesity in humans. *Endocr Rev* 2006;27:710–718.
- 118 Vissers LE, et al: Mutations in a new member of the chromodomain gene family cause CHARGE syndrome. *Nat Genet* 2004;36:955–957.
- 119 Pinto G, et al: CHARGE syndrome includes hypogonadotropic hypogonadism and abnormal olfactory bulb development. *J Clin Endocrinol Metab* 2005;90:5621–5626.
- 120 Kim HG, et al: Mutations in *CHD7*, encoding a chromatin-remodeling protein, cause idiopathic hypogonadotropic hypogonadism and Kallmann syndrome. *Am J Hum Genet* 2008;83:511–519.
- 121 Jongmans MC, et al: *CHD7* mutations in patients initially diagnosed with Kallmann syndrome – the clinical overlap with CHARGE syndrome. *Clin Genet* 2009;75:65–71.
- 122 Pennisi E: What determines species diversity? *Science* 2005;309:90.