Abdominal Adhesion Prevention: Still a Sticky Subject?

Chris I.W. Lauder Giuseppe Garcea Andrew Strickland Guy J. Maddern

University of Adelaide Discipline of Surgery, The Queen Elizabeth Hospital, Woodville, S.A., Australia

Key Words

Adhesions • Peritoneum • Adhesion barrier • Adhesion prevention • Abdominal surgery

Abstract

Background: Adhesion formation remains an almost inevitable consequence of abdominal procedures, potentially resulting in significant morbidity and mortality. There is an ongoing need to evaluate current understanding of adhesion formation and products aimed at prevention. Failure to keep up to date with adhesion treatment may subject clinicians to a greater medico-legal risk. **Design:** Review of published studies exploring the problem of peritoneal adhesion formation. This encompasses the underlying processes of adhesion formation combined with general approaches to reduce formation. An overview of products trialled to prevent formation in both the animal model and clinical setting describes products of scientific interest and commercial success. **Results:** Advances in surgical technique, such as laparoscopic surgery, can help minimize the probability of adhesion formation. Currently barrier products, whilst reducing adhesion formation, have not been shown to reduce the risk of readmission with complications related to adhesions. Hybrid products may improve upon this situation. **Conclusions:** No single approach has been wholly satisfactory in reducing adhesions. Research into the processes driving adhesion formation is providing exciting new targets for therapeutic agents. It would seem plausible that with many promising avenues of research a revolutionary agent to reduce the incidence of adhesional small bowel obstruction may result.

Copyright © 2010 S. Karger AG, Basel

Introduction

Adhesion formation following surgery remains an almost inevitable consequence of most abdominal procedures. Studies have found the incidence of adhesions to be as high as 95% [1]. Though surgeons are aware of adhesions and the problems that can result, it seems that few routinely take measures to reduce their formation. Advances in surgical technique, such as the use of laparoscopic surgery, can help minimize the probability of adhesion formation [2]. However, the significant morbidity and cost associated with adhesion-related disorders continues to be highly prevalent. Estimates on the workload for the treatment of adhesion-related disorders have put the annual cost in the USA at around USD 1.3 billion [3]. Undoubtedly, these figures will continue to increase with the growing cost of health care and medico-legal repercussions of adhesion-related complications. Whilst many methods have been employed in an attempt to reduce the formation of adhesions, no single approach has been wholly satisfactory. A concerted effort is required from all branches of medical science and surgery if the problem of adhesions is to be conquered. Continued ignorance of the subject, or a failure to adopt new practices aimed at adhesion prevention is simply not acceptable in the current litigious healthcare environment.

KARGER

Fax +41 61 306 12 34
E-Mail karger@karger.ch
www.karger.com

© 2010 S. Karger AG, Basel
0253-4886/10/0275-0347\$26.00/0

Accessible online at:
www.karger.com/dsu

Prof. G.J. Maddern
University of Adelaide Discipline of Surgery, The Queen Elizabeth Hospital
28 Woodville Road
Woodville South, S.A. 5011 (Australia)
Tel. +61 8 8222 6756, Fax +61 8 8222 6028, E-Mail guy.maddern@adelaide.edu.au

Consequences of Abdominal Adhesions

The development of intra-abdominal adhesions may manifest in a number of ways and can be categorized as primary or secondary. In primary instances, patients who would not have otherwise required operative intervention require surgery to treat adhesive small bowel obstruction [4, 5], volvulus or chronic abdominal pain [6]. In patients with secondary problems from adhesions, future surgery (for abdominal pathology other than adhesions) may be complicated [7] or prolonged [8, 9]. In such cases, the number and density of adhesions will have a direct impact on the second surgical procedure. In female patients, adhesions within the pelvis may result in infertility [10, 11], ectopic gestation [12] and chronic pelvic pain [13, 14].

Patients undergoing surgery for symptomatic adhesions may also require repeat operations, with each instance associated with an increasing risk of morbidity. Repeated bowel resection or enterotomy as a result of adhesion surgery exposes patients to the risk of anastomotic leak and can increase the chances of developing a variety of other chronic conditions such as short bowel syndrome [15] or enterocutaneous fistulae [16]. Recurrent adhesion-forming patients are perhaps the group with the most to gain from adhesion prevention to break the cycle of re-formation.

Although there is a wide spectrum of adhesive response, the clinically relevant outcome is dependent on patient symptoms or the need for operative intervention. It is difficult to identify individual patients with an increased risk of developing symptomatic adhesions since the relationship between adhesion formation and the surgical pathologies that occur as a consequence is somewhat variable. Logic would dictate that a greater number of adhesions should cause more symptoms; however, this is not always the case. Surgeons will be familiar with the scenario of a single band adhesion causing small bowel obstruction in an otherwise unremarkable abdomen.

Pathogenesis of Adhesion Formation

Adhesions should be considered as highly cellular, vascularized and dynamic structures under the influence of complex signaling pathways. The idea of an adhesion being a simple nonfunctioning scar is an outdated concept [17]. It is now understood why many of the processes occur in response to an initial adhesion-stimulating event. Damage to the peritoneal mesothelium promotes

an acute inflammatory response, with subsequent pathways dictating the number and strength of the fibrous adhesions that form (fig. 1). The mechanisms responsible for a change in normal peritoneal healing to those where adhesions propagate are complex.

Fibrinolysis and extracellular matrix remodeling may be as important in regulating the adhesive process as controlling inflammation in the early stages. The difficulty in modulating adhesion formation lies in separating the process of adhesion formation from that of normal wound healing. Studies examining adhesion formation on a genetic level in an animal model have demonstrated that a large number of genes are upregulated at the wound site postoperatively [18]. Further work is required to identify those genes with a specific role in adhesion formation within the profiles already identified.

On a molecular level, the factors determining cell proliferation, migration, differentiation, angiogenesis, apoptosis and host defense will all have an impact on adhesion formation. Targets for the control of these processes include chemokines [19, 20] and matrix metalloproteinases [21]. Vascular endothelial growth factor has also been implicated in having an early role in adhesion initiation [22]. Chemokines trigger a signaling cascade which is responsible for adherence of leukocytes to specific sites in the blood vessels and extravasation into different tissue sites. As such, they can recruit leukocytes into inflammatory areas following surgery. Several cytokines are continuously produced by macrophages and fibroblasts entrapped within the fibrin meshwork [23, 24], including TNF- α , IL-1, PDGF, EGF and MCP-1, which in turn recruit leukocytes and increase collagen synthesis [25]. As long as these conditions persist, adhesions will proliferate. The driving forces for this are the presence of foreign material, persistent infection or trauma.

Incidence of Adhesions

Assessing the true incidence of adhesion formation is a very difficult task. Asymptomatic adhesions have little chance of being discovered unless the patient has further abdominal surgery, or is noted to have adhesions at autopsy. Numerous studies have looked at the incidence of adhesion formation in isolation; however, the key outcome measure in terms of health care provision is the risk of readmission following the original procedure. This will include patients who may not have undergone further surgery, but have repeated admissions with adhesion-related complications. Of the studies looking at this

Fig. 1. Pathways of adhesion formation or resolution.

risk, the SCAR-3 study [26] is perhaps the best, following a large cohort of patients for an extended period of time. A 5% risk of readmission directly related to adhesions in the 5 years following open abdominal surgery (excluding appendectomy) was demonstrated. In a retrospective American study investigating readmission rates following open colorectal or general surgical procedures, the readmission rate (with a diagnosis of intestinal obstruction) within 2 years ranged from 12.4 to 17% [4]. Another retrospective study reported proven adhesion-related readmission rates following gynecological surgery at 1.5 to 2% over a 4-year period [27]. When admission data included patients with symptoms not proven to be due to adhesions, the figure is substantially higher ranging from 14.5 to 16.1%. This discrepancy highlights the difficulty in isolating iatrogenic adhesion formation in the presence of non-specific symptoms like pain or with coexisting disease such as pelvic inflammatory disease which can produce de novo adhesions. Coding techniques often differ amongst studies which will also have an effect on reporting of patient morbidities.

A final factor in considering the workload associated with the treatment of adhesions is the ageing population.

As the mean survival age increases, the probability of performing more than one surgical procedure on any given patient may also increase.

Models of Adhesion Formation

The majority of studies investigating adhesion formation utilize an animal model to reproduce surgical insults, although they vary greatly depending on the interests of the investigating party. A lack of uniformity makes collation and interpretation of results between studies difficult. Most models utilize small animals such as the rat or mouse [20, 28–53]; however, the desire to test laparoscopic procedures has necessitated large animal studies [54–56].

The methods used to promote adhesion formation are similarly disparate. Some studies mimic standard surgical procedures on a smaller scale [57], while others use a variety of means to produce much more pronounced adhesions [58]. Although the underlying processes are the same, some stimuli provide a reaction that is unlikely to be replicated in the clinical setting, for example insertion

of large areas of exposed polypropylene mesh [44, 56, 59]. Results from these 'extreme' models therefore need to be interpreted with caution and their use limited solely to a research tool.

A key issue in assessing adhesions in any context (particularly when assessing therapies to reduce adhesions) is the severity scoring criteria used. Although there has been some agreement between committees on the use of a standardized scoring system for the evaluation of adhesions [60, 61], no single method has received widespread acceptance. The two fundamental principles at odds in a method to classify adhesions are those of simplicity versus reproducibility; a scoring system with an infinite number of classifications is unlikely to be popular since it will be difficult to remember, apply and analyze. Conversely, a system that is too simple may fail to demonstrate the difference between treatment arms in adhesion reduction studies, under reporting the true spectrum of disease. Certain criteria would seem essential in the design of adhesion scoring system. The number of sites at which adhesions are present must be recorded as well as noting adhesion length. Further assessment of adhesion strength is necessary, although this can be somewhat subjective. Studies have negated this problem to some degree by using video of procedures to allow an independent observer's interpretation of adhesion strength to be compared with the operating surgeon [62–64]. Typically, the expression of adhesion density starts with filmy adhesions easily taken down by blunt dissection, progressing in density where sharp dissection is required and finally firmly adherent, where no discernable plane is evident.

There are numerous systems based upon these concepts [65–67], although little agreement between studies makes accurate comparison of results difficult. The lack of uniformity between study design and result reporting also makes a meta-analysis of existing data impossible.

Clinical Studies of Adhesion Formation

Examining adhesion reduction at the surgical site provides a direct measure of the effectiveness of an adhesion prevention regimen; however, few studies have recorded rates of readmission after treatment. To collect accurate data for adhesion-related readmission, the time period needs to be long and the cohort large to demonstrate statistical significance. A study investigating this principle estimated that around 15,000 patients would need to be screened to show a reduced risk of readmission in a trial

lasting at least 3.5 years [68]. This is with the assumption that the adhesion reduction product has a 25% efficacy. Clearly, if a product is more effective, then the numbers of patients required to demonstrate statistical significance would be lower.

Adhesion assessment in the clinical setting does not necessarily require an extra procedure as it can be part of a phased reconstruction such as the reversal of a loop ileostomy [69]. Some gynecological treatment algorithms may also involve a second-look laparoscopy [63, 64, 70–72]. In these circumstances, it is possible to directly examine the extent of any adhesions formed.

With the initial use of any new adhesion reduction strategy relook assessment is essential given the quiescent nature of many adhesions. Once the effectiveness of an anti-adhesion agent has been established for a particular surgical operation, longer-term follow-up should confirm the expected effect of a reduced risk of adhesion-related readmission.

Methods of Adhesion Reduction

The process of adhesion formation can be thought of as a cycle since failed attempts at adhesion prevention may well lead to further surgical intervention (fig. 2). At each point in this cycle, there is the potential to minimize the formation of adhesions which may be influenced by the following methods.

Surgical Technique

Since the impact of adhesion-related disorders was realized, there have been numerous attempts to reduce their formation. Changes in surgical technique aim to reduce the trauma associated with a procedure by minimizing tissue handling and practicing meticulous hemostasis. The nidus for persistent inflammation must also be curtailed, including synthetic material such as sutures as well as bacterial soiling, blood or other necrotic material. Aids to achieving these surgical goals include the use of hemostatic agents as well as electrosurgical instruments and laparoscopy. Research investigating the use of specific surgical instruments in isolation have not shown a statistically significant beneficial effect in the reduction of adhesions [73–77]. However, any technique or instrument reducing procedure time, or blood loss, should decrease the physiological insult for the patient, with a reduction in adhesions as a consequence.

Topical products to improve hemostasis have also been shown to provide a beneficial role in reducing adhe-

Fig. 2. Adhesion cycle and points of manipulation.

sion formation [78]. Many of the early products used to promote hemostasis show similarities in their composition to adhesion prevention treatments. For example Surgicel® (Johnson & Johnson, Somerville, N.J., USA) consists of oxidized regenerated cellulose. Topical thrombin [79–81] and fibrin [81–83] sealants have evidence to show their efficacy in aiding hemostasis and as such can be thought of as adjuncts in an adhesion reduction strategy.

Laparoscopic versus Open Surgery and Natural Orifice Transluminal Endoscopic Surgery

There are many benefits to performing a procedure by the minimally invasive method, though adhesion reduction is often not considered. Good evidence exists supporting a reduction in adhesion formation following laparoscopic surgery for a large number of procedures [54, 55, 84, 85]. There are potential disadvantages associated with laparoscopic procedures such as a longer duration of surgery with resulting damage to peritoneal mesothelium from the pneumoperitoneum [28, 29, 86, 87]. This is due to a combination of ischemia, acidosis and desiccation. Amelioration of peritoneal damage may be achieved by heating and humidifying the CO₂ as well as reducing

gas flow rate and insufflation pressures [88–90]. Although some damage to the peritoneum is inevitable, protective measures such as these may allow quicker resolution of any insult.

A variation in the application of minimally invasive surgery has led to the development of natural orifice transluminal endoscopic surgery [91]. It is conceivable that this method may lead to reduced adhesion formation by virtue of the fact that there are no transabdominal ports. However, there is insufficient evidence to support this concept at present [92].

Adhesion Reduction Agents – An Overview

Adhesion reduction agents can be broadly separated into two categories. The first are the pharmacological therapies given around the time of the patient's operation. The second encompasses topical products applied directly to the operative site.

Pharmacological Therapy

Many different pharmacological agents have been tried to achieve adhesion reduction [30–43, 93–103]. There is sound logic behind the use of such agents, although their effectiveness has been somewhat limited.

Reports in the literature describe the effective use of a large number of pharmacological agents in experimental animal studies [30–43, 93, 94, 96, 97, 99–103]. However, few agents progress to clinical trials. The focus of this review is to explore the generic classes of drugs based upon their mode of action within the adhesion formation pathways.

Pharmacological agents have also been applied topically in an attempt to enhance the local effects of the drug. Some systemic absorption is inevitable with drugs administered intraperitoneally, although this is a complex process which depends on the properties of the agent in question. The peritoneum is highly permeable to water, small solutes, and proteins and therefore is not a physical barrier [104]. This porosity can lead to rapid absorption of the agent thus limiting any benefit to intraperitoneal application. No strong evidence exists to suggest that the addition of pharmacological agents to fluid left within the abdomen leads to a reduction in adhesions.

Drugs to mediate the inflammatory response following surgery have been trialled extensively. Initial attempts used steroids [30, 93], and nonsteroidal anti-inflammatory drugs [31–33, 94, 95] (NSAIDs). The balance between systemic side effects and adhesion reduction has proved difficult to overcome for this group of agents. The risk of bleeding associated with NSAIDs, or impaired wound healing with steroid use, has limited their use in the clinical setting. More recently, free radical scavengers [34, 96, 97] including methylene blue [105] as well as inhibitors of pro-inflammatory cytokines [35] and antihistamines [36, 37, 72–74] have been trialled. Although there have been promising results from animal studies, no single agent has progressed to widespread clinical use.

Secondary targets include the processes determining fibrin deposition [38, 98] and fibrinolysis [39, 99–103]. Despite low-dose anticoagulation being the accepted standard for deep vein thrombosis prophylaxis in surgery, the use of therapeutic anticoagulation to reduce fibrin deposition has not been shown to have a statistically significant effect on adhesion reduction. Furthermore, this approach is associated with an increased risk of postoperative bleeding. The alternative approach of promoting fibrinolysis has also been met with concern due to the potential for postoperative bleeding [106].

Adhesion reduction by the inhibition of fibroblast proliferation [40, 41] with agents such as mitomycin C has also been trialled, though toxic side effects have limited its use.

Other emerging products include those targeted towards the sites of adhesion formation by inhibition of

chemokines. Experimental studies of broad-spectrum chemokine blockade have demonstrated a reduction in adhesion formation in the murine model [42]. Further selective chemokine inhibition such as the prevention of CCL1-CCR8 interaction [43] has also been shown to be effective in experimental models, although clinical trials are awaited.

Topical Products

The topical products utilized in prevention of intra-abdominal adhesions can be divided into two basic categories. The first consists of liquids which are instilled to the abdominal cavity at the end of the operation. Liquids rely on the principal of flotation – since the bowel is freely floating within the abdominal cavity and thus separated to some degree. The simplest form of this method was the application of Ringer's lactate or other crystalloid fluid [107]. There is no significant evidence to support adhesion reduction with this method since the fluid is rapidly absorbed before any surgical injury has healed. Modification of this principle led to the addition of solutes to the fluid to decrease the rate at which absorption occurs. The resultant products are similar to peritoneal dialysis solutions, and have shown some promise in adhesion prevention [108, 109].

The second category of topical products consists of gels or films. The basic principle is the separation of the operative surfaces with a mechanical barrier. The barrier must be retained for a suitably long period of time for the surgical insult to heal sufficiently so that adhesions will not form. In order to achieve this, there are certain properties that the product must display. For reabsorption to occur, the material must be easily degraded without a fibrous reaction. Popular constituents which meet this criteria include viscous polysaccharides such as cellulose, chitosan, dextran or hyaluronic acid. Interestingly, chitosan, a naturally occurring polysaccharide formed from the deacetylation of chitin, has also been shown to have other beneficial properties that make it a desirable constituent for use in an adhesion prevention product. One aspect is the hemostatic nature of chitosan, with successful use in dressings [110, 111] which are now issued to US army personnel. Another is the antimicrobial action of chitosan to a variety of bacteria, fungi and viruses [112]. Chitosan has shown promising results in animal studies as an adhesion prevention product [51, 58]. Synthetic alternatives include polylactic acid or polyethylene glycol. There is little to choose between these basic materials, although subtle modification of their form may have a significant effect on the polymer properties.

Some crossover exists between gels and films, many of the films taking on more of a gel type consistency during reabsorption. The exception is the application of a PTFE (Goretex®) film between the injured peritoneal surfaces. Whilst results describing the use of PTFE films have been encouraging [113–115], the application of a nonabsorbable foreign body within the abdomen has been criticized. Not only because of the need for later removal (although this has been debated) but also the potential for chronic infection and difficulty associated with fixation of the film to prevent migration.

Modification of Surgical Materials

Another challenge to adhesion prevention is the presence of a foreign body such as nonabsorbable sutures or mesh. Coating of the latter has shown some success [44, 56, 59] in applications where a side of the mesh will potentially come into contact with the bowel. Evaluation of six commercially available meshes in an experimental model showed the absorbable layers of Parietex Composite® and C-Qur® were effective at reducing adhesions in the short term. However, as the absorbable layer was phagocytosed from the mesh, the adhesions reduction was seen to diminish [118]. The simple solution is to avoid or minimizes the use of nonabsorbable materials within the abdomen whenever possible. When this is unavoidable, using adhesion reduction products to isolate the operative site may help to reduce adhesions in the immediate postoperative period. However, suture material is frequently found at the site of important vascular or enteric anastomosis, where an adhesion response is essential in order to reduce the probability of subsequent anastomotic leakage. As such, isolation of the area may be the best compromise.

Evidence-Based Use of Adhesion Prevention Products

An abundance of published literature describing the use of adhesion reduction products exists; however, the number of good quality randomized control trials is somewhat less. Products that have had the most documented success in humans are the absorbable gels and films. Although the use of some agents is not without certain caveats (such as the avoidance of wrapping the product around any anastomosis, and ensuring complete hemostasis), randomized control trials have shown statistically significant benefits in their ability to reduce adhesion formation [117–125]. As such, the predominance of evidence would suggest that unless there are any contraindications, the surgeon should consider using hyaluronic acid/carboxymethyl membranes. This is of particular rel-

evance in patients undergoing staged surgery. Unfortunately, despite a proven reduction in the incidence, extent and severity of adhesions, a decrease in the incidence of intestinal obstruction or need for operative intervention has yet to be shown [126].

A tabulated summary of those products which have been commercially marketed with proven clinical effectiveness is shown in table 1. This gives the opportunity for a rapid review of a product's constituents, any adverse outcomes, and the evidence supporting its use.

Considerations for the Use of Adhesion Prevention Products

The choice of product used may be influenced by the mode of application. Films/solid membranes may prove difficult to apply laparoscopically. An alternative is an in situ cross-linkable hydrogel that can be applied as a liquid which rapidly gels at the desired site of action. There is no one product that is currently available without certain drawbacks. This may be a result of a sensitivity to the product itself, for example with Hyskon® where patients reacted to the antigenic properties of the constituents [127]. Alternatively, the product may increase the incidence of anastomotic leakage or wound dehiscence such as with InterGel®. Adverse outcomes during clinical studies were considered serious enough to cause a product recall [124, 128]. Other instances where an increased risk of anastomotic leakage has been observed are with the use of Seprafilm®. Although the increase in leak rate was relatively low [114], the manufacturers now recommend avoiding wrapping the product around anastomoses. Clinical trials using Interceed® observed that unless complete hemostasis was achieved prior to the application of the product, it may actually precipitate adhesion formation [129].

More minor problems associated with the use of adhesion prevention products center around their application. Many of the more solid sheet preparations require fixation. This makes laparoscopic use a challenge and may necessitate the use of suture material which itself can act as a nidus for adhesion formation. The opposite is true for liquid preparations where the challenge is to keep the solution within the abdomen without leakage from the surgical site. Product storage can be difficult if refrigeration or heating is required prior to use, with specialized applicators using compressed gas or the need for a dedicated light source also deterring routine use.

Table 1. Adhesion-reducing products

Brand name (Company)	Composition and properties	Adverse issues	Current availability	Evidence [ref.]
Adcon-P (Gliatech, Inc.)	Gelatin/proteoglycan Hydrophilic, bioresorbable	FDA recalled related product Adcon-L	Animal studies only as of 2003	46, 47
Adept (Innovata)	4% icodextrin solution Hydrophilic, peritoneal residence time >4 days	Leakage of fluid from the surgical wound, and abdominal distension/discomfort, pulmonary effusion	FDA approved for gynecological laparoscopic procedures in August 2006, marketed in Europe for general and gynecological surgery	108, 109
FloGel (Alliance Pharm, Co.)	Poloxamer 407 Hydrophilic, thermosensitive, bioresorbable	Requires refrigeration	Early clinical trials	48, 49
FocalGel (Focal)	Polyethylene glycol and polylactic acid Photo-cross-linkable; bioresorbable	Requires dedicated light source	Marketed for general surgical use	50
Hyskon (Medisan Pharmaceuticals)	Dextran 70 Hydrophilic, bioresorbable	Local and systemic side effects due to osmotic and antigenic properties Not sufficient evidence of clinical effectiveness	Indicated as hysteroscopy fluid	127, 131–133
Incert (Anika)	Chemically crosslinked hyaluronic acid Hydrophilic, bioresorbable	Difficult laparoscopic application	Pilot human trials in the UK from 2004, Incert-S marketed for spinal surgery	134
Interceed (Ethicon)	Oxidized regenerated cellulose membrane Hydrophilic, absorbed within 3–10 days	Limited effectiveness in the presence of blood and peritoneal fluids	Marketed for general surgical use	66, 67, 135, 136
Intergel (Lifecore)	0.5% ferric hyaluronate gel Hydrophilic, elimination half-life within the peritoneum 51 h	Withdrawn from clinical studies due to high morbidity associated with postoperative peritonitis/anastomotic leak. Clinical study discontinued	Discontinued	124, 128, 137
NOCC (Kytogenics)	N,O-carboxymethyl chitosan Hydrophilic, bioresorbable	Pilot clinical trial demonstrated no significant difference between treatment and control groups	Phase III clinical trial (2002)	51, 52, 64
Preclude (W.L. Gore)	Expanded polytetrafluoroethylene membrane Hydrophobic, not biodegradable	Requires strict sterility and fixation; difficult to use laparoscopically and leaves foreign body in the peritoneum	Marketed mainly for gynecological surgery	113–115
Repel (Life Medical Sciences)	Polyethylene glycol and polylactic acid block copolymer membrane Bioresorbable	Needs fixation; difficult to use laparoscopically	Repel-CV FDA approved in March 2009 for cardiac surgery	138, 139
Sepracoat (Genzyme)	Sodium hyaluronate solution Hydrophilic, elimination half-life within the peritoneum: 26 h	Insufficient evidence of clinical effectiveness	Marketed in Europe for general surgical use (Not FDA approved)	140
Seprafilm (Genzyme)	Hyaluronic acid-carboxymethylcellulose membrane Hydrophilic, absorbed within 2 weeks	Requires careful handling at open operation; difficult to use laparoscopically	Marketed for general surgery use	117–123, 125
Sepragel (Genzyme)	Hyaluronic acid-carboxymethylcellulose gel Hydrophilic, bioresorbable	Sepragel spine withdrawn from clinical trials 2002	Indicated for ENT surgery as a space-occupying gel	67
SprayGel (Confluent surgical)	In situ cross-linkable polyethylene glycol gel Hydrophilic, bioresorbable in 6 days	Requires a specialized applicator and air supply	Marketed for general surgery use	63, 68
Surgiwrap sheet (MAST Biosurgery)	Polylactide membrane Hydrophobic, bioresorbable in 24 weeks	Needs fixation	FDA approved in May 2005, also marketed in Europe for general surgical use	53

Discussion

Surgery in its essence is a rather 'blunt tool', injuring the tissues and organs at the operative site. In order for the body to recover from this injury, there must be an adequate healing response. For the most part, this response is a normal part of wound healing with a number of factors differentiating between resolution and pathological adhesion formation. Targeted molecular-level therapy provides an attractive panacea to the problem of adhesions; however, the reality may be somewhat different. Overmodulation of the mechanisms driving this response may lead to an inadequate reaction resulting in complications such as bleeding, anastomotic leak or even local cancer recurrence.

Any product designed to reduce adhesions may be overwhelmed if the stimulating adhesive process is on a massive scale, for example gross fecal contamination of the abdominal cavity. In this context, any product used to modulate the formation of adhesions may be ineffective despite attempts to minimize bacterial soiling at the time of the operation. Patients who are systemically unwell often have a number of other clinical problems, such as coagulopathies, which contribute to the difficulty of adhesion reduction in the acute setting. Such extreme circumstances are difficult to replicate in the research setting since animal models will often not survive them.

No product will be a substitute for good surgical technique, and the best approach to adhesion reduction will be a combination of treatments designed to minimize the adhesion-forming process from the outset. Whilst it is not possible to recommend one specific product for every situation, a number of therapies are currently available that have a good evidence base for their use and should be considered by the operating surgeon. Foresight is required to ensure that products are available at the time of elective surgery so that an opportunity to reduce adhesions is not missed. It is a duty of care that surgeons should be aware of a suitable adhesion reduction strategy relevant to their practice. Failing to discuss adhesions with patients during the consent process is to pave the way for potential negligence claims [130].

Continuing research is providing advances in the field which promise more effective products in adhesion reduction. Barrier gels or films can now be modified to include new pharmacological agents [45] further modulating the adhesion-forming process. Such hybrid solutions may provide a synergistic effect to overcome adhesion formation in most settings. As these new therapies emerge, they will clearly require thorough validation in their use to ensure that patient safety is not compromised.

References

- 1 Menzies D, Ellis H: Intestinal obstruction from adhesions – how big is the problem? *Ann R Coll Surg Engl* 1990;72:60–63
- 2 Gutt CN, Oniu T, Schemmer P, et al: Fewer adhesions induced by laparoscopic surgery? *Surg Endosc* 2004;18:898–906.
- 3 Ray NF, Denton WG, Thamer M, et al: Abdominal adhesiolysis: inpatient care and expenditures in the United States in 1994. *J Am Coll Surg* 1998;186:1–9.
- 4 Beck DE, Opelka FG, Bailey HR, et al: Incidence of small-bowel obstruction and adhesiolysis after open colorectal and general surgery. *Dis Colon Rectum* 1999;42:241–248; erratum in *Dis Colon Rectum* 1999;42:578.
- 5 Ellis H: The clinical significance of adhesions: focus on intestinal obstruction. *Eur J Surg Suppl* 1997;577:5–9.
- 6 Sulaiman H, Gabella G, Davis C, et al: Presence and distribution of sensory nerve fibers in human peritoneal adhesions. *Ann Surg* 2001;234:256–261.
- 7 Van Der Krabben AA, Dijkstra FR, Nieuwenhuijzen M, et al: Morbidity and mortality of inadvertent enterotomy during adhesiotomy. *Br J Surg* 2000;87:467–471.
- 8 Beck DE, Ferguson MA, Opelka FG, et al: Effect of previous surgery on abdominal opening time. *Dis Colon Rectum* 2000;43:1749–1753.
- 9 Coleman MG, McLain AD, Moran BJ: Impact of previous surgery on time taken for incision and division of adhesions during laparotomy. *Dis Colon Rectum* 2000;43:1297–1299.
- 10 Marana R, Rizzi M, Muzii L, et al: Correlation between the American Fertility Society classifications of adnexal adhesions and distal tubal occlusion and reproductive outcome in tubal surgery. *Fertil Steril* 1995;64:924–929.
- 11 Stovall TG, Elder RF, Ling FW: Predictors of pelvic adhesions. *J Reprod Med* 1989;34:345–348.
- 12 Liakakos T, Thomakos N, Fine PM, et al: Peritoneal Adhesions: Etiology, Pathophysiology, and Clinical Significance. *Dig Surg* 2001;18:260–273.
- 13 Duffy DM, diZerega GS: Adhesion controversies: pelvic pain as a cause of adhesions, crystalloids in preventing them. *J Reprod Med* 1996;41:19–26.
- 14 Howard FM: The role of laparoscopy in chronic pelvic pain: promise and pitfalls. *Obstet Gynecol Surv* 1993;48:357–387.
- 15 Thompson JS, DiBaise JK, Iyer KR, et al: Postoperative short bowel syndrome. *J Am Coll Surg* 2005;201:85–89.
- 16 van Goor H: Consequences and complications of peritoneal adhesions. *Colorectal Dis* 2007;9(suppl 2):25–34.
- 17 Cheong YC, Laird SM, Li TC, et al: Peritoneal healing and adhesion formation/reformation. *Hum Reprod Update* 2001;7:556–566.
- 18 Yu SL, Singh S, Chen HW, et al: Intra-abdominal adhesion formation induces anti-oxidative injury, enhances cell proliferation, and prevents complement-mediated lysis. *Wound Repair Regen* 2008;16:388–398.
- 19 Chegini N, Rong H, Bennett B, et al: Peritoneal fluid cytokine and eicosanoid levels and their relation to the incidence of peritoneal adhesion. *J Soc Gynecol Investig* 1999;6:153–157.
- 20 Hoshino A, Kawamura YI, Yasuhara M, et al: Inhibition of CCL1-CCR8 interaction prevents aggregation of macrophages and development of peritoneal adhesions. *J Immunol* 2007;178:5296–5304.

- 21 Chegini N, Kotseos K, Zhao Y, et al: Expression of matrix metalloproteinase (MMP-1) and tissue inhibitor of MMP in serosal tissue of intraperitoneal organs and adhesions. *Fertil Steril* 2001;76:1212–1219.
- 22 Cahill RA, Wang JH, Soohkai S, Redmond HP: Mast cells facilitate local VEGF release as an early event in the pathogenesis of postoperative peritoneal adhesions. *Surgery* 2006;140:108–112.
- 23 Kovacs EJ, DiPietro LA: Fibrogenic cytokines and connective tissue production. *FASEB J* 1994;8:854–861.
- 24 Zeyneloglu HB, Senturk LM, Seli E, et al: The role of monocyte chemotactic protein-1 in intraperitoneal adhesion formation. *Hum Reprod* 1998;13:1194–1199.
- 25 Kovacs EJ: Fibrogenic cytokines: the role of immune mediators in the development of scar tissue. *Immunol Today* 1991;12:17–23.
- 26 Parker MC, Wilson MS, Menzies D, et al: The SCAR-3 study: 5-year adhesion-related readmission risk following lower abdominal surgical procedures. *Colorectal Dis* 2005;7:551–558.
- 27 Lower AM, Hawthorn RJ, Ellis H, et al: The impact of adhesions on hospital readmissions over ten years after 8,849 open gynaecological operations: an assessment from the Surgical and Clinical Adhesions Research Study. *BJOG* 2000;107:855–862.
- 28 Molinas CR, Mynbaev O, Pauwels A, et al: Peritoneal mesothelial hypoxia during pneumoperitoneum is a cofactor in adhesion formation in a laparoscopic mouse model. *Fertil Steril* 2001;76:560–567.
- 29 Rosário MT, Ribeiro U Jr, Corbett CE, et al: Does CO₂ pneumoperitoneum alter the ultra-structure of the mesothelium? *J Surg Res* 2006;133:84–88.
- 30 Höckel M, Ott S, Siemann U, et al: Prevention of peritoneal adhesions in the rat with sustained intraperitoneal dexamethasone delivered by a novel therapeutic system. *Ann Chir Gynaecol* 1987;76:306–313.
- 31 LeGrand EK, Rodgers KE, Girgis W, et al: Efficacy of tolmetin sodium for adhesion prevention in rabbit and rat models. *J Surg Res* 1994;56:67–71.
- 32 Guvenal T, Cetin A, Ozdemir H, et al: Prevention of postoperative adhesion formation in rat uterine horn model by nimesulide: a selective COX-2 inhibitor. *Hum Reprod* 2001;16:1732–1735.
- 33 De Leon F, Toledo A, Sanfilippo J, et al: The prevention of adhesion formation by nonsteroidal antiinflammatory drugs: an animal study comparing ibuprofen and indomethacin. *Fertil Steril* 1984;41:639–642.
- 34 de la Portilla F, Ynfante I, Bejarano D, et al: Prevention of peritoneal adhesions by intraperitoneal administration of vitamin E: an experimental study in rats. *Dis Colon Rectum* 2005;47:2157–2161.
- 35 Lucas P, Warejcka D, Young H, et al: Formation of abdominal adhesions is inhibited by antibodies to transforming growth factor-beta1. *J Surg Res* 1996;65:135–138.
- 36 Avsar F, Sahin M, Aksoy F, et al: Effects of diphenhydramine HCl and methylprednisolone in the prevention of abdominal adhesions. *Am J Surg* 2001;181:512–515.
- 37 Replogle RL, Johnson R, Gross RE: Prevention of postoperative intestinal adhesions with combined promethazine and dexamethasone therapy: experimental and clinical studies. *Ann Surg* 1966;163:580–588.
- 38 Chowdhury SM, Hubbell JA: Adhesion prevention with anicrod released via a tissue-adherent hydrogel. *J Surg Res* 1996;61:58–64.
- 39 Hill-West JL, Dunn RC, Hubbell JA: Local release of fibrinolytic agents for adhesion prevention. *J Surg Res* 1995;59:759–763.
- 40 Liu Y, Li H, Shu XZ, SD, et al: Crosslinked hyaluronan hydrogels containing mitomycin C reduce postoperative abdominal adhesions. *Fertil Steril* 2005;83(suppl 1):1275–1283.
- 41 Cubukcu A, Alponat A, Gonullu N: An experimental study evaluating the effect of Mitomycin C on the prevention of postoperative intraabdominal adhesions. *J Surg Res* 2001;96:163–166.
- 42 Berkkanoglu M, Zhang L, Ulukus M, et al: Inhibition of chemokines prevents intraperitoneal adhesions in mice. *Hum Reprod* 2005;20:3047–3052.
- 43 Hoshino A, Kawamura YI, Yasuhara M, et al: Inhibition of CCL1-CCR8 interaction prevents aggregation of macrophages and development of peritoneal adhesions. *J Immunol* 2007;178:5296–5304.
- 44 Ansaloni L, Catena F, Coccolini F, et al: Peritoneal adhesions to prosthetic materials: An experimental comparative study of treated and untreated polypropylene meshes placed in the abdominal cavity. *J Laparoendosc Adv Surg Tech A* 2009;19:369–374.
- 45 Yeo Y, Ito T, Bellas E, et al: In situ cross-linkable hyaluronan hydrogels containing polymeric nanoparticles for preventing postsurgical adhesions. *Ann Surg* 2007;245:819–824.
- 46 Oncel M, Remzi FH, Senagore AJ, et al: Liquid antiadhesive product (Adcon-p) prevents post-operative adhesions within the intra-abdominal organs in a rat model. *Int J Colorectal Dis* 2003;18:514–517.
- 47 Oncel M, Remzi FH, Senagore AJ, et al: Application of Adcon-P or Seprafilm in consecutive laparotomies using a murine model. *Am J Surg* 2004;187:304–308.
- 48 Leach RE, Henry RL: Reduction of postoperative adhesions in the rat uterine horn model with poloxamer 407. *Am J Obstet Gynecol* 1990;162:1317–1319.
- 49 Steinleitner A, Lambert H, Kazensky C, et al: Poloxamer 407 as an intraperitoneal barrier material for the prevention of postsurgical adhesion formation and reformation in rodent models for reproductive surgery. *Obstet Gynecol* 1991;77:48–52.
- 50 Hill-West JL, Chowdhury SM, Sawhney AS, et al: Prevention of postoperative adhesions in the rat by in situ photopolymerization of bioresorbable hydrogel barriers. *Obstet Gynecol* 1994;83:59–64.
- 51 Zhou J, Elson C, Lee TD: Reduction in postoperative adhesion formation and re-formation after an abdominal operation with the use of N,O-carboxymethyl chitosan. *Surgery* 2004;135:307–312.
- 52 Kennedy R, Costain DJ, McAlister VC, et al: Prevention of experimental postoperative peritoneal adhesions by N,O-carboxymethyl chitosan. *Surgery* 1996;120:866–870.
- 53 Avital S, Bollinger TJ, Wilkinson JD, et al: Preventing intra-abdominal adhesions with polylactic acid film: an animal study. *Dis Colon Rectum* 2005;48:153–157.
- 54 Chen MD, Teigen GA, Reynolds HT, et al: Laparoscopy versus laparotomy: an evaluation of adhesion formation after pelvic and paraaortic lymphadenectomy in a porcine model. *Am J Obstet Gynecol* 1998;178:499–503.
- 55 Moore RG, Partin AW, Adams JB, et al: Adhesion formation after transperitoneal nephrectomy: laparoscopic vs open approach. *J Endourol* 1995;9:277–280.
- 56 Jacob BP, Hogle NJ, Durak E, et al: Tissue ingrowth and bowel adhesion formation in an animal comparative study: polypropylene versus Proceed versus Parietex Composite. *Surg Endosc* 2007;21:629–633.
- 57 Krähenbühl L, Schäfer M, Kuzinkovas V, et al: Experimental study of adhesion formation in open and laparoscopic fundoplication. *Br J Surg* 1998;85:826–830.
- 58 Zhang ZL, Xu SW, Zhou XL: Preventive effects of chitosan on peritoneal adhesion in rats. *World J Gastroenterol* 2006;12:4572–4577.
- 59 Bellón JM, Rodríguez M, García-Hondurilla N, et al: Peritoneal effects of prosthetic meshes used to repair abdominal wall defects: monitoring adhesions by sequential laparoscopy. *J Laparoendosc Adv Surg Tech A* 2007;17:160–166.
- 60 Adhesion Scoring Group. Improvement of interobserver reproducibility of adhesion scoring system. *Fertil Steril* 1994;62:984–988.
- 61 American Fertility Society. The AFS classification of adnexal adhesions, distal tubal occlusion, tubal occlusion secondary to tubal ligation, tubal pregnancies, Müllerian anomalies and intrauterine adhesions. *Fertil Steril* 1988;49:944–955.
- 62 Corson SL, Batzer FR, Gocial B, et al: Intra-observer and inter-observer variability in scoring laparoscopic diagnosis of pelvic adhesions. *Hum Reprod* 1995;10:161–164.
- 63 Johns DA, Ferland R, Dunn R: Initial feasibility study of a sprayable hydrogel adhesion barrier system in patients undergoing laparoscopic ovarian surgery. *J Am Assoc Gynecol Laparosc* 2003;10:334–338.

- 64 Diamond MP, Luciano A, Johns DA, et al: Reduction of postoperative adhesions by N,O-carboxymethylchitosan: a pilot study. *Fertil Steril* 2003;80:631–636.
- 65 Hulka JF: Adnexal adhesions, a prognostic staging and classification system based on a 5-year survey of fertility surgery results at Chapel Hill, North Carolina. *Am J Obstet and Gynecol* 1982;144:141–148.
- 66 Linsky CB, Diamond MP, Cunningham T, et al: Adhesion reduction in the rabbit uterine horn model using an absorbable barrier, TC-7. *J Reprod Med* 1987;32:17–20.
- 67 Leach RE, Burns JW, Dawe EJ, et al: Reduction of postsurgical adhesion formation in the rabbit uterine horn model with use of hyaluronate/carboxymethylcellulose gel. *Fertil Steril* 1998;69:415–418.
- 68 Wilson MS: Practicalities and costs of adhesions. *Colorectal Dis* 2007;9(suppl 2):60–65.
- 69 Tjandra JJ, Chan MK: A sprayable hydrogel adhesion barrier facilitates closure of de-functioning loop ileostomy: a randomized trial. *Dis Colon Rectum* 2008;51:956–960.
- 70 Takeuchi H, Kitade M, Kikuchi I, et al: Adhesion prevention effects of fibrin sealants after laparoscopic myomectomy as determined by second look laparoscopy: a prospective randomized, controlled study. *J Reprod Med* 2005;50:571–577.
- 71 Hellebrekers BW, Trimbos-Kemper TC, Boesten L, et al: Preoperative predictors of postsurgical adhesion formation and the Prevention of Adhesions with Plasminogen Activator (PAPA-study): results of a clinical pilot study. *Fertil Steril* 2009;91:1204–1214.
- 72 Mercorio F, Mercorio A, Di Spizio Sardo A, et al: Evaluation of ovarian adhesion formation after laparoscopic ovarian drilling by second-look minilaparoscopy. *Fertil Steril* 2008;89:1229–1233.
- 73 Montgomery TC, Sharp JB, Bellina JH, et al: Comparative gross and histological study of the effects of scalpel, electric knife, and carbon dioxide laser on skin and uterine incisions in dogs. *Lasers Surg Med* 1983;3:9–22.
- 74 Bellina JH, Hemmings R, Voros JJ, et al: Carbon dioxide laser and electrosurgical wound study with an animal model: a comparison of tissue damage and healing patterns in peritoneal tissue. *Am J Obstet Gynecol* 1984;148:327–334.
- 75 Filmar S, Jetha N, McComb P, et al: A comparative histologic study on the healing process after tissue transection. I. Carbon dioxide laser and electromicrosurgery. *Am J Obstet Gynecol* 1989;160:1062–1067.
- 76 Filmar S, Jetha N, McComb P, et al: A comparative histologic study on the healing process after tissue transection. II. Carbon dioxide laser and surgical microscissors. *Am J Obstet Gynecol* 1989;160:1068–1072.
- 77 Forestier D, Slim K, Joubert-Zakey J, et al: Do bipolar scissors increase postoperative adhesions? An experimental double-blind randomized trial. *Ann Chir* 2002;127:680–684.
- 78 Weaver FA, Lew W, Granke K, et al: A comparison of recombinant thrombin to bovine thrombin as a hemostatic ancillary in patients undergoing peripheral arterial bypass and arteriovenous graft procedures. *J Vasc Surg* 2008;47:1266–1273.
- 79 Chapman WC, Lockstadt H, Singla N, et al: Phase 2, randomized, double-blind, placebo-controlled, multicenter clinical evaluation of recombinant human thrombin in multiple surgical indications. *J Thromb Haemost* 2006;4:2083–2085.
- 80 Doria C, Fischer CP, Wood CG, et al: Phase 3, randomized, double-blind study of plasma-derived human thrombin versus bovine thrombin in achieving hemostasis in patients undergoing surgery. *Curr Med Res Opin* 2008;24:785–794.
- 81 Levy O, Martinowitz U, Oran A, et al: The use of fibrin tissue adhesive to reduce blood loss and the need for blood transfusion after total knee arthroplasty. A prospective, randomized, multicenter study. *J Bone Joint Surg Am* 1999;81:1580–1588.
- 82 Wang GJ, Goldthwaite CA Jr, Burks S, et al: Fibrin sealant reduces perioperative blood loss in total hip replacement. *J Long Term Eff Med Implants* 2003;13:399–411.
- 83 Milne AA, Murphy WG, Reading SJ, et al: Fibrin sealant reduces suture line bleeding during carotid endarterectomy: a randomised trial. *Eur J Vasc Endovasc Surg* 1995;10:91–94.
- 84 Schäfer M, Krähenb L, Büchler MW: Comparison of adhesion formation in open and laparoscopic surgery. *Dig Surg* 1998;15:148–152.
- 85 Garrard CL, Clements RH, Nanney L, et al: Adhesion formation is reduced after laparoscopic surgery. *Surg Endosc* 1999;13:10–13.
- 86 Molinas CR, Koninckx PR: Hypoxaemia induced by CO₂ or helium pneumoperitoneum is a co-factor in adhesion formation in rabbits. *Hum Reprod* 2000;15:1758–1763.
- 87 Kopernik G, Avinoach E, Grossman Y, et al: The effect of a high partial pressure of carbon dioxide environment on metabolism and immune functions of human peritoneal cells—relevance to carbon dioxide pneumoperitoneum. *Am J Obstet Gynecol* 1998;179:1503–1510.
- 88 Sajid MS, Mallick AS, Rimpel J, et al: Effect of heated and humidified carbon dioxide on patients after laparoscopic procedures: a meta-analysis. *Surg Laparosc Endosc Percutan Tech* 2008;18:539–546.
- 89 Savel RH, Balasubramanya S, Lasheen S, et al: Beneficial effects of humidified, warmed carbon dioxide insufflation during laparoscopic bariatric surgery: a randomized clinical trial. *Obes Surg* 2005;15:64–69.
- 90 Farley DR, Greenlee SM, Larson DR, et al: Double-blind, prospective, randomized study of warmed, humidified carbon dioxide insufflation vs standard carbon dioxide for patients undergoing laparoscopic cholecystectomy. *Arch Surg* 2004;139:739–743; discussion 743–744.
- 91 Flora ED, Wilson TG, Martin IJ, O'Rourke NA, Maddern GJ: A review of natural orifice transluminal endoscopic surgery (NOTES) for intra-abdominal surgery: experimental models, techniques, and applicability to the clinical setting. *Ann Surg* 2008;247:583–602.
- 92 Maddern GJ: NOTES: progress or marketing? *ANZ J Surg* 2009;79:318–319.
- 93 Kucukozkan T, Ersoy B, Uygur D, et al: Prevention of adhesions by sodium chromoglycate, dexamethasone, saline and aprotinin after pelvic surgery. *ANZ J Surg* 2004;74:1111–1115.
- 94 Nishimura K, Nakamura R, diZerega G: Ibuprofen inhibition of postsurgical adhesion formation: a time and dose response biochemical evaluation in rabbits. *J Surg Res* 1984;36:115–124.
- 95 Holtz G: Failure of a nonsteroidal anti-inflammatory agent (ibuprofen) to inhibit peritoneal adhesion reformation after lysis. *Fertil Steril* 1982;37:582–583.
- 96 Binda MM, Molinas CR, Koninckx PR: Reactive oxygen species and adhesion formation: clinical implications in adhesion prevention. *Hum Reprod* 2003;18:2503–2507.
- 97 Portz D, Elkins T, White R, et al: Oxygen free radicals and pelvic adhesions formation: I. Blocking oxygen free radical toxicity to prevent adhesion formation in an endometriosis model. *Int J Fertil* 1991;36:39–42.
- 98 Reid RL, Lie K, Spence JE, et al: Clinical evaluation of the efficacy of heparin-saturated interceed for prevention of adhesion reformation in the pelvic sidewall of the human; in Diamond MP, diZerega GS, Linsky CB, et al (eds): *Gynecologic Surgery and Adhesion Prevention*. New York, Wiley, 1993, pp 261–264.
- 99 Dunn RC, Mohler M: Effect of varying days of tissue plasminogen activator therapy on the prevention of postsurgical adhesions in a rabbit model. *J Surg Res* 1993;54:242–245.
- 100 Orita H, Fukasawa M, Girgis W, diZerega GS: Inhibition of postsurgical adhesions in a standardized rabbit model: intraperitoneal treatment with tissue plasminogen activator. *Int J Fertil* 1991;36:172–177.
- 101 Menzies D, Ellis H: The role of plasminogen activator in adhesion prevention. *Surg Gynecol Obstet* 1991;172:362–366.
- 102 Dunn RC, Mohler M: Effect of varying days of tissue plasminogen activator therapy on the prevention of post-surgical adhesions in a rabbit model. *J Surg Res* 1993;54:242–245.
- 103 Holmdahl LE, Al-Jabreen M, Risberg B: Role of fibrinolysis in the formation of postoperative adhesions. *Wound Repair Regen* 1994;2:171–176.
- 104 Flessner MF: The transport barrier in intraperitoneal therapy. *Am J Physiol Renal Physiol* 2005;288:F433–F442.

- 105 Mahdy T, Mohamed G, Elhawary A: Effect of methylene blue on intra-abdominal adhesion formation in rats. *Int J Surg* 2008;6: 452–455.
- 106 Chalela JA, Katzan I, Liebeskind DS, et al: Safety of intra-arterial thrombolysis in the postoperative period. *Stroke* 2001;32:1365–1369.
- 107 Wiseman DM, Trout JR, Diamond MP: The rates of adhesion development and the effects of crystalloid solutions on adhesion development in pelvic surgery. *Fertil Steril* 1998;70:702–711.
- 108 Menzies D, Pascual MH, Walz MK, et al: Use of icodextrin 4% solution in the prevention of adhesion formation following general surgery: from the multicentre ARIEL Registry. *Ann R Coll Surg Engl* 2006;88: 375–382.
- 109 diZerega GS, Verco SJ, Young P, et al: A randomized, controlled pilot study of the safety and efficacy of 4% icodextrin solution in the reduction of adhesions following laparoscopic gynaecological surgery. *Hum Reprod* 2002;17:1031–1038.
- 110 Ong SY, Wu J, Mochhala SM, et al: Development of a chitosan-based wound dressing with improved hemostatic and antimicrobial properties. *Biomaterials* 2008;29: 4323–4332.
- 111 Gustafson SB, Fulkerson P, Bildfell R, et al: Chitosan dressing provides hemostasis in swine femoral arterial injury model. *Prehosp Emerg Care* 2007;11:172–178.
- 112 Rabea EI, Badawy ME, Stevens CV, et al: Chitosan as antimicrobial agent: applications and mode of action. *Biomacromolecules* 2003;4:1457–1465.
- 113 Haney A, Hesla J, Hurst B, et al: Expanded polytetrafluoroethylene (Gore-Tex Surgical Membrane) is superior to oxidized regenerated cellulose (Interceed TC7+) in preventing adhesions. *Fertil Steril* 1995;63: 1021–1026.
- 114 Haney A, Hesla J, Hurst B, et al: Prevention of pelvic side wall adhesion reformation using surgical barriers: expanded polytetrafluoroethylene (Gore-Tex Surgical Membrane) is superior to oxidized regenerated cellulose (Interceed TC7+) (abstract). *Fertil Steril* 1994;S210:265.
- 115 Korell M: Reduction of adhesion by INTERCEED Barrier and Gortex Surgical Membrane after laparoscopic myomectomy; presented at 'Möglichkeiten der Adhäsionsprophylaxe', Munich, April 1994.
- 116 Schreinemacher MH, Emans PJ, Gijbels MJ, et al: Degradation of mesh coatings and intraperitoneal adhesion formation in an experimental model. *Br J Surg* 2009;96:305–313.
- 117 Diamond MP: Reduction of adhesions after uterine myomectomy by Seprafilm membrane (HAL-F): a blinded, prospective, randomized, multicenter clinical study. *Seprafilm Adhesion Study Group. Fertil Steril* 1996;66:904–910.
- 118 Beck DE, Cohen Z, Fleshman JW, et al: A prospective, randomised, multicenter, controlled study of the safety of seprafilm adhesion barrier in abdominopelvic surgery of the intestine. *Dis Colon Rectum* 2003;46: 1310–1319.
- 119 Becker JM, Dayton MT, Fazio VW, et al: Prevention of postoperative abdominal adhesions by a sodium hyaluronate based bioresorbable membrane: a prospective, randomised, double-blind multicenter study. *J Am Coll Surg* 1996;183:297–306.
- 120 Cohen Z, Senagore A, Dayton M, et al: Prevention of postoperative abdominal adhesions by a novel, glycerol/sodium hyaluronate/carboxymethylcellulose-based bioresorbable membrane: a prospective, randomised, evaluator-blinded multicenter study. *Dis Colon Rectum* 2005;48: 1130–1139.
- 121 Fazio VW, Cohen Z, Fleshman JW, et al: Reduction in adhesive small-bowel obstruction by seprafilm adhesion barrier after intestinal resection. *Dis Colon Rectum* 2006;49:1–11.
- 122 Kusunoki M, Ikeuchi H, Yanagi H, et al: Bioresorbable hyaluronate-carboxymethylcellulose membrane (Seprafilm) in surgery for rectal carcinoma: a prospective randomised clinical trial. *Surg Today* 2005; 35:940–945.
- 123 Tang CL, Seow-Choen F, Fook-Chong S, et al: Bioresorbable adhesion barrier facilitates early closure of the defunctioning ileostomy after rectal excision: a prospective, randomised trial. *Dis Colon Rectum* 2003; 46:1200–1207.
- 124 Tang CL, Jayne DG, Seow-Choen F, et al: A randomised controlled trial of 0.5% ferric hyaluronate gel (Intergel) in the prevention of adhesions following abdominal surgery. *Ann Surg* 2006;243:449–455.
- 125 Vrijland WW, Tseng LNL, Eijkman HJM, et al: Fewer intraperitoneal adhesions with use of hyaluronic acid-carboxymethylcellulose membrane: a randomised clinical trial. *Ann Surg* 2002;235:193–199.
- 126 Kumar S, Wong PF, Leaper DJ: Intra-peritoneal prophylactic agents for preventing adhesions and adhesive intestinal obstruction after non-gynaecological abdominal surgery. *Cochrane Database Syst Rev* 2009;CD005080.
- 127 Perlitz Y, Oettinger M, Karam K, et al: Anaphylactic shock during hysteroscopy using Hyskon solution: case report and review of adverse reactions and their treatment. *Gynecol Obstet Invest* 1996;41:67–69.
- 128 Johns DB, Keyport GM, Hoehler F, et al: Reduction of postsurgical adhesions with Intergel adhesion prevention solution: a multicenter study of safety and efficacy after conservative gynecologic surgery. *Fertil Steril* 2001;76:595–604.
- 129 DeCherney AH, diZerega GS: Clinical problem of intraperitoneal postsurgical adhesion formation following general surgery and the use of adhesion prevention barriers. *Surg Clin North Am* 1997;77:671–688.
- 130 Ellis H: Medicolegal consequences of adhesions. *Hosp Med* 2004;65:348–350.
- 131 Gauwerky JF, Heinrich D, Kubli F: Complications of intraperitoneal dextran application for prevention of adhesions. *Biol Res Pregnancy Perinatol* 1986;7:93–97.
- 132 Larsson B, Lalos O, Marsk L, et al: Effect of intraperitoneal instillation of 32% dextran 70 on postoperative adhesion formation after tubal surgery. *Acta Obstet Gynecol Scand* 1985;64:437–441.
- 133 Sites CK, Jensen BA, Glock JL, et al: Transvaginal ultrasonographic assessment of transvaginal ultrasonographic assessment of Hyskon or lactated Ringer's solution instillation after laparoscopy: randomized, controlled study. *J Ultrasound Med* 1997; 16:195–199.
- 134 Haney AF, Doty E: A barrier composed of chemically cross-linked hyaluronic acid (Incert) reduces postoperative adhesion formation. *Fertil Steril* 1998;70:145–151.
- 135 Nordic Adhesion Prevention Study Group. The efficacy of Interceed (TC7)* for prevention of reformation of postoperative adhesions on ovaries, fallopian tubes, and fimbriae in microsurgical operations for fertility: a multicenter study. *Fertil Steril* 1995;63:709–714.
- 136 Adhesion Barrier Study Group. Prevention of postsurgical adhesions by INTERCEED (TC7), an absorbable adhesion barrier: a prospective randomized multicenter clinical study. *Fertil Steril* 1989;51:933–938.
- 137 Wiseman DM: Possible intergel reaction syndrome (pIRS). *Ann Surg* 2006;244:630–632.
- 138 Okuyama N, Rodgers KE, Wang CY, et al: Prevention of retrosternal adhesion formation in a rabbit model using bioresorbable films of polyethylene glycol and polylactic acid. *J Surg Res* 1998;78:118–122.
- 139 Rodgers K, Cohn D, Hotovely A, et al: Evaluation of polyethylene glycol/polylactic acid films in the prevention of adhesions in the rabbit adhesion formation and reformation sidewall models. *Fertil Steril* 1998;69: 403–408.
- 140 Diamond MP: Reduction of de novo postsurgical adhesions by intraoperative pre-coating with Sepracoat (HAL-C) solution: a prospective, randomized, blinded, placebo-controlled multicenter study. The Sepracoat Adhesion Study Group. *Fertil Steril* 1998;69:1067–1074.