The Autoinducer-2 Exporter Superfamily

Rachael E. Rettner Milton H. Saier, Jr.

Division of Biological Sciences, University of California at San Diego, La Jolla, Calif., USA

Key Words

Transport • Metabolite exporters • Autoinducer •
Phylogeny • Topology • Evolution

Abstract

The TqsA (YdgG) protein of *Escherichia coli* has been shown to export the autoinducer-2 (AI-2) molecule, a furanosyl borate diester that bears little resemblance to previously characterized biological molecules. TqsA belongs to a large superfamily, the AI-2 exporter (AI-2E) superfamily, of putative transporters with no other functionally characterized members. These proteins derive exclusively from bacteria. Many different bacterial kingdoms contain them, although several kingdoms do not. These proteins exhibit a uniform topology with 8 putative transmembrane segments (TMSs) which we show probably arose from a 4-TMS precursor in a process that involved at least one and possibly two intragenic duplication event(s). The first halves of these proteins are more diverse in sequence than the second halves, suggesting that the first halves may serve substrate-specific functions while the second halves serve family-specific functions. Conserved residues and motifs in these proteins are identified. Some homologues include extra catalytic domains including those involved in purine nucleotide biosynthesis, ATP and GTP binding, and molecular signaling. The results presented provide guides for future functional studies on members of this superfamily of bacterial transporters.

Copyright © 2010 S. Karger AG, Basel

Bacteria use a variety of means to communicate with one another and with their eukaryotic hosts [1–6]. In some cases, social interactions in complex bacterial colonies allow bacteria to synchronize the behavior of all members of the group and thereby act like multicellular organisms [7]. By contrast, some bacterial social engagements promote individuality among members within the group and thereby foster diversity [8, 9]. Bacterial communication systems include long- and short-range chemical signaling [10]; one-way, two-way, and multi-way communication [11]; contact-mediated and contact-inhibited signaling [12, 13], and the use and spread of misinformation, even deadly information [8, 14].

Bacteria use a diversity of small molecules for extra- and intracellular signaling [8]. They scan small-molecule mixtures to access information about both their extracellular environment and their intracellular physiological status. Based on the integrated information, they continuously interpret their circumstances and react rapidly to changes [15]. They must integrate extra- and intracellular signaling information to mount appropriate responses to environmental changes.

Several small-molecule bacterial signaling pathways have been identified. These include extracellular ‘quorum-sensing’ signaling and intracellular cyclic dinucleotide signaling. Possibly, these two pathways converge to control complex processes including multicellularity, biofilm formation, and virulence [15].

The exchange of extracellular signaling molecules must be mediated by specific transporters, some involved

in molecular uptake, others catalyzing efflux [16–18]. Unlike other autoinducers, which are specific to a particular species of bacteria, autoinducer-2 (AI-2) is produced by a large number of bacterial species. AI-2 has been proposed to serve as a ‘universal’ signal for inter-species communication [8]. The crystal structure of an AI-2 sensor protein, LuxP, in complexation with the autoinducer, revealed the bound ligand to be a furanosyl borate diester that bears no resemblance to previously characterized autoinducers [19].

Recently, the *Escherichia coli* YdgG (TqsA) protein was shown to control biofilm formation in *E. coli* K-12 by mediating AI-2 transport. YdgG had been known to be induced in *E. coli* biofilms. Deletion of *ydgG* decreased extracellular and increased intracellular concentrations of AI-2, suggesting that YdgG enhances export of AI-2 [20]. Consistent with this hypothesis, deletion of the *ydgG*

gene increased cell motility by increasing transcription of flagellar genes that were known to be induced by AI-2. By expressing *ydgG* in *trans*, the wild-type phenotypes for extracellular AI-2 activity, motility, and biofilm formation were restored.

YdgG is a membrane-spanning protein that is conserved in many bacteria. It influences resistance to several antimicrobials, including crystal violet and streptomycin [20]. Deletion of *ydgG* caused 31% of the bacterial chromosome to be differentially expressed in biofilms. This fact apparently resulted because AI-2 controls the transcription of hundreds of genes. Since YdgG catalyzes export of the quorum-sensing signal, AI-2, the gene name *tqsA* (transporter of quorum sensing-A) was suggested [20].

In this communication, we analyze the family of putative transporters to which TqsA belongs. We name this

Fig. 1. Phylogenetic trees for the 5 families of proteins in the AI-2E superfamily (a–e) as well as for the complete superfamily (f). **a–e** Families 1–5, respectively. **f** The full superfamily tree showing the positioning of families 1–5 relative to each other. The dendrogram corresponding to the tree in **f** is presented in figure S2. It is based on the multiple alignment shown in figure S1. Programs used to derive the trees have been described (CLUSTAL X [24] and TreeView [25]). Protein abbreviations for all members of the AI-2E superfamily can be found in table S1 (alphabetically arranged) and tables S3–S7 for the 5 families (1–5, respectively). The standard abbreviations used in our laboratory and many others include three letters often followed by a number. The first capitalized letter refers to the genus name while the second two small letters are the first two letters of the species name. Thus, Eco2 refers to the second paralogue in *E. coli*. (For figure 1 **b–f** see next pages.)

family the AI-2 exporter (AI-2E) superfamily, based on the function of its first characterized member. We show that it is a huge superfamily (Pfam family PF01594) with members derived from a wide range of bacteria. However, no member of the family was identified from an archaeon, a eukaryote, or a member of certain bacterial kingdoms that include sequenced organisms with reduced genome sizes. Phylogenetic analyses revealed the relationships of these homologues to each other with groupings largely according to organismal type. However, this occurred in an unusual fashion suggestive of several functional types with occasional lateral gene transfer responsible for some of the anomalies. Thus, the analyses reveal groups of apparent orthologues that presumably serve the same function. Conserved motifs are identified which must play specific functional/structural roles. An evolutionary pathway for the appearance of these proteins, based on intragenic duplication of an ancestral 4-transmembrane segment (TMS) unit is established. Finally,

extra domains with recognized functions in some of these proteins are identified, and these provide clues as to the associated transport functions.

Methods and Results

Initial Phylogenetic Analyses of the AI-2E Superfamily

The PerM protein of *E. coli* (TC #2.A.86.1.1) was used as the query sequence in PSI-BLAST searches (default settings) with six iterations [21]. A modified CD hit program [22] was used to eliminate redundancies and closely similar sequences of greater than 90% identity (the default setting). This program randomly selects one protein of several that are of greater than 90% identity with each other. Membership to the AI-2E superfamily was assigned based on sequence similarity throughout the transmembrane regions. All members of the superfamily thus exhibited at least 9 SD (probability that the observed degree of similarity arose by chance is less than 10^{-19} [23]). 391 non-redundant homologues from the NCBI protein databank remained and served as our data set for all analyses reported here (see table S1 on our website; <http://www.biology.ucsd.edu/~msaier/supmat/AI-2>). A neighbor-join-

ing phylogenetic tree (fig. 1f), based on a CLUSTAL X multiple alignment [24] (see fig. S1 on our website), and a dendrogram (fig. S2), drawn using the TreeView program [25], were generated.

The proteins generally fell into 5 distinct clusters or families of fairly similar size (fig. 1f). Cluster 1 (family 1) contains 78 proteins; family 2, 58 proteins; family 3, 89 proteins; family 4, 103 proteins, and family 5, 63 proteins. The multiple alignment of all 391 proteins (fig. S1) revealed two fully conserved prolyl residues, one at alignment position 532, and the other at alignment position 627 within hydrophobic peaks 6 and 8, respectively (see below). Several gaps were observed within the hydrophilic regions of these aligned sequences (positions 249–263, 336–384, 404–416, 441–446, 463–468, 508–515, 523–526, and 559–569 of the 1,112 residue position alignment presented in fig. S1).

Topological and Sequence Conservation Analyses

Each of the 5 families within the AI-2E superfamily was analyzed separately for sequence conservation, topology, and phylogeny. The proteins, their organismal sources, abbreviations, sizes, numbers of putative TMSs and Genbank (gi) identification numbers for the 5 clusters are provided in tables S3–S7. The 5 neighbor-joining phylogenetic trees are shown in figure 1a–e, while the average hydropathy, amphipathicity and similarity plots, based on the AveHAS program [26] are shown in figure 2a–e. The AveHAS program averages the hydropathy plots for all aligned sequences, providing much greater accuracy than would otherwise be possible. These figures provide the bases for several of the conclusions cited below. They were derived using the multiple alignments presented in figure S3–S7 on our website. Hydropathy and amphipathicity plots for individual proteins were generated using the WHAT program [26].

Family 1 proteins show eight peaks of hydrophobicity (fig. 2a) and several fully conserved residues (fig. S3). These include seven prolyl residues (alignment positions 104, peak 3; 195, end of peak 4; 217, between peaks 4 and 5; 279, peak 6; 326 and 337, beginning of peak 8; and 361, end of peak 8). The second halves of these proteins proved to be better conserved than the first halves.

We derived consensus sequences for the two best conserved regions of these proteins for all 5 families. In peak 6, the consensus sequence was:

```
* : : : : : *
GLSVLPY (LIV) GA (LIV)3 TVP
```

while in peak 8, the consensus sequence was:

```
* : : : : : * * : : : :
P (LIV) FSEAVNLHP (LIV)4 (SA) (LIV)3 FGGLWGFVGFVFAIP
```

where asterisks indicate identities only, colons close similarities, and dots more distant similarities as defined by the CLUSTAL X program; alternative consensus residues at any one position are indicated in parentheses.

Figure 3 shows how these two consensus sequences differ between the proteins of the 5 phylogenetic families. Of these 5 families, family 1 appears to be the best conserved with respect to both identities and conservative substitutions. The two sets of consensus sequences show similarities in all 5 families. For example, the P at position 7 in motif 1 and the P in the terminal position of motif 2 are fully conserved in all proteins of the AI-2E superfamily as noted above, suggesting an essential structural or functional role. The G at position 10 in the first consensus se-

quence and those at positions 21 and 25 in the second consensus sequence are conserved in nearly all families. While the dominant residue may differ in the different families at specific positions, these are not fully conserved in any of these families. Thus, when the consensus residues differ, they are never fully conserved. We conclude that in none of these 5 families is a distinctive function, common to all members in that family, but different from those of the other four families, dictated by a specific residue or set of residues within these two conserved motifs. It appears that these motifs provide structural features or a function that is common to all members of the AI-2E superfamily.

The average hydropathy plots shown in figure 2a–e reveal the characteristics of each of the 5 families in the AI-2E superfamily. Family 1 is most compact, with no protein showing N- or C-terminal extensions. In this plot, peaks 1 and 2 are close to each other, peak 3 is more distant, and peak 4 is still more distant. Peaks 5–7 are all overlapping while peak 8 is distinct but close to peak 7. This pattern is observed for all 5 families (compare figure 2a–e).

The average amphipathicity plots revealed that all 5 families show similar characteristics: amphipathic peaks occur following hydrophobic peaks 3, 4 and 8 and coinciding with peak 6. Motif 1, also coinciding with peak 6, reveals a general pattern of a semipolar residue followed by three hydrophobic residues. A helical wheel depiction showed that strongly hydrophobic residues occur on one side of helix 6, while semipolar residues predominate on the other side (data not shown), thus accounting for the amphipathicity of putative TMS 6.

As noted above, families 2–5 show poorer conservation than observed for family 1 (figure 2a–e and 3). Further, families 2–4 contain proteins with short N-terminal and long C-terminal hydrophilic sequences, while family 5 contains proteins with both N- and C-terminal extensions maximally of about 160 residues each. The domains in these longer proteins will be analyzed and described below.

Phylogenetic Analyses of the 5 Families of the AI-2E Superfamily

The phylogenetic trees for the 5 AI-2E families are presented in figure 1a–e. Family 1 proteins derive exclusively from the proteobacterial phylum of the γ - and δ -orders. Family 2 is largely from α -, β -, γ - and δ -proteobacteria, but three proteins are from a single species each of the *Bacteroides*, Firmicute and *Planctomycetes* phyla, respectively. Family 3 includes proteins from the same proteobacteria as for family 2, but *Chlorobi* and *Chloroflexi* proteins are also present. Family 4 proteins are exclusively from firmicutes except for one actinobacterial protein. Finally, family 5 proteins are mostly from actinobacteria and cyanobacteria with a few proteins from *Deinococcus*, *Chloroflexi* and ϵ -proteobacterial species. It is therefore clear that phylogenetic clustering is predominantly according to organismal type with a few interesting exceptions. Also of interest is the absence of homologues in specific bacterial kingdoms such as the spirochetes, chlamydia, and mycoplasma. Most of the sequenced genomes of bacteria in these kingdoms are of reduced size due to their obligate parasitic lifestyles. We detected no homologues outside of the bacterial domain.

Repeat Elements in AI-2E Superfamily Proteins

The GAP and IC programs, with default settings and 500 random shuffles (to establish statistical significance) [27, 28] were

Fig. 2. Relative average hydropathy (solid dark lines, top), average amphipathicity (dotted gray lines, top) and average similarity (solid but thin lines, bottom) plots for the 5 families of proteins in the AI-2E superfamily. **a–e** Families 1–5, respectively. The Ave-HAS program was used to derive the plots using all members of each of the 5 families included in this study [47]. Peaks of hydro-

phobicity that are believed to correspond to the 8 putative TMSs are numbered above the hydropathy plot. Assignments were confirmed using the WHAT [26] and TMHMM [48] programs for individual proteins. These plots are based on the multiple alignments shown in figure S3–S7.

The two fully conserved prolyl residues in all members of the AI-2E superfamily are worthy of note. Full conservation in a wide range of sequence-divergent homologues suggests one or more essential functional or structural roles. No molecular genetic analyses have been reported that provide a clue as to what this role or these roles may be. However, cis/trans-prolyl isomerization has been reported to control ion gating channel activity [38]. It is possible that one or both of the conserved prolyl residues in members of the AI-2E superfamily undergo cis/trans-induced conformational changes that accompany transport. Other possibilities, such as essential structural features, can also be considered.

With only one member of the AI-2E superfamily characterized, the functions of most members remain an open question. However, there are clues. The clustering of members of this superfamily into at least 5 families, each derived from a different but often overlapping set of bacterial kingdoms, leads to the suggestion that each family has a set of characteristic properties. Usually, members of a superfamily catalyze only uptake or efflux, but occasionally large superfamilies (e.g. MFS, ABC, DMT) can catalyze both. In almost all cases where both sym- and antiporters occur within the same superfamily, the symporters cluster separately from the antiporters on a phylogenetic tree [36, 39–46]. Thus, if all AI-2E superfamily members catalyze efflux, as we tentatively predict, they are likely to transport a variety of compounds, possibly all related in structure.

What might these substrates be? For clues, we examined members of the AI-2E superfamily that are fused to other domains. We found that various hydrophilic domains of recognizable function may be fused to them, either N-terminally or C-terminally. Based on homology, these domains appeared to be involved in purine nucleotide biosynthesis, purine nucleotide binding and/or hydrolysis, and molecular signaling. On this basis, as well as the known function of TqsA, we tentatively suggest that many or all members of the AI-2E superfamily catalyze the pmf-dependent efflux of both intracellular and extracellular signaling molecules. If so, they are generally involved in transmembrane signaling. Substrates may include cyclic nucleotides such as cyclic AMP, cyclic GMP and bis-3',5'-cyclic diguanosine monophosphate (cyclic di-GMP) [27, 32] as well as quorum sensors such as AI-2 [17]. Further studies will be required to establish the functional diversity of this large superfamily of secondary carriers.

Acknowledgments

This work was supported by NIH grant GM077402 from the National Institute of General Medical Sciences. We thank Mary Beth Hiller and Jeeni Criscenzo for assistance in the preparation of the manuscript.

References

- 1 Ansaldi M, Marolt D, Stebe T, Mandic-Mulec I, Dubnau D: Specific activation of the *Bacillus* quorum-sensing systems by isoprenylated pheromone variants. *Mol Microbiol* 2002;44:1561–1573.
- 2 Bassler BL: Small talk. Cell-to-cell communication in bacteria. *Cell* 2002;109:421–424.
- 3 Ben Jacob E, Becker I, Shapira Y, Levine H: Bacterial linguistic communication and social intelligence. *Trends Microbiol* 2004;12:366–372.
- 4 Ben-Jacob E, Shmueli H, Shochet O, Tenenbaum A: Adaptive self-organization during growth of bacterial colonies. *Physica A* 1992;187:378–424.
- 5 Fux CA, Costerton JW, Stewart PS, Stoodley P: Survival strategies of infectious biofilms. *Trends Microbiol* 2005;13:34–40.
- 6 Visick KL: Layers of signaling in a bacterium-host association. *J Bacteriol* 2005;187:3603–3606.
- 7 Taga ME, Bassler BL: Chemical communication among bacteria. *Proc Natl Acad Sci USA* 2003;100(suppl 2):14549–14554.
- 8 Bassler BL, Losick R: Bacterially speaking. *Cell* 2006;125:237–246.
- 9 Bhinu VS: Insight into biofilm-associated microbial life. *J Mol Microbiol Biotechnol* 2005;10:15–21.
- 10 Kolter R: Surfacing views of biofilm biology. *Trends Microbiol* 2005;13:1–2.
- 11 Parsek MR, Greenberg EP: Sociomicrobiology: the connections between quorum sensing and biofilms. *Trends Microbiol* 2005;13:27–33.
- 12 Kaiser D: Signaling in myxobacteria. *Annu Rev Microbiol* 2004;58:75–98.
- 13 Tiller JC, Liao CJ, Lewis K, Klibanov AM: Designing surfaces that kill bacteria on contact. *Proc Natl Acad Sci USA* 2001;98:5981–5985.
- 14 Lewis K: Programmed death in bacteria. *Microbiol Mol Biol Rev* 2000;64:503–514.
- 15 Camilli A, Bassler BL: Bacterial small-molecule signaling pathways. *Science* 2006;311:1113–1116.
- 16 Evans K, Passador L, Srikumar R, Tsang E, Nezezon J, Poole K: Influence of the MexAB-OprM multidrug efflux system on quorum sensing in *Pseudomonas aeruginosa*. *J Bacteriol* 1998;180:5443–5447.
- 17 Taga ME, Miller ST, Bassler BL: Lsr-mediated transport and processing of AI-2 in *Salmonella typhimurium*. *Mol Microbiol* 2003;50:1411–1427.
- 18 Walters M, Sircili MP, Sperandio V: AI-3 synthesis is not dependent on luxS in *Escherichia coli*. *J Bacteriol* 2006;188:5668–5681.
- 19 Chen X, Schauder S, Potier N, Van Dorsselaer A, Pelczar I, Bassler BL, Hughson FM: Structural identification of a bacterial quorum-sensing signal containing boron. *Nature* 2002;415:545–549.

- 20 Herzberg M, Kaye IK, Peti W, Wood TK: YdgG (TqsA) controls biofilm formation in *Escherichia coli* K-12 through autoinducer 2 transport. *J Bacteriol* 2006;188:587–598.
- 21 Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ: Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 1997;25:3389–3402.
- 22 Li W, Godzik A: Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics* 2006;22:1658–1659.
- 23 Dayhoff MO, Barker WC, Hunt LT: Establishing homologies in protein sequences. *Methods Enzymol* 1983;91:524–545.
- 24 Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG: The CLUSTAL X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* 1997;25:4876–4882.
- 25 Zhai Y, Tchieu J, Saier MH Jr: A web-based Tree View (TV) program for the visualization of phylogenetic trees. *J Mol Microbiol Biotechnol* 2002;4:69–70.
- 26 Zhai Y, Saier MH Jr: A web-based program (WHAT) for the simultaneous prediction of hydropathy, amphipathicity, secondary structure and transmembrane topology for a single protein sequence. *J Mol Microbiol Biotechnol* 2001;3:501–502.
- 27 Devereux J, Haeblerli P, Smithies O: A comprehensive set of sequence analysis programs for the VAX. *Nucleic Acids Res* 1984;12:387–395.
- 28 Zhai Y, Saier MH Jr: A simple sensitive program for detecting internal repeats in sets of multiply aligned homologous proteins. *J Mol Microbiol Biotechnol* 2002;4:375–377.
- 29 Galperin MY: Bacterial signal transduction network in a genomic perspective. *Environ Microbiol* 2004;6:552–567.
- 30 Ho YS, Burden LM, Hurley JH: Structure of the GAF domain, a ubiquitous signaling motif and a new class of cyclic GMP receptor. *EMBO J* 2000;19:5288–5299.
- 31 Saier MH Jr, Feucht BU, McCaman MT: Regulation of intracellular adenosine cyclic 3':5'-monophosphate levels in *Escherichia coli* and *Salmonella typhimurium*. Evidence for energy-dependent excretion of the cyclic nucleotide. *J Biol Chem* 1975;250:7593–7601.
- 32 D'Andrea LD, Regan L: TPR proteins: the versatile helix. *Trends Biochem Sci* 2003;28:655–662.
- 33 Galperin MY: Structural classification of bacterial response regulators: diversity of output domains and domain combinations. *J Bacteriol* 2006;188:4169–4182.
- 34 Saier MH Jr, Tran CV, Barabote RD: TCDB: the Transporter Classification Database for membrane transport protein analyses and information. *Nucleic Acids Res* 2006;34:D181–D186.
- 35 Saier MH Jr: Computer-aided analyses of transport protein sequences: gleaned evidence concerning function, structure, biogenesis, and evolution. *Microbiol Rev* 1994;58:71–93.
- 36 Busch W, Saier MH Jr: The transporter classification (TC) system, 2002. *Crit Rev Biochem Mol Biol* 2002;37:287–337.
- 37 Saier MH Jr: A functional-phylogenetic classification system for transmembrane solute transporters. *Microbiol Mol Biol Rev* 2000;64:354–411.
- 38 Ye S, Li Y, Chen L, Jiang Y: Crystal structures of a ligand-free MthK gating ring: insights into the ligand gating mechanism of K⁺ channels. *Cell* 2006;126:1161–1173.
- 39 Jack DL, Paulsen IT, Saier MH: The amino acid/polyamine/organocation (APC) superfamily of transporters specific for amino acids, polyamines and organocations. *Microbiology* 2000;146:1797–1814.
- 40 Jack DL, Yang NM, Saier MH Jr: The drug/metabolite transporter superfamily. *Eur J Biochem* 2001;268:3620–3639.
- 41 Kennerknecht N, Sahm H, Yen MR, Patek M, Saier MH Jr, Eggeling L: Export of L-isoleucine from *Corynebacterium glutamicum*: a two-gene-encoded member of a new translocator family. *J Bacteriol* 2002;184:3947–3956.
- 42 Pao SS, Paulsen IT, Saier MH Jr: Major facilitator superfamily. *Microbiol Mol Biol Rev* 1998;62:1–34.
- 43 Saier MH Jr, Beatty JT, Goffeau A, Harley KT, Heijne WH, Huang SC, Jack DL, Jahn PS, Lew K, Liu J, Pao SS, Paulsen IT, Tseng TT, Virk PS: The major facilitator superfamily. *J Mol Microbiol Biotechnol* 1999;1:257–279.
- 44 Saier MH Jr, Eng BH, Fard S, Garg J, Haggerty DA, Hutchinson WJ, Jack DL, Lai EC, Liu HJ, Nusinew DP, Omar AM, Pao SS, Paulsen IT, Quan JA, Sliwinski M, Tseng TT, Wachi S, Young GB: Phylogenetic characterization of novel transport protein families revealed by genome analyses. *Biochim Biophys Acta* 1999;1422:1–56.
- 45 Tseng TT, Gratwick KS, Kollman J, Park D, Nies DH, Goffeau A, Saier MH Jr: The RND permease superfamily: an ancient, ubiquitous and diverse family that includes human disease and development proteins. *J Mol Microbiol Biotechnol* 1999;1:107–125.
- 46 Vrljic M, Garg J, Bellmann A, Wachi S, Freudl R, Malecki MJ, Sahm H, Kozina VJ, Eggeling L, Saier MH Jr: The LysE superfamily: topology of the lysine exporter LysE of *Corynebacterium glutamicum*, a paradigm for a novel superfamily of transmembrane solute translocators. *J Mol Microbiol Biotechnol* 1999;1:327–336.
- 47 Zhai Y, Saier MH Jr: A web-based program for the prediction of average hydropathy, average amphipathicity and average similarity of multiply aligned homologous proteins. *J Mol Microbiol Biotechnol* 2001;3:285–286.
- 48 Moller S, Croning MD, Apweiler R: Evaluation of methods for the prediction of membrane spanning regions. *Bioinformatics* 2001;17:646–653.