Is This Stricture Inflammatory?

Gerhard Rogler

Division of Gastroenterology and Hepatology, University Hospital of Zürich, Zürich, Switzerland

About 75% of all patients suffering from Crohn's disease (CD) have to undergo surgery at least once during the course of their disease [1–4]. In half of these patients, intestinal obstructions and strictures are the indication for surgery. This means that intestinal obstructions cause the need for abdominal surgery in approximately one third of all CD patients. In more than 45% of those patients the obstructions are recurrent.

These data indicate that a stricturing disease course is still a major challenge in the clinical care of CD patients [1]. As therapeutic options differ, it is of major importance for the clinical management if a stenosis is fibrotic, inflammatory or both (and in the latter case more fibrotic or more inflammatory). An inflammatory stenosis would primarily be treated conservatively with (topical) steroids, immunosuppressants or biologicals. A fibrotic stenosis will not respond to such a treatment and should be operated early as a conservative approach may raise the risk for complete bowel obstruction.

In clinical practice we usually apply different diagnostic tools to discriminate between fibrotic and inflammatory strictures. We use imaging techniques (such as MRI or CT and ultrasound), evaluate systemic parameters of inflammation (such as CRP or ESR), and quantify fecal markers of inflammation (such as calprotectin or lactoferrin) [5]. Sometimes we obtain sufficient information from such diagnostic techniques to make a decision, sometimes we do not have sufficient evidence for one or the other option. Therefore, frequently a probatory steroid treatment will be performed. However, it has to be kept in mind that a steroid dose of more than 20 mg/day may increase the frequency of perioperative complications [6].

Recent data indicate that ultrasound has similar specificity and sensitivity as compared to MRI for the evaluation of CD activity [7, 8]. Ultrasound has been shown to be useful for the clinical management of CD [8–12]. Ultrasound results have been shown to be predictive for the disease behavior [10–16], which is of importance in the context of the study presented in this issue of *Digestion*.

Bowel wall thickness has been the most important readout in studies on the diagnostic value of ultrasound in IBD patients [10, 11, 16–19]. In addition to simple morphometry, contrast-enhanced ultrasound or splanchnic artery blood flow measurements have been applied with mixed results [12, 20–25].

Ultrasound has on one hand the great advantage of being readily available and non-invasive. It can be used repeatedly to monitor therapeutic success. On the other hand the subjective nature of ultrasound has been regarded as its major disadvantage. Therefore, the aim of the study by Schirin-Sokhan and co-workers [26] was to define objective quantitative measures of stricture characteristics (fibrostenotic vs. inflammatory) using contrast-enhanced ultrasound in CD patients with small bowel stenosis. The authors prospectively recruited 18 consecutive CD patients with a localized significant small bowel stenosis during a period of 18 months. They performed a standardized ultrasound examination, color-coded duplex sonography and contrast-enhanced ultrasound using SonoVue®. Bowel wall vascularity was quantified using computerized algorithms.

Schirin-Sokhan et al. [26] found no significant correlations between the vascularity parameters and the outcome in the clinical follow-up (which is the most important parameter) that would indicate a method for a clear sono-

graphic discrimination between fibrostenotic and inflammatory stenoses in CD patients. Certainly the results of this study have to be interpreted carefully due to the limited number of patients investigated. With a number of only 18 patients that may have had mixed (fibrotic and inflammatory) strictures it is very hard to find sufficient positive predictive values for a diagnostic technique.

Whether the establishment of computerized algorithms for objective quantification of bowel wall vascu-

larity will be helpful has to be evaluated by further studies. The study presented in this issue of *Digestion* should stimulate such research.

Disclosure Statement

G.R. discloses grant support from Abbot, Ardeypharm, Essex, FALK, Flamentera, Novartis, Tillots, UCB and Zeller.

References

- Peyrin-Biroulet L, Loftus EV Jr, Colombel JF, Sandborn WJ: The natural history of adult Crohn's disease in population-based cohorts. *Am J Gastroenterol* 2010;105:289–97.
- Doherty G, Bennett G, Patil S, Cheifetz A, Moss AC: Interventions for prevention of post-operative recurrence of Crohn's disease. *Cochrane Database Syst Rev* 2009;CD006873.
- Cosnes J: Can we modulate the clinical course of inflammatory bowel diseases by our current treatment strategies? *Dig Dis* 2009;27:516–521.
- Van Assche G, Vermeire S, Rutgeerts P: The potential for disease modification in Crohn's disease. *Nat Rev Gastroenterol Hepatol* 2010;7:79–85.
- Lakatos PL, Kiss LS: Is the disease course predictable in inflammatory bowel diseases? *World J Gastroenterol* 2010;16:2591–2599.
- Aberra FN, Lewis JD, Hass D, Rombeau JL, Osborne B, Lichtenstein GR: Corticosteroids and immunomodulators: postoperative infectious complication risk in inflammatory bowel disease patients. *Gastroenterology* 2003;125:320–327.
- Martinez MJ, Ripolles T, Paredes JM, Blanc E, Marti-Bonmati L: Assessment of the extension and the inflammatory activity in Crohn's disease: comparison of ultrasound and MRI. *Abdom Imaging* 2009;34:141–148.
- Pascu M, Roznowski AB, Muller HP, Adler A, Wiedenmann B, Dignass AU: Clinical relevance of transabdominal ultrasonography and magnetic resonance imaging in patients with inflammatory bowel disease of the terminal ileum and large bowel. *Inflamm Bowel Dis* 2004;10:373–382.
- Swoger JM, Regueiro M: Preventive therapy in postoperative Crohn's disease. *Curr Opin Gastroenterol* 2010;26:337–343.
- Rigazio C, Ercole E, Laudi C, Daperno M, Lavagna A, Crocella L, et al: Abdominal bowel ultrasound can predict the risk of surgery in Crohn's disease: proposal of an ultrasonographic score. *Scand J Gastroenterol* 2009;44:585–593.
- Dietrich CF: Significance of abdominal ultrasound in inflammatory bowel disease. *Dig Dis* 2009;27:482–493.
- Ripolles T, Martinez MJ, Barrachina MM: Crohn's disease and color Doppler sonography: response to treatment and its relationship with long-term prognosis. *J Clin Ultrasound* 2008;36:267–272.
- Castiglione F, de Sio I, Cozzolino A, Rispo A, Manguso F, Del Vecchio Blanco G, et al: Bowel wall thickness at abdominal ultrasound and the one-year-risk of surgery in patients with Crohn's disease. *Am J Gastroenterol* 2004;99:1977–1983.
- Kunihiro K, Hata J, Manabe N, Mitsuoka Y, Tanaka S, Haruma K, et al: Predicting the need for surgery in Crohn's disease with contrast harmonic ultrasound. *Scand J Gastroenterol* 2007;42:577–585.
- Maconi G, Parente F, Bollani S, Cesana B, Bianchi Porro G: Abdominal ultrasound in the assessment of extent and activity of Crohn's disease: clinical significance and implication of bowel wall thickening. *Am J Gastroenterol* 1996;91:1604–1609.
- Parente F, Sampietro GM, Molteni M, Greco S, Anderloni A, Sposito C, et al: Behaviour of the bowel wall during the first year after surgery is a strong predictor of symptomatic recurrence of Crohn's disease: a prospective study. *Aliment Pharmacol Ther* 2004;20:959–968.
- Paredes JM, Ripolles T, Cortes X, Martinez MJ, Barrachina M, Gomez F, et al: Abdominal sonographic changes after antibody to tumor necrosis factor (anti-TNF) α therapy in Crohn's Disease. *Dig Dis Sci* 2010;55:404–410.
- Calabrese E, Petruzzello C, Onali S, Condino G, Zorzi F, Pallone F, et al: Severity of postoperative recurrence in Crohn's disease: correlation between endoscopic and sonographic findings. *Inflamm Bowel Dis* 2009;15:1635–1642.
- Fraquelli M, Sarno A, Girelli C, Laudi C, Buscarini E, Villa C, et al: Reproducibility of bowel ultrasonography in the evaluation of Crohn's disease. *Dig Liver Dis* 2008;40:860–866.
- Quaia E, Migaleddu V, Baratella E, Pizzolato R, Rossi A, Grotto M, et al: The diagnostic value of small bowel wall vascularity after sulfur hexafluoride-filled microbubble injection in patients with Crohn's disease. Correlation with the therapeutic effectiveness of specific anti-inflammatory treatment. *Eur J Radiol* 2009;69:438–444.
- Migaleddu V, Scanu AM, Quaia E, Rocca PC, Dore MP, Scanu D, et al: Contrast-enhanced ultrasonographic evaluation of inflammatory activity in Crohn's disease. *Gastroenterology* 2009;137:43–52.
- Castiglione F, Bucci L, Pesce G, De Palma GD, Camera L, Cipolletta F, et al: Oral contrast-enhanced sonography for the diagnosis and grading of postsurgical recurrence of Crohn's disease. *Inflamm Bowel Dis* 2008;14:1240–1245.
- Serra C, Menozzi G, Labate AM, Giangregorio F, Gionchetti P, Beltrami M, et al: Ultrasound assessment of vascularization of the thickened terminal ileum wall in Crohn's disease patients using a low-mechanical index real-time scanning technique with a second-generation ultrasound contrast agent. *Eur J Radiol* 2007;62:114–121.
- Dietrich CF, Jedrzejczyk M, Ignee A: Sonographic assessment of splanchnic arteries and the bowel wall. *Eur J Radiol* 2007;64:202–212.
- Kratzer W, Schmidt SA, Mittrach C, Haenle MM, Mason RA, Von Tirpitz C, et al: Contrast-enhanced wideband harmonic imaging ultrasound (SonoVue®): a new technique for quantifying bowel wall vascularity in Crohn's disease. *Scand J Gastroenterol* 2005;40:985–991.
- Schirin-Sokhan R, Winograd R, Tischendorf S, Wasmuth HE, Streetz K, Tacke F, Trautwein C, Tischendorf JJW: Assessment of inflammatory and fibrotic stenoses in patients with Crohn's disease using contrast-enhanced ultrasound and computerized algorithm: a pilot study. *Digestion* 2011;83:263–268.