A Novel Approach to Assess Oral Feeding Skills of Preterm Infants

C. Lau E.O. Smith

Department of Pediatrics, Baylor College of Medicine, Houston Tex., USA

Key Words

Very low birth weight · Bottle feeding · Prematurity · Oral feeding evaluation

Abstract

Background: There is no well-defined means to identify the level of oral feeding skills (OFS) in preterm infants. **Objective:** To determine whether OFS as reflected by the combination of proficiency (PRO, %ml taken during the first 5 min/ml prescribed) and rate of milk transfer (RT, ml/min) correlates with gestational age (GA), oral feeding performance (OT, %ml taken during a feeding/ml prescribed) and days from start to independent oral feeding (SOF-IOF). Our working premises are that PRO is reflective of infants' *actual* feeding skills *when fatigue is minimal* and RT, monitored over an entire feeding session, reflects their overall skills when fatigue comes into play. **Methods:** Infants (26–36 weeks GA) with prematurity as their principal diagnosis were recruited and monitored at their first oral feeding. GA was divided into 3 strata, 26–29, 30–33, and 34–36 weeks GA. OFS was divided into 4 levels delineated by PRO ($\geq$ or $<30\%$) and RT ($\geq$ or <1.5 ml/min). ANOVA with post-hoc Bonferroni and multiple regression analyses were used. **Results:** OFS levels were correlated with GA. OT, PRO, and days from SOF-IOF were associated with OFS and GA, whereas RT was only correlated with OFS levels. **Conclusions:** OFS is a novel objective indicator of infants' feeding ability that takes into account infants' skills

and endurance. As a clinical tool, it can help caretakers monitor infants' skills as they transition to oral feeding and identify oral feeding issues arising from immature skills and/or poor endurance.

Copyright © 2011 S. Karger AG, Basel

Introduction

Attainment of independent oral feeding is one of the criteria recommended by the American Academy of Pediatrics for hospital discharge of preterm infants [1]. As such, an infant's inability to wean from tube feeding likely will delay hospital discharge and mother-infant reunion, while increasing medical cost and maternal stress [2, 3].

There are two dilemmas caregivers face when addressing oral feeding difficulties, i.e. infant ability to complete their feedings safely and the appropriate rate of advancement to independent oral feeding. Assessing infants' oral feeding skills has been difficult due to the lack of well-defined outcomes. Descriptive and objective scales have been developed. A descriptive scale uses visual, sensory and behavioral observations to assess the appropriateness of oral feeding skills, e.g. the Neonatal Oral-Motor Assessment Scale (NOMAS) developed by Palmer et al. [4] and the Early Feeding Skill assessment (EFS) by Thoyre et al. [5]. The accuracy of this type of approach has been

debated due to its subjective nature and lack of direct measure of specific outcomes. Recently, a debate was raised regarding the validity of NOMAS when used in preterm infants insofar as this scale was developed from term infants [6, 7]. Als et al. [8] have recommended the use of a developmental care approach to promote transition to oral feeding with the reasoning that if an infant's stability, organization and competence could be enhanced, his/her physiologic and behavioral expression would be optimized. Multidisciplinary approaches have also been proposed. An integrative assessment of an infant's potential ability to feed orally would be more accurate if feedback from the varied caregivers involved in his/her care were taken into account [9, 10]. More objective and quantitative evaluations of feeding skills are illustrated by tools developed to measure outcomes such as nutritive and nonnutritive sucking patterns and their rhythmicity, sucking force and coordination of suck-swallow-respiration [11–15]. However, these approaches necessitate special equipment, not readily available in neonatal intensive care units (NICU). Therefore, due to the difficulty in identifying the level of oral feeding skills (OFS), introduction and advancement of oral feeding remain in the hands of caregivers, with oral feeding customarily introduced around 33–35 weeks postmenstrual age (PMA).

In an earlier study [16] conducted at the first oral feeding of infants born between 26 and 30 weeks gestation (GA), we defined infants' OFS as a function of their combined proficiency (PRO, % volume taken during the first 5 min/total volume prescribed) and rate of milk transfer over an entire feeding (RT, ml/min). PRO, being monitored during the first 5 min of a feeding, was used as an index of infants' *actual* feeding ability when fatigue is minimal. RT, being monitored over an entire feeding session, was used as an index of endurance. It was reasoned that during a feeding the rate of transfer would decrease if infants' *actual* skills were held back by increasing fatigue. We defined feeding as successful or not if infants completed $\geq 80\%$ or $< 80\%$ of their prescribed feeding, respectively. Based on their performance, we distinguished 4 OFS levels as delineated by $\text{PRO} > 30\%$ or $< 30\%$ and $\text{RT} > 1.5$ or < 1.5 ml/min. The cutoffs for these two outcomes were based on the observations that infants demonstrating $\text{RT} \geq 1.5$ ml/min and $\text{PRO} \geq 30\%$ were successful at feeding, i.e. taking $\geq 80\%$ of their prescribed volume and attained independent oral feeding at an earlier PMA than counterparts whose PRO were $< 30\%$. Compared with this first group, infants with $\text{RT} \geq 1.5$ ml/min, but $\text{PRO} < 30\%$, also fed successfully ($\geq 80\%$), but

attained independent oral feeding at a later PMA, whereas infants with $\text{PRO} < 30\%$ and $\text{RT} < 1.5$ ml/min were not successful at feeding ($< 80\%$) and reached independent oral feeding at a later PMA.

This prospective study is a follow-up to our earlier work. Its goal was to determine whether the defined OFS levels can be used as an *objective* tool for the assessment of preterm infants' oral feeding skills. We hypothesized that (1) the more mature an infant's OFS level, the better his/her OT at that feeding; (2) the more premature an infant (GA), the more immature his/her OFS level, and (3) the better the OFS levels, the faster independent oral feeding will be attained.

Methods

Clinical Practice

Although breast-feeding is the optimal approach to infant feeding, in developed countries exclusive breast-feeding of hospitalized preterm infants is difficult to achieve. Thus, in the absence of mothers and when preterm infants are deemed ready to start oral feeding, bottle feeding is common practice rather than maintaining tube feeding. As is the practice at our hospital, our subjects' first oral feeding experience is with bottle rather than breast. Introduction and advancement of oral feeding are left to the discretion of the attending neonatologists. With no evidence-based determinant(s) of oral feeding readiness available, our attending neonatologists' decision to advance oral feeding generally takes into account their patients' clinical status, PMA, and the infants' oral feeding performance the day before if oral feeding started. It is also our hospital's practice that neonatal nutritionists monitor infants' daily weight to ensure at least a gain of 15 g/kg body weight/day.

Subjects

Infants born between 26 and 36 weeks GA were recruited from the level II intermediate care nurseries at Texas Children's Hospital (Houston, Tex., USA). They were stratified into 3 GA strata: 26–29, 30–33 and 34–36 weeks GA. Exclusion criteria included gastrointestinal complications and various medical conditions, e.g. congenital cytomegalovirus, intraventricular hemorrhage grades III and IV. No infants with bronchopulmonary dysplasia were recruited as they remained in the level III NICU as per hospital protocol. Only 'feeders and growers' with a primary diagnosis of prematurity were enrolled. This diagnosis defined infants who were clinically stable, demonstrated clinical signs of immature systems, e.g. lung function, that resolved with maturation and whose discharge from the hospital was primarily based on their ability to feed by mouth. This study was approved by the Baylor College of Medicine Institutional Review Board for Human Research. Parental consent was obtained following approval by attending neonatologists.

Outcome Measures

Infants were monitored at the time of their first oral feeding. They were identified by charge nurses who provided the infants'

nurses with a study form to fill out immediately following the feeding session with the following information: total volume prescribed (ml), total volume taken during the feeding (ml), volume taken during the first 5 min of the feeding (ml), duration of oral feeding (min) and any episodes of adverse events, namely oxygen desaturation, apnea and/or bradycardia. Infants were fed for a maximum of 20 min as per nursery protocol, with nurses deciding on how long to feed based on their perception of infants' performance and endurance. Infants were not disturbed for at least 30 min prior to this feeding session in order to minimize fatigue prior to oral feeding. From the measures collected, the following outcomes were computed: overall transfer (OT, % volume taken/total volume to be taken); PRO (% volume taken during the first 5 min/total volume prescribed); RT (ml/min). The number of days taken from the start to independent oral feeding (SOF-IOF) was recorded. Four OFS levels were delineated by their respective PRO ($\geq 30\%$ or $<30\%$) and RT (≥ 1.5 or <1.5 ml/min) as described in our earlier study [16]. Thus, level 1, the most immature, was defined by PRO $<30\%$ and RT <1.5 ml/min, level 2, by PRO $<30\%$ and RT ≥ 1.5 ml/min, level 3, by PRO $\geq 30\%$ and RT <1.5 ml/min, and level 4, the most mature, by PRO $\geq 30\%$ and RT ≥ 1.5 ml/min. As in our earlier study, our working premises were as follows. PRO, being monitored during the first 5 min of a feeding, is representative of infants' *actual* nutritive feeding skills as fatigue was deemed minimal (infants being undisturbed for 30 min prior to feeding). RT, being monitored over an entire feeding session, reflected infant overall skill when fatigue came into play and was used as an index of overall endurance (fig. 1).

Statistical analyses included univariate analyses of variance using Bonferroni for post-hoc comparisons when appropriate. Potential associations between outcomes and multiple factors were determined by multiple linear regression analyses (SPSS, v.17, Chicago, Ill., USA).

Results

Sixty-six subjects classified as 'feeders and growers' with a primary diagnosis of prematurity were recruited between March and September 2008. Their ethnic/racial and gender distribution are presented in table 1. Table 2 shows infants' characteristics by GA strata. As expected, significant differences in GA, birth weight and days of life at first and independent oral feeding were noted between GA strata ($p < 0.001$). Infants less than 34 weeks GA were introduced to oral feeding around 34–35 weeks PMA and within the first week of life for those born between 34 and 36 weeks GA. PMA at IOF was similar between GA strata (table 2), but significantly younger with more mature OFS levels (table 3). On the other hand, whereas PMA at discharge was significantly greater the more premature the infants (table 2), it was not significant between OFS levels (table 3). Multiple linear regression demonstrated that OFS levels were correlated with GA ($p = 0.017$) but not with PMA ($p = 0.234$). With increasing GA, a trend

Fig. 1. Four OFS levels as defined by PRO and RT.

Table 1. Subjects' ethnic/racial and gender distribution

	Number	Percent
Ethnicity/Race		
Hispanic	14	21
Caucasian	32	49
African-American	19	29
Asian	1	1
Gender		
Male	36	55
Female	30	45

was noted towards a gradual decrease in the predominance of the most immature OFS (level 1) with a concurrent increase in the more mature levels 2–4 (fig. 2). Subjects' oral feeding outcomes, i.e. OT, PRO, RT and feeding duration by GA strata and OFS levels, are presented in tables 2 and 3, respectively. None of these infants demonstrated any significant adverse events at their first oral feeding. OT and PRO were positively correlated with GA strata and OFS levels ($p \leq 0.004$), and a negative association was noted with feeding duration ($p \leq 0.014$). RT did not differ between GA strata ($p = 0.172$) but did between OFS levels ($p < 0.001$). Multiple linear regression analyses showed associations between oral feeding outcomes with OFS levels, GA and PMA ($p \leq 0.022$; table 4). Post-hoc analyses demonstrated that OT, PRO and days from SOF to IOF were primarily associated with OFS levels and GA. In turn, RT was only with OFS levels and feeding duration with GA strata. Average days from SOF-IOF by OFS levels and GA strata are presented in table 5.

In an attempt to determine the individual effect of infants' *actual* feeding skills (as measured by PRO when

Table 2. Characteristics and oral feeding outcomes of subjects who attained independent oral feeding prior to discharge

	GA strata, weeks			p ¹
	26–29	30–33	34–36	
Number	20	39	7	
GA, weeks	27.9 ± 1.2	31.8 ± 1.1	34.8 ± 0.8	<0.001
Birth weight, g	1,047.9 ± 194.8 ^{a, b}	1,545.2 ± 305.4 ^{a, c}	2,275.1 ± 444.8 ^{b, c}	<0.001
DOL at SOF	53 ± 33 ^{a, b}	17 ± 9 ^a	5 ± 5 ^b	<0.001
PMA at SOF	35.4 ± 4.3	34.2 ± 0.9	35.5 ± 1.0	0.149
DOL at IOF	62 ± 17 ^{a, b}	28 ± 11 ^{a, c}	10 ± 7 ^{b, c}	<0.001
PMA at IOF	36.9 ± 2.5	35.8 ± 1.2	36.2 ± 1.1	0.112
PMA at discharge	40.1 ± 4.9 ^{a, b}	37.3 ± 1.8 ^a	36.9 ± 1.0 ^b	0.004
OT, %	46.6 ± 35.5 ^a	53.0 ± 33.4 ^b	95.2 ± 12.6 ^{a, b}	0.004
PRO, %	21.3 ± 18.8 ^a	28.4 ± 24.8 ^b	57.1 ± 26.3 ^{a, b}	0.004
RT, ml/min	1.0 ± 1.1	1.1 ± 0.7	1.7 ± 1.1	0.172
Feeding duration, min	17 ± 5 ^{a, b}	14 ± 5 ^a	11 ± 5 ^b	0.014

Means ± SD. GA = Gestational age; DOL = days of life; SOF = start of oral feeding; PMA = postmenstrual age (weeks); IOF = independent oral feeding (8 oral feedings/day).

¹ One-way ANOVA; post-hoc Bonferroni between different superscript letters p ≤ 0.05.

Table 3. Oral feeding performance at first oral feeding by OFS level (mean ± SD)

	OFS level				p ¹
	1	2	3	4	
Number	36	5	14	11	
Infants at each level, %	54.5	7.6	21.2	16.7	
OT, %	30.3 ± 23.6 ^{a–c}	84.8 ± 16.5 ^a	77.9 ± 21.1 ^{b, d}	96.3 ± 12.3 ^{c, d}	<0.001
PRO, %	12.5 ± 7.3 ^{a, b}	23.1 ± 3.6 ^{c, d}	47.3 ± 20.2 ^{a, c, e}	64.4 ± 22.8 ^{b, d, e}	<0.001
RT, ml/min	0.6 ± 0.4 ^{a–c}	1.9 ± 0.3 ^{a, d, e}	1.0 ± 0.2 ^{b, d, f}	2.6 ± 1.0 ^{c, e, f}	<0.001
Feeding duration, min	14.9 ± 4.9 ^a	17.4 ± 2.5 ^b	15.9 ± 5.4 ^c	9.8 ± 4.7 ^{a–c}	0.007
PMA at IOF, weeks	36.8 ± 1.9 ^{a, b}	35.9 ± 1.1	35.6 ± 1.0 ^a	35.2 ± 1.4 ^b	0.020
PMA at discharge, weeks	38.9 ± 3.9	37.3 ± 1.1	37.6 ± 2.0	36.4 ± 1.4	0.102

¹ One-way ANOVA; post-hoc Bonferroni between different superscript letters p ≤ 0.05.

fatigue was minimal) and endurance (as measured by RT) on overall transfer (OT), infants were divided into two groups, i.e. ‘high’ and ‘low’ *actual* skill level as defined by PRO ≥ 30% and <30%, respectively (fig. 3). Two-way ANOVA showed that significant PRO and RT effects along with a significant interaction between PRO·RT on OT (p < 0.001). OT of infants in the high-PRO groups (OFS levels 3 and 4) were significantly greater than of those of their low-PRO counterparts (OFS levels 1 and 2). Similarly, infants in the high RT (≥ 1.5 ml/min, OFS levels 2 and 4) performed better than counterparts with low RT (< 1.5 ml/min, OFS levels 1 and 3). Post-hoc analyses

of PRO·RT demonstrated that infants at an OFS level 1 (low PRO, low RT) performed significantly worse than counterparts at levels 2–4 (p < 0.001). No differences were observed between infants at OFS levels ≥ 2 (p ≥ 0.210).

Discussion

Caregivers frequently question the ‘oral feeding readiness’ of preterm infants when considering introduction of oral feeding. As no accurate tool is available to determine whether an infant is ‘ready’ to wean from tube feed-

Fig. 2. Percent distribution of OFS levels by GA stratum in infants born between 26 and 36 weeks gestation.

Fig. 3. OT of infants with high versus low *actual* feeding skills (PRO >30%/<30%) vs. endurance (RT >1.5/<1.5 ml/min).

Table 4. Multiple regression analyses of oral feeding outcomes by OFS levels, GA strata and PMA category

	OFS level	GA strata	PMA category	p
OT, %	<0.001	0.053	0.431	<0.001 (64.4) ^a
PRO, %	<0.001	0.034	0.180	<0.001 (70.0)
RT, ml/min	<0.001	0.714	0.928	<0.001 (43.9)
Feeding duration, min	0.274	0.013	0.440	<0.022 (10.5)
SOF-IOF, days	0.003	<0.001	0.080	<0.001 (39.5)

^a Figures in parentheses are adj. R2 (expressed in percentages).

Table 5. Time from introduction to attainment of independent oral feeding

	GA strata, weeks				OFS levels				
	26-29	30-33	34-36	p ¹	1	2	3	4	p ¹
SOF to IOF, days	20 ± 11 ^{a, b}	13 ± 5 ^a	5 ± 3 ^b	<0.001	17 ± 9 ^{a, b}	14 ± 7	12 ± 6 ^a	7 ± 4 ^b	0.002

¹ One-way ANOVA; post-hoc Bonferroni between different superscript letters p ≤ 0.05.

ing, introduction and advancement of oral feedings are commonly ordered by attending physicians, with some practices leaving the decision to nurses or following prefeeding assessment programs such as those of Thoyre et al. [5] and McCain [17]. Infants with oral feeding issues are routinely referred to feeding therapists. However, if any improvement occurs, it is uncertain whether it resulted from normal maturation or the therapies provided

as evidence-based data lack in supporting the latter's benefit(s).

This study was conducted to determine whether OFS levels as defined by the combination of PRO and RT may be used as an objective indicator of infants' feeding skills. Our hypotheses were confirmed insofar as the OFS levels were (1) positively correlated with an infant's feeding performance, i.e. the better the levels, the greater the OT and

Table 6. OFS levels as *potential* screening tool for assessment of infants' *actual* feeding skills and endurance

OFS level (OT)	Actual feeding skills (PRO)	Endurance (RT)	Potential intervention(s)
1	low	low	nonnutritive oral motor stimulation + endurance training
2	low	high	nonnutritive oral motor stimulation
3	high	low	endurance training
4	high	high	none

the shorter the feeding duration; (2) positively correlated with GA strata, i.e. the less premature the infant, the more mature his/her skills, and (3) inversely associated with days from SOF to IOF, i.e. the better the skills, the faster the attainment of independent oral feeding.

It is unclear why infants within a particular GA stratum, even the most premature one, exhibited all 4 levels of OFS when tested at the same PMA. Prenatal factors may have been involved. The lack of correlation between oral feeding outcomes and PMA was expected insofar as infants were introduced to oral feeding at similar PMA. The associations of *OT* and *PRO* with OFS levels helps explain why some infants of similar gestational and/or postnatal ages readily transition from tube to oral feeding while others do not. Indeed, we showed that infants within each GA stratum exhibited OFS levels ranging from 1 to 4. Consequently, the use of GA may not be the optimal indicator for assisting the decision-making of caregivers in the initiation and advancement of oral feeding in preterm infants. The correlations of *OT* and *PRO* with GA can be explained by the greater proportion of infants in the older GA strata that naturally demonstrated more mature OFS levels. The observation that *RT* was associated with OFS, but not GA, suggests that *RT* is primarily regulated by an infant's feeding aptitude, e.g. suck, swallow, suck-swallow-respiration coordination and/or endurance. This is supported by an earlier work where we showed that infants feeding with nipples of varied flow rates within a 24-hour period could modify their sucking skills, e.g. sucking pattern and suction amplitude, in order to maintain the same *RT* [18]. The correlation between *feeding duration* and GA is consistent with the lower *PRO*, *OT* and the trend towards slower *RT* observed at the younger GA. The association of *days from SOF-IOF*

with both OFS levels and GA supports the notion that feeding performance resulting from infants' OFS levels and GA are criteria used by our attending physicians in advancing oral feeding. Infants with *high PRO* ($\geq 30\%$, OFS levels 3 and 4) perform better than their counterparts with low *PRO* ($<30\%$, OFS levels 1 and 2). Similarly, infants with *high RT* (OFS levels 2 and 4) also performed better than infants with low *RT* (OFS levels 1 and 3). The observation that infants at OFS levels 2 and 3 had similar *OT* suggests that both *actual* feeding skill and endurance are equally important in determining oral feeding success. From this, one may speculate that enhancing both factors would optimize outcome. This is supported by the observations that *OT*, *PRO*, *RT* and feeding duration of infants at OFS level 4 were superior to those of their counterparts at levels 1–3.

From this study, it is proposed that the use of OFS levels can offer a more objective indicator of infants' ability to feed by mouth than GA or other tools currently available. It does not claim to provide the ultimate answer for solving infants' oral feeding difficulties as the latter are multifactorial. However, it offers the possibility of differentiating between feeding aptitude and endurance/fatigue which, as we showed, are both equally important for oral feeding success. Its use offers several advantages. (1) It is easy to measure as caregivers need only collect the volume reading at 5 min into the feeding session in addition to the routine information collected, i.e. volume prescribed, volume taken and feeding duration. No special equipment is required. (2) It provides an objective rather than subjective assessment of infants' feeding skills during a feeding session. (3) As infants of similar GA differ in OFS, evaluating their levels prior to the introduction of oral feeding can help identify infants' ability when oral feeding is initiated. (4) Measuring OFS levels does not only pertain to infants' first oral feeding. Monitoring OFS longitudinally as infants wean from tube feeding provides information on their maturation process. It can be used as an indicator of whether oral feeding should be advanced or held back. Additionally, if an infant is receiving a particular intervention, monitoring over time can help determine its efficacy. (5) OFS levels may also assist caregivers in identifying whether infants' oral feeding issues relate to skill levels or endurance (table 6). For instance, if an infant exhibits an OFS level of 1, with low skill and endurance, he/she may benefit from oral feeding therapy and 'endurance training' (see below). A level 2 infant with low skill and high endurance would likely only require oral feeding therapy, whereas a level 3 infant with high skill and low endurance would benefit from

‘endurance training’. An infant at level 4 would need no intervention. Oral feeding therapy to enhance *actual* feeding skills may consist of a nonnutritive oral motor stimulation program, which has been shown to accelerate attainment of IOF [19–21]. In order to prevent negative oral feeding experiences and/or excessive fatigue [22], it is proposed that an ‘endurance training’ program be set up. This may consist of daily shortened feeding sessions, *the total duration of which equals the duration corresponding to the number of oral feedings per day ordered*. For instance, if an infant is allowed to feed once a day for a maximum of 20 min, but at the first feeding on that day, after 5 min he/she exhibits signs of fatigue, disorganization and/or unstable behavioral state, the ‘endurance training’ may consist of four 5-min feedings on that particular day. Feeding duration can be gradually increased on a daily basis as the above symptoms decrease. Although no study, to the authors’ knowledge, has yet been conducted to show that such training program is efficacious, it is based on the general acceptance that ‘practice makes perfect’ in the absence of adverse events.

In summary, we propose that the use of OFS levels is an additional indicator to assist caregivers in determining

infants’ oral feeding aptitude. It is novel and offers an objective assessment of oral feeding skills whether they are monitored at a first oral feeding experience or on a regular basis as they advance towards independent oral feeding.

Acknowledgments

The authors are grateful to all the nurses in the level II intermediate care nurseries at Texas Children’s Hospital for their enthusiasm in assisting with data collection. Special recognition is given to Clarito Arbilo, RN, who supervised the entire study and Averie Swanson, undergraduate research assistant for data management. This work was supported by a grant from the National Institute of Child Health and Human Development (R01-HD044469). The contents of this publication are solely the responsibility of the authors and do not necessarily represent the official views of the National Institute of Child Health and Human Development or the National Institutes of Health.

Disclosure Statement

The authors have no financial disclosure or conflict of interest to disclose relevant to this article.

References

- 1 American Academy of Pediatrics: Hospital discharge of the high-risk neonate. *Pediatrics* 2008;122:1119–1126.
- 2 Lau C, Hurst N: Oral feeding in infants. *Curr Probl Pediatr* 1999;29:105–124.
- 3 Burklow KA, McGrath AM, Kaul A: Management and prevention of feeding problems in young children with prematurity and very low birth weight. *Infants Young Children* 2002;14:19–30.
- 4 Palmer MM, Crawley K, Blanco IA: Neonatal Oral-Motor Assessment scale: a reliability study. *J Perinatol* 1993;13:28–35.
- 5 Thoyre SM, Shaker CS, Pridham KF: The early feeding skills assessment for preterm infants. *Neonatal Netw* 2005;24:7–16.
- 6 da Costa SP, van der Schans CP: The reliability of the Neonatal Oral-Motor Assessment Scale. *Acta Paediatr* 2008;97:21–26.
- 7 da Costa SP, van den Engel-Hoek L, Bos AF: Sucking and swallowing in infants and diagnostic tools. *J Perinatol* 2008;28:247–257.
- 8 Als H, Lawhon G, Duffy FH, McAnulty GB, Gibes-Grossman R, Blickman JG: Individualized developmental care for the very low-birth-weight preterm infant. Medical and neurofunctional effects (see comments). *JAMA* 1994;272:853–858.
- 9 Rommel N, De Meyer AM, Feenstra L, Veerman-Wauters G: The complexity of feeding problems in 700 infants and young children presenting to a tertiary care institution. *J Pediatr Gastroenterol Nutr* 2003;37:75–84.
- 10 Premji SS, McNeil DA, Scotland J: Regional neonatal oral feeding protocol: changing the ethos of feeding preterm infants. *J Perinat Neonatal Nurs* 2004;18:371–384.
- 11 Medoff-Cooper B, Verklan T, Carlson S: The development of sucking patterns and physiologic correlates in very-low-birth-weight infants. *Nurs Res* 1993;42:100–105.
- 12 Lau C, Alagurusamy R, Schanler RJ, Smith EO, Shulman RJ: Characterization of the developmental stages of sucking in preterm infants during bottle feeding. *Acta Paediatr* 2000;89:846–852.
- 13 Poore M, Zimmerman E, Barlow SM, Wang J, Gu F: Patterned orocutaneous therapy improves sucking and oral feeding in preterm infants. *Acta Paediatr* 2008;97:920–927.
- 14 Gewolb IH, Vice FL: Maturation changes in the rhythms, patterning, and coordination of respiration and swallow during feeding in preterm and term infants. *Dev Med Child Neurol* 2006;48:589–594.
- 15 Lau C, Smith EO, Schanler RJ: Coordination of suck-swallow and swallow respiration in preterm infants. *Acta Paediatr* 2003;92:721–727.
- 16 Lau C, Sheena HR, Shulman RJ, Schanler RJ: Oral feeding in low birth weight infants. *J Pediatr* 1997;130:561–569.
- 17 McCain GC: An evidence-based guideline for introducing oral feeding to healthy preterm infants. *Neonatal Netw* 2003;22:45–50.
- 18 Scheel CE, Schanler RJ, Lau C: Does the choice of bottle nipple affect the oral feeding performance of very-low-birthweight (VLBW) infants? *Acta Paediatr* 2005;94:1266–1272.
- 19 Fucile S, Gisel E, Lau C: Oral stimulation accelerates the transition from tube to oral feeding in preterm infants. *J Pediatr* 2002;141:230–236.
- 20 Fucile S, Gisel EG, Lau C: Effect of an oral stimulation program on sucking skill maturation of preterm infants. *Dev Med Child Neurol* 2005;47:158–162.
- 21 Rocha AD, Moreira ME, Pimenta HP, Ramos JR, Lucena SL: A randomized study of the efficacy of sensory-motor-oral stimulation and non-nutritive sucking in very low birth-weight infant. *Early Hum Dev* 2007;83:385–388.
- 22 Lau C: Oral feeding in the preterm infant. *NeoReviews* 2006;7:e19–e27.