Functional Neuroimaging in Obsessive-Compulsive Disorder

A. Del Casale^a G.D. Kotzalidis^a C. Rapinesi^a D. Serata^a E. Ambrosi^a
A. Simonetti^a M. Pompili^{a,b} S. Ferracuti^a R. Tatarelli^a P. Girardi^a

^aNESMOS (Neurosciences, Mental Health, and Sensory Organs) Department, School of Medicine and Psychology, Sapienza University and Sant'Andrea Hospital, Rome, Italy; ^bMcLean Hospital, Harvard Medical School, Belmont, Mass., USA

Key Words

Obsessive-compulsive disorder • Functional neuroimaging • Striatum • Orbitofrontal cortex • Cognitive functions

Abstract

Background and Aim: Obsessive-compulsive disorder (OCD) is a severe, highly prevalent and chronically disabling psychiatric disorder that usually emerges during childhood or adolescence. This paper aims to review the literature on functional neuroimaging in OCD, analysing the reported dysfunctional connectivity in the corticostriatothalamocortical circuitry. **Method:** This study included papers published in peer-reviewed journals dealing with functional imaging in OCD. **Results:** Striatal dysfunction, mainly of the caudate nucleus, leads to inefficient thalamic gating, resulting in hyperactivity within the orbitofrontal cortex (intrusive thoughts) and the anterior cingulate cortex (non-specific anxiety). Compulsions consist of ritualistic behaviours performed to recruit the inefficient striatum and neutralise unwanted thoughts and anxiety. Functional neuroimaging findings are discussed against the background of specific cognitive impairments, mainly regarding visuospatial processing, executive functioning and motor speed. Cognitive deficits are partial and specific, matching imaging data. **Conclusions:** Sev-

eral studies have targeted brain regions hypothesised to be involved in the pathogenesis of OCD, showing the existence of dysfunctional connectivity in the corticostriatothalamocortical circuitry. Improvements in spatial resolution of neuroimaging techniques may contribute to a better understanding of the neurocircuitry of OCD and other anxiety disorders.

Copyright © 2011 S. Karger AG, Basel

Introduction

Obsessive-Compulsive Disorder: Diagnostic Criteria and Epidemiology

Obsessive-compulsive disorder (OCD) is a common psychiatric disorder affecting more than 1% of the population worldwide [1]. It is included in the group of anxiety disorders in the DSM-IV [2], but there is considerable debate whether this should continue in the DSM-5 [3–7]. The hallmark symptoms of OCD include intrusive thoughts (obsessions) as well as ritualistic behaviour (compulsions). The lifetime prevalence of OCD is 2–3% in the general population [8], with no difference in gender distribution; however, juvenile-onset OCD tends more often to be familial and to have a higher prevalence in

boys [9, 10]. Its clinical presentation in children and adults is generally similar [2], but comorbidities differ [10]. Its usual onset is in late adolescence or early adulthood, although it may occur from childhood to advanced age [11]. Onset is usually gradual. The course follows a chronic but fluctuating pattern, often related to stressful life events [12, 13]. Epidemiological surveys suggest that 50% or more people with OCD have at least one comorbid psychiatric disorder, most commonly an anxiety disorder or unipolar depression. Alcohol abuse/dependence is higher in OCD than in the general population [14].

Functional Neuroimaging Techniques Used to Study OCD

Functional imaging studies are most useful to clarify the neural substrate underlying the pathophysiology of OCD. Positron emission tomography (PET), single photon emission computed tomography (SPECT), proton magnetic resonance spectroscopy (^1H -MRS) and functional magnetic resonance imaging (fMRI) are the most frequently used techniques for this purpose.

In the late 1980s, various groups [15–17] carried out PET studies and showed differences between OCD patients and healthy volunteers (table 1). SPECT has been first used in OCD by Machlin et al. [18]. They found alterations in the medial frontal cortex (MFC). This technique uses conventional nuclear medicine planar imaging and a gamma camera to provide three-dimensional rendering (table 2). fMRI first used the paramagnetic contrast agent gadolinium [19], and subsequently non-contrast-based techniques [20–22]. Since then, fMRI has been increasingly employed in psychiatric research and neuroscience. To evaluate brain functions after exposure to stimuli, fMRI is often combined with one or more tasks (i.e. verbal word fluency test, spatial item-recognition tasks, go/no go task, Stroop task, etc.). The use of cognitive or behavioural paradigms as circuitry-specific probes highlights the advantages of fMRI in pursuing this goal.

Studies comparing baseline scans with scans after effective treatment suggest that alterations are OCD state dependent as improvement in symptoms is associated with reestablishing a normal activation pattern [23–25]. Imaging studies of brain response to symptom provocation suggested direct frontal-subcortical circuit involvement in obsessions and compulsions [26–30]. However, these functional alterations could represent only part of a more widespread, latent, prefrontal dysfunction requiring different behavioural challenges to be detected [31, 32]. Our aim has been to review functional neuroimaging

studies of OCD, and to analyse the dysfunctional connectivity reported for the corticostriatalthalamocortical circuitry (table 3).

Method

We searched the Medline and PsycINFO databases combining ‘obsessive-compulsive disorder’ with other keywords like ‘positron emission tomography’, ‘functional magnetic resonance imaging’, ‘single photon emission computerized tomography’ and ‘magnetic resonance spectroscopy’. Papers were included if they satisfied standards for adequate methodology, diagnostics (DSM or ICD) and population inclusion (patients with OCD and at least healthy controls). Papers published in peer-reviewed journals dealing with functional imaging in OCD were included. Papers were excluded: if they did not clarify a method for diagnosing OCD, or if OCD was not the principal diagnosis; if severe psychiatric comorbidity was present that could have influenced OCD findings; if the imaging methods were unspecified or inadequately described, and if they were reviews or meta-analyses (but their reference lists were searched anyway). Further papers that did not appear in the above databases were searched from reference lists of papers retrieved.

Data Analysis

Our search yielded a total of 714 papers in the National Library Database/PubMed and 187 papers in PsycLIT as of June 1, 2010; the latter overlapped with the former by more than 95%. A large number of papers were irrelevant and emerged from the search due to its overinclusiveness. By restricting the search for keywords to the title of papers, the yield in PubMed was limited to 82 relevant papers, which we considered too restrictive. We therefore decided to scan all papers found, and eventually kept for analysis and discussion 143 of them that were actually neuroimaging studies, 125 of which were fMRI studies, 6 were PET, 7 were SPECT and 5 were MRS studies. Most irrelevant studies were excluded by simply focusing on their title and abstract; 53 were not immediately excluded from our analysis and our above criteria were applied to exclude them. Thirty-five did not focus on OCD (i.e. the focus lay on other diagnoses and obsessive-compulsive symptoms or OCD were only comorbid), 13 included patients with severe comorbidity, 4 did not use an adequate classification of symptoms, and 1 was excluded for multiple reasons.

The 143 included studies reported data on 755 patients with OCD. Original articles assessed several neuroimaging techniques (MRS: 3.5%; fMRI: 87.42%; SPECT: 4.89%; PET: 4.19%). Most studies (81.6%) compared OCD patients with healthy or other controls; some focused on intragroup comparisons (28.6%), others compared OCD with other disorders (10.2%). In OCD patients, these studies showed alterations: in the cingulate cortex (32.6%); in the ventrolateral prefrontal (10.2%), dorsolateral prefrontal (32.6%) and orbitofrontal (51%) cortices; in other prefrontal areas (10.2%); in the insular (10.2%), frontal, parietal and temporal (63.3%) cortices; in the thalamus (20.4%); in the cerebellum (8.2%); in the striatum (40.8%) including the caudate nuclei (18.4%) and the putamen (4%); in the amygdala (8.2%); in the hippocampal/parahippocampal region (6.1%), and in other brain regions (8.2%).

Table 1. Main PET studies in OCD

Study Ref. No.	Technique	Age group	Participants	Medication	Comorbidity	General findings in OCD patients
Baxter et al. [15]	PET	Adults	14 OCD patients: 9 m, 5 f; mean age: 31.6 years; 14 unipolar depression: 5 m, 9 f; 14 healthy controls: 7 m, 7 f	5 on medication	9 of OCD patients met criteria for major depression; no neurological disorder; head injury or serious medical condition	Increased bilateral head of caudate; increased left orbitofrontal gyrus
Baxter et al. [16]	PET	Adults	10 OCD patients: 5 m, 5 f; mean age: 35.5 years; illness duration: at least 1 year; 10 healthy controls: 5 m, 5 f	Drug free for at least 2 weeks before study	No neurological disorder; head injury or serious medical condition; 8 had major depression; 3 met criteria for social phobia, 2 of whom met criteria for simple phobia	Increased bilateral orbital gyri
Swedo et al. [17]	PET	Adults	18 OCD patients: 9 m, 9 f; severe OCD symptoms; adults with childhood-onset OCD; mean age: 27.8 years; mean age at onset: 8.9 years; illness duration: 19 years; 18 healthy controls: 9 m, 9 f	Free of psychoactive drugs for at least 4 weeks before the study	No neurological disorder; head injury or serious medical condition; major depression; anxiety disorder	Increased right thalamus, left orbitofrontal bilateral prefrontal cortex and bilateral anterior cingulate gyri
Martinot et al. [177]	PET	Adults	16 OCD patients: 9 m, 7 f; mean age: 44 years; adults with mixed childhood/adult onset; 8 healthy controls: 5 m, 3 f	10 on medication; 5 had no medications for at least 2 weeks before study; 1 was drug free	No major depressive episode; no neurological disorder; head injury or serious medical condition	Decreased metabolism in lateral prefrontal cortex
Sawle et al. [178]	PET	Adults	6 OCD patients; mixed childhood/adult onset; mean age: 34.3 years; illness duration: 15.8 years; 6 healthy controls	3 on medication	Depression; no neurological disorder; head injury or serious medical condition	Hypermetabolism in bilateral orbitofrontal, premotor and midfrontal cortices
Rauch et al. [29]	PET; symptom provocation paradigm	Adults	8 OCD patients: 5 m, 3 f; handedness (R:L) 8:0; mean age: 36.1 years; mean age at onset: 18.1 years	Free of psychoactive drugs for at least 2 weeks before the study	No neurological disorder; head injury or serious medical condition	Increase in relative regional cerebral blood flow during the OCD symptomatic state vs. the resting state in right caudate nucleus, left anterior cingulate cortex and bilateral orbitofrontal cortex
Perani et al. [35]	PET	Adults	11 OCD patients: 8 f, 3 m; 7 checkers, 4 washers; mean age: 26.1 years; illness duration: 7.5 years; 15 healthy controls: 11 m, 4 f	Free of psychoactive drugs for at least 2 weeks before the study	No axis I comorbid disorders; no neurological disorder; head injury or serious medical condition	Increased pallidum/putamen complex; increased thalamus; bilateral anterior, middle and posterior cingulate cortex increase
Cottraux et al. [26]	PET; symptom provocation paradigm	Adults	10 OCD patients: 5 m, 5 f; all with predominant checking rituals; handedness (R:L) 10:0; mean duration of symptoms: ≥ 2 h/day; illness duration: ≥ 1 year; mean age: 31.6 years; 10 HC: 5 m, 5 f; handedness (R:L) 10:0; mean age: 28.3 years	Free of psychoactive drugs for at least 2 weeks before the study	No comorbid axis I or II disorders; no neurological disorder; head injury or serious medical condition	Control subjects had significantly higher regional rCBF levels in the thalamus and putamen; a trend towards higher rCBF in OCD patients was found in the superior temporal regions; obsessive stimulation was associated with higher rCBF than neutral stimulation in orbitofrontal regions in both groups of subjects; under obsessive stimulation, hyperactivation in temporal and orbitofrontal regions

Table 1 (continued)

Study Ref. No.	Technique	Age group	Participants	Medication	Comorbidity	General findings in OCD patients
Rauch and Savage [110]	PET	Adults	9 OCD patients: 9 f; mean age: 31.7 years; mean years of education: 14.1; 9 HC: 9 f; mean age: 32.3 years; mean years of education: 15.7	Free of psychoactive drugs for at least 4 weeks before the study	No comorbid axis I or II disorders or substance abuse; no neurological disorder, head injury or serious medical condition	Bilateral activation of the hippocampal/parahippocampal region; non-bilateral activation within an inferior territory of the striatum
Brody et al. [136]	PET	Adults	27 OCD patients divided into: – fluoxetine group: 18 patients: 6 m, 9 f; handedness (R:L) 17:1; mean age: 31.2 years – behavioral therapy group: 9 patients: 3 m, 6 f; handedness (R:L) 8:1; mean age: 34.1 years	All subjects received FDG-PET scans before treatment with either BT (n = 18) or 60 mg/day of fluoxetine (n = 9)	In the medication treatment group: concomitant diagnoses of cyclothymic disorder, panic disorder, Tourette's disorder mild, and social phobia in 1 subject each In the BT treatment group: comorbid diagnoses of cyclothymic disorder, panic disorder and acrophobia in one subject each; no patients with comorbid current depressive episode at the time of the study; no neurological disorder, head injury or serious medical condition	Higher normalised metabolism in left OFC was associated with greater improvement in the BT-treated group, but with worse outcome in the fluoxetine-treated group
Rauch et al. [179]	PET; symptom provocation paradigm	Adults	9 OCD patients: 4 m, 5 f; handedness (R:L) 9:0; all with contamination symptoms as primary OCD manifestation; mean age: 28.8 years	Free of psychotropic medications for at least 4 weeks prior to the initial PET session (at least 6 weeks for fluoxetine); subjects were ineligible for participation if they had a history of prior treatment with fluvoxamine; 9 subjects with contamination-related OCD underwent a 12-week open trial of treatment with fluvoxamine	No comorbid axis I or II disorders; no history of drug or alcohol addiction; no neurological disorder, head injury or serious medical condition	Lower rCBF values in the OFC and higher rCBF values in the PCC predicted better treatment response; this same pattern of associations was present regardless of whether the imaging data were acquired during a provoked or neutral state
van den Heuvel et al. [72]	PET; symptom provocation paradigm	Adults	11 OCD patients: 8 m, 3 f; handedness (R:L) 11:0; mean age: 40.5 years; 10 HC: 7 m, 3 f; handedness (R:L) 10:0; mean age: 36.5 years	No psychotropic medication	No comorbid axis I or II disorders; no neurological disorder, head injury or serious medical condition	Hyperactivation in the left amygdala in response to contamination stimuli; sensitisation effects in the right amygdala; these paralleled an increase in levels of distress and obsessivity as well as a decrease in dorsolateral prefrontal activity

BT = Behavioural therapy; HC = healthy controls; OFC = orbitofrontal cortex; PCC = posterior cingulate cortex; rCBF = regional cerebral blood flow.

Table 2. SPECT and MRS Studies in OCD (PTSD = Posttraumatic stress disorder)

Study Ref. No.	Tech- nique	Age group	Participants	Medication	Comorbidity	General findings in OCD patients
Machlin et al. [18]	SPECT	Adults	10 OCD patients; varying severity; mixed childhood/adult onset; mean age: 34.1 years; age at onset: 15.8 years; 8 healthy controls	Medication free for 4 weeks to 2 years before study	No major depressive episode; no neurological disorder, head injury or serious medical condition	Increased medial frontal cortex
Rubin et al. [107]	SPECT	Adults	10 OCD patients: 10 m; mean age: 34.9 years; 10 healthy controls	Drug free for at least 4 weeks before study	No axis I comorbid disorders; no neurological disorder, head injury or serious medical condition	Decreased heads of bilateral caudate nuclei; increased orbitofrontal cortex bilaterally; increased left posterofrontal cortex
Ebert et al. [34]	¹ H-MRS	Adults	12 OCD patients: 8 m, 4 f; mean age: 27 years; illness duration: 11 years; 6 healthy controls: 5 m, 1 f	10 were never medicated or medication free for at least 6 months before study; 1 receiving clomipramine; 1 receiving fluvoxamine	N/A	Lowered metabolism in right striatum
Lucey et al. [180]	SPECT	Adults	15 OCD patients: 8 m, 7 f; mean age: 36 years; illness duration: 14 years; 16 PTSD patients: 14 m, 2 f; 15 patients with panic disorder and agoraphobia: 8 m, 7 f; 15 healthy controls: 8 m, 7 f	N/A	No axis I comorbid disorders; no neurological disorder, head injury or serious medical condition	Decreased superior frontal cortex bilaterally; decreased right caudate volume
Bartha et al. [181]	¹ H-MRS	Adults	13 OCD patients: 7 m, 6 f; 13 healthy controls	Medication free for 6 weeks before study	No neurological disorder, head injury or serious medical condition; 1 patient had comorbid depression	Lowered metabolism in the left striatum
Crespo-Facorro et al. [77]	SPECT	Adults	21 OCD patients: 14 m, 7 f without motor tics; mean age: 32 years; 8 OCD patients: 7 m, 1 f with motor tics; mean age: 26 years; 16 healthy controls: 10 m, 6 f	Drug free for a 2-week period before study (or 4 weeks if they were using fluoxetine), or never been treated	Current or past history of Gilles de la Tourette syndrome was ruled out; 7 patients had chronic tic disorder; no neurological disorder, head injury or serious medical condition	Decrease in the right orbitofrontal cortex in the OCD group without chronic tic; no difference for OCD patients with and without motor tics
Busatto et al. [71]	SPECT	Adults	26 OCD patients: 15 m, 11 f; mean age: 32.1 years; mean age at onset: 15.2 years; illness duration: 16.9 years; 22 healthy controls: 12 m, 10 f	11 drug naïve; 15 had no medication for 3 weeks (or 6 weeks in case of fluoxetine)	5 patients had current motor/vocal tics; no neurological disorder, head injury or serious medical condition	Reduction in right orbitofrontal cortex and in left anterior cingulate cortex
Alptekin et al. [182]	SPECT	Adults	9 OCD patients: 3 m, 6 f; mean age: 30.8 years; 6 healthy controls: 2 m, 4 f	5 were never medicated; 4 drug free for at least 6 months before study	N/A	Increased right thalamus and bilateral orbitofrontal cortex
Lacerda et al. [70]	SPECT	Adults	16 OCD patients: 8 m, 8 f; mean age: 29.5 years; age at onset: 15.8 years; illness duration: 13.7 years; 17 healthy controls: 10 m, 8 f	Drug free for at least 30 days before the study; 10 patients were drug naïve	N/A	Increased right superior frontal cortex; increases in right and left thalamus; increased inferior frontal cortex

Table 3. Main fMRI studies in OCD

Study	Technique	Age group	Participants	Medication	Comorbidity	General findings in OCD patients
Levine et al. [58]	fMRI; word fluency verbal paradigm	Adults	11 OCD patients: 10 m, 1 f; mean age: 34.4 years; 11 HC: 6 m, 5 f; mean age: 28.2 years	Outpatients receiving neuroleptic treatment, clinically stable for at least 1 year before the study	All patients with comorbid schizophrenia; no neurological disorder, head injury or serious medical condition	Negative relationship between OCD symptomatology and activation of the left dorsolateral prefrontal cortex in schizophrenic patients vs. controls
Pujol et al. [175]	fMRI; word generation task	Adults	20 OCD patients: 15 m, 5 f; handedness (R:L) 18:2; mean age: 29.4 years; 20 HC: 14 m, 6 f; mean age: 26.2 years	11 patients receiving diazepam (10–25 mg/day); 9 patients free from psychoactive treatment for at least 4 weeks before the study	No comorbid axis I or II disorders; no neurological disorder, head injury or serious medical condition	Focal activation of the left frontal cortex during word generation, significantly correlated with the severity of the clinical process
Adler et al. [43]	fMRI; symptom provocation paradigm; 'off-on' exposure	Adults	7 OCD outpatients: 3 m, 4 f; all right-handed; mean age: 27 years; mean years of education: 14; mean age at onset: 17 years; had illness for at least 4 years	4 drug naïve; 3 drug free for 2 years before the study; no subjects were taking medications with central nervous system effects	No current comorbid diagnoses; no substance use disorders; 5 subjects had at least 1 episode of MDD; 1 subject had dysthymia in the past; no subjects met criteria for a tic disorder; no neurological disorder, head injury or serious medical condition	Activation in frontal cortex (orbitofrontal, superior frontal, dorsolateral prefrontal), anterior medial, lateral temporal cortex and right anterior cingulate; right superior frontal activation inversely correlated with baseline compulsion symptomatology, and left orbitofrontal cortical activation inversely associated with changes in OCD self-ratings following provocative stimuli
Ursu et al. [37]	fMRI; continuous-performance task	Adults	13 OCD patients: 8 m, 5 f; handedness (R:L) 9:4; mean age: 30.85 years; 11 HC: 5 m, 6 f; handedness (R:L) 11:0; mean age: 35.45 years	1 patient free of medication; 10 patients on medications; 3 were taking sertraline, 2 fluvoxamine, 2 fluoxetine, 1 clomipramine, 1 citalopram, 1 venlafaxine, 1 gabapentin, 1 valproate	5 patients with history of MDD; 1 with history of anorexia; 3 with comorbid generalised anxiety disorder; 1 with comorbid social phobia; no lifetime psychotic symptoms, substance abuse within the previous 6 months, and presence of first-degree relatives with psychotic disorders; no neurological disorder, head injury or serious medical condition	ACC hyperactivity in OCD may play a more significant role in the pathogenesis of this disorder than previously thought; more general account of the ACC hyperactivity in OCD
Fitzgerald et al. [38]	fMRI; flanker interference task	Adults	8 OCD patients: 6 m, 2 f; mean age: 27.4 years; mean years of education: 15.5; 7 HC: 5 m, 2 f; mean age: 30 years	2 patients receiving fluoxetine; 1 receiving fluoxetine + clonazepam; 1 receiving fluoxetine + risperidone; 3 unmedicated	3 patients with comorbid major depression; 2 with comorbid dysthymia; no neurological disorder, head injury or serious medical condition	Greater error-related activation of the rostral ACC in OCD patients positively correlated with symptom severity
Maltby et al. [39]	fMRI; speeded reaction time task (go/no go task)	Adults	11 OCD patients: 4 m, 7 f; mean age: 39.36 years; 11 HC: 4 m, 7 f; mean age: 36.55 years	N/A	No comorbid psychotic disorder, neurologic disorder, substance abuse, serious suicidal ideation, head injury or serious medical condition; OCD was patients' primary diagnosis	Only correctly rejected, high-conflict trials produced excessive activation in both action monitoring (rostral and caudal ACC, LPFC), frontal striatal regions (lateral orbitofrontal cortex, caudate and thalamus) and posterior cingulate cortex

Study	Technique	Age group	Participants	Medication	Comorbidity	General findings in OCD patients
Nakao et al. [40]	fMRI; Stroop task	Adults	24 OCD patients: 9 m, 15 f; handedness (R:L) 24:0; mean age: 33.9 years; 14 HC: 5 m, 9 f; handedness (R:L) 13:1; mean age: 30.2 years	Free of psychoactive drugs for at least 2 weeks before the study	No comorbid axis I disorders; free of drug or alcohol addiction; no neurological disorder; head injury or serious medical condition	Weaker activation in the ACC and in the right caudate nucleus
van den Heuvel et al. [53]	fMRI; Tower of London task	Adults	22 OCD patients: 7 m, 15 f; handedness (R:L) 22:0; mean age: 34.4 years; 22 HC: 11 m, 11 f; handedness (R:L) 22:0; mean age: 29.9 years	Free of psychotropic medication or not receiving medication for at least 4 weeks	No comorbid axis I or II disorders; no neurological disorder; head injury or serious medical condition	Significant planning impairments: during planning, decreased frontostriatal responsiveness, mainly in dorsolateral prefrontal cortex and caudate nucleus; increased – presumably compensatory – anterior cingulate, ventrolateral prefrontal and parahippocampal cortices
Viard et al. [42]	fMRI; conflict task	Adults	20 OCD patients: 7m, 5f; handedness (R:L) 19:1; mean onset age: 8.5 years; mean duration of illness: 13.1 years; 9 with obsessions related to numbers or counting compulsions; mean age: 21 years; 15 HC: 11 m, 4 f; handedness (R:L) 14:1; mean age: 25 years	Free of benzodiazepine or neuroleptics; 5 patients receiving sertraline, 3 fluoxetine, 2 paroxetine, 1 fluoxetine and clomipramine	No comorbid axis I or II disorders; no neurological disorder; head injury or serious medical condition	In prime target repetition condition, hyperactivation of anterior cingulate gyrus and left parietal lobe; 'non-resistant' OCD subjects activated a bilateral network including the precuneus and the pulvinar and paracentral lobules
Remijnse et al. [73]	fMRI; reversal learning task	Adults	20 OCD patients: 6 m, 14 f; mean age: 34 years; 27 HC: 8 m, 19 f; mean age: 32 years	Free of psychotropic medication for at least 2 weeks and, in case of fluoxetine or antipsychotic medication, for at least 1 month; no patients involved in a BT program	7 patients with comorbid MDD, 4 with dysthymia, 3 with social phobia, 3 with generalised anxiety disorder, 2 with panic disorder, 1 with agoraphobia, 1 with posttraumatic stress disorder, 2 with comorbid Tourette disorder, 5 patients were diagnosed with 'pure' OCD; no patients with alcohol or substance abuse; no neurological disorder; head injury or serious medical condition	Decreased responsiveness in right medial and lateral OFC, in the right caudate (border zone ventral striatum); during affective switching, hypoactivation of left posterior OFC, bilateral insular cortex, and bilateral dorsolateral and bilateral anterior prefrontal cortex; no results of hyperactivation areas
Rauch et al. [74]	fMRI; serial reaction time task modified for fMRI	Adults	12 OCD patients: 4 m, 8 f; handedness (R:L) 12:0; mean age: 28 years; mean years of education: 15.33; 12 HC: 4 m, 8 f; handedness (R:L) 12:0; mean age: 27 years; mean years of education: 16.67	2 patients receiving sertraline; all other subjects were free of psychotropic medication use at the time of study; no subjects received benzodiazepines within 4 weeks or neuroleptics within 1 year before image acquisition	No comorbid axis I disorder, except for 1 patient with comorbid generalised anxiety disorder; no neurological disorder; head injury or serious medical condition	For the implicit learning vs. random contrast, group-by-condition interactions revealed aberrant recruitment within the hippocampus as well as orbitofrontal cortex, but no striatal group differences; an inverse correlation was found between striatal activation and specific symptom factors

Table 3 (continued)

Study	Technique	Age group	Participants	Medication	Comorbidity	General findings in OCD patients
Roth et al. [79]	fMRI; go/no go task	Adults	12 OCD patients: 5 m, 7 f; mean age: 37.8 years; 14 HC: 6 m, 8 f; mean age: 34.9 years	6 OCD patients treated with SSRI; 6 had not taken any psychotropic medication for at least 6 weeks before scanning; no one in psychotherapy at the time of the study	2 patients with MDD, 1 of whom with current comorbid social phobia; no history of substance use disorder; no neurological disorder, head injury or serious medical condition	During response inhibition, healthy adults showed predominantly right-hemisphere activation including the right inferior frontal gyrus, whereas the patient group showed a more diffuse, bilateral pattern of activation; the OCD group demonstrated less activation than the comparison group in several right-hemisphere regions during response inhibition, including inferior and medial frontal gyri; symptom severity was inversely correlated with activation in right orbitofrontal and anterior cingulate gyri, positively correlated with thalamic and posterior cortical activations
Yücel et al. [82]	fMRI; Multi-Source Interference Task	Adults	19 OCD patients: 10 m, 9 f; current IQ >80; handedness (R/L) 19/0; mean age: 33.7 years; mean duration of illness: 13.4 years; 19 HC: 10 m, 9 f; handedness (R/L) 19/0; mean age: 30.6 years	Medication free or stable on their medication dose for at least 1 month; no corticosteroid use	No comorbid axis I or II disorders; no electroconvulsive therapy; no impaired thyroid function; no neurological disorder, head injury or serious medical condition	Greater relative activation of the supplementary motor area and deactivation of the rostral anterior cingulate during high- vs. low-conflict trials; reduced neuronal N-acetylaspartate levels in the dorsal anterior cingulate region, negatively correlated with their blood oxygen level-dependent activation of the region
Gu et al. [54]	fMRI; task-switching paradigm	Adults	21 OCD patients: 18 m, 3 f; mean age: 23.6 years; 21 HC: 18 m, 3 f; mean age: 24.8 years	8 patients drug naïve, 2 unmedicated, 3 on SSRI, 2 on SSRI + anxiolytic, 1 on SSRI + atypical antipsychotics, 5 on SSRI + anxiolytic + antipsychotic	1 patient with comorbid panic disorder, 1 with obsessive-compulsive personality disorder, 11 patients without comorbid axis I or II disorders; no neurological disorder, head injury or serious medical condition	Healthy controls showed significant activation in various areas, including dorsal frontal-striatal regions, during task switching, whereas OCD patients showed no activation in these areas; significant differences were also observed in the dorsal frontal-striatal regions and ventromedial prefrontal and right orbitofrontal cortices between OCD patients and healthy controls; activations of orbitofrontal cortex were related to the performance in both groups and also to activation of the anterior cingulate cortex in the OCD group
Henseler et al. [92]	fMRI; 3 verbal and spatial item-recognition tasks	Adults	11 OCD patients; mean age: 32.64 years; mean age at onset: 18.91 years; 11 HC; mean age: 33.73 years	8 patients receiving SSRI; 1 receiving TCA; 2 medication free	No substance abuse, acute depression, acute suicidal tendency or diagnosis of psychosis; no neurological disorder, head injury or serious medical condition	Significantly greater task-related activation in several frontal and parietal brain areas known to underlie WM: during verbal and spatial WM tasks, increased activation showed in the left precentral sulcus, left intraparietal cortex, left inferior frontal gyrus, left inferior frontal sulcus, and middle third and right intraparietal cortex
Lázaro et al. [93]	fMRI; performance of simple and complex sequences	Children and adolescents	20 OCD patients: 7 m, 5 f; mean age: 13.1 years; 20 HC; mean age: 13.7 years	Patients were drug naïve; after initial assessment have received SSRI and behavioural counselling; no other pharmacological treatment was used	No comorbid affective disorder, attention deficit hyperactivity disorder, learning and writing disabilities, schizophrenia and concurrent eating disorder; no neurological disorder, head injury or serious medical condition	Both patients and controls showed a pattern of cerebral activation involving the frontoparietal cortex and basal ganglia; OCD patients presented significantly higher brain activation bilaterally in the middle frontal gyrus; after 6 months of pharmacological treatment and with clear clinical improvement, activation in the left insula and left putamen decreased significantly

Study	Technique	Age group	Participants	Medication	Comorbidity	General findings in OCD patients
Nabeyama et al. [41]	fMRI; Stroop Task	Adults	11 OCD patients: 4 m, 7 f; handedness (R:L) 11:0; mean age: 32.4 years; 19 HC: 8 m, 11 f; handedness (R:L) 18:1; mean age: 32.7 years	Drug free for at least 2 weeks before their initial fMRI scan; 2 were drug free for 18 days, 2 for 30 days, 3 for 60 days; 4 had never taken medication and remained drug free throughout the study; the patients received 12 weeks of BT	N/A	Hypoactivation in the anterior cingulate gyrus and cerebellum; following significant improvement in obsessive-compulsive symptoms, the cerebellum and parietal lobe showed increased activation, and the orbitofrontal cortex, middle frontal gyrus and temporal regions showed decreased activation during the Stroop task, and the performance of the task itself improved
Nakao et al. [57]	fMRI; Stroop and N-back tasks	Adults	32 OCD patients; mean age at onset: 14 years; mean duration of illness before a medical examination: 10 years; divided into 2 groups: – short-term group (examined before 10 years of illness): 5 m, 12 f, mean age at onset: 14.5 years, mean age: 30.6 years, mean duration of illness: 5.5 years, mean duration of treatment: 6.3 months; – long-term group (examined after 10 years of illness): 7 m, 8 f, mean age at onset: 14.4 years, mean age: 36.1 years, mean duration of illness: 20.3 years, mean duration of treatment: 34.9 months; 16 HC: 9 m, 7 f; mean age: 30.8 years	Drug free for a 2-week period before study; 8 patients drug naïve; 24 on medications: 16 received SSRI, fluvoxamine or paroxetine, 4 clomipramine, 4 other psychotropics; 9 of these 24 patients had stopped medication 14–28 days before entering the study, 15 had stopped more than 29 days before	No current comorbid depressive episode; no comorbid axis I disorder, or history of drug or alcohol addiction; no neurological disorder, head injury or serious medical condition	The short-term group showed weaker activation of the right caudate during the Stroop task and stronger activation of the right dorsolateral prefrontal cortex during the N-back task than the long-term and normal control groups; the long-term group showed attention deficit and nonverbal memory dysfunction in the neuropsychological tests
Woolley et al. [153]	fMRI; Stop task; motor Stroop task; Switch task	Children and adolescents	10 OCD patients: 10 m; mean age: 14.3 years; mean duration of illness: 37.7 months; 9 HC: 9 m; mean age: 14.5 years	8 patients receiving SSRI; mean duration of treatment: 5 months; 5 patients had completed a CBT (mean: 8 sessions)	No comorbid axis I or II disorders; no neurological disorder, head injury or serious medical condition	During the Stop task, there was reduced activation in the right orbitofrontal cortex, thalamus and basal ganglia; inhibition failure elicited mesial frontal underactivation; task switching and interference inhibition associated with attenuated activation in frontal, temporoparietal and cerebellar regions
Gilbert et al. [151]	fMRI; symptom provocation paradigm	Children and adolescents	18 OCD patients: 11 m, 7 f; mean age: 13.1 years; 18 HC: 11 m, 7 f; mean age: 13.6 years	9 patients were taking psychotropic medications	Exclusion criteria included a history of head injury, bipolar disorder, psychosis, substance abuse/dependence, debilitating medical or neurological conditions, pervasive developmental disorders, mental retardation or learning disorders	Reduced activity in the right insula, putamen, thalamus, dorsolateral prefrontal cortex and left orbitofrontal cortex (contamination experiment), and in the right thalamus and right insula (symmetry experiment); higher scores on OCD symptom-related measures (contamination and total severity) were significantly predictive of reduced neural activity in the right dorsolateral prefrontal cortex during the contamination experiment

Table 3 (continued)

Study	Technique	Age group	Participants	Medication	Comorbidity	General findings in OCD patients
Nakao et al. [56]	fMRI; N-back task using a block design paradigm	Adults	40 OCD patients: 16 m, 24 f; handedness (R:L) 38:2; mean age: 33.3 years; 25 HC: 10 m, 15 f; handedness (R:L) 23:2; mean age: 30.9 years	Drug free for a 2-week period before study; 8 patients drug naïve; 24 on medications: 16 received SSRI, fluoxetine or paroxetine, 4 clomipramine, 4 other psychotropics; 9 of these 24 patients had stopped medication 14–28 days before entering the study, 15 had stopped more than 29 days before	No comorbid axis I or II disorder, or history of drug or alcohol addiction; no neurological disorder; head injury or serious medical condition	Greater activation in the right dorsolateral prefrontal cortex, left superior temporal gyrus, left insula and cuneus during 2-back task; right orbitofrontal cortex activity showed a significant positive correlation with Y-BOCS scores in OCD; patients with obsessions/checking rituals showed severer memory deficits and decreased activity in the postcentral gyrus compared with patients with cleanliness/washing rituals
Page et al. [131]	fMRI; go/no go task; motor Stroop task; Switch task	Adults	10 OCD patients; mean age: 39.1 years; 11 HC; mean age: 34.1 years	No medications	2 patients with OCD had a current comorbid diagnosis of dysthymic disorder, 3 had a past history of major depressive disorder and 1 of past alcohol dependence	During the go/no go and Switch experiments, people with OCD had underactivation in task-relevant orbitofrontal/dorsolateral prefrontal, striatal and thalamic regions; during the motor Stroop and Switch tasks, people with OCD also displayed underactivation in temporoparietal areas; in the go/no go and motor Stroop tasks, the OCD group showed increased activation compared with controls in the cerebellum and predominantly posterior brain regions
Remijnse et al. [76]	fMRI; self-paced reversal learning task	Adults	20 OCD patients: 5 m, 15 f; handedness (R:L) 17:3; mean age: 34 years; 20 MDD patients: 12 m, 8 f; handedness (R:L) 16:4; 8 OCD patients with prior MDD; mean length of time since remission of MDD: 36 months; 27 HC: 8 m, 19 f; handedness (R:L) 23:4	Drug free for at least 2 weeks before study and, in case of fluoxetine or antipsychotic medication, for at least 1 month; 8 MDD patients and 7 OCD patients were drug naïve; mean length of drug-free interval: 16 months in the MDD sample, 30 months in the OCD group	No history of drug or alcohol addiction; – MDD group: 3 patients with comorbid social anxiety disorder, 1 with generalised anxiety disorder, 1 with panic disorder without agoraphobia, 1 with pain disorder, 1 with cannabis abuse in early adulthood, 1 with sustained full remission, 12 of them were free from comorbidity, 6 patients had their first lifetime depressive episode; – OCD group: 9 patients were diagnosed with ‘pure’ OCD, the following disorders were comorbid: 1 posttraumatic stress disorder, 2 panic disorder, 4 generalised anxiety disorder, 4 dysthymic disorder, 4 social anxiety disorder, 1 opioid abuse in sustained full remission, 1 Tourette disorder; no neurological disorder; head injury or serious medical condition	Both MDD and OCD patients displayed prolonged mean reaction times, but normal accuracy; the OCD group showed blunted responsiveness of the orbitofrontal-striatal loop during reward, and in the OFC and anterior insula during affective switching

Study	Technique	Age group	Participants	Medication	Comorbidity	General findings in OCD patients
Rubia et al. [152]	fMRI; Stop task; Switch task	Adolescents	18 ADHD patients; mean age: 13.9 years; 10 OCD patients; mean age: 14.3 years; 20 HC; mean age: 14.5 years	All patients with ADHD were medication naïve; patients with OCD were treated and in partial remission	Exclusion criteria for both patient groups were drug and substance abuse and a history of a general or specific learning disability or comorbidity with any other major psychiatric disorder; the exception was comorbidity with OCD for the ADHD group, which was present in 1 patient	Both patient groups shared brain dysfunction compared with healthy controls in the right orbitofrontal (successful inhibition) and left dorsolateral prefrontal cortices (failed inhibition); right inferior prefrontal dysfunction, however, was disorder specific to ADHD during both tasks; left inferior prefrontal dysfunction during the Switch task was significant in children with ADHD relative to controls, but only reached a trend in patients with OCD; patients with ADHD furthermore showed disorder-specific underactivation in a cluster comprising the caudate, putamen, and anterior and posterior cingulate gyri during the Switch task
Sanematsu et al. [132]	fMRI; symptom provocation task	Adults	15 OCD patients: 7 m, 8 f; mean age: 33.6 years	9 patients had been on psychotropic medication; 8 patients were drug free	All patients with axis I disorders, neurological disorder, head injury, serious medical condition, or history of drug/alcohol addiction were excluded	Patients showed brain activation in the left superior temporal gyrus (BA 39), left precuneus (BA 7), left frontal cortices (BA 10, 47), right cerebellum and right frontal cortices (BA 47); no region showed a significant correlation between brain activation and OCD symptoms
Ursu and Carter [176]	fMRI BOLD; conflict-generating task	Adults	15 OCD patients: 7 m, 8 f; handedness (R:L) 13:2; mean age: 32.06 years; mean years of education: 15.8; 15 HC: 8 m, 7 f; handedness (R:L) 13:2; mean age: 30.85 years; mean years of education: 16.56	13 patients medicated	None reported; no exclusion criteria reported	Hyperactivation in the right lateral orbitofrontal cortex in OCD patients relative to controls during conflict-laden tasks; anxiety but not OCD symptoms correlated with OFC activation; relative hyperactivity has also been found in the left lateral prefrontal cortex and bilaterally in the superior temporal gyrus
Jung et al. [94]	fMRI; biological and scrambled motion; stimuli defined by point lights	Adults	15 OCD patients: 12 m, 3 f; mean age: 23.4 years; mean age at onset: 17.33 years; mean duration of illness: 6.13 years	7 patients drug naïve; 8 receiving medications: 2 MAOI, 3 SSRI with antianxiety medication, 3 SSRI + anti-anxiety medication + antipsychotics	No comorbid axis I disorder; 1 patient with comorbid panic disorder, 1 with obsessive-compulsive personality disorder, 1 with schizotypal personality disorder; the remaining 12 patients had not been diagnosed with any comorbid axis I or II disorders; no neurological disorder, head injury or serious medical condition	Increased activation in the right superior and middle temporal gyri as well as in the left inferior temporal and fusiform gyri; reduced activation in the right postcentral gyrus (BA 40); increased activation in the ventral visual system including the inferior temporal and fusiform gyri

ACC = Anterior cingulate cortex; ADHD = attention deficit hyperactivity disorder; BA = Brodmann area; BOLD = blood oxygen level dependency; BT = behavioural therapy; CBT = cognitive-behavioural therapy; HC = healthy controls; IQ = intelligence quotient; LPFC = lateral prefrontal cortex; MAOI = monoamine oxidase inhibitor; MDD = major depressive disorder; OFC = orbitofrontal cortex; PCC = posterior cingulate cortex; SSRI = selective serotonin reuptake inhibitor; TCA = tricyclic antidepressant; WM = working memory; Y-BOCS = Yale-Brown Obsessive Compulsive Scale.

When suspecting population overlap, for example in patients and controls whose details appeared in more than one study by the same research group, we kept the results of the most recent study and with the higher sample size, provided that the same methodology was used. In case the older or smaller-sized studies contained more data than the more recent ones, we kept the older papers.

Results

Cingulate Cortex Dysfunction

Structural MRI had shown the involvement of the cingulate cortex in OCD in the context of a more general corticolimbic-striatal circuitry alteration [33]. Functional studies were in the same direction; MRS showed the existence of neuronal loss or impairment in the anterior cingulate cortex [34]. A SPECT study found increased cingulate perfusion ratios relative to the whole brain [18], whereas PET studies found cingulate hypermetabolism in patients with OCD [17, 35].

The hyperactivity of the anterior cingulate cortex that has been identified by functional studies has been interpreted as a non-specific index of heightened alertness and anxiety [17, 18] as the anterior cingulate cortex is believed to signal to the prefrontal decisional ('executive') cortex anticipatory mismatching between a desired and an actual state and calls for implementation of effective strategies. Hyperactivity in the anterior cingulate cortex is aimed at preventing errors; the anterior cingulate cortex is active during cognitive verification processes that control that everything is going all right, but in OCD patients this activity is not set off when the desired result has been achieved. Despite the fact that the performance of OCD patients is unimpaired in many cognitive tasks [36], anterior cingulate hyperactivity ensues in signalling a conflict when there is none, thus explaining some symptoms like constant doubt and need for repetition although one recognises the correctness of one's own performance. Hyperactivation in the anterior cingulate cortex of OCD patients relative to healthy controls was found in an fMRI study while subjects were tested in a conflict-generating, continuous performance test graded for degree of conflict, but only under the higher-conflict conditions, and this was independent from task-related anxiety [37]. Overall, these findings are consistent with hyperactivity and inability to dampen activity after success of the conflict-monitoring system in OCD.

Fitzgerald et al. [38] subjected OCD patients and healthy controls to fMRI during a flanker interference test that was unable to trigger active OCD symptoms and

found greater hyperactivation of the rostral anterior cingulate cortex during conflict in the patient group, which correlated with background OCD symptom severity in the patients. It appears that a basic alteration in conflict-processing stimulus processing in the anterior cingulate cortex is present in OCD patients even when their symptoms are not manifest.

The role of the cingulate cortex in OCD has been explored by Maltby et al. [39] using a 'go/no go' inhibition task. In this task, the subject is required to abstain from responding to a certain cue while he/she has to respond as fast as possible to another cue that is presented at a much higher frequency, thus constituting a sort of habit that is learned (a prepotent response). The no go cue is a highly conflicting condition. The subject may commit an error either by responding to the no go cue (commission error) or by not responding to the go cue (correct reject). The result was that OCD patients responded with hyperactivation, relative to controls, of both rostral and caudal portions of the anterior cingulate cortex only to no go errors, i.e. when they were required to inhibit a prepotent learned action, lending further support to the hypothesis of hyperactivity of action-monitoring brain areas in OCD. The caudal anterior cingulate cortex is believed to be involved in the recognition of the conflicting nature of a stimulus, whereas the rostral anterior cingulate portion would be linked to the affective response to a conflict. Based on these data, Maltby et al. [39] concluded that in OCD there is a relative inability to inhibit a learned response, and an exaggerated response to it, rather than the incapacity to complete an initiated action.

However, using a different inhibition task, the Stroop colour word naming test, which involves the inhibition of a spontaneous prepotent learned response, both Nakao et al. [40] and Nabeyama et al. [41] found decreased anterior cingulate activation in OCD patients relative to controls during task performance.

Viard et al. [42], using event-related fMRI, compared regional brain activity of healthy controls with adolescents and young adults with childhood-onset OCD. They investigated conflict task performance, involving the presentation of two consecutive and possibly conflicting prime and target numbers. The image dataset of the patients was further analysed according to whether they resisted or yielded to symptoms during the scans. They identified a subregion of the anterior cingulate gyrus where OCD patients showed more activation than healthy controls. Symptom-resisting and symptom-surrendering OCD patients did not differ in activation of the anterior cingulate cortex, but rather bilaterally activated the pre-

cuneus-pulvinar-paracentral lobule circuitry. It appears from this study that OCD patients abnormally amplify the brain circuitry involved in repetitive visual stimulus discrimination, and that these activations may depend on a patient's ability to resist obsessions [42].

Adler et al. [43] subjected patients with OCD to patient-targeted innocuous and OCD symptom-provoking stimuli (like supposedly infected material to compulsive washers) and found higher activation of the right anterior cingulate cortex under the symptom-provoking condition.

These fMRI findings match those of several previous structural, neuropsychological and functional studies suggesting that dysfunctions of the anterior cingulate cortex [17, 29, 44, 45] play a role in OCD symptomatology. However, the direction and extent of activation depends upon the function explored and the task employed.

Dysfunctions of the Prefrontal Cortex: Ventrolateral Prefrontal Cortex and Dorsolateral Prefrontal Cortex

The functions of the lateral prefrontal cortex (LPFC) were studied according to stimulus type (verbal vs. object or spatial) [46, 47] and process type (maintenance or manipulation) [48, 49]. The ventrolateral prefrontal cortex (VLPFC), which includes Broca's area (Brodmann areas, BA, 44, 45 and 47), supports processes that convey, sustain and match information during working memory tasks. The dorsolateral prefrontal cortex (DLPFC), which includes BA 9 and 46, is involved in the processing of spatial and non-spatial information maintained in working memory, i.e. monitoring, manipulation and higher-level planning. A third executive control process, after maintenance in the VLPFC and manipulation in the DLPFC, is run by the anterior prefrontal cortex (frontopolar) including BA 8 and 10 [50].

Models of LPFC interactions with the anterior cingulate cortex may suggest that the LPFC serves to reduce conflict by filtering responses unrelated to task demands, although further research is needed to evaluate the relationship between the LPFC and the anterior cingulate cortex in action monitoring [39]. An increase in VLPFC activity may be a sign of working memory loading (the number of items kept available) and of recovery of abstract rules used for problem solving [51]. Moreover, VLPFC activity is selectively increased during arithmetic computations, mainly when people attempt to resolve a conflict between externally presented answers and internally computed solutions [52]. This network may explain the increased bilateral activity of the VLPFC associated with task load in OCD patients compared with control

subjects [39, 53]. In OCD patients, fMRI under the reversed-contrast condition (such as task repeat minus task switch) showed significant differences in ventromedial prefrontal cortex activities compared with healthy subjects [54].

In an fMRI study, van der Wee et al. [55] found decreased performance only at the highest task level in OCD patients compared with controls during performance of a spatial N-back task. Functional imaging showed similar bilateral DLPFC and parietal cortex activities in both groups, from which the authors concluded that spatial working memory in OCD was not abnormal.

On the other hand, several fMRI studies using the N-back task [56], Stroop and N-back tasks [57] or an off/on exposure paradigm [43] showed greater activation of the DLPFC in OCD patients than in control subjects. It seems that DLPFC activation is greater in recent-onset OCD patients compared with patients with chronic, stabilised disease. In general, OCD patients seem to have some kind of disorder of spatial cognition, attention and non-verbal memory.

In OCD patients, Nakao et al. [56, 57] found greater activation in the right DLPFC, left superior temporal gyrus, left insula and cuneus than in control subjects. Furthermore, the patients with cleaning/washing compulsions showed greater activation in the right thalamus and left postcentral gyrus during the N-back task. According to these data, there might be a close relationship between checking rituals and memory-related neuropsychological dysfunction [56, 57].

For several investigators, different brain networks might be involved in checking and washing rituals. Symptom severity and symptom subtypes such as obsessions/checking might affect neuropsychological function and related brain activities.

Levine et al. [58] studied the neurobiological differences between patients with schizophrenia and high or low levels of obsessive-compulsive symptomatology; they did not find significant differences in signal activation of the left DLPFC. However, they found a subgroup of patients with schizophrenia in whom MRI activation of the left DLPFC during a word fluency challenge task was significantly associated with severity of OCD symptomatology. In this subgroup, as OCD symptomatology increased, the activation of the DLPFC decreased.

Dysfunctions of the Orbitofrontal Cortex

The network involving orbitofrontal cortex (OFC) and striatal structures is important for motivational behaviour, and may be involved in the pathophysiology of OCD

[45, 59]. Several functional imaging studies [26, 28, 29, 43, 60], using symptom provocation designs, have reported orbitofrontal-striatal functional abnormalities in OCD, hypothesised to reflect its role in ritualistic behaviour.

Selective serotonin reuptake inhibitors and dopamine antagonists showed some efficacy in OCD [30, 61], and intact transmissions of serotonin (5-hydroxytryptamine) and dopamine have been associated with normal OFC functioning [62] and reward processing in the ventral striatum [63], respectively.

Reward and punishment perception appears to be abnormal in OCD. When experiencing obsessions, patients are always conscious that they are committing a kind of error. Moreover, they give the impression of feeling insufficiently relieved by compulsive behaviour that serves a rewarding goal [59].

A number of neuropsychological tasks addressing particular OFC functions have shown impaired performance in patients with OCD compared with healthy controls [64–67]. Structural neuroimaging studies of OCD showed inconsistent OFC abnormalities in OCD patients compared with healthy controls. OFC volumes were reported to be increased [68] or decreased [69]. Inconsistency was also confirmed by functional imaging studies showing increased [16, 70] or decreased [71] OFC activity in OCD patients compared with healthy volunteers. Symptom provocation studies showed increased OFC activity [60] along with both increased [29, 60] and decreased [72] caudate activity.

fMRI studies showed decreased responsiveness in the right medial and lateral OFC during a reversal learning task [73], exaggerated OFC activation during implicit sequence learning [74] and after provocative stimuli [43], and asymmetrical patterns of activation in lateral OFC during high-conflict trials [39]. In the first of these studies, the reduction in the number of correct responses relative to healthy controls in OCD patients was paralleled by adequate behaviour on receiving punishment and with regard to affective switching [73]. On reward outcome, patients showed decreased responsiveness in the right medial and lateral OFC as well as in the right caudate nucleus (border area of the ventral striatum) when compared with controls. During affective switching, OCD patients recruited the left posterior OFC, bilateral insular cortex, and bilateral dorsolateral and bilateral anterior prefrontal cortex to a lesser extent than controls [73]. The inverse correlation existing between OCD symptoms and extent of activation in the left OFC appears to suggest that orbitofrontal structures may be active in inhibiting symptom pro-

vocation, which is consistent with previous findings [29, 43].

During high-conflict trials, fMRI showed in some studies that activation patterns in the lateral OFC were asymmetrical, favouring the left side during both error and correctly rejected, high-conflict trials [39]. Other studies reported that OFC activation was generally bilateral in OCD, but might tend towards increased activation on the left side [75]. It was advanced that the asymmetries observed represented excessive tone in contralateral OFC-caudate projections [39]. However, animal studies support ipsilateral preponderance of OFC-caudate projections [75].

Remijnse et al. [76] analysed 20 unmedicated OCD-free patients with major depressive disorder (MDD), 20 unmedicated MDD-free patients with OCD, and 27 healthy controls. Patients underwent a self-paced reversal learning task according to an event-related design during fMRI. OCD patients showed blunted response of the OFC-striatal loop during reward, and in the OFC and anterior insula during affective switching. This is consistent with differential neural patterns in MDD and OCD in frontal-striatal and paralimbic structures in this task. These findings are consistent with those of several previous structural, neuropsychological and functional studies suggesting that dysfunctions of the OFC [25, 26, 29, 44, 45, 64, 77] play a role in OCD symptomatology.

Dysfunctions of the Frontal, Parietal and Temporal Cortices

Decreased baseline cerebral blood flow in the superior frontal cortex [60, 78] may be associated, in OCD patients, with activation after exposure to provocative stimuli [29, 43]. Activation of the right hemisphere, mostly of the right inferior frontal gyrus, is physiological during response inhibition. In OCD patients this activation is of poorer quality both in the inferior and medial frontal gyri [79]. Adaptive and inhibitory control of behaviour is critically regulated by the MFC, which includes the dorsal anterior cingulate cortex and the supplementary motor area (SMA), which is just about 25 mm posterior to the dorsal anterior cingulate cortex. Its activity might reflect the processing of response to conflict, as suggested by studies using both flanker-type interference tasks [80] and response inhibition tasks [81]. Abnormally high MFC activity has been a consistent finding in functional neuroimaging studies of OCD. However, the precise regions and the neural alterations associated with this abnormality remain unclear.

Yücel et al. [82] conducted a cross-sectional study combining volume localised proton MRS and fMRI with a task encompassing inhibitory control processes (the Multi-Source Interference Task), designed to activate the MFC. In this study, compared with controls, OCD patients had a greater relative activation of the SMA and deactivation of the rostral anterior cingulate cortex during high- versus low-conflict (incongruent > congruent) trials. Patients with OCD also showed reduced levels of neuronal *N*-acetylaspartate in the dorsal anterior cingulate region, which correlated inversely with their blood oxygen level-dependent activation of the MFC.

Hyperactivation of the MFC during high/low-conflict conditions in patients with OCD may be a compensatory reaction to neuronal deficiency in the region. This association may somewhat clarify the nature of inhibitory control deficits that are commonly seen in OCD [82]. Fitzgerald et al. [38] found conflict-related SMA/pre-SMA activation in controls, but not in OCD patients. As this greater activation in controls was not predicted, the finding is difficult to interpret.

The frontal cortex, especially Broca's language area and the surrounding cortex, is physiologically activated during word generation tasks, mostly when those tasks are phonologically guided (i.e. when subjects are required to generate words beginning with a designated letter) [83–85]. Several studies have confirmed the utility of fMRI in studying frontal activation during this type of verbal fluency in normal individuals [85–89]. In OCD patients, previous neuropsychological studies reported verbal fluency ranging from normal to only mildly altered [90, 91].

People showing activation in frontal and parietal brain areas may suffer working memory deficits. Henseler et al. [92] scanned 11 patients and 11 matched controls while they performed three verbal and spatial item-recognition tasks. OCD patients exhibited significantly greater task-related activation in several frontal and parietal brain areas involved in working memory. Working memory in OCD patients may be weakened by interference from regions supporting OCD-related processing. On the other hand, working memory disturbances may contribute to the expression of the typical behaviours of OCD patients [92]. It is interesting that hyperactivity of the superior right parietal lobe also correlates with severity of disorder [93].

Some findings may indicate that OCD patients have functional differences related to the perception of biological motion. In OCD patients, alterations in visual association areas were found after exposure to provocative

stimuli [43]; abnormal activation was found in temporal areas and in the right postcentral gyrus [94]. Jung et al. [94] studied 15 patients with OCD and 15 age- and IQ-matched healthy volunteers. All subjects participated in a biological motion task in which they performed a 1-back conflict task. Patients with OCD showed increased activation in the right superior and middle temporal gyri and in the left inferior temporal and fusiform gyri, as well as reduced activation in the right postcentral gyrus (BA 40), compared with healthy controls. They also exhibited increased activation in the ventral visual system including the inferior temporal and fusiform gyri.

Furthermore, during fMRI while performing a planning task, OCD patients showed increased posterior temporal and parietal cortical activity relative to controls, explained as compensatory mechanisms [53]. Exploratory analysis of individual fMRI scans employing an 'off/on' exposure paradigm showed activation within the lateral temporal cortex and the left medial temporal aspect of the hippocampus and insula in over 50% of subjects [43]. Aberrant hippocampal recruitment during implicit sequence learning in OCD patients versus healthy controls was reported in an fMRI study using the serial reaction time (SRT) task [74]. The insula is a paralimbic region that plays a role in mediating emotional states including anxiety and disgust. It is possible that clinical improvement after treatment, accompanied by decreased anxiety, leads to decreased activation in this area [93].

Taken together, these findings are consistent with those of a number of earlier structural, neuropsychological and functional studies suggesting that dysfunction of the temporal cortex [78, 95], in particular of the anterior temporal cortex [96, 97] and of superior regions of the temporal cortex [26], is involved in OCD symptomatology.

Basal Ganglia

Dopamine-serotonin interplay is important in both the OFC and basal ganglia. Serotonin regulates executive functions in the OFC, like inhibition and flexibility, by preventing possible interfering salient stimuli from being responded to during tasks targeted at other goals [98], thus allowing dopamine-mediated, rewarded behaviour to proceed. Dopaminergic activity in the ventral striatum controls reward processing [63, 73, 99], whereas striatal serotonergic impairments or OFC lesions cause compulsive behaviour that is inhibited by increasing intrasynaptic serotonin content [100]. Recent SPECT studies reported abnormal dopamine transporter density and irregular D₂ receptor binding in the basal ganglia in OCD [101, 102].

Basal ganglion disinhibition due to an altered balance between indirect, inhibitory and direct excitatory corticostriatalthalamocortical circuits has been advanced to explain OCD symptoms [45, 103]. There is evidence that the basal ganglia are critical for implicit sequence learning [44]. Patients with Huntington's or Parkinson's disease show poor performance in the SRT task [104], which provides a measure of implicit sequence learning.

Striatum

Some morphometric studies reported enlarged [69], normal [105] or diminished [106] striatal volumes in OCD patients, whereas functional imaging studies during the resting state showed increased [16, 70] or decreased [71] activity in the OFC, and decreased [107] or increased [16, 108] activity in the caudate nucleus. PET studies, using an implicit learning task, showed striatal dysfunction in OCD [109, 110].

The caudate nucleus projects to the OFC, the anterior cingulate cortex and the thalamus. It has been proposed to play a central role in the corticolimbic-basal ganglia-thalamus network; network dysfunction may be associated with OCD symptoms [45]. During Stroop test performance, OCD patients showed weaker activation than healthy controls both in the anterior cingulate cortex and in the right caudate nucleus [111] in the absence of neuropsychological impairment. In a subgroup of patients with recent-onset OCD, the same group of researchers found caudate nucleus hypoactivity to be related to attention, and right DLPFC hyperactivity to be related to delayed visual recall [56, 57]. Higher caudate nucleus activation in OCD patients during conflict processing has been reported in one study [38]. Caudate nucleus hyperactivity in OCD [15, 16, 112] could be related to excitatory projections from the pre-SMA/SMA region.

Patients with OCD fail to recruit brain systems that are typically responsible for unconscious information processing [44]. This hyperactivation involves the frontal and temporal cortices as well as the cingulate cortex, insula, caudate and lenticular nuclei and amygdala [44], supporting that the neurophysiology of OCD involves hyperactivation of a neural feedback circuit that includes cortical, paralimbic, limbic and striatal structures [45, 79, 113].

Adults with OCD show frontostriatalthalamocortical circuitry hypoactivation during response inhibition. The thalamus and related circuitry may play a role in the expression or intensity of OCD symptoms, whereas right frontal subregions may be involved in symptom suppression [79]. In an fMRI study, patients with recent OCD

onset showed weaker activation of the right caudate nucleus than healthy controls during Stroop task performance [57].

Both children and adolescent OCD patients, studied by fMRI during the performance of simple and complex sequences and compared with healthy controls, showed significant bilateral hyperactivation in the middle frontal gyrus. After 6 months of pharmacological treatment and with clear clinical improvement, activation in the left insula and left putamen decreased significantly [93]. Other studies reported a decreased responsiveness of dorsal prefrontal-striatal circuits during planning in OCD patients compared with healthy controls, reflecting basal ganglion dysfunction in OCD in implicit learning [109, 114].

Some investigators reported delayed responding [115, 116] or reduced performance [53, 117] in planning tasks, whereas others found no differences in planning between OCD patients and healthy controls [118, 119]. Inconsistencies may reflect methodological differences in task implementation, such as mental performance versus the use of a touch screen, or providing feedback versus no feedback [115, 120]. During planning tasks, fMRI showed that OCD patients failed to recruit dorsal prefrontal-striatal regions as compared with controls [53].

Amygdala

Among other functions, the amygdala is involved in visual recognition of fearful expressions, whereas the fusiform gyrus is implicated in recognition of facial identity. Use of the fearful versus neutral expression contrast during fMRI allows studying the physiological function of the amygdala. Recent PET findings in normal people suggest that the amygdala has a role in emotional memory [121].

Alternative conceptualisations, however, emphasise the role of the amygdala in rapid threat assessment. The activation of the amygdala in symptom provocation studies may reflect a quantitative versus qualitative distinction related to the magnitude of the threat perceived. It seems that paralimbic-limbic activation, particularly concerning the amygdala, is much more prominent in fMRI studies than in PET investigations of OCD [44].

Cerebellum

The cerebellum is crucial for movement coordination and motor learning. Recent anatomical and functional studies support its involvement in a variety of cognitive functions such as attention, verbal learning and memory, and cognitive planning [122–125].

Nonhuman primates possess pontine projections to the cerebellum from association areas in the prefrontal cortex [126], posterior parietal region [127], parahippocampal region [128] and cingulate gyrus [129]. The existence of a 'cerebellar cognitive affective syndrome' was postulated by Schmahmann and Sherman [130]. This syndrome shows deficits of executive function, visual spatial ability and memory. Such cognitive impairments in patients with cerebellar lesions may be due to alterations to the network linking prefrontal, temporal, posterior parietal and limbic cortices with the cerebellum. Apparently, regions believed to be crucial for interference processes are connected anatomically with the cerebellum. Specifically, fMRI activation was found in OCD in cerebellar regions during a 'focused-attention' task, independent of motor activation, and during a Stroop task [131]. Like the Stroop task, this task requires cognitive control to ignore irrelevant information. Cerebellar hyperactivity [40, 132] or hypoactivity [41] was found in OCD patients compared with a group of healthy controls. After symptom improvement, increased cerebellar activation was found [41], even when preceded by hypoactivity [111].

Few studies have described the relationships between obsessive-compulsive symptom improvement, neuropsychological performance and alterations in a wide range of brain activities involving the cerebellum. Metabolic changes in the putamen, cerebellum and hippocampus significantly correlated with symptom improvement [133]. Symptom improvement after behaviour therapy (BT) seems to be significantly associated with increased activation of the cerebellum and parietal lobe, and decreased activation of the OFC, middle frontal gyrus and temporal regions during fMRI while performing the Stroop task. The performance of the task itself improves after BT [41]. Briefly, dysfunctions of the posterior brain regions, especially the cerebellum, are involved in the pathogenesis of OCD; these functions may recover with obsessive-compulsive symptom improvement.

Discussion

Corticostriatal Pathophysiological Model

According to a model focusing on the corticostriato-thalamocortical circuitry [134], primary OCD alteration lies within the striatum (specifically, the caudate nucleus). This leads to inefficient gating at the level of the thalamus, which results in hyperactivity within the OFC (associated with intrusive thoughts) and hyperactivity

within the anterior cingulate cortex (corresponding to non-specific anxiety). Compulsions are viewed as ritualistic behaviours that are performed to recruit the inefficient striatum so as to finally achieve thalamic gating, and consequently neutralise unwanted thoughts and anxiety.

PET and SPECT studies in OCD patients compared with healthy controls consistently found increased activity within the orbitofrontal and anterior cingulate cortices [17, 18, 23, 135], and showed differences in caudate regional activity [16, 18]. Successful treatment of OCD patients with drugs, BT or their combination is associated with attenuation of abnormal regional brain activity within the OFC, anterior cingulate cortex and caudate nucleus [17, 35, 136, 137]. Some evidence suggests that lower pretreatment activity in the OFC predicts better response to serotonergic reuptake inhibitors [23, 28, 138, 139]. Imaging studies showed that regional cerebral blood flow and glucose metabolic rates within the posterior cingulate cortex positively correlate with subsequent response to treatment with fluvoxamine [138] or cingulotomy [140], respectively. Symptom provocation studies with PET [29] or with fMRI [114] have also shown increased brain activity within the anterior/lateral OFC and anterior cingulate cortex in OCD. Cognitive activation studies using PET and fMRI that probed the functional integrity of the striatum in OCD found that OCD patients failed to sufficiently recruit the striatum, which is physiologically activated during completion of the SRT task [110, 114]. Patients with OCD may utilise fronto-temporal systems as well as explicit information processing strategies to compensate for frontal-striatal dysfunction and corresponding implicit information processing deficits, thereby potentially contributing to the phenomenon of intrusive thoughts [140]. Aberrant hippocampal activation has recently been observed in the absence of deficient striatal recruitment in OCD [74], suggesting the possibility that aberrant hippocampal function plays a primary role in OCD.

The heterogeneity of OCD prompts a decomposition of the OCD phenotype, so to respect the distinct contributions of various symptom dimensions [74, 141, 142]. Hence, general models of OCD are not likely to apply to all cases. Data gathered on the various symptom dimensions or subtypes of OCD call for multifaceted models that reflect these various aspects of the disorder. A four-factor model of OCD symptom dimensions [143] represents a refinement of the model first presented by Baer [144]. The model by Leckman et al. [143] has now been essentially replicated across an impressive array of subse-

Fig. 1. Simplified drawing of corticostriatothalamocortical circuitry involved in obsessive-compulsive symptoms (continuous lines), with a hypothetical role for the cerebellum (dashed lines).

quent studies [12, 117, 145]. This model envisages four dimensions/subtypes, i.e. obsession and checking, symmetry and ordering, cleanliness and washing, and hoarding. Using this four-factor model, investigators have found that symptom dimensions are relatively stable over time [146] and that they are associated with differential response rates to treatment [117, 146], possibly reflecting different genetic underpinnings [147–149]. Neuroimaging data suggests that distinct profiles of regional brain function may be associated with each of the various symptom factors [74, 108, 150]. Corticostriatal network dysfunction was also reported in children and adolescents affected by OCD [151, 152]. Woolley et al. [153] studied a group of adolescents by fMRI during the Stop task, an inhibition task similar to a go/no go task, showing in OCD adolescents reduced activation with respect to matched controls in brain areas connected via the inferior and orbital frontostriatothalamocortical pathways. In the OCD group, stop failures were associated with reduced activation in the mesial and dorsolateral prefrontal cortex, including the anterior cingulate gyrus. During the more cognitive inhibition tasks, the OCD group showed reduced activation in inferior frontal (Switch task) and temporoparietocerebellar regions (motor Stroop and Switch tasks). The findings confirm frontostriatal network abnormalities during motor response inhibition, but in addition show that tasks requiring more cognitive

forms of inhibitory control (such as selective and flexible attention) were associated with abnormal functioning in extrafrontal temporal, parietal and cerebellar brain regions.

These findings of frontostriatothalamocortical hypoactivation during motor inhibition support that a dysregulation of orbitofrontostriatal pathways mediating inhibitory control functions underlies alterations in OCD [153]. They are also in line with reported structural and biochemical abnormalities in paediatric OCD in the frontal lobes, basal ganglia and thalamus [154, 155], which have been shown to correlate with severity of OCD symptoms [156]. The complexity of the circuitry involved in OCD (fig. 1) makes heterogeneity possible in that a deficit may be present at any point of the pathway and may trigger dysfunction in the other areas connected.

An issue for future research is whether the dorsal prefrontal-striatal dysfunction is specific to OCD or extends to other neuropsychiatric disorders such as anxiety disorders, basal ganglia disorders, Tourette syndrome and MDD. The latter is characterised by impairment of various executive functions such as verbal fluency and attentional set shifting [157].

However, in MDD, dorsal prefrontal baseline perfusion is decreased rather than increased [158]. In addition, MDD is not associated with striatal alterations [159]. Whereas depressive symptoms frequently cooccur in

OCD, cognitive deficits in OCD are not associated with comorbid depression [115, 116], suggesting different pathophysiological mechanisms in OCD with respect to MDD [53].

Cognitive Impairments in OCD

Neuropsychological studies [115, 116, 160] showed cognitive impairments in OCD, mainly regarding visuospatial processing, executive functioning and motor speed. Other cognitive domains appear to remain intact, demonstrating a specific, rather than a general, cognitive deficit.

Executive functioning implies different subdomains of higher-order cognitive functioning. Planning, intended as the ability to achieve a goal via a series of intermediate steps, is an essential component of higher-order cognitive processing such as problem solving. By a neuronal network model, Dehaene and Changeux [161] proposed several hierarchical levels coding for specialised subprocesses of planning such as plan generation, working memory, and internal evaluation and reward. Some subprocesses seem to be relatively independent of task load, whereas other subprocesses are mainly involved at higher levels of planning behaviour.

A test frequently used to probe planning processes is the Tower of London task, adapted from the Tower of Hanoi task [162, 163]. This task has been used in several studies to investigate planning in healthy controls, using PET [164–167], SPECT [168, 169] and fMRI [170–174]. The results of these imaging studies confirm the involvement of the DLPFC and parietal-occipital regions during planning.

No differences were found in accuracy between OCD patients and controls [118, 119]. However, when OCD patients provided erroneous responses, they were spending more time than controls in generating alternative solutions or in checking the next response [119]. The results of two studies by Purcell et al. [115, 116] – who compared the neuropsychological profiles of patients with OCD, panic disorder and MDD with healthy controls, finding specific deficits of executive functions only in OCD patients, namely spatial working memory, spatial recognition and motor initiation and execution – highlight the importance of task implementation.

Mataix-Cols et al. [117] found impaired performance relative to controls also in subclinical obsessive-compulsive individuals; performance impairment correlated with symptom severity, particularly regarding checking behaviour. Although motor speed was decreased, OCD patients showed a normal ability to manage and execute

a sequence of goal-directed moves in a planning task when using a touch screen which provided external validation of ongoing performance. In contrast, when the task had to be performed mentally, OCD patients were significantly impaired [53]. It is still not clear whether impaired executive functioning is a specific OCD trait or whether it is secondary to mood-related or anxiety-related symptoms, but the fact that disorders with such symptoms do not show such impairments [115, 116] prompts us to consider the first hypothesis as more likely.

Conclusions

Several studies have been targeting brain regions hypothesised to be involved in the pathogenesis of OCD, showing the existence of dysfunctional connectivity in the corticostriatalthalamocortical circuitry. However, when people with OCD are tested by cognitive tasks exploring domains other than executive functions such as inhibition or flexibility, alterations also in memory function and in structures associated with it come to the fore. Hence, future studies should focus on other brain regions as well to determine their involvement in OCD. Other major foci should comprise the disentangling of OCD from other frequently occurring comorbid conditions like MDD and attention deficit hyperactivity disorder, and the establishment of similarities and differences, as well as the subtyping of OCD according to neuroimaging data, which would help in classifying and treating the disorder better.

Improvements in spatial resolution of neuroimaging techniques may contribute to a better understanding of the neurocircuitry of OCD and other anxiety disorders. However, currently available data allow to conclude that there is an overlap between the pathophysiology of other anxiety disorders and OCD. Hence, on the grounds of the suggestions coming from functional neuroimaging, it would be premature to separate OCD from other anxiety disorders in the DSM-5 and create a new spectrum.

Focusing upon separate symptoms of OCD is not appropriate when investigating alterations in OCD as these symptoms may also be present in other mental disorders, being supported by different biological mechanisms and bearing different psychopathological meanings. Individual symptoms in OCD patients may reveal something about functioning of the OCD while such symptoms are experienced, but they are unable to provide clues to general patterns of OCD brain functioning. Our analysis – differently from meta-analyses that by their nature have

to focus on methodologically similar studies, thus limiting the number of includible studies – shows that patients with OCD have some consistent hyper- or hypoactivations underlying the true nature of brain circuitry alterations in OCD, whereas inconsistencies are more likely to reflect heterogeneity within the context of different OCD subtypes that should be addressed by future studies focusing on clinically homogeneous populations according to Leckman's subtypes [143].

Acknowledgements

We acknowledge the aid provided by the librarians of the School of Medicine and Psychology of Sapienza University, Ms. Mimma Ariano, Ms. Tiziana Mattei and Ms. Felicia Proietti for the localisation of relevant literature.

References

- Rasmussen SA, Eisen JL: The epidemiology and differential diagnosis of obsessive compulsive disorder. *J Clin Psychiatry* 1994; 55(suppl):5–10, discussion 11–14.
- American Psychiatric Association: Diagnostic and Statistical Manual of Mental Disorders (DSM-IV). Washington, American Psychiatric Association, 1994.
- Hollander E: Obsessive-compulsive spectrum phenomena and the DSM-V developmental process. *CNS Spectr* 2008;13:107–108.
- Hollander E, Braun A, Simeon D: Should OCD leave the anxiety disorders in DSM-V? The case for obsessive compulsive-related disorders. *Depress Anxiety* 2008;25:317–329.
- Mataix-Cols D, Pertusa A, Leckman JF: Issues for DSM-V: how should obsessive-compulsive and related disorders be classified? *Am J Psychiatry* 2007;164:1313–1314.
- Regier DA: Obsessive-compulsive behavior spectrum: refining the research agenda for DSM-V. *Psychiatry Res* 2009;170:1–2.
- Storch EA, Abramowitz J, Goodman WK: Where does obsessive-compulsive disorder belong in DSM-V? *Depress Anxiety* 2008;25:336–347.
- Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE: Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 2005;62:593–602.
- Geller DA: Obsessive-compulsive and spectrum disorders in children and adolescents. *Psychiatr Clin North Am* 2006;29:353–370.
- Geller DA, Biederman J, Jones J, Shapiro S, Schwartz S, Park KS: Obsessive-compulsive disorder in children and adolescents: a review. *Harv Rev Psychiatry* 1998;5:260–273.
- Heyman I, Fombonne E, Simmons H, Ford T, Meltzer H, Goodman R: Prevalence of obsessive-compulsive disorder in the British nationwide survey of child mental health. *Int Rev Psychiatry* 2003;15:178–184.
- Mataix-Cols D, Marks IM, Greist JH, Kobak KA, Baer L: Obsessive-compulsive symptom dimensions as predictors of compliance with and response to behaviour therapy: results from a controlled trial. *Psychother Psychosom* 2002;71:255–262.
- Stewart SE, Geller DA, Jenike M, Pauls D, Shaw D, Mullin B, Faraone SV: Long-term outcome of pediatric obsessive-compulsive disorder: a meta-analysis and qualitative review of the literature. *Acta Psychiatr Scand* 2004;110:4–13.
- Torres AR, Prince MJ, Bebbington PE, Bhugra D, Brugha TS, Farrell M, Jenkins R, Lewis G, Meltzer H, Singleton N: Obsessive-compulsive disorder: prevalence, comorbidity, impact, and help-seeking in the British National Psychiatric Morbidity Survey of 2000. *Am J Psychiatry* 2006;163:1978–1985.
- Baxter LR Jr, Phelps ME, Mazziotta JC, Guze BH, Schwartz JM, Selin CE: Local cerebral glucose metabolic rates in obsessive-compulsive disorder: a comparison with rates in unipolar depression and in normal controls. *Arch Gen Psychiatry* 1987;44:211–218.
- Baxter LR Jr, Schwartz JM, Mazziotta JC, Phelps ME, Pahl JJ, Guze BH, Fairbanks L: Cerebral glucose metabolic rates in nondepressed patients with obsessive-compulsive disorder. *Am J Psychiatry* 1988;145:1560–1563.
- Swedo SE, Schapiro MB, Grady CL, Cheslow DL, Leonard HL, Kumar A, Friedland R, Rapoport SI, Rapoport JL: Cerebral glucose metabolism in childhood-onset obsessive-compulsive disorder. *Arch Gen Psychiatry* 1989;46:518–523.
- Machlin SR, Harris GJ, Pearlson GD, Hoehn-Saric R, Jeffery P, Camargo EE: Elevated medial-frontal cerebral blood flow in obsessive-compulsive patients: a SPECT study. *Am J Psychiatry* 1991;148:1240–1242.
- Belliveau JW, Kennedy DN Jr, McKinstry RC, Buchbinder BR, Weisskoff RM, Cohen MS, Vevea JM, Brady TJ, Rosen BR: Functional mapping of the human visual cortex by magnetic resonance imaging. *Science* 1991;254:716–719.
- Bandettini PA, Wong EC, Hinks RS, Tikofsky RS, Hyde JS: Time course EPI of human brain function during task activation. *Magn Reson Med* 1992;25:390–397.
- Kwong KK, Belliveau JW, Chesler DA, Goldberg IE, Weisskoff RM, Poncelet BP, Kennedy DN, Hoppel BE, Cohen MS, Turner R: Dynamic magnetic resonance imaging of human brain activity during primary sensory stimulation. *Proc Natl Acad Sci USA* 1992; 89:5675–5679.
- Ogawa S, Tank DW, Menon R, Ellermann JM, Kim SG, Merkle H, Ugurbil K: Intrinsic signal changes accompanying sensory stimulation: functional brain mapping with magnetic resonance imaging. *Proc Natl Acad Sci USA* 1992;89:5951–5955.

Disclosure Statement

M.P. has served as a consultant to, or was engaged in research collaborations with Eli Lilly and Organon Corporations in Italy. S.F. has in the past 3 years participated on the advisory boards of Pfizer and Lilly and received honoraria from Lilly, Bristol-Meyers Squibb, Sigma Tau, Schering and Pfizer. P.G. has in the past 3 years received research support from Lilly and Janssen, has participated on the advisory boards of Lilly, Organon, Pfizer and Schering, and received honoraria from Lilly and Organon. R.T. has in the past 3 years participated on the advisory boards of Schering, Servier and Pfizer, and received honoraria from Schering, Servier and Pfizer.

All other authors of this paper have no relevant affiliations or financial involvement with any organisation or entity with a financial interest in, or financial conflict with, the subject matter or materials discussed in the manuscript. This includes employment, consultancies, honoraria, stock ownership or options, expert testimony, grants or patents received or pending, or royalties.

- 23 Baxter LR Jr, Schwartz JM, Bergman KS, Szuba MP, Guze BH, Mazzotta JC, Alazraki A, Selin CE, Ferng HK, Munford P: Caudate glucose metabolic rate changes with both drug and behavior therapy for obsessive-compulsive disorder. *Arch Gen Psychiatry* 1992;49:681–689.
- 24 Hoehn-Saric R, Pearlson GD, Harris GJ, Machlin SR, Camargo EE: Effects of fluoxetine on regional cerebral blood flow in obsessive-compulsive patients. *Am J Psychiatry* 1991;148:1243–1245.
- 25 Swedo SE, Pietrini P, Leonard HL, Schapiro MB, Rettew DC, Goldberger EL, Rapoport SI, Rapoport JL, Grady CL: Cerebral glucose metabolism in childhood-onset obsessive-compulsive disorder: revisualization during pharmacotherapy. *Arch Gen Psychiatry* 1992;49:690–694.
- 26 Cottraux J, Gérard D, Cinotti L, Froment JC, Deiber MP, le Bars D, Galy G, Millet P, Labbé C, Lavenne F, Bouvard M, Mauguère F: A controlled positron emission tomography study of obsessive and neutral auditory stimulation in obsessive-compulsive disorder with checking rituals. *Psychiatry Res* 1996; 60:101–112.
- 27 Jenike MA, Breiter HC, Baer L, Kennedy DN, Savage CR, Olivares MJ, O'Sullivan RL, Shera DM, Rauch SL, Keuthen N, Rosen BR, Caviness VS, Filipek PA: Cerebral structural abnormalities in obsessive-compulsive disorder: a quantitative morphometric magnetic resonance imaging study. *Arch Gen Psychiatry* 1996;53:625–632.
- 28 McGuire PK, Bench CJ, Frith CD, Marks IM, Frackowiak RS, Dolan RJ: Functional anatomy of obsessive-compulsive phenomena. *Br J Psychiatry* 1994;164:459–468.
- 29 Rauch SL, Jenike MA, Alpert NM, Baer L, Breiter HC, Savage CR, Fischman AJ: Regional cerebral blood flow measured during symptom provocation in obsessive-compulsive disorder using oxygen 15-labeled carbon dioxide and positron emission tomography. *Arch Gen Psychiatry* 1994;51:62–70.
- 30 Zohar J, Insel TR, Berman KF, Foa EB, Hill JL, Weinberger DR: Anxiety and cerebral blood flow during behavioral challenge: dissociation of central from peripheral and subjective measures. *Arch Gen Psychiatry* 1989; 46:505–510.
- 31 Savage CR, Weilburg JB, Duffy FH, Baer L, Shera DM, Jenike MA: Low-level sensory processing in obsessive-compulsive disorder: an evoked potential study. *Biol Psychiatry* 1994;35:247–252.
- 32 Towey J, Bruder G, Tenke C, Leite P, DeCaria C, Friedman D, Hollander E: Event-related potential and clinical correlates of neurodysfunction in obsessive-compulsive disorder. *Psychiatry Res* 1993;49:167–181.
- 33 Garber HJ, Ananth JV, Chiu LC, Griswold VJ, Oldendorf WH: Nuclear magnetic resonance study of obsessive-compulsive disorder. *Am J Psychiatry* 1989;146:1001–1005.
- 34 Ebert D, Speck O, König A, Berger M, Hennig J, Hohagen F: ¹H-magnetic resonance spectroscopy in obsessive-compulsive disorder: evidence for neuronal loss in the cingulate gyrus and the right striatum. *Psychiatry Res* 1997;74:173–176.
- 35 Perani D, Colombo C, Bressi S, Bonfanti A, Grassi F, Scarone S, Bellodi L, Smeraldi E, Fazio F: [¹⁸F]FDG PET study in obsessive-compulsive disorder: a clinical/metabolic correlation study after treatment. *Br J Psychiatry* 1995;166:244–250.
- 36 Galderisi S, Mucci A, Catapano F, D'Amato AC, Maj M: Neuropsychological slowness in obsessive-compulsive patients: is it confined to tests involving the fronto-subcortical systems? *Br J Psychiatry* 1995;167:394–398.
- 37 Ursu S, Stenger VA, Shear MK, Jones MR, Carter CS: Overactive action monitoring in obsessive-compulsive disorder: evidence from functional magnetic resonance imaging. *Psychol Sci* 2003;14:347–353.
- 38 Fitzgerald KD, Welsh RC, Gehring WJ, Abelson JL, Himle JA, Liberzon I, Taylor SF: Error-related hyperactivity of the anterior cingulate cortex in obsessive-compulsive disorder. *Biol Psychiatry* 2005;57:287–294.
- 39 Maltby N, Tolin DF, Worhunsky P, O'Keefe TM, Kiehl KA: Dysfunctional action monitoring hyperactivates frontal-striatal circuits in obsessive-compulsive disorder: an event-related fMRI study. *Neuroimage* 2005; 24:495–503.
- 40 Nakao T, Nakagawa A, Yoshiura T, Nakatani E, Nabeyama M, Yoshizato C, Kudoh A, Tada K, Yoshioka K, Kawamoto M: A functional MRI comparison of patients with obsessive-compulsive disorder and normal controls during a Chinese character Stroop task. *Psychiatry Res* 2005;139:101–114.
- 41 Nabeyama M, Nakagawa A, Yoshiura T, Nakao T, Nakatani E, Togao O, Yoshizato C, Yoshioka K, Tomita M, Kanba S: Functional MRI study of brain activation alterations in patients with obsessive-compulsive disorder after symptom improvement. *Psychiatry Res* 2008;163:236–247.
- 42 Viard A, Flament MF, Artiges E, Dehaene S, Naccache L, Cohen D, Mazet P, Mouren MC, Martinot JL: Cognitive control in childhood-onset obsessive-compulsive disorder: a functional MRI study. *Psychol Med* 2005; 35:1007–1017.
- 43 Adler CM, McDonough-Ryan P, Sax KW, Holland SK, Arndt S, Strakowski SM: fMRI of neuronal activation with symptom provocation in unmedicated patients with obsessive compulsive disorder. *J Psychiatr Res* 2000;34:317–324.
- 44 Breiter HC, Rauch SL: Functional MRI and the study of OCD: from symptom provocation to cognitive-behavioral probes of cortico-striatal systems and the amygdala. *Neuroimage* 1996;4:S127–S138.
- 45 Saxena S, Brody AL, Schwartz JM, Baxter LR: Neuroimaging and frontal-subcortical circuitry in obsessive-compulsive disorder. *Br J Psychiatry Suppl* 1998;35:26–37.
- 46 Levy R, Goldman-Rakic PS: Segregation of working memory functions within the dorsolateral prefrontal cortex. *Exp Brain Res* 2000;133:23–32.
- 47 Petrides M: Specialized systems for the processing of mnemonic information within the primate frontal cortex. *Philos Trans R Soc Lond B Biol Sci* 1996;351:1455–1461, discussion 1461–1462.
- 48 Manoach DS, White NS, Lindgren KA, Heckers S, Coleman MJ, Dubal S, Holzman PS: Hemispheric specialization of the lateral prefrontal cortex for strategic processing during spatial and shape working memory. *Neuroimage* 2004;21:894–903.
- 49 Petrides M: Frontal lobes and behaviour. *Curr Opin Neurobiol* 1994;4:207–211.
- 50 Burgess PW, Veitch E, de Lacy Costello A, Shallice T: The cognitive and neuroanatomical correlates of multitasking. *Neuropsychologia* 2000;38:848–863.
- 51 Bunge SA, Kahn I, Wallis JD, Miller EK, Wagner AD: Neural circuits subserving the retrieval and maintenance of abstract rules. *J Neurophysiol* 2003;90:3419–3428.
- 52 Menon V, Mackenzie K, Rivera SM, Reiss AL: Prefrontal cortex involvement in processing incorrect arithmetic equations: evidence from event-related fMRI. *Hum Brain Mapp* 2002;16:119–130.
- 53 van den Heuvel OA, Veltman DJ, Groenewegen HJ, Cath DC, van Balkom AJLM, van Hartkamp J, Barkhof F, van Dyck R: Frontal-striatal dysfunction during planning in obsessive-compulsive disorder. *Arch Gen Psychiatry* 2005;62:301–309.
- 54 Gu BM, Park JY, Kang DH, Lee SJ, Yoo SY, Jo HJ, Choi CH, Lee JM, Kwon JS: Neural correlates of cognitive inflexibility during task-switching in obsessive-compulsive disorder. *Brain* 2008;131:155–164.
- 55 van der Wee NJA, Ramsey NF, Jansma JM, Denys DA, van Megen HJGM, Westenberg HMG, Kahn RS: Spatial working memory deficits in obsessive compulsive disorder are associated with excessive engagement of the medial frontal cortex. *Neuroimage* 2003;20: 2271–2280.
- 56 Nakao T, Nakagawa A, Nakatani E, Nabeyama M, Sanematsu H, Yoshiura T, Togao O, Tomita M, Masuda Y, Yoshioka K, Kuroki T, Kanba S: Working memory dysfunction in obsessive-compulsive disorder: a neuropsychological and functional MRI study. *J Psychiatr Res* 2009;43:784–791.
- 57 Nakao T, Nakagawa A, Yoshiura T, Nakatani E, Nabeyama M, Sanematsu H, Togao O, Yoshioka K, Tomita M, Kuroki T, Kanba S: Duration effect of obsessive-compulsive disorder on cognitive function: a functional MRI study. *Depress Anxiety* 2009;26:814–823.

- 58 Levine JB, Gruber SA, Baird AA, Yurgelun-Todd D: Obsessive-compulsive disorder among schizophrenic patients: an exploratory study using functional magnetic resonance imaging data. *Compr Psychiatry* 1998; 39:308–311.
- 59 Aouizerate B, Guehl D, Cuny E, Rougier A, Bioulac B, Tignol J, Burbaud P: Pathophysiology of obsessive-compulsive disorder: a necessary link between phenomenology, neuropsychology, imagery and physiology. *Prog Neurobiol* 2004;72:195–221.
- 60 Breiter HC, Rauch SL, Kwong KK, Baker JR, Weisskoff RM, Kennedy DN, Kendrick AD, Davis TL, Jiang A, Cohen MS, Stern CE, Belliveau JW, Baer L, O'Sullivan RL, Savage CR, Jenike MA, Rosen BR: Functional magnetic resonance imaging of symptom provocation in obsessive-compulsive disorder. *Arch Gen Psychiatry* 1996;53:595–606.
- 61 McDougle CJ, Epperson CN, Pelton GH, Wasyluk S, Price LH: A double-blind, placebo-controlled study of risperidone addition in serotonin reuptake inhibitor-refractory obsessive-compulsive disorder. *Arch Gen Psychiatry* 2000;57:794–801.
- 62 Rogers RD, Blackshaw AJ, Middleton HC, Matthews K, Hawtin K, Crowley C, Hopwood A, Wallace C, Deakin JF, Sahakian BJ, Robbins TW: Tryptophan depletion impairs stimulus-reward learning while methylphenidate disrupts attentional control in healthy young adults: implications for the monoaminergic basis of impulsive behaviour. *Psychopharmacology* 1999;146:482–491.
- 63 Schultz W: Predictive reward signal of dopamine neurons. *J Neurophysiol* 1998;80:1–27.
- 64 Abbruzzese M, Ferri S, Scarone S: The selective breakdown of frontal functions in patients with obsessive-compulsive disorder and in patients with schizophrenia: a double dissociation experimental finding. *Neuropsychologia* 1997;35:907–912.
- 65 Cavedini P, Riboldi G, D'Annunzio A, Belotti P, Cisima M, Bellodi L: Decision-making heterogeneity in obsessive-compulsive disorder: ventromedial prefrontal cortex function predicts different treatment outcomes. *Neuropsychologia* 2002;40:205–211.
- 66 Hermesh H, Zohar J, Weizman A, Voet H, Gross-Isseroff R: Orbitofrontal cortex dysfunction in obsessive-compulsive disorder? 2. Olfactory quality discrimination in obsessive-compulsive disorder. *Eur Neuropsychopharmacol* 1999;9:415–420.
- 67 Nielen MMA, Veltman DJ, de Jong R, Mulder G, den Boer JA: Decision making performance in obsessive compulsive disorder. *J Affect Disord* 2002;69:257–260.
- 68 Kim JJ, Lee MC, Kim J, Kim IY, Kim SI, Han MH, Chang KH, Kwon JS: Grey matter abnormalities in obsessive-compulsive disorder: statistical parametric mapping of segmented magnetic resonance images. *Br J Psychiatry* 2001;179:330–334.
- 69 Pujol J, Soriano-Mas C, Alonso P, Cardoner N, Menchon JM, Deus J, Vallejo J: Mapping structural brain alterations in obsessive-compulsive disorder. *Arch Gen Psychiatry* 2004;61:720–730.
- 70 Lacerda ALT, Dalgalarondo P, Caetano D, Camargo EE, Etchebehere ECSC, Soares JC: Elevated thalamic and prefrontal regional cerebral blood flow in obsessive-compulsive disorder: a SPECT study. *Psychiatry Res* 2003;123:125–134.
- 71 Busatto GF, Zamignani DR, Buchpiguel CA, Garrido GE, Glabus MF, Rocha ET, Maia AF, Rosario-Campos MC, Campi Castro C, Furuie SS, Gutierrez MA, McGuire PK, Miguel EC: A voxel-based investigation of regional cerebral blood flow abnormalities in obsessive-compulsive disorder using single photon emission computed tomography (SPECT). *Psychiatry Res* 2000;99:15–27.
- 72 van den Heuvel OA, Veltman DJ, Groenewegen HJ, Dolan RJ, Cath DC, Boellaard R, Messina CT, van Balkom AJLM, van Oppen P, Witter MP, Lammertsma AA, van Dyck R: Amygdala activity in obsessive-compulsive disorder with contamination fear: a study with oxygen-15 water positron emission tomography. *Psychiatry Res* 2004;132:225–237.
- 73 Remijnse PL, Nielen MMA, van Balkom AJLM, Cath DC, van Oppen P, Uylings HBM, Veltman DJ: Reduced orbitofrontal activity on a reversal learning task in obsessive-compulsive disorder. *Arch Gen Psychiatry* 2006;63:1225–1236.
- 74 Rauch SL, Wedig MM, Wright CI, Martis B, McMullin KG, Shin LM, Cannistraro PA, Wilhelm S: Functional magnetic resonance imaging study of regional brain activation during implicit sequence learning in obsessive-compulsive disorder. *Biol Psychiatry* 2007;61:330–336.
- 75 Baxter LR Jr: Basal ganglia systems in ritualistic social displays: reptiles and humans – function and illness. *Physiol Behav* 2003;79:451–460.
- 76 Remijnse PL, Nielen MMA, van Balkom AJLM, Hendriks GJ, Hoogendijk WJ, Uylings HBM, Veltman DJ: Differential frontal-striatal and paralimbic activity during reversal learning in major depressive disorder and obsessive-compulsive disorder. *Psychol Med* 2009;39:1503–1518.
- 77 Crespo-Facorro B, Cabranes JA, Lopez-Ibor Alcocer MI, Paya B, Fernandez Perez C, Encinas M, Ayuso Mateos JL, Lopez-Ibor JJ Jr: Regional cerebral blood flow in obsessive-compulsive patients with and without a chronic tic disorder: a SPECT study. *Eur Arch Psychiatry Clin Neurosci* 1999;249:156–161.
- 78 Lucey JV, Costa DC, Blanes T, Busatto GF, Pilowsky LS, Takei N, Marks IM, Ell PJ, Kerwin RW: Regional cerebral blood flow in obsessive-compulsive disordered patients at rest: differential correlates with obsessive-compulsive and anxious-avoidant dimensions. *Br J Psychiatry* 1995;167:629–634.
- 79 Roth RM, Saykin AJ, Flashman LA, Pixley HS, West JD, Mamourian AC: Event-related functional magnetic resonance imaging of response inhibition in obsessive-compulsive disorder. *Biol Psychiatry* 2007;62:901–909.
- 80 Ullsperger M, von Cramon DY: Subprocesses of performance monitoring: a dissociation of error processing and response competition revealed by event-related fMRI and ERPs. *Neuroimage* 2001;14:1387–1401.
- 81 Garavan H, Ross TJ, Kaufman J, Stein EA: A midline dissociation between error-processing and response-conflict monitoring. *Neuroimage* 2003;20:1132–1139.
- 82 Yücel M, Harrison BJ, Wood SJ, Fornito A, Wellard RM, Pujol J, Clarke K, Phillips ML, Kyrios M, Velakoulis D, Pantelis C: Functional and biochemical alterations of the medial frontal cortex in obsessive-compulsive disorder. *Arch Gen Psychiatry* 2007;64:946–955.
- 83 Buckner RL, Raichle ME, Petersen SE: Dissociation of human prefrontal cortical areas across different speech production tasks and gender groups. *J Neurophysiol* 1995;74:2163–2173.
- 84 Mummery CJ, Patterson K, Hodges JR, Wise RJ: Generating 'tiger' as an animal name or a word beginning with T: differences in brain activation. *Proc Biol Sci* 1996;263:989–995.
- 85 Pujol J, Vendrell P, Deus J, Kulisevsky J, Martí-Vilalta JL, García C, Junqué C, Capdevila A: Frontal lobe activation during word generation studied by functional MRI. *Acta Neurol Scand* 1996;93:403–410.
- 86 Cuenod CA, Bookheimer SY, Hertz-Pannier L, Zeffiro TA, Theodore WH, le Bihan D: Functional MRI during word generation, using conventional equipment: a potential tool for language localization in the clinical environment. *Neurology* 1995;45:1821–1827.
- 87 Rueckert L, Appollonio I, Grafman J, Jezzard P, Johnson R Jr, le Bihan D, Turner R: Magnetic resonance imaging functional activation of left frontal cortex during covert word production. *J Neuroimaging* 1994;4:67–70.
- 88 Yetkin FZ, Hammeke TA, Swanson SJ, Morris GL, Mueller WM, McAuliffe TL, Haughton VM: A comparison of functional MR activation patterns during silent and audible language tasks. *AJNR Am J Neuroradiol* 1995;16:1087–1092.
- 89 Yurgelun-Todd DA, Wateraux CM, Cohen BM, Gruber SA, English CD, Renshaw PF: Functional magnetic resonance imaging of schizophrenic patients and comparison subjects during word production. *Am J Psychiatry* 1996;153:200–205.
- 90 Christensen KJ, Kim SW, Dysken MW, Hoover KM: Neuropsychological performance in obsessive-compulsive disorder. *Biol Psychiatry* 1992;31:4–18.
- 91 Otto MW: Normal and abnormal information processing: a neuropsychological perspective on obsessive-compulsive disorder. *Psychiatr Clin North Am* 1992;15:825–848.

- 92 Henseler I, Gruber O, Kraft S, Krick C, Reith W, Falkai P: Compensatory hyperactivations as markers of latent working memory dysfunctions in patients with obsessive-compulsive disorder: an fMRI study. *J Psychiatry Neurosci* 2008;33:209–215.
- 93 Lázaro L, Caldú X, Junqué C, Bargalló N, Andrés S, Morer A, Castro-Fornieles J: Cerebral activation in children and adolescents with obsessive-compulsive disorder before and after treatment: a functional MRI study. *J Psychiatr Res* 2008;42:1051–1059.
- 94 Jung WH, Gu BM, Kang DH, Park JY, Yoo SY, Choi CH, Lee JM, Kwon JS: BOLD response during visual perception of biological motion in obsessive-compulsive disorder: an fMRI study using the dynamic point-light animation paradigm. *Eur Arch Psychiatry Clin Neurosci* 2009;259:46–54.
- 95 Simpson S, Baldwin B: Neuropsychiatry and SPECT of an acute obsessive-compulsive syndrome patient. *Br J Psychiatry* 1995; 166:390–392.
- 96 Berthier ML, Kulisevsky J, Gironell A, Heras JA: Obsessive-compulsive disorder associated with brain lesions: clinical phenomenology, cognitive function, and anatomic correlates. *Neurology* 1996;47:353–361.
- 97 Hendler T, Goshen E, Tadmor R, Lustig M, Zwas ST, Zohar J: Evidence for striatal modulation in the presence of fixed cortical injury in obsessive-compulsive disorder (OCD). *Eur Neuropsychopharmacol* 1999; 9:371–376.
- 98 Walker SC, Robbins TW, Roberts AC: Differential contributions of dopamine and serotonin to orbitofrontal cortex function in the marmoset. *Cereb Cortex* 2009;19:889–898.
- 99 Koeppe MJ, Gunn RN, Lawrence AD, Cunningham VJ, Dagher A, Jones T, Brooks DJ, Bench CJ, Grasby PM: Evidence for striatal dopamine release during a video game. *Nature* 1998;393:266–268.
- 100 Schilman EA, Klavir O, Winter C, Sohr R, Joel D: The role of the striatum in compulsive behavior in intact and orbitofrontal-cortex-lesioned rats: possible involvement of the serotonergic system. *Neuropsychopharmacology* 2010;35:1026–1039.
- 101 van der Wee NJ, Stevens H, Hardeman JA, Mandl RC, Denys DA, van Megen HJ, Kahn RS, Westenberg HM: Enhanced dopamine transporter density in psychotropic-naïve patients with obsessive-compulsive disorder shown by [¹²³I]β-CIT SPECT. *Am J Psychiatry* 2004;161:2201–2206.
- 102 Denys D, van der Wee N, Janssen J, de Geus F, Westenberg HGM: Low level of dopaminergic D₂ receptor binding in obsessive-compulsive disorder. *Biol Psychiatry* 2004; 55:1041–1045.
- 103 Alexander GE, DeLong MR, Strick PL: Parallel organization of functionally segregated circuits linking basal ganglia and cortex. *Annu Rev Neurosci* 1986;9:357–381.
- 104 Knopman D, Nissen MJ: Procedural learning is impaired in Huntington's disease: evidence from the serial reaction time task. *Neuropsychologia* 1991;29:245–254.
- 105 Aylward EH, Harris GJ, Hoehn-Saric R, Barta PE, Machlin SR, Pearson GD: Normal caudate nucleus in obsessive-compulsive disorder assessed by quantitative neuroimaging. *Arch Gen Psychiatry* 1996;53: 577–584.
- 106 Robinson D, Wu H, Munne RA, Ashtari M, Alvir JM, Lerner G, Koreen A, Cole K, Bogerts B: Reduced caudate nucleus volume in obsessive-compulsive disorder. *Arch Gen Psychiatry* 1995;52:393–398.
- 107 Rubin RT, Villanueva-Meyer J, Ananth J, Trajmar PG, Mena I: Regional xenon 133 cerebral blood flow and cerebral technetium 99m HMPAO uptake in unmedicated patients with obsessive-compulsive disorder and matched normal control subjects: determination by high-resolution single-photon emission computed tomography. *Arch Gen Psychiatry* 1992;49:695–702.
- 108 Saxena S, Brody AL, Maidment KM, Smith EC, Zohrabi N, Katz E, Baker SK, Baxter LR Jr: Cerebral glucose metabolism in obsessive-compulsive hoarding. *Am J Psychiatry* 2004;161:1038–1048.
- 109 Deckersbach T, Savage CR, Curran T, Bohne A, Wilhelm S, Baer L, Jenike MA, Rauch SL: A study of parallel implicit and explicit information processing in patients with obsessive-compulsive disorder. *Am J Psychiatry* 2002;159:1780–1782.
- 110 Rauch SL, Savage CR: Neuroimaging and neuropsychology of the striatum: bridging basic science and clinical practice. *Psychiatr Clin North Am* 1997;20:741–768.
- 111 Nakao T, Nakagawa A, Yoshiura T, Nakatani E, Nabeyama M, Yoshizato C, Kudoh A, Tada K, Yoshioka K, Kawamoto M, Togao O, Kanba S: Brain activation of patients with obsessive-compulsive disorder during neuropsychological and symptom provocation tasks before and after symptom improvement: a functional magnetic resonance imaging study. *Biol Psychiatry* 2005;57:901–910.
- 112 Benkelfat C, Nordahl TE, Semple WE, King AC, Murphy DL, Cohen RM: Local cerebral glucose metabolic rates in obsessive-compulsive disorder: patients treated with clomipramine. *Arch Gen Psychiatry* 1990;47: 840–848.
- 113 Rauch SL, Jenike MA: Neurobiological models of obsessive-compulsive disorder. *Psychosomatics* 1993;34:20–32.
- 114 Rauch SL, Savage CR, Alpert NM, Dougherty D, Kendrick A, Curran T, Brown HD, Manzo P, Fischman AJ, Jenike MA: Probing striatal function in obsessive-compulsive disorder: a PET study of implicit sequence learning. *J Neuropsychiatry Clin Neurosci* 1997;9:568–573.
- 115 Purcell R, Maruff P, Kyrios M, Pantelis C: Cognitive deficits in obsessive-compulsive disorder on tests of frontal-striatal function. *Biol Psychiatry* 1998;43:348–357.
- 116 Purcell R, Maruff P, Kyrios M, Pantelis C: Neuropsychological deficits in obsessive-compulsive disorder: a comparison with unipolar depression, panic disorder, and normal controls. *Arch Gen Psychiatry* 1998;55:415–423.
- 117 Mataix-Cols D, Junqué C, Sánchez-Turet M, Vallejo J, Verger K, Barrios M: Neuropsychological functioning in a subclinical obsessive-compulsive sample. *Biol Psychiatry* 1999;45:898–904.
- 118 Schmidtke K, Schorb A, Winkelmann G, Hohagen F: Cognitive frontal lobe dysfunction in obsessive-compulsive disorder. *Biol Psychiatry* 1998;43:666–673.
- 119 Veale DM, Sahakian BJ, Owen AM, Marks IM: Specific cognitive deficits in tests sensitive to frontal lobe dysfunction in obsessive-compulsive disorder. *Psychol Med* 1996;26:1261–1269.
- 120 Keith Berg W, Byrd D: The Tower of London spatial problem-solving task: enhancing clinical and research implementation. *J Clin Exp Neuropsychol* 2002;24:586–604.
- 121 Cahill L, Haier RJ, Fallon J, Alkire MT, Tang C, Keator D, Wu J, McLaugh JL: Amygdala activity at encoding correlated with long-term, free recall of emotional information. *Proc Natl Acad Sci USA* 1996; 93:8016–8021.
- 122 Corbetta M, Shulman GL, Miezin FM, Petersen SE: Superior parietal cortex activation during spatial attention shifts and visual feature conjunction. *Science* 1995;270: 802–805.
- 123 Courchesne E, Townsend J, Akshoomoff NA, Saitoh O, Yeung-Courchesne R, Lincoln AJ, James HE, Haas RH, Schreibman L, Lau L: Impairment in shifting attention in autistic and cerebellar patients. *Behav Neurosci* 1994;108:848–865.
- 124 Kim SG, Ugurbil K, Strick PL: Activation of a cerebellar output nucleus during cognitive processing. *Science* 1994;265:949–951.
- 125 Le TH, Pardo JV, Hu X: 4 T-fMRI study of nonspatial shifting of selective attention: cerebellar and parietal contributions. *J Neurophysiol* 1998;79:1535–1548.
- 126 Schmahmann JD, Pandya DN: Prefrontal cortex projections to the basilar pons in rhesus monkey: implications for the cerebellar contribution to higher function. *Neurosci Lett* 1995;199:175–178.
- 127 Schmahmann JD, Pandya DN: Anatomical investigation of projections to the basis pontis from posterior parietal association cortices in rhesus monkey. *J Comp Neurol* 1989;289:53–73.
- 128 Schmahmann JD, Pandya DN: Prelunate, occipitotemporal, and parahippocampal projections to the basis pontis in rhesus monkey. *J Comp Neurol* 1993;337:94–112.

- 129 Vilensky JA, van Hoesen GW: Corticopontine projections from the cingulate cortex in the rhesus monkey. *Brain Res* 1981;205:391–395.
- 130 Schmahmann JD, Sherman JC: The cerebellar cognitive affective syndrome. *Brain* 1998;121(pt 4):561–579.
- 131 Page LA, Rubia K, Deeley Q, Daly E, Toal F, Mataix-Cols D, Giampietro V, Schmitz N, Murphy DGM: A functional magnetic resonance imaging study of inhibitory control in obsessive-compulsive disorder. *Psychiatry Res* 2009;174:202–209.
- 132 Sanematsu H, Nakao T, Yoshiura T, Nabeyama M, Togao O, Tomita M, Masuda Y, Nakatani E, Nakagawa A, Kanba S: Predictors of treatment response to fluvoxamine in obsessive-compulsive disorder: an fMRI study. *J Psychiatr Res* 2010;44:193–200.
- 133 Kang DH, Kwon JS, Kim JJ, Youn T, Park HJ, Kim MS, Lee DS, Lee MC: Brain glucose metabolic changes associated with neuropsychological improvements after 4 months of treatment in patients with obsessive-compulsive disorder. *Acta Psychiatr Scand* 2003;107:291–297.
- 134 Rauch SL, Baxter LR: Neuroimaging of OCD and related disorders; in Jenike MA, Baer L, Minichiello WE (eds): *Obsessive-Compulsive Disorders. Theory and Management*. St. Louis, Mosby, 1998, pp 289–317.
- 135 Nordahl TE, Benkelfat C, Semple WE, Gross M, King AC, Cohen RM: Cerebral glucose metabolic rates in obsessive compulsive disorder. *Neuropsychopharmacology* 1989;2:23–28.
- 136 Brody AL, Saxena S, Schwartz JM, Stoessel PW, Maidment K, Phelps ME, Baxter LR Jr: FDG-PET predictors of response to behavioral therapy and pharmacotherapy in obsessive-compulsive disorder. *Psychiatry Res* 1998;84:1–6.
- 137 Schwartz JM, Stoessel PW, Baxter LR Jr, Martin KM, Phelps ME: Systematic changes in cerebral glucose metabolic rate after successful behavior modification treatment of obsessive-compulsive disorder. *Arch Gen Psychiatry* 1996;53:109–113.
- 138 Rauch SL, Dougherty DD, Cosgrove GR, Cassem EH, Alpert NM, Price BH, Nierenberg AA, Mayberg HS, Baer L, Jenike MA, Fischman AJ: Cerebral metabolic correlates as potential predictors of response to anterior cingulotomy for obsessive compulsive disorder. *Biol Psychiatry* 2001;50:659–667.
- 139 Saxena S, Brody AL, Maidment KM, Dunkin JJ, Colgan M, Alborzian S, Phelps ME, Baxter LR Jr: Localized orbitofrontal and subcortical metabolic changes and predictors of response to paroxetine treatment in obsessive-compulsive disorder. *Neuropsychopharmacology* 1999;21:683–693.
- 140 Deckersbach T, Dougherty DD, Rauch SL: Functional imaging of mood and anxiety disorders. *J Neuroimaging* 2006;16:1–10.
- 141 Mataix-Cols D, Rosario-Campos MC, Leckman JF: A multidimensional model of obsessive-compulsive disorder. *Am J Psychiatry* 2005;162:228–238.
- 142 Miguel EC, Leckman JF, Rauch S, do Rosario-Campos MC, Hounie AG, Mercadante MT, Chacon P, Pauls DL: Obsessive-compulsive disorder phenotypes: implications for genetic studies. *Mol Psychiatry* 2005;10:258–275.
- 143 Leckman JF, Grice DE, Boardman J, Zhang H, Vitale A, Bondi C, Alsobrook J, Peterson BS, Cohen DJ, Rasmussen SA, Goodman WK, McDougle CJ, Pauls DL: Symptoms of obsessive-compulsive disorder. *Am J Psychiatry* 1997;154:911–917.
- 144 Baer L: Factor analysis of symptom subtypes of obsessive-compulsive disorder and their relation to personality and tic disorders. *J Clin Psychiatry* 1994;55(suppl):18–23.
- 145 Summerfeldt LJ, Richter MA, Antony MM, Swinson RP: Symptom structure in obsessive-compulsive disorder: a confirmatory factor-analytic study. *Behav Res Ther* 1999;37:297–311.
- 146 Mataix-Cols D, Rauch SL, Baer L, Eisen JL, Shera DM, Goodman WK, Rasmussen SA, Jenike MA: Symptom stability in adult obsessive-compulsive disorder: data from a naturalistic two-year follow-up study. *Am J Psychiatry* 2002;159:263–268.
- 147 Alsobrook JP 2nd, Leckman JF, Goodman WK, Rasmussen SA, Pauls DL: Segregation analysis of obsessive-compulsive disorder using symptom-based factor scores. *Am J Med Genet* 1999;88:669–675.
- 148 Leckman JF, Pauls DL, Zhang H, Rosario-Campos MC, Katsochis L, Kidd KK, Pakstis AJ, Alsobrook JP, Robertson MM, McMahon WM, Walkup JT, van de Wetering BJM, King RA, Cohen DJ: Obsessive-compulsive symptom dimensions in affected sibling pairs diagnosed with Gilles de la Tourette syndrome. *Am J Med Genet B Neuropsychiatr Genet* 2003;116B:60–68.
- 149 Zhang H, Leckman JF, Pauls DL, Tsai CP, Kidd KK, Campos MR: Genomewide scan of hoarding in sib pairs in which both sibs have Gilles de la Tourette syndrome. *Am J Hum Genet* 2002;70:896–904.
- 150 Mataix-Cols D, Wooderson S, Lawrence N, Brammer MJ, Speckens A, Phillips ML: Distinct neural correlates of washing, checking, and hoarding symptom dimensions in obsessive-compulsive disorder. *Arch Gen Psychiatry* 2004;61:564–576.
- 151 Gilbert AR, Akkal D, Almeida JRC, Mataix-Cols D, Kalas C, Devlin B, Birmaher B, Phillips ML: Neural correlates of symptom dimensions in pediatric obsessive-compulsive disorder: a functional magnetic resonance imaging study. *J Am Acad Child Adolesc Psychiatry* 2009;48:936–944.
- 152 Rubia K, Cubillo A, Smith AB, Woolley J, Heyman I, Brammer MJ: Disorder-specific dysfunction in right inferior prefrontal cortex during two inhibition tasks in boys with attention-deficit hyperactivity disorder compared to boys with obsessive-compulsive disorder. *Hum Brain Mapp* 2010;31:287–299.
- 153 Woolley J, Heyman I, Brammer M, Framp-ton I, McGuire PK, Rubia K: Brain activation in paediatric obsessive compulsive disorder during tasks of inhibitory control. *Br J Psychiatry* 2008;192:25–31.
- 154 Rosenberg DR, Keshavan MS, O'Hearn KM, Dick EL, Bagwell WW, Seymour AB, Montrose DM, Pierri JN, Birmaher B: Frontostriatal measurement in treatment-naive children with obsessive-compulsive disorder. *Arch Gen Psychiatry* 1997;54:824–830.
- 155 Szeszko PR, MacMillan S, McMeniman M, Chen S, Baribault K, Lim KO, Ivey J, Rose M, Banerjee SP, Bhandari R, Moore GJ, Rosenberg DR: Brain structural abnormalities in psychotropic drug-naive pediatric patients with obsessive-compulsive disorder. *Am J Psychiatry* 2004;161:1049–1056.
- 156 Gilbert AR, Moore GJ, Keshavan MS, Paulson LA, Narula V, Mac Master FP, Stewart CM, Rosenberg DR: Decrease in thalamic volumes of pediatric patients with obsessive-compulsive disorder who are taking paroxetine. *Arch Gen Psychiatry* 2000;57:449–456.
- 157 Austin MP, Mitchell P, Goodwin GM: Cognitive deficits in depression: possible implications for functional neuropathology. *Br J Psychiatry* 2001;178:200–206.
- 158 Swaab DF, Fliers E, Hoogendijk WJ, Veltman DJ, Zhou JN: Interaction of prefrontal cortical and hypothalamic systems in the pathogenesis of depression. *Prog Brain Res* 2000;126:369–396.
- 159 Saxena S, Brody AL, Ho ML, Alborzian S, Ho MK, Maidment KM, Huang SC, Wu HM, Au SC, Baxter LR Jr: Cerebral metabolism in major depression and obsessive-compulsive disorder occurring separately and concurrently. *Biol Psychiatry* 2001;50:159–170.
- 160 Greisberg S, McKay D: Neuropsychology of obsessive-compulsive disorder: a review and treatment implications. *Clin Psychol Rev* 2003;23:95–117.
- 161 Dehaene S, Changeux JP: A hierarchical neuronal network for planning behavior. *Proc Natl Acad Sci USA* 1997;94:13293–13298.
- 162 Anzai Y, Simon HA: The theory of learning by doing. *Psychol Rev* 1979;86:124–140.
- 163 Shallice T: Specific impairments of planning. *Philos Trans R Soc Lond B Biol Sci* 1982;298:199–209.
- 164 Baker SC, Rogers RD, Owen AM, Frith CD, Dolan RJ, Frackowiak RS, Robbins TW: Neural systems engaged by planning: a PET study of the Tower of London task. *Neuropsychologia* 1996;34:515–526.

- 165 Dagher A, Owen AM, Boecker H, Brooks DJ: Mapping the network for planning: a correlational PET activation study with the Tower of London task. *Brain* 1999;122(pt 10):1973–1987.
- 166 Owen AM, Doyon J, Petrides M, Evans AC: Planning and spatial working memory: a positron emission tomography study in humans. *Eur J Neurosci* 1996;8:353–364.
- 167 Rowe JB, Owen AM, Johnsrude IS, Passingham RE: Imaging the mental components of a planning task. *Neuropsychologia* 2001;39:315–327.
- 168 Morris RG, Ahmed S, Syed GM, Toone BK: Neural correlates of planning ability: frontal lobe activation during the Tower of London test. *Neuropsychologia* 1993;31:1367–1378.
- 169 Rezai K, Andreasen NC, Alliger R, Cohen G, Swayze V 2nd, O'Leary DS: The neuropsychology of the prefrontal cortex. *Arch Neurol* 1993;50:636–642.
- 170 Cazalis F, Valabregue R, Pelegrini-Issac M, Asloun S, Robbins TW, Granon S: Individual differences in prefrontal cortical activation on the Tower of London planning task: implication for effortful processing. *Eur J Neurosci* 2003;17:2219–2225.
- 171 Fincham JM, Carter CS, van Veen V, Stenger VA, Anderson JR: Neural mechanisms of planning: a computational analysis using event-related fMRI. *Proc Natl Acad Sci USA* 2002;99:3346–3351.
- 172 Lazeron RH, Rombouts SA, Machielsen WC, Scheltens P, Witter MP, Uylings HB, Barkhof F: Visualizing brain activation during planning: the Tower of London test adapted for functional MR imaging. *AJNR Am J Neuroradiol* 2000;21:1407–1414.
- 173 Newman SD, Carpenter PA, Varma S, Just MA: Frontal and parietal participation in problem solving in the Tower of London: fMRI and computational modeling of planning and high-level perception. *Neuropsychologia* 2003;41:1668–1682.
- 174 van den Heuvel OA, Groenewegen HJ, Barkhof F, Lazeron RHC, van Dyck R, Veltman DJ: Frontostriatal system in planning complexity: a parametric functional magnetic resonance version of Tower of London task. *Neuroimage* 2003;18:367–374.
- 175 Pujol J, Torres L, Deus J, Cardoner N, Pi-farré J, Capdevila A, Vallejo J: Functional magnetic resonance imaging study of frontal lobe activation during word generation in obsessive-compulsive disorder. *Biol Psychiatry* 1999;45:891–897.
- 176 Ursu S, Carter CS: An initial investigation of the orbitofrontal cortex hyperactivity in obsessive-compulsive disorder: exaggerated representations of anticipated aversive events? *Neuropsychologia* 2009;47:2145–2148.
- 177 Martinot JL, Hardy P, Feline A, Huret JD, Mazoyer B, Attar-Levy D, Pappata S, Syrota A: Left prefrontal glucose hypometabolism in the depressed state: a confirmation. *Am J Psychiatry* 1990;147:1313–1317.
- 178 Sawle GV, Hymas NF, Lees AJ, Frackowiak RS: Obsessional slowness: functional studies with positron emission tomography. *Brain* 1991;114:2191–2202.
- 179 Rauch SL, Shin LM, Dougherty DD, Alpert NM, Fischman AJ, Jenike MA: Predictors of fluvoxamine response in contamination-related obsessive compulsive disorder: a PET symptom provocation study. *Neuropsychopharmacology* 2002;27:782–791.
- 180 Lucey JV, Costa DC, Busatto G, Pilowsky LS, Marks IM, Ell PJ, Kerwin RW: Caudate regional cerebral blood flow in obsessive-compulsive disorder, panic disorder and healthy controls on single photon emission computerised tomography. *Psychiatry Res* 1997;74:25–33.
- 181 Bartha R, Stein MB, Williamson PC, Drost DJ, Neufeld RW, Carr TJ, Canaran G, Densmore M, Anderson G, Siddiqui AR: A short echo ¹H spectroscopy and volumetric MRI study of the corpus striatum in patients with obsessive-compulsive disorder and comparison subjects. *Am J Psychiatry* 1998;155:1584–1591.
- 182 Alptekin K, Degirmenci B, Kivircik B, Durak H, Yemez B, Derebek E, Tunca Z: Tc-99m HMPAO brain perfusion SPECT in drug-free obsessive-compulsive patients without depression. *Psychiatry Res* 2001;107:51–56.