Bronchoepithelial Expression of CXCR1 and CXCR2 Does Not Facilitate Transepithelial Migration of Neutrophils

Christian Schulz Benjamin Stoelcker Brigitte Ruhland Nico Jentsch
Andreas Steege

Pneumologie, Klinik und Poliklinik für Innere Medizin II, Klinikum der Universität Regensburg,
Regensburg, Germany

Key Words

Airway inflammation • Chemokine receptors • Chronic obstructive pulmonary disease • Epithelial cells • Inflammation • Neutrophils • Transepithelial migration

Abstract

Background: Neutrophilic airway inflammation is one of the key features of chronic obstructive pulmonary disease (COPD). The chemokine receptors 1 (CXCR1) and 2 (CXCR2) are expressed in the bronchial mucosa during chronic inflammation and might be of importance for transepithelial migration of neutrophils. **Objectives:** This study addressed the role of bronchoepithelial CXCR1 and CXCR2 expression with respect to transepithelial migration of neutrophils. **Methods:** Primary bronchial epithelial cells (PBEs) derived from COPD patients and healthy controls as well as transiently CXCR1- and CXCR2-transfected Calu-6 cells were used for transepithelial migration assays of neutrophils under various conditions. Epithelial CXCR1 and CXCR2 expression was verified by means of flow cytometry. **Results:** Transepithelial migration of neutrophils was significantly increased following lipopolysaccharide pretreatment of epithelial cells. Tran-

sient transfection of CXCR1 and CXCR2 neither augmented the transepithelial migration of neutrophils, nor did the selective blockade of CXCR1 and CXCR2 have any significant effect on neutrophilic transepithelial migration. In addition, no differences were found in PBEs and neutrophils derived from healthy controls and COPD patients. **Conclusions:** The data of the present study do not support the hypothesis that bronchoepithelial expression of CXCR1 and/or CXCR2 facilitate transepithelial migration of neutrophils.

Copyright © 2011 S. Karger AG, Basel

Introduction

The pathophysiology of chronic obstructive pulmonary disease (COPD) includes persistent inflammation of the airways and lung parenchyma, which is caused by an abnormal response of the lung to noxious gases and inhaled particles [1]. Typically, the cellular composition of the airway infiltrates consists of neutrophils, macrophages, and CD8+ T cells [1, 2]. There is convincing evidence that neutrophils are one of the primary effector cells in COPD [3–7]. The number of neutrophils is in-

creased in bronchoalveolar lavage fluid as well as in induced sputum samples and airways from COPD patients [6, 8, 9]. The amount of neutrophils correlates with disease severity and progression in these patients [3, 8]. Neutrophil recruitment to the airways is induced by chemotactic factors such as CXC ligand 8 (CXCL8) and leukotriene B₄ (LTB₄), and both mediators are often elevated in the airways of patients with COPD [10–13]. In addition, neutrophils were predominant in the airway lumen, but a lack of increase in these cells in the subepithelium can be noted in bronchial biopsies of COPD patients [14]. This observation may be due to accelerated transepithelial migration of neutrophils from the circulation into the airway lumen, although this has not been investigated in detail. It is well known that increased LTB₄ and CXCL8 levels derive from alveolar macrophages and epithelial cells, respectively. This, however, should also result in increased levels of neutrophils in the subepithelium, which is not the case, and points towards potential local mechanisms that facilitate and regulate transepithelial migration of neutrophils to the site of inflammation.

CXCL8 and other neutrophil chemoattractants belong to the ELR-CXC chemokines, which are ligands of the CXC chemokine receptor 1 (CXCR1) or 2 (CXCR2). CXCR1 is the receptor for CXCL8 and CXCL6 (granulocyte chemoattractant protein-2), and CXCR2 binds these as well as all other ELR-CXC chemokines; CXCL8 binds with high affinity to both receptors, while the others are ligands of lower affinity [15–18]. Recent studies focusing on urinary tract infections demonstrated that transepithelial neutrophil migration is CXCR1 dependent *in vitro* and defective in CXCR1-knockout mice [19]. Along this line, it was hypothesized that CXCR1 and CXCR2 expression on bronchial epithelial cells is relevant for the interaction between neutrophils and the airway mucosal barrier. It was speculated that these receptors are of importance for neutrophil migration across airway epithelial layers, and, consequently, a potential future therapeutic target for the treatment of neutrophilic airway inflammation, e.g. in COPD patients.

Materials and Methods

Flow Cytometry

CXCR1 and CXCR2 receptor proteins were detected on the cell surface of primary cells and cell lines using specific antibodies (anti-CXCR1 R&D Systems, MAB330, and anti-CXCR2 R&D Systems, MAB331). Primary cells or transfected cell lines were harvested and washed once and resuspended in PBS. Specific pri-

mary antibodies were added (1:50) and the suspension was incubated for 30 min on ice. After additional washing with PBS, labeled secondary antibodies were added to the suspension (1:50). The suspension was incubated for 30 min at 4°C in the dark. After incubation, cells were washed in PBS and resuspended in PBS prior to analysis. Fluorescence was measured with a BD FACS-Calibur flow cytometer (BD, Franklin Lakes, N.J., USA) using a 488-nm excitation laser. The number of cells stained was calculated using CellQuest software (BD).

Cloning of CXCR1 and CXCR2 and Transfection of Calu-6 Cells

For transient and stable transfection, the coding sequences of CXCR1 and CXCR2 were cloned in the pDsred-N1 backbone (Clontech, Mountain View, Calif., USA) or pcGFP-N1 (Clontech). The coding sequences from CXCR1 (NM_000634.2) and CXCR2 (NM_001557.3) were amplified using CXCR1_fw 5'-GCTAGC-GCTGAACTGAAGAGGACATG-3' and CXCR1_rev 5'-CTC-GAGGTTGGAAGAGACATTGACA-3' with attached *NheI* and *XhoI* restriction sites. A specific CXCR2 sequence was amplified using CXCR2_fw 5'-CTCGAGAGTAGTGGAAGTGTGCC-3' and CXCR2_rev 5'-CTCGAGAGTAGTGGAAGTGTGCC-3' with attached *NheI* and *XhoI* restriction sites. cDNA from neutrophilic granulocytes served as template DNA. Amplicons were subcloned into the pCR2.1 backbone using the TOPO-Cloning kit (Invitrogen, Darmstadt, Germany). After digestion with *NheI* and *XhoI*, the fragments were cloned into the expression vectors. The vectors are named CXCR1-pDsred-N1 and CXCR2-pcGFP-N1. The accuracy of inserts was checked by sequencing.

For transient overexpression of CXCR1 and CXCR2, Calu-6 cells were transfected with the CXCR1-pDsred-N1 and CXCR2-pcGFP-N1. After reaching 80% confluency, cells were transfected with 4 µg plasmid DNA in a cell culture flask. DNA was mixed in a 1:2 ratio with FuGENE6 transfection reagent (Roche, Mannheim, Germany). After 30 min of incubation in serum-free media, the mix was added to the cells. After 24 h, cells were checked for CXCR1 and CXCR2 expression using flow cytometry. Positive cells were used for transmigration assays.

Granulocyte Isolation

Informed consent was obtained from all patients and healthy volunteers according to the procedures approved by the local ethics committee. Neutrophilic granulocytes were isolated from healthy volunteers and COPD patients; 20 ml of whole blood was transferred to a dextran gradient [18 ml Deltadex 60 (DeltaSelect) plus 2 ml citrate dextrose solution; Sigma, Taufkirchen, Germany]. Erythrocytes were removed by centrifugation. The supernatant containing lymphocytes and sera was transferred to 20 ml Ficoll (Ficoll-Paque Plus). After 20 min of centrifugation at 600 g (no break), the neutrophilic granulocytes were at the bottom of the gradient. The supernatant containing the plasma and lymphocytes was removed. Remaining erythrocytes were removed from the pellet with erythrocyte lysis buffer (Buffer EL; Qiagen, Hilden, Germany) plus a 5-min incubation at 4°C and centrifugation. The neutrophilic granulocytes were resuspended in normal growth media. Cell numbers were determined using trypan blue exclusion (Sigma) and a Neubauer chamber.

Transmigration Assay

To test granulocyte migration, transmigration assays were performed. Tissue culture (TC) inserts with 3.0- μ m pore size (Greiner Bio-One, Frickenhausen, Germany) were coated with rat tail collagen I (Sigma-Aldrich). CXCR1- and CXCR2-positive Calu-6 cells were seeded on the outer side of the TC insert membrane, which represents the apical side in the migration assays. The TC inserts were placed into 24-well plates filled with normal growth media (Calu-6 medium). A volt ohmmeter (World Precision Instruments, Berlin, Germany) was used to verify the tightness of the cell layer; 12 h before the assay, half of the TC inserts were pretreated from the apical side with 5 μ g/ml lipopolysaccharide (LPS; source: *Escherichia coli* 055:B5; Sigma-Aldrich). LPS was removed completely before the migration assay was performed. Isolated granulocytes (250,000) were added to the basal side of each TC insert membrane. After 3 h of incubation at 37°C, the TC inserts were removed from the wells. The remaining growth medium was transferred to fresh cups. The number of migrated cells was assessed using the CyQuant Cell Proliferation Kit (Invitrogen, Darmstadt, Germany) according to the manufacturer's protocol. Specific antibodies (anti-CXCR1 R&D Systems, MAB330, and anti-CXCR2 R&D Systems, MAB331) were used to block the receptors. Antibodies were applied 12 h before the assay was performed to the apical side of the TC inserts. The assay antibodies had been completely removed in advance. Untransfected Calu-6 cells or naive primary bronchial epithelial cells (PBECs) served as controls. Experiments were performed twice for each condition and granulocyte isolation.

Subjects

Patients who were referred to our clinic for flexible bronchoscopy for various reasons were screened to be included in the present study. Selection of COPD patients was based on the definition and classification given by the Global Initiative for Chronic Obstructive Lung Disease (GOLD) [1]. In brief, the following inclusion criteria had to be met: stable airflow limitation, FEV₁ <70% of predicted, reversibility <10% of predicted FEV₁ after 200 μ g inhaled salbutamol and a smoking history of >10 pack years. None of the patients with COPD had a history of atopy or evidence of atopy on skin prick testing for common aeroallergens. Control subjects were nonsmokers with normal lung function and no history of airway disease. Exclusion criteria were a history of respiratory tract infection or exacerbation of airway disease within the previous 8 weeks, and systemic or topic corticosteroid treatment during the previous 12 weeks. All patients gave written informed consent, and the study was approved by the local ethics committee (approval No. 03/180).

Patients underwent fiberoptic bronchoscopy under light sedation. Bronchial epithelium was obtained by gentle brushing of segmental and subsegmental bronchi under direct visual guidance by means of a protected brush. Each patient had 20–30 brushes of the bronchial epithelium. Brushing was accomplished by a very light gliding along the surfaces of the airways. Brushes were immediately placed in ice-cold bronchial epithelial growth medium and transported directly to the laboratory for further processing.

Culture of PBECs

Brushes were vortexed vigorously, and the cell suspension harvested was filtered through a 200- μ m filter (Millipore; Billerica,

Mass., USA) to remove mucus and cellular debris, and then treated with 4.8 U/ml Dispase II (Roche, Mannheim, Germany) to eliminate cell clumping. The cells were then centrifuged at 150 g for 10 min, and the pellet was resuspended in cell culture medium. Cells were counted using a hemocytometer. Cell viability was determined by trypan blue exclusion. Cells were resuspended in serum-free bronchial epithelial cell growth medium (Promocell, Freiburg, Germany) supplemented with a variety of growth factors, including bovine pituitary extract (0.052 mg/ml), recombinant human epidermal growth factor (0.5 ng/ml), insulin (5 μ g/ml), hydrocortisone (0.5 μ g/ml), epinephrine (0.5 μ g/ml), triiodothyronine (6.5 ng/ml), transferrin (0.01 mg/ml) and retinoic acid (0.1 ng/ml). In addition, the medium contained penicillin G (100 U/ml), streptomycin (100 μ g/ml) and amphotericin B (0.25 μ g/ml; Invitrogen, Darmstadt, Germany). Cultures were maintained in a humidified atmosphere at 37°C in air/carbon dioxide (95/5% vol/vol). The identity of the epithelial cells was confirmed in all cultures by light microscopy and in randomly selected cultures by immunocytochemical staining for cytokeratin expression using a pancytokeratin antibody (clone KL1) directed against cytokeratin types 1, 2, 5–8, 11, 14, 16, and 18 (Immunotec, Marl, Germany). The monoclonal antifibroblast antibody FibAS02 (Dianova, Hamburg, Germany) was used to exclude contamination by fibroblasts [20]. The staining was performed according to Kunz-Schughart et al. [21], as described in detail previously. In general, cell culture experiments were performed after the first passage and no higher passage cell cultures were used.

Calu-6 Cell Culture

Calu-6 cells (human lung fibroblasts) were maintained in MEM with EBSS and L-glutamine (PAN Biotech) supplemented with sodium pyruvate, non-essential amino acids (1 mM; Biochrome, Berlin, Germany), penicillin (50 U/ml), streptomycin (50 μ g/ml) and 10% FCS (PAA Laboratories, Pasching, Austria). Cells were grown at 37°C in humidified air with 5% CO₂.

Statistical Analysis

For statistical analyses, the Kruskal-Wallis rank sum test was used to compare data between groups, followed by the Mann-Whitney rank sum test using SigmaStat for Windows V2.03. Differences were considered to be statistically different if $p < 0.05$. x-fold migration was calculated for each value compared to the mean of the control group.

Results

Subjects

Nine COPD and 21 control subjects without airflow limitation were included in the study. All patients were referred for bronchoscopy for clinical reasons. All patients underwent preoperative diagnostic procedures for suspected peripheral or central carcinoma of the bronchus. In addition, based on the patients' history and files, no other conditions known or suggested to be associated with altered immune responses were detected. In particular, no signs of diabetes, rheumatoid arthritis, in-

Fig. 1. Representative FACS analyses of CXCR1 (**a, b**) and CXCR2 (**c, d**) expression on the surface of naive PBECs from a COPD patient (**a, c, e, g**) and a non-COPD donor (**b, d, f, h**). Note that there is an even higher expression of CXCR1 in PBECs in both cases. Isotype staining (ISO) and unstained cells served as controls. FSC = Forward scatter; Quad = quadrant; UL = upper left; UR = upper right; LL = lower left; LR = lower right.

flammatory bowel diseases, or myeloproliferative syndromes were found. The COPD patients were classified as stage I (n = 3), stage II (n = 2), stage III (n = 1) and stage IV (n = 3) based on the GOLD classification [1]. All were exsmokers and aged between 45 and 70 years.

COPD patients were clinically stable for at least 8 weeks prior to bronchoscopy without any changes in medication. Control subjects, aged between 42 and 68 years, had no evidence of chronic disease and were free of medication.

Fig. 2. Representative FACS analysis of CXCR1 and CXCR2 expression on the surface of Calu-6 cells following transient transfection. **a, c** CXCR1 and CXCR2 expression on untransfected Calu-6 cells. **b, d** CXCR1 and CXCR2 expression after transient transfection. Note that no expression was detected on naive Calu-6 cells. Iso-type staining (ISO; **e**) and unstained cells (**f**) served as controls. FSC = Forward scatter; Quad = quadrant; UL = upper left; UR = upper right; LL = lower left; LR = lower right.

Surface Expression of CXCR1 and CXCR2

First we confirmed the expression of CXCR1 and CXCR2 on the surface of PBECs using flow cytometry. PBECs from COPD patients and controls (non-COPD patients) were stained for CXCR1 and CXCR2 using specific antibodies. In figure 1a, c, representative FACS analyses are shown. On PBECs from COPD patients, expression of CXCR1 was increased compared to CXCR2 expression on PBECs: 60% of PBECs analyzed were positive for CXCR1 whereas only 3% of the cells were CXCR2 positive. In PBECs from control patients, 37% of the cells expressed CXCR1, whereas CXCR2 expression amounted to 3%. All of the PBECs showed similar mean fluorescence intensity ratios; the calculated ratio for CXCR1 on PBECs was 22.92 (control 1) and 25.53 (control 2) com-

pared to 26.1 on PBECs from a COPD patient. For CXCR2, the respective ratios were 4.26 (control 1) and 4.53 (control 2) compared to 4.82 (COPD patient).

Surface Expression of CXCR1 and CXCR2 in Calu-6 Cells following Transient Transfection

Calu-6 cells were transiently transfected with CXCR1 or CXCR2 expression vectors. Before using these cells in the migration assay, surface expression of CXCR1 and CXCR2 was confirmed by means of FACS analysis. The analysis was performed three times with cells from independent transfections with similar results. In figure 2, representative FACS analyses of the expression of both receptors on naive Calu-6 and following transient transfection are shown. Naive Calu-6 cells show neither

Fig. 3, 4. Transmigration assay of neutrophilic granulocytes from healthy donors (3) and COPD patients (4) in combination with transiently transfected Calu-6 cells. To determine the transmigration ability of neutrophilic granulocytes from healthy donors (3) and COPD patients (4), these cells were added to the basal side of transiently transfected Calu-6 cells. Shown is the x-fold migration of the neutrophilic granulocytes after 3 h of coculture. Un-

transfected and untreated Calu-6 cells served as controls. Experiments were performed twice for each condition and granulocyte isolation. Each box represents the x-fold change compared to control; the ends of the boxes define the 25th and 75th percentiles. Error bars define the 5th and 95th percentiles. Outliers outside the 5th and 95th percentiles are also shown. NS = Nonsignificant.

CXCR1 nor CXCR2 expression (fig. 2a, c). CXCR1 expression was higher on the Calu-6 cells compared to CXCR2. After transient transfection, 40% of the Calu-6 cells showed CXCR1 expression, with 18% of the cells showing CXCR2 expression. This expression pattern is comparable with the expression on the surface of naive PBECs. Subsequently, the transiently transfected Calu-6 cells were used in transmigration assays in combination with neutrophilic granulocytes from either healthy controls or COPD patients.

Transmigration Assays of Neutrophilic Granulocytes in Combination with Transiently CXCR1- and CXCR2-Transfected Calu-6 Cells

The resistance of the transiently transfected Calu-6 cell layer was measured using a voltmeter to confirm the tightness of the cell layer (data not shown). Neutrophilic granulocytes from COPD patients or healthy volunteers were added to the basal side of the transfected cells and cocultured for 3 h. In figure 3, the x-fold migration of neutrophils from control patients is shown. LPS stimulation significantly increased the transmigration of neutro-

phils resulting in a 1.4-fold (range 1.05–1.87) increase. Similar findings were seen for CXCR1 overexpression following LPS stimulation (1.3-fold; range 1.08–2.11) and unstimulated conditions. For CXCR2, transmigration of neutrophils was significantly lower following LPS treatment (median = 0.98; range 0.89–1.48) compared to control cells and unstimulated conditions. When neutrophils from COPD patients were used instead, no significant increase in the transmigration of neutrophils was observed with respect to CXCR1 and CXCR2 overexpression under either condition (fig. 4). In summary, as shown in figures 3 and 4, there is no augmentation in the migration of neutrophilic granulocytes from either healthy donors or COPD patients following CXCR1 and CXCR2 overexpression.

Transmigration Assays of Neutrophilic Granulocytes Using PBECs

PBECs derived from control subjects and COPD patients were used in combination with neutrophils from healthy controls and COPD patients to perform transmigration assays in both the absence and presence of spe-

cific CXCR1 and CXCR2 antibodies. The antibodies were applied to PBECs 12 h prior to incubation with neutrophilic granulocytes.

As shown in figure 5a, LPS stimulation of PBECs derived from non-COPD patients resulted in a 1.9-fold (median = 1.89, range 0.24–4.70) increase in transepithelial migration of neutrophils. In the presence of anti-CXCR1 and anti-CXCR2 antibodies, the increase in LPS-stimulated transepithelial migration of neutrophils derived from control subjects was 1.5-fold (range 0.62–3.26; non-significant compared to LPS stimulation). In the presence of anti-CXCR1 or anti-CXCR2, the increase was 1.9-fold (range 0.88–3.42) and 1.08-fold (range 1.0–1.14), respectively (data not shown).

Similar results (fig. 5b) were obtained when PBECs derived from COPD patients were used. LPS stimulation resulted in a 1.8-fold (range 0.88–4.62) increase in the transmigration of neutrophils derived from COPD patients ($p < 0.05$ vs. unstimulated control). Blockade of both receptors revealed a 2.1-fold increase in the transepithelial migration of neutrophils (range 1.47–3.13; non-significant compared to LPS stimulation), whereas a 1.4-fold (range 1.22–1.96) increase for CXCR1 and a 2.1-fold (range 1.47–3.13) increase were observed when only CXCR1 or CXCR2 were blocked (data not shown).

Figure 5 summarizes the results demonstrating that the CXCR1 and/or CXCR2 blockade had no significant effect on transepithelial migration of neutrophilic granulocytes under either condition.

Discussion

In the present report, the potentially CXCR1- and CXCR2-dependent transepithelial migration of neutrophils was investigated by using different cell culture models as well as airway epithelial cell lines and PBECs derived from healthy nonsmokers and COPD patients. Recent studies suggest an accelerated transepithelial migration of neutrophils from the circulation to the airway lumen, which could, at least in part, explain why airway inflammation dominated by neutrophils is often found in COPD patients.

Significant CXCR1 and CXCR2 expression has already been demonstrated in bronchial epithelial cells [22–24]. Qiu et al. [23] showed upregulation of epithelial CXCR2 mRNA expression in COPD patients with severe exacerbation. However, in the same paper, neither in stable COPD patients nor in patients with severe COPD exacerbation CXCR1 expression was detected. In a more

Fig. 5. **a** Transmigration assay of neutrophilic granulocytes from healthy donors in combination with blocking of CXCR1 and CXCR2 in PBECs isolated from healthy donors (control $n = 33$; LPS $n = 35$; aCXCR1 + aCXCR2 + LPS $n = 14$). **b** Transmigration assay of neutrophilic granulocytes from COPD patients in combination with blocking of CXCR1 and CXCR2 in PBECs isolated from COPD patients (control $n = 23$; LPS $n = 24$; aCXCR1 + aCXCR2 + LPS $n = 7$). x-fold migration of the neutrophilic granulocytes after 3 h of coculture. For further information, see legend to figures 3, 4. aCXCR1 = Anti-CXCR1; aCXCR2 = anti-CXCR2.

recent paper, a robust bronchoepithelial expression of both receptors was seen in COPD patients across all stages of severity [22]. This is in line with the results of the present study demonstrating cell surface expression of CXCR1 and CXCR2 in PBECs from healthy donors and COPD patients, respectively. Therefore, it is evident that the bronchial mucosa expresses both chemokine receptors, which may be involved in the transepithelial migration of neutrophils.

Godaly et al. [19] were able to demonstrate that the expression of functional IL-8 receptors is crucial for neutrophils to cross the epithelial barrier. This was shown in IL-8 receptor-knockout mice with urinary tract infection as well as in human uroepithelial cell lines. In addition, it was speculated that IL-8 dimer formation, which is required for IL-8 receptor activation, forms a bridge between the neutrophils and the epithelial cells, thereby enabling transepithelial cell migration. This hypothesis was supported by a recently published paper using molecular dynamic simulations, thereby showing that electrostatic interactions are favorable for CXCL8 homodimer formation, which could have substantial consequences biologically [25].

However, the results of the present study do not reinforce this view with respect to the transepithelial migra-

tion of neutrophils across the bronchoepithelial barrier. Different experimental approaches were realized to substantiate the hypothesis. Transient CXCR1 and CXCR2 transfections of Calu-6 cells, neither of which express CXCR1 or CXCR2 in the control situation, were performed and used for transmigration experiments. In addition, PBECs from healthy nonsmokers and COPD patients, with confirmed IL-8 receptor expression, as well as neutrophils from both cohorts were differentially applied for migration assays. Anti-CXCR1 and anti-CXCR2 control experiments confirmed the negative results found. Therefore, several lines of evidence were generated demonstrating that the migration of neutrophils through airway epithelial cell layers is not controlled and/or facilitated by CXCR1 and CXCR2 expression.

Based on the results presented here and available data in the literature, the effects of CXCR1 and CXCR2 on respiratory epithelial cells remain obscure. According to the evidence presented for IL-8 receptor-expressing intestinal epithelial cells, one might speculate that CXCR1 and/or CXCR2 play an important role in the maintenance of the mucosal epithelial surface barrier, thus affecting the infiltration of inflammatory cells into the host that may maintain airway inflammation. This was at least demonstrated in an *in vitro* migration model using intestinal Caco-2 and HT-29 cells [26]. Whether the same holds true for respiratory epithelial cells requires further investigation and was beyond the scope of the present study.

Neutrophil recruitment during COPD exacerbations appears to be partially mediated by the two important neutrophil chemoattractants CXCL5 (ENA-78) and CXCL8 (IL-8). Upregulation of their receptors, CXCR1 and CXCR2, has been observed in bronchial biopsy specimens in severe COPD exacerbations [23], which is in line with our results. In addition, CXCR1 and CXCR2 are expressed on a broad range of leukocytes, most notably neutrophilic granulocytes. The physiological role of these receptors is mediating neutrophil recruitment, which has been described in humans as well as in primates, rodents and rabbits. Recent data provide evidence that the expression of CXCR1 and CXCR2 in neutrophils is maintained in COPD patients when compared to healthy controls or healthy smokers, indicating that COPD *per se* does not alter the expression of chemoattractant receptors [27].

Importantly, treatment with a CXCR2 antagonist (SCH527123) blocked pulmonary neutrophilia associated with the repeat bronchoscopy procedure [28], suggesting that CXCR2 inhibition is important for the inflammatory events occurring in the distal lungs of human

COPD patients. Similar results were found in ozone-induced neutrophil recruitment to the lungs in healthy humans [29]. Thus, there is overwhelming evidence that IL-8 receptor expression on neutrophils is important for the development of the neutrophilic lung inflammation present in COPD. The aforementioned studies in animals and humans have clearly demonstrated that CXCR1/CXCR2 antagonism inhibits pulmonary and airway neutrophilia [28, 29]. Since the present study shows that the blockade of bronchoepithelial CXCR1/CXCR2 expression has no impact on transepithelial neutrophil migration, the important site of CXCR1/CXCR2 blockade has to be the neutrophil itself. This hypothesis is further supported by the observation that the CXCR2 antagonist SCH527123 reversibly decreases peripheral neutrophil counts in humans. This implies that the CXCR2 antagonism inhibits neutrophil egress from the bone marrow and migration from peripheral blood to tissue [29]. Neutrophils are generated in large numbers in the bone marrow; mature neutrophils circulate in the bloodstream for several hours and are recruited to tissues and sites of infection via a multistep process [30, 31]. There is convincing evidence that the mechanisms of neutrophilic trafficking are complex and probably disease and stimulus specific in connection with a certain degree of redundancy in the mechanisms involved [30]. To further address this issue in the future, it would be important to use disease-specific models in conjunction with disease- and phenotype-specific neutrophils.

There are some limitations of the present study. Firstly, the data presented are from *in vitro* cell culture experiments, which do not necessarily reflect the complex *in vivo* situation seen in COPD patients. Secondly, CXCR1 and CXCR2 expression of neutrophils was not assessed on a regular basis before each experiment. Although unlikely, this could also have influenced our results.

In summary, the present study generated substantial evidence indicating that airway epithelial expression of both IL-8 receptors CXCR1 and CXCR2 does not facilitate transepithelial migration of neutrophilic granulocytes. This, however, does not contradict the importance of CXCR1 and CXCR2 in neutrophilic lung inflammation when expressed on polymorphonuclear granulocytes.

Acknowledgment

This study was supported by a grant (SCHU-1606/2-1) from the Deutsche Forschungsgemeinschaft (German Research Foundation).

References

- 1 Global Initiative for Chronic Obstructive Lung Disease: Global Strategy for the Diagnosis, Management and Prevention of Chronic Obstructive Pulmonary Disease. NHLBI/WHO Workshop Report, updated 2009 (www.goldcopd.org).
- 2 Barnes PJ, Shapiro SD, Pauwels RA: Chronic obstructive pulmonary disease: molecular and cellular mechanisms. *Eur Respir J* 2003; 22:672–688.
- 3 Stancescu D, Sanna A, Veriter C, Kostianev S, Calcagni PG, Fabbri LM, Maestrelli P: Airways obstruction, chronic expectoration, and rapid decline of FEV₁ in smokers are associated with increased levels of sputum neutrophils. *Thorax* 1996;51:267–271.
- 4 Hunninghake GW, Crystal RG: Cigarette smoking and lung destruction. Accumulation of neutrophils in the lungs of cigarette smokers. *Am Rev Respir Dis* 1983;128:833–838.
- 5 Thompson AB, Daughton D, Robbins RA, Ghafouri MA, Oehlerking M, Rennard SI: Intraluminal airway inflammation in chronic bronchitis. Characterization and correlation with clinical parameters. *Am Rev Respir Dis* 1989;140:1527–1537.
- 6 Rutgers SR, Timens W, Kaufmann HF, van der Mark TW, Koeter GH, Postma DS: Comparison of induced sputum with bronchial wash, bronchoalveolar lavage and bronchial biopsies in COPD. *Eur Respir J* 2000;15:109–115.
- 7 Lapperre TS, Willems LN, Timens W, Rabe KF, Hiemstra PS, Postma DS, Sterk PJ: Small airways dysfunction and neutrophilic inflammation in bronchial biopsies and BAL in COPD. *Chest* 2007;131:53–59.
- 8 Hodge SJ, Hodge GL, Holmes M, Reynolds PN: Flow cytometric characterization of cell populations in bronchoalveolar lavage and bronchial brushings from patients with chronic obstructive pulmonary disease. *Cytometry B Clin Cytom* 2004;61:27–34.
- 9 Isajevs S, Taivans I, Svirina D, Strazda G, Kopeika U: Patterns of inflammatory responses in large and small airways in smokers with and without chronic obstructive pulmonary disease. *Respiration* 2011;81:362–371.
- 10 Corhay JL, Henket M, Nguyen D, Duysinx B, Sele J, Louis R: Leukotriene B₄ contributes to exhaled breath condensate and sputum neutrophil chemotaxis in COPD. *Chest* 2009; 136:1047–1054.
- 11 Keatings VM, Collins PD, Scott DM, Barnes PJ: Differences in interleukin-8 and tumor necrosis factor- α in induced sputum from patients with chronic obstructive pulmonary disease or asthma. *Am J Respir Crit Care Med* 1996;153:530–534.
- 12 Yamamoto C, Yoneda T, Yoshikawa M, Fu A, Tokuyama T, Tsukaguchi K, Narita N: Airway inflammation in COPD assessed by sputum levels of interleukin-8. *Chest* 1997;112: 505–510.
- 13 Schulz C, Wolf K, Harth M, Kratzel K, Kunz-Schughart L, Pfeifer M: Expression and release of interleukin-8 by human bronchial epithelial cells from patients with chronic obstructive pulmonary disease, smokers, and never-smokers. *Respiration* 2003;70: 254–261.
- 14 Saetta M, Di Stefano A, Maestrelli P, Ferrarresso A, Drigo R, Potena A, Ciaccia A, Fabbri LM: Activated T-lymphocytes and macrophages in bronchial mucosa of subjects with chronic bronchitis. *Am Rev Respir Dis* 1993; 147:301–306.
- 15 Schumacher C, Clark-Lewis I, Baggiolini M, Moser B: High- and low-affinity binding of GRO α and neutrophil-activating peptide 2 to interleukin 8 receptors on human neutrophils. *Proc Natl Acad Sci USA* 1992; 89:10542–10546.
- 16 Holmes WE, Lee J, Kuang WJ, Rice GC, Wood WI: Structure and functional expression of a human interleukin-8 receptor. *Science* 1991;253:1278–1280. *J Immunol* 2009; 183:2895–2897.
- 17 Wuyts A, Proost P, Lenaerts JP, Ben Baruch A, Van Damme J, Wang JM: Differential usage of the CXCR chemokine receptors 1 and 2 by interleukin-8, granulocyte chemotactic protein-2 and epithelial-cell-derived neutrophil attractant-78. *Eur J Biochem* 1998;255: 67–73.
- 18 Baggiolini M: Chemokines in pathology and medicine. *J Intern Med* 2001;250:91–104.
- 19 Godaly G, Hang L, Frendus B, Svanborg C: Transepithelial neutrophil migration is CXCR1 dependent in vitro and is defective in IL-8 receptor knockout mice. *J Immunol* 2000;165:5287–5294.
- 20 Saalbach A, Aneregg U, Bruns M, Schnabel E, Herrmann K, Hausteiner UF: Novel fibroblast-specific monoclonal antibodies: properties and specificities. *J Invest Dermatol* 1996;106:1314–1319.
- 21 Kunz-Schughart LA, Heyder P, Schroeder J, Kneuchel R: A heterologous 3-D coculture model of breast tumor cells and fibroblasts to study tumor-associated fibroblast differentiation. *Exp Cell Res* 2001;266:74–86.
- 22 Di Stefano A, Caramori G, Gnemmi I, Conzoli M, Bristot L, Capelli A, Ricciardolo FL, Magno F, D'Anna SE, Zanini A, Carbone M, Sabatini F, Usai C, Brun P, Chung KF, Barnes PJ, Papi A, Adcock IM, Balbi B: Association of increased CCL5 and CXCL7 chemokine expression with neutrophil activation in severe stable COPD. *Thorax* 2009;64:968–975.
- 23 Qiu Y, Zhu J, Bandi V, Atmar RL, Hattotuwa K, Guntupalli KK, Jeffery PK: Biopsy neutrophilia, neutrophil chemokine and receptor gene expression in severe exacerbations of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2003;168:968–975.
- 24 Farkas L, Hahn MC, Schmoezer M, Jentsch N, Kratzel K, Pfeifer M, Schulz C: Expression of CXCR chemokine receptors 1 and 2 in human bronchial epithelial cells. *Chest* 2005; 128:3724–3734.
- 25 Nesmelova IV, Sham Y, Gao J, Mayo KH: CXCR and CC chemokines form mixed heterodimers: association free energies from molecular dynamics simulations and experimental correlations. *J Biol Chem* 2008;283: 24155–24166.
- 26 Sturm A, Baumgart DC, d'Heureuse JH, Hotz A, Wiedenmann B, Dignass AU: CXCL8 modulates human intestinal epithelial cells through a CXCR1 dependent pathway. *Cytokine* 2005;29:42–48.
- 27 Sapely E, Stockley JA, Greenwood H, Ahmad A, Bayley D, Lord JM, Insall RH, Stockley RA: Behavioral and structural differences in migrating peripheral neutrophils from patients with COPD. *Am J Respir Crit Care Med* 2011;183:1176–1186.
- 28 Chapman RW, Minniccozi M, Celly CS, Phillips JE, Kung TT, Hipkin RW, Fan X, Rindgen D, Deno G, Bond R, Gonsiorek W, Billah MM, Fine JS, Hey JA: A novel, orally active CXCR1/2 receptor antagonist, SCH527123, inhibits neutrophil recruitment, mucus production, and goblet cell hyperplasia in animal models of pulmonary inflammation. *J Pharmacol Exp Ther* 2007;322: 486–493.
- 29 Holz O, Khalilieh S, Ludwig-Sengpiel A, Watz H, Stryzak P, Soni P, Tsai M, Sadeh J, Magnussen H: SCH527123, a novel CXCR2 antagonist, inhibits ozone-induced neutrophilia in healthy subjects. *Eur Respir J* 2010; 35:564–570.
- 30 Gane J, Stockley R: Mechanisms of neutrophil transmigration across the vascular endothelium in COPD. *Thorax* 2011, E-pub ahead of print.
- 31 Borregaard N: Neutrophils, from marrow to microbes. *Immunity* 2010;33:657–670.