22q11.2 Distal Deletion Syndrome: Description of a New Case with Truncus Arteriosus Type 2 and Review

L. Garavelli^a S. Rosato^a A. Wischmeijer^{a, c} C. Gelmini^a A. Esposito^a
L. Mazzanti^d F. Franchi^b A. De Crescenzo^e O. Palumbo^f M. Carella^f
A. Riccio^{e, g}

^aClinical Genetics Unit, Obstetric and Paediatric Department, and ^bLaboratory of Genetics, S. Maria Nuova IRCCS Hospital, Reggio Emilia, ^cDepartment of Medical Genetics, Policlinico Sant'Orsola-Malpighi, ^dRare Disease Unit, Paediatric Department, University of Bologna, Bologna, ^eDepartment of Environmental Science, 2nd University of Naples, Caserta, ^fGenetics Unit, IRCCS Casa Sollievo della Sofferenza, San Giovanni Rotondo, and ^gInstitute of Genetics and Biophysics 'A. Buzzati-Traverso', CNR, Naples, Italy

Key Words

22q11.2 distal deletion syndrome · Pre- and post-natal growth retardation · *TBX1* · Truncus arteriosus

Abstract

22q11.2 deletion syndrome is mainly characterized by conotruncal congenital heart defects, velopharyngeal insufficiency, hypocalcemia and a characteristic craniofacial appearance. The etiology in the majority of patients is a 3-Mb recurrent deletion in region 22q11.2. Nevertheless, recently some cases of infrequent deletions with various sizes have been reported with a different phenotype. We report on a patient with congenital heart disease (truncus arteriosus type 2) in whom a de novo 1.3-Mb 22q11.2 deletion was detected by array comparative genomic hybridization. The deletion described corresponds to an atypical and distal deletion which spans low copy repeat (LCR) 4 and is associated with breakpoint sites that do not correspond to known LCRs of 22q11.2. We examine the clinical phenotype of our case and compare our findings with those published in the litera-

ture. The most prevalent clinical features in this type of deletion are a history of prematurity, pre-natal and post-natal growth retardation, slight facial dysmorphic features, microcephaly and developmental delay, with a speech defect in particular. These are clearly different from those found in the classic 22q11.2 deletion syndrome, and we believe that the main differential diagnosis should be with Silver-Russell syndrome. In our case we observe the cardiac phenotype with truncus arteriosus communis usually seen in the classic 22q11.2 deletion syndrome, and so far associated with the *TBX1* gene. Significantly, however, *TBX1* is not included in our patient's deletion. The possible roles of a position effect or other genes are discussed.

Copyright © 2011 S. Karger AG, Basel

Interstitial deletions of chromosome 22q11.2 are the most commonly seen microdeletions observed in humans, with a frequency of approximately 1:4,000–1:8,000 live births. The main clinical phenotypes associated with the 22q11.2 deletion syndrome are the velocardiofacial

syndrome (OMIM 192430) and DiGeorge syndrome (OMIM 188400). Patients with the 22q11.2 deletion present various combinations of the following clinical features: congenital heart disease, in particular conotruncal heart anomalies, typical facial phenotype, palatal abnormalities with velopharyngeal insufficiency, congenital hypocalcemia and immune deficiency. The vast majority of patients with the 22q11.2 microdeletion exhibit a common ~3-Mb deletion while 7% of the individuals have a smaller 1.5-Mb deletion [Carlson et al., 1997]. It was discovered that the 3-Mb as well as the 1.5-Mb deletions occur as a result of nonallelic homologous recombination [Stankiewicz and Lupski, 2002]. The majority of deletions were located within the common ~3-Mb deletion region or were found to overlap with it. 'Atypical deletions' correspond to deletions located in the typical deletion region with varied breakpoints, while 'distal deletions' are located distal to the 3-Mb common deletion region.

So far, only a small number of 'atypical deletions' [Kurahashi et al., 1996, 1997; Garcia-Minaur et al., 2002; Rauch et al., 2005; D'Angelo et al., 2007; Fernandez et al., 2009] and 'distal deletions' [Rauch et al., 1999, 2005; Saitta et al., 1999; Mikhail et al., 2007; Ben-Shachar et al., 2008; Rodningen et al., 2008; Xu et al., 2008; Bruce et al., 2009; Lafay-Cousin et al., 2009] were discovered. Here, we describe the case of a 5-year-old girl with an atypical and distal 22q11.2 deletion slightly overlapping with the 3-Mb recurrent deletion and compare the clinical features of our case with those reported in the literature.

Materials and Methods

DNA Methylation Analysis

DNA methylation of *KCNQ1OT1* and *H19* imprinting control regions of 11p15.5 was analyzed in DNA derived from peripheral blood leukocytes by combined bisulfite restriction analysis, as previously described [Sparago et al., 2007]. Two micrograms of genomic DNA were treated with sodium bisulfite, amplified by polymerase chain reaction (PCR), the PCR product digested with a restriction enzyme containing a CpG dinucleotide in its target sequence and the fragments separated on a polyacrylamide gel.

Microsatellite Analysis

11p15 microsatellite markers *TH*, D11S4046 and D11S922 and chromosome 7 microsatellite markers D7S1818, D7S484, D7S1830, D7S686, D7S1824 and D7S669 were amplified using FAM-labeled and HEX-labeled primers and detected using an ABI 3100 capillary electrophoresis instrument. Data were analyzed using GeneMapper Software. Primers and PCR conditions for the microsatellite markers were obtained from the NCBI UniSTS Database (<http://www.ncbi.nlm.nih.gov/sites/entrez?db=unists>).

Array Comparative Genomic Hybridization

Single nucleotide polymorphism (SNP) array analysis was performed using the GeneChip Human Mapping 250K *NspI* Array (Affymetrix, Santa Clara, Calif., USA), which contains a 25-mer oligonucleotide representing a total of 262,264 SNPs with an average resolution of ~12 kb. Labeling and hybridization were performed in accordance with the manufacturer's protocol while SNP copy number was assessed using the program Genotyping Console Software 4.0 (Affymetrix) by a standard Hidden Markov Model method.

Clinical Case and Results

The girl is the second of 2 children of healthy, nonconsanguineous parents (40-year-old mother and 41-year-old father). She has a healthy brother. Family history was otherwise unremarkable. Pre-natal ultrasound at 18 weeks' gestation showed congenital heart disease (truncus arteriosus). The fetal movements were normal. The karyotype analysis at 400 band resolution on amniocytes was normal, 46,XX. She was born at 35 weeks' gestation with Caesarean section, a birth weight of 1,680 g (<3rd centile), length of 41 cm (<3rd centile) and head circumference of 29 cm (<3rd centile). APGAR scores were 9 and 9 at 1 and 5 min, respectively. She was transferred from the neonatal service to the pediatric cardiac surgery at the age of 32 days and underwent surgery for truncus arteriosus type 2. The psychomotor development was delayed both on the mental and motor scales. She was able to sit at 6 months, started walking at 18 months and produced her first words at 18 months. A developmental assessment during the preschool years revealed mild mental retardation. She showed a harmonic cognitive profile with impairments in verbal and performance tasks. Delays in expressive language were worse than those expected by cognitive level, and were not always related to structural anomalies of the palate. Within the verbal domain, receptive language seemed to be the strongest component. Executive function and attention tests showed impairments of several types.

Because of the growth retardation, she was treated with growth hormone, and is also being treated for hypothyroidism. On examination at the age of 2 years and 3 months and prior to the beginning of the treatment with growth hormone, her head circumference was 43 cm (<-2 SD), height 77 cm (<<3rd centile) and weight 7.050 kg (<<3rd centile). At 4 years and 11 months of age, her head circumference was 46.5 cm (<-2 SD), height 99 cm (3rd-10th centile) and weight 12 kg (<<3th centile). She had microcephaly, triangular face, high forehead, frontal

Fig. 1. Craniofacial phenotype of the probanda.

Fig. 2. Hands of the patient. Note the long fingers.

bossing, horizontal palpebral fissures, epicanthic folds, broad nose, long and flat philtrum, thin upper lip, downturned mouth, micrognathia, pointed chin, posteriorly rotated ears and long fingers (fig. 1, 2). Bone age at 1 year 11 months corresponded to that of 1 year 6 months. Renal ultrasound was normal.

Concerning the molecular investigations performed on this case, microsatellite and combined bisulfite restriction analyses excluded the typical defects of Silver-Russell syndrome (SRS), e.g. chromosome 7 UPD and 11p15.5 copy number or DNA methylation abnormalities, while FISH analysis (probe D22S75) excluded the classical 22q11.2 deletion (data not shown). The DNAs of the probanda and her parents were then analyzed by array comparative genomic hybridization using the GeneChip Human Mapping 250K *Nsp* I Array. This study disclosed a de novo 1.3-Mb 22q11.2 distal deletion from SNP A-2255904 (19,433,551 bp) to SNP A-2110718 (20,747,829 bp) probes, based on UCSC Genome Browser (<http://genome.ucsc.edu/>; March 2006 release), in the probanda. The deletion marginally and distally overlapped the common 22q11.2 deletion (fig. 3, 4).

Discussion

The recent diffusion of genome-wide analytical methods for the identification of copy number variations resulted in an increasing number of reports on rare dele-

tions differing from the more commonly observed ones and those associated with variant phenotypes.

In this report, we describe a patient with atypical and distal 22q11.2 deletion and congenital heart disease (truncus arteriosus type 2) and compare the molecular data and clinical phenotype with those previously reported on this genomic region [Kurahashi et al., 1996, 1997; Rauch et al., 1999, 2005; Saitta et al., 1999; Garcia-Minaur et al., 2002; D'Angelo et al., 2007; Mikhail et al., 2007; Ben-Shachar et al., 2008; Rodningen et al., 2008; Xu et al., 2008; Bruce et al., 2009; Fernandez et al., 2009; Lafay-Cousin et al., 2009; Ogilvie et al., 2009].

Apparently, this is the second case, after the first one described by Ogilvie et al. [2009], with an atypical and distal deletion spanning LCR4, the distal point of the common deleted region and the proximal point of many of the distal deletions. Thus, this present deletion, spanning atypical and distal deletions, represents a new variant deletion, associated with breakpoint sites that do not correspond to known LCRs of 22q11.2. Both deletions are novel deletions and are overlapping some of the reported atypical or distal deletions. In both cases, location of the breakpoints does not coincide with known LCRs, indicating alternative mechanisms in the origin of these deletions, which did not utilize LCRs as recombination substrates.

The most common clinical features associated with the atypical and the distal 22q11.2 deletions are a history of prematurity, pre-natal and post-natal growth retardation as well as developmental delay (tables 1–4). This clinical aspect was already described by Ben-Shachar et al. [2008]

Fig. 3. Schematic overview of the 22q11.2 deletion region [modified from Rauch et al., 2005]. Low copy repeats are depicted as green squares and labeled with A2 to H8, according to Edelmann et al. [1999], McDermid and Morrow [2002], and Shaikh et al. [2000]. Bars below the map depict deletions found in this study and in reported cases. The exact deletion borders are sometimes only estimated in reported cases.

with pre- and post-natal growth restriction found in 83% of his patients. In contrast, pre-natal growth retardation is not a typical feature of the classic 22q11.2 deletion syndrome and post-natal growth retardation has been noted in only 36% of the cases [Ben-Shachar et al., 2008].

We subdivided the reported cases with atypical and distal 22q11.2 deletions into 2 groups: those associated with a 'distal deletion' in the more telomeric region (interval VI–VII, fig. 3 and tables 1, 2) [Rauch et al., 1999, 2005; Saitta et al., 1999; Mikhail et al., 2007; Ben-Shachar et al., 2008; Rodningen et al., 2008; Xu et al., 2008; Bruce et al., 2009; Lafay-Cousin et al., 2009] and those with an 'atypical deletion' in the more centromeric region (interval IV–V, fig. 3 and tables 3, 4) [Kurahashi et al., 1996, 1997; Garcia-Minaur et al., 2002; Rauch et al., 2005; D'Angelo et al., 2007; Fernandez et al., 2009]. The case reported by Ogilvie et al. [2009] is very similar to the present patient, with an 'atypical' and 'distal' 22q11.2 deletion, with the same breakpoints, but it is difficult to do a comparison between the facial phenotypes because these clinical aspects have not been described in detail by Ogilvie et al. We did not consider in our review the 5 patients reported by Jackson et al. [2007], with germline 22q11.2 deletion, because these children had a more distal

deletion including the *IN11* gene and predisposing to development of malignant rhabdoid tumors.

In each of the 2 groups, the clinical features and morphologic phenotypes are similar, although the sample of the second group is too small to allow statistically significant conclusions. In both groups, we observe the presence of prematurity (13 out of 18 patients), with an average of 35 weeks gestation, and pre-natal growth retardation (8 out of 11 newborns). Post-natal growth retardation is present in 11 out of 20 cases.

The facial phenotype of the atypical and distal 22q11.2 deletion patients is characterized by slight dysmorphic features with arched eyebrows, deep-set eyes, broad nose, hypoplastic alae nasi, smooth philtrum, down-turned mouth, micrognathia and pointed chin. In our experience, the phenotype of the atypical and distal 22q11.2 deletion patients is not too evident and tricky to suspect, but clearly distinguishable from that of the classic 22q11.2 deletion syndrome. Instead, the main differential diagnosis is with SRS. Indeed, we and others [Bruce et al., 2009] had initially suspected SRS in distal 22q11.2 cases. However, the molecular analyses for chromosome 7 UPD and 11p15.5 copy number or methylation defects have not revealed any abnormality. Moreover, patients with SRS do

Table 1. Clinical features and review of the literature (interval VI–VII)

	Saitta et al., 1999	Xu et al., 2008	Ben-Shachar et al., 2008	Rodnigen et al., 2008	Bruce et al., 2009	Rauch et al., 2005	Lafay-Cousin et al., 2009	Rauch et al., 2005	Mikhail et al., 2007	Ben-Shachar et al., 2008	Patient	Total
Deleted interval and genes (see fig. 4)	VI IGLV	VI IGLV	VI IGLV	VI IGLV	VI IGLV	VII GNAZ	VI+VII GNAZ	VI+VII GNAZ	VI+VII	VI+VII	VI+VII	
22q11.2 size deletion Mb	?	1.12	1.4	1.4	1.4	1.1	2	2	2.1	1.55–1.92	2.1	1.3
Parental analysis for deletion	P-NL M-NL	P-NL M-NL	P-NL M-NL	P-NL M-NL	P-NL M-NL	P-NL M-NL	P-NL M-NL	P-NL M-NL	P-NL M-NL	P-NL M-NL	P-NL M-NL	P-NL M-NL
Gender	m	m	m	m	m	f	f	f	f	f	m	f
Karyotype	46,XY	46,XY	46,XY	46,XY	46,XY	46,XX	46,XX	46,XX	46,XX	46,XX	46,XX	46,XX
Age, years	2	11/12	6	5	3	4	7	8/12	11/12	2/12	31	12
Premature death	–	–	–	–	–	–	–	–	–	–	–	–
Gestational age, weeks	35	37	38	36	35	37	38	36	36	36	36	35
Birth weight, kg	?	1.800	?	2.210	1.250	1.700	2.100	1.930	?	?	?	?
centile	<3rd	<3rd	3rd–10th	<3rd	<3rd	<3rd	<3rd	<3rd	<3rd	<3rd	<3rd	<3rd
Postnatal growth retardation	+	?	+	–	+	+	+	+	+	+	+	+
Weight at presentation (<3rd centile)	?	?	<3rd	60th	<3rd	<3rd	2.5th–10th	5th	<3rd	?	?	?
Height at presentation (<3rd centile)	?	?	?	60th	<3rd	<3rd	2.5th–10th	5th	<3rd	?	?	?
Head circumference (<3rd centile/<2 SD)	? MC	? MC	?	?	?	?	<2.5th	<2.5th	?	?	?	?
Developmental delay	+sd	+	+sd	–	±	+sd	motor	+	+sd	+sd	+sd	+sd
Behavioural problems	?	NA	–	+	–	–	±	–	–	–	–	–
Congenital heart defect	+TAC2 VSD	VSD	+BAV	–	–	–	VSD	PAS VSD	NA	NA	NA	NA
Hypocalcaemia	+	?	?	?	?	?	?	?	?	?	?	?
Urogenital anomalies	+HYSP	?	?	?	?	?	?	?	?	?	?	?
Tracheomalacia/laryngomalacia	–	–	–	–	–	–	–	–	–	–	–	–
Immunodeficiency	–	?	?	?	?	?	?	?	?	?	?	?
Recurrent infections	–	NA	–	–	–	–	+	+	+	+	+	+

BAV = Bicuspid aortic valve; del = deletion; f = female; HYSP = hypospadias; IAAB = interrupted aortic arch type B; m = male; M = maternal; MC = microcephaly; NA = not applicable; NL = normal; P = paternal; PAS = pulmonary artery stenosis; Pat = patient; PDA = persistent ductus arteriosus Botalli; sd = speech delay; TAC = truncus arteriosus communis; TAC2/A4 = truncus arteriosus communis type 2 or A4; VSD = ventricular septal defect.

BU = Bifid uvula; CP = cleft palate; del = deletion; EDC = epibulbar dermoid cyst; f = female; HL = hearing loss; HM = hemifacial microsomia; IATRT = intracranial atypical teratoid rhabdoid tumor; m = male; M = maternal; NL = normal; OAVS = oculo-auriculo-vertebral spectrum; P = paternal; Pat = patient; PT = preauricular tag; SA = various skeletal anomalies (1 scoliosis, 1 lumbar lordosis, 1 upper limb asymmetry, 1 anomaly in upper extremity); ST = strabismus; T = tooth; T1 = tooth 1; T2 = tooth 2; T3 = tooth 3; T4 = tooth 4; T5 = tooth 5; T6 = tooth 6; T7 = tooth 7; T8 = tooth 8; T9 = tooth 9; T10 = tooth 10; T11 = tooth 11; T12 = tooth 12; T13 = tooth 13; T14 = tooth 14; T15 = tooth 15; T16 = tooth 16; T17 = tooth 17; T18 = tooth 18; T19 = tooth 19; T20 = tooth 20; T21 = tooth 21; T22 = tooth 22; T23 = tooth 23; T24 = tooth 24; T25 = tooth 25; T26 = tooth 26; T27 = tooth 27; T28 = tooth 28; T29 = tooth 29; T30 = tooth 30; T31 = tooth 31; T32 = tooth 32; T33 = tooth 33; T34 = tooth 34; T35 = tooth 35; T36 = tooth 36; T37 = tooth 37; T38 = tooth 38; T39 = tooth 39; T40 = tooth 40; T41 = tooth 41; T42 = tooth 42; T43 = tooth 43; T44 = tooth 44; T45 = tooth 45; T46 = tooth 46; T47 = tooth 47; T48 = tooth 48; T49 = tooth 49; T50 = tooth 50; T51 = tooth 51; T52 = tooth 52; T53 = tooth 53; T54 = tooth 54; T55 = tooth 55; T56 = tooth 56; T57 = tooth 57; T58 = tooth 58; T59 = tooth 59; T60 = tooth 60; T61 = tooth 61; T62 = tooth 62; T63 = tooth 63; T64 = tooth 64; T65 = tooth 65; T66 = tooth 66; T67 = tooth 67; T68 = tooth 68; T69 = tooth 69; T70 = tooth 70; T71 = tooth 71; T72 = tooth 72; T73 = tooth 73; T74 = tooth 74; T75 = tooth 75; T76 = tooth 76; T77 = tooth 77; T78 = tooth 78; T79 = tooth 79; T80 = tooth 80; T81 = tooth 81; T82 = tooth 82; T83 = tooth 83; T84 = tooth 84; T85 = tooth 85; T86 = tooth 86; T87 = tooth 87; T88 = tooth 88; T89 = tooth 89; T90 = tooth 90; T91 = tooth 91; T92 = tooth 92; T93 = tooth 93; T94 = tooth 94; T95 = tooth 95; T96 = tooth 96; T97 = tooth 97; T98 = tooth 98; T99 = tooth 99; T100 = tooth 100; T101 = tooth 101; T102 = tooth 102; T103 = tooth 103; T104 = tooth 104; T105 = tooth 105; T106 = tooth 106; T107 = tooth 107; T108 = tooth 108; T109 = tooth 109; T110 = tooth 110; T111 = tooth 111; T112 = tooth 112; T113 = tooth 113; T114 = tooth 114; T115 = tooth 115; T116 = tooth 116; T117 = tooth 117; T118 = tooth 118; T119 = tooth 119; T120 = tooth 120; T121 = tooth 121; T122 = tooth 122; T123 = tooth 123; T124 = tooth 124; T125 = tooth 125; T126 = tooth 126; T127 = tooth 127; T128 = tooth 128; T129 = tooth 129; T130 = tooth 130; T131 = tooth 131; T132 = tooth 132; T133 = tooth 133; T134 = tooth 134; T135 = tooth 135; T136 = tooth 136; T137 = tooth 137; T138 = tooth 138; T139 = tooth 139; T140 = tooth 140; T141 = tooth 141; T142 = tooth 142; T143 = tooth 143; T144 = tooth 144; T145 = tooth 145; T146 = tooth 146; T147 = tooth 147; T148 = tooth 148; T149 = tooth 149; T150 = tooth 150; T151 = tooth 151; T152 = tooth 152; T153 = tooth 153; T154 = tooth 154; T155 = tooth 155; T156 = tooth 156; T157 = tooth 157; T158 = tooth 158; T159 = tooth 159; T160 = tooth 160; T161 = tooth 161; T162 = tooth 162; T163 = tooth 163; T164 = tooth 164; T165 = tooth 165; T166 = tooth 166; T167 = tooth 167; T168 = tooth 168; T169 = tooth 169; T170 = tooth 170; T171 = tooth 171; T172 = tooth 172; T173 = tooth 173; T174 = tooth 174; T175 = tooth 175; T176 = tooth 176; T177 = tooth 177; T178 = tooth 178; T179 = tooth 179; T180 = tooth 180; T181 = tooth 181; T182 = tooth 182; T183 = tooth 183; T184 = tooth 184; T185 = tooth 185; T186 = tooth 186; T187 = tooth 187; T188 = tooth 188; T189 = tooth 189; T190 = tooth 190; T191 = tooth 191; T192 = tooth 192; T193 = tooth 193; T194 = tooth 194; T195 = tooth 195; T196 = tooth 196; T197 = tooth 197; T198 = tooth 198; T199 = tooth 199; T200 = tooth 200; T201 = tooth 201; T202 = tooth 202; T203 = tooth 203; T204 = tooth 204; T205 = tooth 205; T206 = tooth 206; T207 = tooth 207; T208 = tooth 208; T209 = tooth 209; T210 = tooth 210; T211 = tooth 211; T212 = tooth 212; T213 = tooth 213; T214 = tooth 214; T215 = tooth 215; T216 = tooth 216; T217 = tooth 217; T218 = tooth 218; T219 = tooth 219; T220 = tooth 220; T221 = tooth 221; T222 = tooth 222; T223 = tooth 223; T224 = tooth 224; T225 = tooth 225; T226 = tooth 226; T227 = tooth 227; T228 = tooth 228; T229 = tooth 229; T230 = tooth 230; T231 = tooth 231; T232 = tooth 232; T233 = tooth 233; T234 = tooth 234; T235 = tooth 235; T236 = tooth 236; T237 = tooth 237; T238 = tooth 238; T239 = tooth 239; T240 = tooth 240; T241 = tooth 241; T242 = tooth 242; T243 = tooth 243; T244 = tooth 244; T245 = tooth 245; T246 = tooth 246; T247 = tooth 247; T248 = tooth 248; T249 = tooth 249; T250 = tooth 250; T251 = tooth 251; T252 = tooth 252; T253 = tooth 253; T254 = tooth 254; T255 = tooth 255; T256 = tooth 256; T257 = tooth 257; T258 = tooth 258; T259 = tooth 259; T260 = tooth 260; T261 = tooth 261; T262 = tooth 262; T263 = tooth 263; T264 = tooth 264; T265 = tooth 265; T266 = tooth 266; T267 = tooth 267; T268 = tooth 268; T269 = tooth 269; T270 = tooth 270; T271 = tooth 271; T272 = tooth 272; T273 = tooth 273; T274 = tooth 274; T275 = tooth 275; T276 = tooth 276; T277 = tooth 277; T278 = tooth 278; T279 = tooth 279; T280 = tooth 280; T281 = tooth 281; T282 = tooth 282; T283 = tooth 283; T284 = tooth 284; T285 = tooth 285; T286 = tooth 286; T287 = tooth 287; T288 = tooth 288; T289 = tooth 289; T290 = tooth 290; T291 = tooth 291; T292 = tooth 292; T293 = tooth 293; T294 = tooth 294; T295 = tooth 295; T296 = tooth 296; T297 = tooth 297; T298 = tooth 298; T299 = tooth 299; T300 = tooth 300; T301 = tooth 301; T302 = tooth 302; T303 = tooth 303; T304 = tooth 304; T305 = tooth 305; T306 = tooth 306; T307 = tooth 307; T308 = tooth 308; T309 = tooth 309; T310 = tooth 310; T311 = tooth 311; T312 = tooth 312; T313 = tooth 313; T314 = tooth 314; T315 = tooth 315; T316 = tooth 316; T317 = tooth 317; T318 = tooth 318; T319 = tooth 319; T320 = tooth 320; T321 = tooth 321; T322 = tooth 322; T323 = tooth 323; T324 = tooth 324; T325 = tooth 325; T326 = tooth 326; T327 = tooth 327; T328 = tooth 328; T329 = tooth 329; T330 = tooth 330; T331 = tooth 331; T332 = tooth 332; T333 = tooth 333; T334 = tooth 334; T335 = tooth 335; T336 = tooth 336; T337 = tooth 337; T338 = tooth 338; T339 = tooth 339; T340 = tooth 340; T341 = tooth 341; T342 = tooth 342; T343 = tooth 343; T344 = tooth 344; T345 = tooth 345; T346 = tooth 346; T347 = tooth 347; T348 = tooth 348; T349 = tooth 349; T350 = tooth 350; T351 = tooth 351; T352 = tooth 352; T353 = tooth 353; T354 = tooth 354; T355 = tooth 355; T356 = tooth 356; T357 = tooth 357; T358 = tooth 358; T359 = tooth 359; T360 = tooth 360; T361 = tooth 361; T362 = tooth 362; T363 = tooth 363; T364 = tooth 364; T365 = tooth 365; T366 = tooth 366; T367 = tooth 367; T368 = tooth 368; T369 = tooth 369; T370 = tooth 370; T371 = tooth 371; T372 = tooth 372; T373 = tooth 373; T374 = tooth 374; T375 = tooth 375; T376 = tooth 376; T377 = tooth 377; T378 = tooth 378; T37

Table 3. Clinical features and review of the literature (interval IV–V)

	Garcia-Minaur et al., 2002	Rauch et al., 2005	D'Angelo et al., 2007	Fernandez et al., 2009	Kurahashi et al., 1996, 1997	Ogilvie et al., 2009	Patient	Total
Deleted interval and gene (see fig. 4)	IV+V	IV+V	IV+V	IV+V	V <i>CRKL</i>	V+VI	V+VI	
22q11.2 size deletion, Mb	?	?	1	1	1.1	?	1.3	
Parental analysis for deletion	P del	P-NL M-NL	P? M-NL	M del	P-NL M-NL	P-NL M-NL	P-NL M-NL	
Gender	m	m	m	f	m	f	f	4 m/3 f
Karyotype						46,XX	46,XX	
Age, years	5/12	?	6	4	7/12	?	4	
Premature death	–	–	–	–	–	–	–	0/8
Gestational age, weeks	39	?	?	at term	37	?	35	mean 37
Birth weight, kg	3.250	?	?	2.900	2.450	?	1.760	1.680
Centile				25th	10th–25th		<3rd	mean 2.408
Postnatal growth retardation	–	–	–	–	+	?	+	3/7
Weight at presentation (<3rd centile)	<3rd	?	?	>97th	<3rd	?	<<3rd	4/5
Height at presentation (<3rd centile)	10th–25th	?	?	25th–50th	<5th	?	3rd–10th	3/5
Head circumference (<3rd centile/<–2 SD)	<3rd	?	?	50th–98th	5th–10th	?	<3rd	3/5
Developmental delay	–	±	+sd	+sd	+	?	+sd	5/6
Behavioural problems	NA	?	+	+	NA	?	–	2/3
Congenital heart defect	ToF	?	–	–	ASD, VSD	PA, ToF	ASD, VSD	+TAC2
Hypocalcaemia	–	?	–	–	–	?	–	0/6
Urogenital Anomalies	?	?	–	–	?	?	–	0/3
Tracheomalacia/laryngomalacia	–	–	–	–	–	?	–	0/6
Immunodeficiency	?	?	–	–	?	?	–	0/3
Recurrent infections	NA	?	+	?	NA	?	–	1/2

ASD = Atrial septal defect; del = deletion; f = female; m = male; M = maternal; NA = not applicable; NL = normal; P = paternal; PA = pulmonary atresia; sd = speech delay; TAC2 = truncus arteriosus communis type 2; ToF = tetralogy of Fallot; VSD = ventricular septal defect.

Table 4. Clinical features and review of the literature (interval IV–V)

	Garcia-Minaur et al., 2002	Rauch et al., 2005	D'Angelo et al., 2007	Fernandez et al., 2009	Kurahashi et al., 1996 1997	Ogilvie et al., 2009	Patient	Total
Deleted interval and gene (see fig. 4)	IV+V	IV+V	IV+V	IV+V	V <i>CRKL</i>	V+VI	V+VI	
22q11.2 size deletion, Mb	?	?	1	1	1.1	?	1.3	
Parental analysis for deletion	P del	P-NL M-NL	P? M-NL	M del	P-NL M-NL	P-NL M-NL	P-NL M-NL	
Gender	m	m	m	f	m	f	f	4 m
Karyotype						46,XX	46,XX	3 f
Age, years	5/12	?	6	4	7/12	?	4	
Craniofacial								
Arched eye brows	–	–	–	–	+	?	+	2/6
Hypertelorism	+	–	–	–	–	?	–	1/7
Deep-set eyes	–	–	–	+	–	?	–	1/6
Iris coloboma	–	–	–	–	–	?	–	0/6
Choanal stenosis/atresia	–	–	–	–	–	?	–	0/6
Broad nose	–	+	+	+	+	?	+	4/6
Hypoplastic alae nasi	–	–	+	–	–	?	–	1/6
Smooth philtrum	–	–	–	–	–	?	+	1/6
High arched palate	–	?	?	+	?	?	–	1/4
Velopharyngeal insufficiency	–	–	–	–	–	?	–	0/7
Cleft palate/bifid uvula	–	?	–	?	?	?	–	0/4
Micrognathia	–	+	+	+	+	?	+	5/7
Down-turned mouth	+	–	–	–	–	?	+	2/6
Pointed chin	–	–	–	–	–	?	+	1/6
Skeletal								
5th finger clinodactyly	?	?	?	?	?	?	–	1/2
Other clinical features				pes planus OB		+	imperforate anus	

f = Female; m = male; M = maternal; NL = normal; P = paternal; OB = obesity.

reported in other cases. Since a key role of *Tbx1* in heart development has been demonstrated in the mouse, this 22q11.2 gene was also involved in the pathogenesis of the associated human heart defects [Lindsay et al., 1999; Baldini, 2002]. *TBX1* is not included in our as well as other distal 22p11.2 deletion cases (fig. 3). However, genes distal to and functionally interacting with *TBX1* such as *CRKL* and *ERK2/MAPK1* (fig. 3) have been proposed to be responsible for congenital heart defects [Guris et al., 2001; Newbern et al., 2008]. In particular, *ERK2/MAPK1* is included in the 22q11.2 distal deletion interval and its inactivation in the mouse leads to cardiac outflow tract abnormalities similar to those observed in the human patients. Nevertheless, a position effect on *TBX1* expression cannot be excluded. Genes possibly involved in the growth retardation phenotype are *UBE2L3* whose mutation has been associated with retarded pre-natal growth in the mouse [Harbers et al., 1996], and *MIR130B* showing cell growth promoting activity via silencing *TP53INP1* [Ma et al., 2010].

The deleted interval demonstrated in our case overlaps both the 3-Mb proximal deletion in 22q11.2 and the distal deletion intervals and includes both the *CRKL* and

the *ERK2/MAPK1* genes (fig. 3). A similar deletion has been described in another patient, who, like our case, presents preterm delivery, growth retardation, head circumference less than the 3rd centile and congenital heart disease, albeit different from our case, atrial septal defect and ventricular septal defect [Ogilvie et al., 2009]. Moreover, the latter presented imperforate anus, a condition which has never hitherto been reported in the literature.

In conclusion, it is our opinion that pediatric geneticists should take into account this condition when they suspect a SRS and when they have the occasion to examine children with truncus arteriosus.

Acknowledgements

We are thankful to the parents for their cooperation and to Marco Bonazzi for photography work. This work was supported by grants from Telethon-Italia N. GGP07086, GGP11122 (A.R.), MIUR PRIN 2009 (A.R.), Associazione Italiana Ricerca sul Cancro (A.R.) and Ministry of Italian Health grant RC1101GM05 (M.C.).

References

- Baldini A: DiGeorge syndrome: the use of model organisms to dissect complex genetics. *Hum Mol Genet* 11:2363–2369 (2002).
- Ben-Shachar S, Ou Z, Shaw CA, Belmont JW, Patel MS, et al: 22q11.2 distal deletion: a recurrent genomic disorder distinct from DiGeorge syndrome and velocardiofacial syndrome. *Am J Hum Genet* 82:214–221 (2008).
- Bruce S, Hannula-Jouppi K, Puoskari M, Fransson I, Simola KO, et al: Submicroscopic genomic alterations in Silver-Russell syndrome and Silver-Russell-like patients. *J Med Genet* 47:816–822 (2010).
- Carlson C, Sirotkin H, Pandita R, Goldberg R, McKie J, et al: Molecular definition of 22q11 deletions in 151 velo-cardio-facial syndrome patients. *Am J Hum Genet* 61:620–629 (1997).
- D'Angelo CS, Jehee FS, Koiffmann CP: An inherited atypical 1 Mb 22q11.2 deletion within the DGS/VCFS 3 Mb region in a child with obesity and aggressive behavior. *Am J Med Genet A* 143A:1928–1932 (2007).
- Edelmann L, Pandita RK, Spiteri E, Funke B, Goldberg R, et al: A common molecular basis for rearrangement disorders on chromosome 22q11. *Hum Mol Genet* 8:1157–1167 (1999).
- Fernandez L, Nevado J, Santos F, Heine-Suner D, Martinez-Glez V, et al: A deletion and a duplication in distal 22q11.2 deletion syndrome region. Clinical implications and review. *BMC Med Genet* 10:48 (2009).
- Garcia-Minaur S, Fantes J, Murray RS, Porteous ME, Strain L, et al: A novel atypical 22q11.2 distal deletion in father and son. *J Med Genet* 39:e62 (2002).
- Guris DL, Fantes J, Tara D, Druker BJ, Imamoto A: Mice lacking the homologue of the human 22q11.2 gene *CRKL* phenocopy neurocristopathies of DiGeorge syndrome. *Nat Genet* 27:293–298 (2001).
- Harbers K, Müller U, Grams A, Li E, Jaenisch R, Franz T: Provirus integration into a gene encoding a ubiquitin-conjugating enzyme results in a placental defect and embryonic lethality. *Proc Natl Acad Sci USA* 93:12412–12417 (1996).
- Jackson EM, Shaikh TH, Gururangan S, Jones MC, Malkin D, et al: High-density single nucleotide polymorphism array analysis in patients with germline deletions of 22q11.2 and malignant rhabdoid tumor. *Hum Genet* 122:117–127 (2007).
- Kurahashi H, Nakayama T, Osugi Y, Tsuda E, Masuno M, et al: Deletion mapping of 22q11 in CATCH22 syndrome: identification of a second critical region. *Am J Hum Genet* 58:1377–1381 (1996).
- Kurahashi H, Tsuda E, Kohama R, Nakayama T, Masuno M, et al: Another critical region for deletion of 22q11: a study of 100 patients. *Am J Med Genet* 72:180–185 (1997).
- Lafay-Cousin L, Payne E, Strother D, Chernos J, Chan M, Bernier FP: Goldenhar phenotype in a child with distal 22q11.2 deletion and intracranial atypical teratoid rhabdoid tumor. *Am J Med Genet A* 149A:2855–2859 (2009).
- Lindsay EA, Botta A, Jurecic V, Carattini-Rivera S, Cheah YC, et al: Congenital heart disease in mice deficient for the DiGeorge syndrome region. *Nature* 401:379–383 (1999).
- Ma S, Tang KH, Chan YP, Lee TK, Kwan PS, et al: miR-130b Promotes CD133(+) liver tumor-initiating cell growth and self-renewal via tumor protein 53-induced nuclear protein 1. *Cell Stem Cell* 7:694–707 (2010).
- McDermid HE, Morrow BE: Genomic disorders on 22q11. *Am J Hum Genet* 70:1077–1088 (2002).
- Mikhail FM, Descartes M, Piotrowski A, Andersson R, Diaz de Stahl T, et al: A previously unrecognized microdeletion syndrome on chromosome 22 band q11.2 encompassing the *BRC* gene. *Am J Med Genet A* 143A:2178–2184 (2007).

- Newbern J, Zhong J, Wickramasinghe RS, Li X, Wu Y, et al: Mouse and human phenotypes indicate a critical conserved role for ERK2 signaling in neural crest development. *Proc Natl Acad Sci USA* 105:17115–17120 (2008).
- Ogilvie CM, Ahn JW, Mann K, Roberts RG, Flinter F: A novel deletion in proximal 22q associated with cardiac septal defects and microcephaly: a case report. *Mol Cytogenet* 2:9 (2009).
- Rauch A, Pfeiffer RA, Leipold G, Singer H, Tiggles M, Hofbeck M: A novel 22q11.2 microdeletion in DiGeorge syndrome. *Am J Hum Genet* 64:659–667 (1999).
- Rauch A, Zink S, Zweier C, Thiel CT, Koch A, et al: Systematic assessment of atypical deletions reveals genotype-phenotype correlation in 22q11.2. *J Med Genet* 42:871–876 (2005).
- Rodningen OK, Prescott T, Eriksson AS, Rosby O: 1.4 Mb recurrent 22q11.2 distal deletion syndrome, two new cases expand the phenotype. *Eur J Med Genet* 51:646–650 (2008).
- Saitta SC, McGrath JM, Mensch H, Shaikh TH, Zackai EH, Emanuel BS: A 22q11.2 deletion that excludes *UFDIL* and *CDC45L* in a patient with conotruncal and craniofacial defects. *Am J Hum Genet* 65:562–566 (1999).
- Shaikh TH, Kurahashi H, Saitta SC, O'Hare AM, Hu P, et al: Chromosome 22-specific low copy repeats and the 22q11.2 deletion syndrome: genomic organization and deletion endpoint analysis. *Hum Mol Genet* 9:489–501 (2000).
- Sparago A, Russo S, Cerrato F, Ferraiuolo S, Castorina P, et al: Mechanisms causing imprinting defects in familial Beckwith-Wiedemann syndrome with Wilms' tumour. *Hum Mol Genet* 16:254–264 (2007).
- Stankiewicz P, Lupski JR: Genome architecture, rearrangements and genomic disorders. *Trends Genet* 18:74–82 (2002).
- Xu J, Fan YS, Siu VM: A child with features of Goldenhar syndrome and a novel 1.12 Mb deletion in 22q11.2 by cytogenetics and oligonucleotide array CGH: is this a candidate region for the syndrome? *Am J Med Genet A* 146A:1886–1889 (2008).