This is an Open Access article licensed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs 3.0 License (www.karger.com/OA-license), applicable to the online version of the article only. Distribution for non-commercial purposes only.

Familial Hypobetalipoproteinemia-Induced Nonalcoholic Steatohepatitis

Mindy C.W. Lam^a Janakie Singham^a Robert A. Hegele^c
Maziar Riazzy^a Matti A. Hiob^a Gordon Francis^b
Urs P. Steinbrecher^a

Divisions of ^aGastroenterology and ^bEndocrinology and Metabolism, Department of Medicine, University of British Columbia, Vancouver, B.C., and ^cBlackburn Cardiovascular Genetics Laboratory, Robarts Research Institute, University of Western Ontario, London, Ont., Canada

Key Words

Familial hypobetalipoproteinemia · Nonalcoholic steatohepatitis · *APOB*

Abstract

Familial hypobetalipoproteinemia (FHBL) is a rare genetic disorder of lipid metabolism that is associated with abnormally low serum levels of low-density lipoprotein (LDL) cholesterol and apolipoprotein B. It is an autosomal co-dominant disorder, and depending on zygosity, the clinical manifestations may vary from none to neurological, endocrine, hematological or liver dysfunction. Nonalcoholic fatty liver disease is common in persons with FHBL, however progression to nonalcoholic steatohepatitis is unusual. We describe here a patient with a novel *APOB* mutation, V703I, which appears to contribute to the severity of the FHBL phenotype. He had liver enzyme abnormalities, increased echogenicity of the liver consistent with steatosis, very low LDL cholesterol at 0.24 mmol/l (normal 1.8–3.5 mmol/l) and an extremely low apolipoprotein B level of 0.16 g/l (normal 0.6–1.2 g/l). *APOB* gene sequencing revealed him to be a compound heterozygote with two mutations (R463W and V703I). *APOB* R463W has previously been reported to cause FHBL. Genetic sequencing of his first-degree relatives identified the *APOB* V703I mutation in his normolipidemic brother and father and the *APOB* R463W mutation in his mother and sister, both of whom have very low LDL cholesterol levels. These results suggest that the *APOB* V703I mutation alone does not cause the FHBL phenotype. However, it is possible that it has a contributory role to a more aggressive phenotype in the presence of *APOB* R463W.

Introduction

Familial hypobetalipoproteinemia (FHBL) is a rare autosomal co-dominant genetic disorder of lipid metabolism. It is characterized by serum low-density lipoprotein (LDL) cholesterol or apolipoprotein B (apoB) levels ≤ 5 th percentile. The disorder was first described in 1979 by Steinberg et al. [1–3]. ApoB is the largest of the apolipoproteins and is important in the formation and secretion of very-low-density lipoprotein (VLDL) and chylomicrons. There are two forms of apoB found in serum, namely apoB-48 and apoB-100. These two isoforms are derived from differential splicing of RNA from a single *APOB* gene [4]. ApoB-48 is synthesized by enterocytes and is a key component of chylomicrons, while apoB-100 is synthesized by the liver and is the key protein component of VLDL and LDL particles; B-100 also contains the binding site for hepatocyte uptake [5]. The *APOB* gene is found on chromosome 2p24 and is unlinked to other apolipoprotein gene clusters [6].

Since the discovery of the first kindred with FHBL, many subsequent kindreds have been described, leading to the identification of numerous single nucleotide substitutions, gene deletions, splicing variations and linkage abnormalities in the *APOB* gene that can lead to defects in VLDL and chylomicron production and secretion. Persons with FHBL can be heterozygotes or homozygotes, with severity of the clinical phenotype related to the number of defective *APOB* alleles. Heterozygotes often have no clinical manifestations other than very low serum apoB-containing lipoprotein levels; these low levels of atherogenic lipoprotein particles may actually confer cardiovascular protection [7]. In contrast, FHBL homozygotes can have a range of clinical symptoms, including fat malabsorption, neurologic symptoms, red cell acanthocytosis, retinitis pigmentosa and/or nonalcoholic fatty liver disease. In fact, the clinical presentation of homozygous FHBL can be indistinguishable from that of abetalipoproteinemia (ABL) due to mutation in *MTTP*, which is the gene that encodes microsomal triglyceride transfer protein [8].

We describe here a case of nonalcoholic steatohepatitis (NASH) that led to the diagnosis of FHBL.

Case Report

A 44-year-old man was referred to the Vancouver General Hospital Division of Gastroenterology for advice regarding management of NASH. He was a shop owner, a life-long nonsmoker and drinking about half a bottle of wine a week. He was first noted to have elevated liver enzymes in 1996. At that time, he was told that AST (normal 5–35 U/l), ALT (normal 7–56 U/l) and GGT (normal 8–78 U/l) were all about 2-fold above the upper limits of normal. Hepatitis B and C serologies were negative. A liver biopsy was recommended, however the patient declined the procedure and elected to initiate a weight loss program with dietary modification and increased exercise for presumed NASH. About 3 years before presentation, he was found to have elevated transaminases on blood work done for an insurance application, with ALT 130 U/l and AST 111 U/l. These abnormalities were again attributed to NASH. After about 6 months of strict diet and exercise, as well as a 10-pound intentional weight loss, his liver enzymes normalized with ALT 44 U/l and AST 41 U/l. He reintroduced mild alcohol intake and reduced the amount of exercise, and 6 months later his liver enzymes were again elevated with ALT 94 U/l and AST 80 U/l. Other causes for elevated liver enzymes, including anti-nuclear antibody, ceruloplasmin and alpha-1 antitrypsin, were normal or negative. However, he was found to have an elevated ferritin at 737 $\mu\text{g/l}$ (normal 18–250 $\mu\text{g/l}$). Subsequent testing revealed him to be heterozygous for the C282Y mutation in the *HFE* gene for hemochromatosis. There was no other significant past medical history.

On examination, he was a stocky male with a body mass index of 32 kg/m². There were no extrahepatic signs of liver disease. Respiratory and cardiovascular examinations were normal. His abdomen was soft and nontender with no palpable organs or masses. Review of his blood tests from 3 years before revealed LDL cholesterol 0.24 mmol/l (normal 1.8–3.5 mmol/l), high-density lipoprotein cholesterol 1.29 mmol/l (normal 0.9–1.5 mmol/l) and triglyceride 0.37 mmol/l (normal 0.5–1.7 mmol/l). The serum apoB level was extremely low at 0.16 g/l (normal 0.6–1.2 mmol/l). A liver biopsy had been done in 2009 by the referring physician and had shown normal liver architecture. There was very mild portal and pericellular centrilobular fibrosis, mild focal mononuclear cell infiltration and moderate macrovesicular steatosis. There was no abnormal iron deposition and no Mallory hyaline.

Family history revealed that his sister (subject II-3 in [fig. 1](#)) also has elevated liver enzymes ([table 1](#)). She was obese, had type 2 diabetes and had recently been diagnosed with esophageal varices. She had been aware of a low LDL cholesterol level for several years. However, details of her liver disease were unknown. His mother (I-2) and several maternal relatives were known to have a very low cholesterol level. His mother's family was of Christian Lebanese origin. His father (I-1) had a normal cholesterol level. His father's family was from Great Britain.

Family history and laboratory results suggested the diagnosis of FHBL. ApoB immunoblotting on fasting plasma showed only a band at the expected apoB-100 position, arguing against truncation mutations. DNA was extracted from plasma samples from the patient and subsequently from all first-degree relatives and sent to the Cardiovascular Genetics Lab in London, Ontario. Direct automated sequencing revealed two different mutations in the *APOB* gene, namely R463W and V703I. These were found to be on opposite chromosomes, so the proband is a compound heterozygote. Family relationships, cholesterol levels and *APOB* mutation analysis are shown in figure 1 and table 1.

Our patient was advised to return to regular exercise and to maintain a healthy low-fat diet. He was also started on vitamin E and multivitamin supplementation. His weight decreased by about 5 kg over 6 months and his ALT normalized to 25 U/l.

Discussion

Rare monogenic disorders have been very informative for identifying key components of the metabolism of apoB-containing lipoproteins and triglycerides [9]. These disorders include ABL due to mutation of the *MTTP* gene, hypobetalipoproteinemia due to mutation of the *APOB* gene, familial combined hypolipidemia due to mutation of the *ANGPTL3* gene and PCSK9 deficiency due to mutations in the *PCSK9* gene [10, 11].

In addition to being important in metabolism, some of the gene products causing these disorders are now targets for new drugs designed to lower LDL cholesterol. For instance, mipomersen is an anti-sense RNA biologic that reduces LDL cholesterol by targeting the *APOB* gene, while lomitapide reduces LDL cholesterol by inhibiting microsomal triglyceride transfer protein [12, 13]. There are also numerous new biologic drugs in development that lower LDL cholesterol by targeting PCSK9 [14]. While these new drugs seem to be very effective at lowering LDL cholesterol, there is concern about potential long-term side effects, such as hepatosteatosis. Clinical evaluation of 'experiments of nature' such as FHBL patients who carry a germ-line mutation in *APOB* can provide clues of the potential long-term benefits and adverse reaction of pharmacologic therapies that target these key metabolic pathways.

ApoB truncation is due to *APOB* gene mutation, exon deletions and splicing variations. About 45 different truncated forms of apoB have been discovered to date, ranging from apoB-2 to apoB-90 [15–38]. The numbers represent estimated apoB mass

based on polyacrylamide gel electrophoresis, with apoB-100 being normal. Truncated apoB can form varying sizes and densities of lipoprotein particles depending on the length of truncation [39]. Truncated apoB particles have a lower production rate and higher clearance rate, resulting in low plasma levels of apoB [19, 40–42]. ApoB smaller than apoB-27.6 is undetectable in plasma due to rapid clearance [40]. In FHBL heterozygotes with apoB truncation, the average plasma concentration of apoB is around 25–30% of normal [43, 44].

FHBL due to *ANGPTL3* gene linked to chromosome 3p21 is associated with about a 50% reduction in apoB levels compared to normal individuals, as well as an elevated rate of catabolism of VLDL apoB [10, 45–47]. At least some individuals with this subtype of FHBL have gain-of-function mutations of PCSK9 [48]. This is a serine protease that destroys LDL receptors in the liver and thereby controls the level of LDL in plasma. Mutations that increase PCSK9 activity cause hypercholesterolemia, whereas mutations that inactivate PCSK9 have the opposite effect.

In FHBL, the genetic defect in triglyceride export can lead to increased intrahepatic triglyceride content. There have been several reports of nonalcoholic fatty liver disease in FHBL subjects as detected by ultrasound or liver biopsy [28, 49–52]. Schonfeld et al. [51] analyzed liver fat by magnetic resonance spectroscopy, body mass index, waist circumference and glucose tolerance of 21 individuals with FHBL due to apoB truncations compared with 14 controls. Despite similar mean values for obesity and insulin indexes, there was a 5-fold increase in liver fat in FHBL subjects. Similar studies with FHBL subjects due to chromosome 3p21 linkage or PCSK9 mutations revealed normal hepatic fat content, an important phenotypic difference between these mutations and apoB truncation [7, 46].

The proband in this report is a 44-year-old man with NASH secondary to FHBL. Genetic analysis revealed compound heterozygosity for two mutations, each of which results in amino acid substitution. As shown in figure 1, he inherited the R463W mutation from his mother (subject I-2). Heterozygosity for this mutation is associated with hypobetalipoproteinemia, while homozygosity has a more severe phenotype [53]. Interestingly, the R463W mutation was first reported in an extended Christian Lebanese family, which is also the ethnic background of our proband's mother. Our patient inherited a second nontruncating *APOB* mutation, V703I, from his father. This is a novel mutation, and analysis using the PolyPhen program predicts the effect of this amino acid substitution as 'possibly damaging' to apoB function. Although the two V703I heterozygotes in this kindred, subjects I-1 and II-1, had normal LDL cholesterol levels when the second allele encoded normal apoB-100, it is possible that this mutation, in the context of a defective form of apoB-100, such as the R463W mutation, contributes to increased intracellular degradation of apoB. This might act to compound the impairment of VLDL secretion in subject II-2 who had one copy of R463W, as evidenced by his lower LDL cholesterol level compared to the two R463W simple heterozygotes (subjects I-2 and II-3).

Although NASH is common in FHBL patients, the long-term effects of FHBL are still unclear [28, 49, 51, 54]. However, steatosis from FHBL can lead to end-stage liver disease. Interestingly, Harada et al. [54] reported a patient with recurrent NASH and cirrhosis due to FHBL post liver transplantation. This suggests that at least in some forms of FHBL, the intestinal defect may be sufficient to cause the phenotype.

Fortunately, our patient had only mild steatohepatitis. However, his sister (subject II-3) has developed esophageal varices, possibly due to liver cirrhosis from FHBL compounded by diabetes mellitus.

Diet and exercise continue to be the mainstay of treatment for patient with NASH induced by FHBL. As in our patient, exercise and weight loss led to improvement in liver enzyme tests. Fat malabsorption can be managed through diet modification by limiting intake of fats containing long-chain fatty acids as well as supplementing fat-soluble vitamins. Reports have suggested that the cellular membranes of patients with FHBL are relatively deficient in vitamin E; indeed severe vitamin E deficiency in both homozygous FHBL and ABL is responsible for the development of neurologic signs and symptoms [55, 56]. While the most common neurologic symptom is decreased or absent deep tendon reflexes, ataxia, proprioceptive deficits and pain have been described in both homozygous FHBL and ABL [57–59]. There are reports of success with vitamin E treatments for neurologic symptoms in FHBL and ABL [57, 60]. There is some evidence that combination therapy of vitamin A and E can slow the progression of retinal degeneration and may lead to restoration of vision if started early [61–63]. Our patient was started on vitamin E and multivitamin supplementation for primary prevention, as he does not currently have any neurological, hematological or ocular abnormalities.

Conclusion

FHBL is an uncommon autosomal co-dominant disorder of lipid metabolism that is often overlooked because low levels of LDL cholesterol may not attract the attention of health care providers. In the present case liver enzyme abnormalities were the first clue to diagnosis. The understanding of FHBL and liver disease is still in its preliminary stages. Patients with FHBL and NASH can progress to end-stage liver failure, therefore monitoring and risk reduction is important. Finally, patients with such rare disorders can provide insight into the possible long-term consequences of new LDL-lowering biologic therapies.

References

- 1 Steinberg D, Grundy SM, Mok HY, et al: Metabolic studies in an unusual case of asymptomatic familial hypobetalipoproteinemia with hypolipolipoproteinemia and fasting chylomicronemia. *J Clin Invest* 1979;64:292–301.
- 2 Heiss G, Tamir I, Davis CE, et al: Lipoprotein-cholesterol distributions in selected North American populations: the lipid research clinics program prevalence study. *Circulation* 1980;61:302–315.
- 3 Contois J, McNamara JR, Lammi-Keefe C, et al: Reference intervals for plasma apolipoprotein A-1 determined with a standardized commercial immunoturbidimetric assay: results from the Framingham Offspring Study. *Clin Chem* 1996;42:507–514.
- 4 Cladaras C, Hadzopoulou-Cladaras M, Nolte RT, et al: The complete sequence and structural analysis of human apolipoprotein B-100: relationship between apoB-100 and apoB-48 forms. *EMBO J* 1986;5:3495–3507.
- 5 Herbert PN, Hyams JS, Bernier DN, et al: Apolipoprotein B-100 deficiency. Intestinal steatosis despite apolipoprotein B-48 synthesis. *J Clin Invest* 1985;76:403–412.
- 6 Lusis AJ, West R, Mehrabian M, et al: Cloning and expression of apolipoprotein B, the major protein of low and very low density lipoproteins. *Proc Natl Acad Sci USA* 1985;82:4597–4601.
- 7 Tarugi P, Averna M, Di Leo E, et al: Molecular diagnosis of hypobetalipoproteinemia: an ENID review. *Atherosclerosis* 2007;195:e19–e27.
- 8 Hegele RA: Plasma lipoproteins: genetic influences and clinical implications. *Nat Rev Genet* 2009;10:109–121.
- 9 Johansen CT, Kathiresan S, Hegele RA: Genetic determinants of plasma triglycerides. *J Lipid Res* 2011;52:189–206.
- 10 Musunuru K, Pirruccello JP, Do R, et al: Exome sequencing, ANGPTL3 mutations, and familial combined hypolipidemia. *N Engl J Med* 2010;363:2220–2227.
- 11 Zhao Z, Tuakli-Wosornu Y, Lagace TA, et al: Molecular characterization of loss-of-function mutations in PCSK9 and identification of a compound heterozygote. *Am J Hum Genet* 2006;79:514–523.
- 12 Raal FJ, Santos RD, Blom DJ, et al: Mipomersen, an apolipoprotein B synthesis inhibitor, for lowering of LDL cholesterol concentrations in patients with homozygous familial hypercholesterolaemia: a randomised, double-blind, placebo-controlled trial. *Lancet* 2010;375:998–1006.
- 13 Samaha FF, McKenney J, Bloedon LT, et al: Inhibition of microsomal triglyceride transfer protein alone or with ezetimibe in patients with moderate hypercholesterolemia. *Nat Clin Pract Cardiovasc Med* 2008;5:497–505.
- 14 Crunkhorn S: Trial watch: PCSK9 antibody reduces LDL cholesterol. *Nat Rev Drug Discov* 2012;11:11.
- 15 Young SG, Bertics SJ, Curtiss LK, et al: Genetic analysis of a kindred with familial hypobetalipoproteinemia. Evidence for two separate gene defects: one associated with an abnormal apolipoprotein B species, apolipoprotein B-37; and a second associated with low plasma concentrations of apolipoprotein B-100. *J Clin Invest* 1987;79:1842–1851.
- 16 Linton MF, Pierotti V, Young SG: Reading-frame restoration with an apolipoprotein B gene frameshift mutation. *Proc Natl Acad Sci USA* 1992;89:11431–11435.
- 17 Collins DR, Knott TJ, Pease RJ, et al: Truncated variants of apolipoprotein B cause hypobetalipoproteinemia. *Nucleic Acids Res* 1988;16:8361–8375.
- 18 Krul ES, Kinoshita M, Talmud P, et al: Two distinct truncated apolipoprotein B species in a kindred with hypobetalipoproteinemia. *Arteriosclerosis* 1989;9:856–868.
- 19 Parhofer KG, Barrett PH, Bier DM, Schonfeld G: Lipoproteins containing the truncated apolipoprotein, apo B-89, are cleared from human plasma more rapidly than apo B-100-containing lipoproteins in vivo. *J Clin Invest* 1992;89:1931–1937.
- 20 Young SG, Hubl ST, Chappell DA, et al: Familial hypobetalipoproteinemia associated with a mutant species of apolipoprotein B (B-46). *N Engl J Med* 1989;320:1604–1610.
- 21 Young SG, Hubl ST, Smith RS, et al: Familial hypobetalipoproteinemia caused by a mutation in the apolipoprotein B gene that results in a truncated species of apolipoprotein B (B-31). A unique mutation that helps to define the portion of the apolipoprotein B molecule required for the formation of buoyant, triglyceride-rich lipoproteins. *J Clin Invest* 1990;85:933–942.
- 22 Huang LS, Ripps ME, Korman SH, et al: Hypobetalipoproteinemia due to an apolipoprotein B gene exon 21 deletion derived by Alu-Alu recombination. *J Biol Chem* 1989;264:11394–11400.
- 23 Welty FK, Hubl ST, Pierotti VR, Young SG: A truncated species of apolipoprotein B (B67) in a kindred with familial hypobetalipoproteinemia. *J Clin Invest* 1991;87:1748–1754.

- 24 McCormick SP, Fellowes AP, Walmsley TA, George PM: Apolipoprotein B-32: a new truncated mutant of human apolipoprotein B capable of forming particles in the low density lipoprotein range. *Biochim Biophys Acta* 1992;1138:290–296.
- 25 Talmud PJ, Converse C, Krul E, et al: A novel truncated apolipoprotein B (apo B55) in a patient with familial hypobetalipoproteinemia and atypical retinitis pigmentosa. *Clin Genet* 1992;42:62–70.
- 26 Yue P, Yuan B, Gerhard DS, et al: Novel mutations of apoB cause apoB truncations undetectable in plasma and familial hypobetalipoproteinemia. *Hum Mutat* 2002;20:110–116.
- 27 Hegele RA, Miskie BA: Acanthocytosis in a patient with homozygous familial hypobetalipoproteinemia due to a novel apoB splice site mutation. *Clin Genet* 2002;61:101–103.
- 28 Tarugi P, Lonardo A, Ballarini G, et al: Fatty liver in heterozygous hypobetalipoproteinemia caused by a novel truncated form of apolipoprotein B. *Gastroenterology* 1996;111:1125–1133.
- 29 Tarugi P, Lonardo A, Gabelli C, et al: Phenotypic expression of familial hypobetalipoproteinemia in three kindreds with mutations of apolipoprotein B gene. *J Lipid Res* 2001;42:1552–1561.
- 30 Pulai JI, Zakeri H, Kwok PY, et al: Donor splice mutation (665 + 1 G_T) in familial hypobetalipoproteinemia with no detectable apoB truncation. *Am J Med Genet* 1998;80:218–220.
- 31 Talmud P, King-Underwood L, Krul E, et al: The molecular basis of truncated forms of apolipoprotein B in a kindred with compound heterozygous hypobetalipoproteinemia. *J Lipid Res* 1989;30:1773–1779.
- 32 Welty FK, Guida KA, Andersen JJ: Donor splice-site mutation (210 + 1G_C) in the apoB gene causes a very low level of apoB-100 and LDL cholesterol. *Arterioscler Thromb Vasc Biol* 2001;21:1864–1865.
- 33 Welty FK, Ordovas J, Schaefer EJ, et al: Identification and molecular analysis of two apoB gene mutations causing low plasma cholesterol levels. *Circulation* 1995;92:2036–2040.
- 34 Pullinger CR, Hillas E, Hardman DA, et al: Two apolipoprotein B gene defects in a kindred with hypobetalipoproteinemia, one of which results in a truncated variant, apoB-61, in VLDL and LDL. *J Lipid Res* 1992;33:699–710.
- 35 Hardman DA, Pullinger CR, Hamilton RL, et al: Molecular and metabolic basis for the metabolic disorder normotriglyceridemic abetalipoproteinemia. *J Clin Invest* 1991;88:1722–1729.
- 36 Young SG, Pullinger CR, Zysow BR, et al: Four new mutations in the apolipoprotein B gene causing hypobetalipoproteinemia, including two different frameshift mutations that yield truncated apolipoprotein B proteins of identical length. *J Lipid Res* 1993;34:501–507.
- 37 Talmud PJ, Krul ES, Pessah M, et al: Donor splice mutation generates a lipid-associated apolipoprotein B-27.6 in a patient with homozygous hypobetalipoproteinemia. *J Lipid Res* 1994;35:468–477.
- 38 Tarugi P, Aversa M: Hypobetalipoproteinemia: genetics, biochemistry, and clinical spectrum. *Adv Clin Chem* 2011;54:81–107.
- 39 Groenewegen WA, Aversa MR, Pulai J, et al: Apolipoprotein B-38.9 does not associate with apo[a] and forms two distinct HDL density particle populations that are larger than HDL. *J Lipid Res* 1994;35:1012–1025.
- 40 Zhu XF, Noto D, Seip R, et al: Organ loci of catabolism of short truncations of apoB. *Arterioscler Thromb Vasc Biol* 1997;17:1032–1038.
- 41 Chen Z, Saffitz JE, Latour MA, Schonfeld G: Truncated apo B-70.5-containing lipoproteins bind to megalin but not the LDL receptor. *J Clin Invest* 1999;103:1419–1430.
- 42 Parhofer KG, Daugherty A, Kinoshita M, Schonfeld G: Enhanced clearance from plasma of low density lipoproteins containing a truncated apolipoprotein, apoB-89. *J Lipid Res* 1990;31:2001–2007.
- 43 Linton MF, Farese RV Jr, Young SG: Familial hypobetalipoproteinemia. *J Lipid Res* 1993;34:521–541.
- 44 Chen Z, Fitzgerald RL, Li G, et al: Hepatic secretion of apoB-100 is impaired in hypobetalipoproteinemic mice with an apoB-38.9-specifying allele. *J Lipid Res* 2004;45:155–163.
- 45 Elias N, Patterson BW, Schonfeld G: In vivo metabolism of ApoB, ApoA-I, and VLDL triglycerides in a form of hypobetalipoproteinemia not linked to the ApoB gene. *Arterioscler Thromb Vasc Biol* 2000;20:1309–1315.
- 46 Yue P, Tanoli T, Wilhelm O, et al: Absence of fatty liver in familial hypobetalipoproteinemia linked to chromosome 3p21. *Metabolism* 2005;54:682–688.
- 47 Yuan B, Neuman R, Duan SH, et al: Linkage of a gene for familial hypobetalipoproteinemia to chromosome 3p21.1–22. *Am J Hum Genet* 2000;66:1699–1704.
- 48 Horton JD, Cohen JC, Hobbs HH: Molecular biology of PCSK9: its role in LDL metabolism. *Trends Biochem Sci* 2007;32:71–77.
- 49 Tarugi P, Lonardo A: Heterozygous familial hypobetalipoproteinemia associated with fatty liver. *Am J Gastroenterol* 1997;92:1400–1402.

- 50 Lonardo A, Tarugi P, Ballarini G, Bagni A: Familial heterozygous hypobetalipoproteinemia, extrahepatic primary malignancy, and hepatocellular carcinoma. *Dig Dis Sci* 1998;43:2489–2492.
- 51 Schonfeld G, Patterson BW, Yablonskiy DA, et al: Fatty liver in familial hypobetalipoproteinemia: triglyceride assembly into VLDL particles is affected by the extent of hepatic steatosis. *J Lipid Res* 2003;44:470–478.
- 52 Whitfield AJ, Barrett PH, Robertson K, et al: Liver dysfunction and steatosis in familial hypobetalipoproteinemia. *Clin Chem* 2005;51:266–269.
- 53 Burnett JR, Shan J, Miskie BA, et al: A novel nontruncating APOB gene mutation, R463W, causes familial hypobetalipoproteinemia. *J Biol Chem* 2003;278:13442–13452.
- 54 Harada N, Soejima Y, Taketomi A, et al: Recurrent familial hypobetalipoproteinemia-induced nonalcoholic fatty liver disease after living donor liver transplantation. *Liver Transpl* 2009;15:806–809.
- 55 Traber MG, Sokol RJ, Ringel SP, et al: Lack of tocopherol in peripheral nerves of vitamin E-deficient patients with peripheral neuropathy. *N Engl J Med* 1987;317:262–265.
- 56 Kayden HJ, Traber MG: Absorption, lipoprotein transport, and regulation of plasma concentrations of vitamin E in humans. *J Lipid Res* 1993;34:343–358.
- 57 Buonomo PS, Ruggiero A, Valeriani M, Mariotti P: Familial hypobetalipoproteinemia: early neurological, hematological, and ocular manifestations in two affected twins responding to vitamin supplementation. *Curr Opin Pediatr* 2009;21:824–827.
- 58 Granot E, Deckelbaum RJ: Hypocholesterolemia in childhood. *J Pediatr* 1989;115:171–185.
- 59 Scott BB, Miller JP, Losowsky MS: Hypobetalipoproteinaemia – a variant of the Bassen-Kornzweig syndrome. *Gut* 1979;20:163–168.
- 60 Hegele RA, Angel A: Arrest of neuropathy and myopathy in abetalipoproteinemia with high-dose vitamin E therapy. *Can Med Assoc J* 1985;132:41–44.
- 61 Bishara S, Merin S, Cooper M, et al: Combined vitamin A and E therapy prevents retinal electrophysiological deterioration in abetalipoproteinaemia. *Br J Ophthalmol* 1982;66:767–770.
- 62 Chowers I, Banin E, Merin S, Cooper M, Granot E: Long-term assessment of combined vitamin A and E treatment for the prevention of retinal degeneration in abetalipoproteinaemia and hypobetalipoproteinaemia patients. *Eye (Lond)* 2001;15:525–530.
- 63 Burnett JR, Hooper AJ: Common and rare gene variants affecting plasma LDL cholesterol. *Clin Biochem Rev* 2008;29:11–26.