Introduction

Martin J. Kelly

Department of Physiology and Pharmacology, L334, Oregon Health and Science University, Portland, Oreg., USA

The concept of ‘rapid’ nongenomic signaling was foreign to neuroendocrinologists in the last century even though there was a plethora of publications documenting the rapid, almost instantaneous effects of steroids on neuronal firing [1–3]. In fact the short latency for the onset of the effects of steroids (e.g. 17β -estradiol, E_2) delivered directly onto neuronal membranes was equivalent to the rapid actions of neurotransmitters. The 1980’s ushered in the era of ‘neurosteroids’. Steroid hormones and their active metabolites were shown to be synthesized *de novo* in the brain and have direct actions on membrane receptors, the best known were allosteric actions on $GABA_A$ channel activity [4, 5]. However, endocrinologists remained skeptical about the ‘pharmacological’ micromolar concentrations necessary to modulate $GABA_A$ channel activity. Some even dismissed the effects of subnanomolar concentrations of gonadal steroids on hypothalamic and hippocampal neurons because of the lack of identifiable steroid receptors at the membrane for mediating these rapid actions [6, 7]. However, this all began to change in the late 1990s based on several papers that documented the membrane localization of estrogen receptor- α (ER α) in pituitary cells and even primary cultures of hippocampal CA1 neurons [8, 9]. Moreover, a seminal paper reported that nuclear and membrane receptors were encoded by the same estrogen receptor genes [10]. The resulting proteins were shown to complex with G protein signaling cascades. As the new century began, several groups identified membrane (m)ERs that were not derived from ER α or ER β transcripts [11–13] includ-

ing a bona fide G protein-coupled receptor, GPR30/GPER1 [14, 15]. It was evident from the interrogation of the ‘nongenomic’ signaling that while some of these signaling cascades initiated at the membrane are tied to rapid membrane effects on ion channel activity, others lead to the regulation of gene transcription – similar to the membrane-to-nucleus signaling described for many neurotransmitters. In this light, it has been more accurate to differentiate between membrane-initiated signaling and nuclear-initiated signaling when discussing hormone actions in neurons and nonneural cells [16].

In the last decade, there has been an explosion in the number of publications documenting the physiological importance of membrane-initiated signaling of steroids in the CNS. Both the number of steroids – gonadal, adrenal, neurosteroids – and the number of brain functions that they affect are much too numerous to cover in one issue. Therefore, this special issue focuses on the gonadal steroid receptors – estrogen receptors (ERs), androgen receptors (AR) and progesterone receptors (PRs) – and their role in vital mammalian physiological functions. These include the neuroendocrine regulation of reproduction and energy homeostasis (Micevych and Kelly), stress responses (Handa and coworkers) and neuroprotection (Simpkins and coworkers) by ER; male sex behavior, aggressive behavior (Cunningham and coworkers) and anabolic steroid abuse (Oberlander and coworkers) by AR, as well as female sex behavior by the classic PR (Mani and Blaustein) and pulsatile GnRH release by a novel mPR (Thomas and Pang).

References

- 1 Mandelbrod I, Feldman S, Werman R: Inhibition of firing is the primary effect of microelectrophoresis of cortisol to units in the rat tuberal hypothalamus. *Brain Res* 1974;80:303–315.
- 2 Kelly MJ, Moss RL, Dudley CA: Differential sensitivity of preoptic-septal neurons to microelectrophoresed estrogen during the estrous cycle. *Brain Res* 1976;114:152–157.
- 3 Teyler TJ, Vardaris RM, Lewis D, Rawitch AB: Gonadal steroids: effects on excitability of hippocampal pyramidal cells. *Science* 1980;209:1017–1019.
- 4 Majewska MD, Harrison NL, Schwartz RD, Barker JL, Paul SM: Steroid hormone metabolites are barbiturate-like modulators of the GABA receptor. *Science* 1986;232:1004–1007.
- 5 Mellon SH: Neurosteroids: Biochemistry, modes of action, and clinical relevance. *J Clin Endocr Metab* 1994;78:1003–1008.
- 6 Mermelstein PG, Becker JB, Surmeier DJ: Estradiol reduces calcium currents in rat neostriatal neurons via a membrane receptor. *J Neurosci* 1996;16:595–604.
- 7 Lagrange AH, Rønnekleiv OK, Kelly MJ: Modulation of G protein-coupled receptors by an estrogen receptor that activates protein kinase A. *Mol Pharmacol* 1997;51:605–612.
- 8 Pappas TC, Gametchu B, Yannariello-Brown J, Collins TJ, Watson CS: Membrane estrogen receptors in GH3/B6 cells are associated with rapid estrogen-induced release of prolactin. *Endocrine* 1994;2:813–822.
- 9 Clarke CH, Norfleet AM, Clarke MSF, Watson CS, Cunningham KA, Thomas ML: Perimembrane localization of the estrogen receptor α protein in neuronal processes of cultured hippocampal neurons. *Neuroendocrinology* 2000;71:34–42.
- 10 Razandi M, Pedram A, Greene GL, Levin ER: Cell membrane and nuclear estrogen receptors (ERs) originate from a single transcript: studies of ER α and ER β expressed in Chinese hamster ovary cells. *Mol Endocrinol* 1999;13:307–319.
- 11 Toran-Allerand CD, Guan X, MacLusky NJ, Horvath TL, Diano S, Singh M, Connolly ES Jr, Nethrapalli IS, Tinnikov AA: ER-X: a novel, plasma membrane-associated, putative estrogen receptor that is regulated during development and after ischemic brain injury. *J Neurosci* 2002;22:8391–8401.
- 12 Qiu J, Bosch MA, Tobias SC, Grandy DK, Scanlan TS, Rønnekleiv OK, Kelly MJ: Rapid signaling of estrogen in hypothalamic neurons involves a novel G protein-coupled estrogen receptor that activates protein kinase C. *J Neurosci* 2003;23:9529–9540.
- 13 Qiu J, Bosch MA, Jamali K, Xue C, Kelly MJ, Rønnekleiv OK: Estrogen upregulates T-type calcium channels in the hypothalamus and pituitary. *J Neurosci* 2006;26:11072–11082.
- 14 Filardo EJ, Quinn JA, Bland KI, Frackelton ARJ: Estrogen-induced activation of Erk-1 and Erk-2 requires the G protein-coupled receptor homolog, GPR30, and occurs via trans-activation of the epidermal growth factor receptor through release of HB-EGF. *Mol Endocrinol* 2000;14:1649–1660.
- 15 Revankar CM, Cimino DF, Sklar LA, Arterburn JB, Prossnitz ER: A transmembrane intracellular estrogen receptor mediates rapid cell signaling. *Science* 2005;307:1625–1630.
- 16 Hammes SR, Levin ER: Extra-nuclear steroid receptors: nature and actions. *Endocr Rev* 2007;28:726–741.