Thyroid Hormone Actions in Cartilage and Bone

Graham R. Williams

Molecular Endocrinology Group, Department of Medicine, Imperial College London, London, UK

Key Words

Thyroid hormone • Cartilage • Chondrocyte • Ossification • Osteoarthritis • Bone • Bone turnover • Osteoblast • Osteoclast • Osteoporosis

Abstract

Thyroid hormones exert widespread and complex actions in almost all tissues during development, throughout childhood and in adults. The skeleton is an important T3-target tissue that exemplifies these processes, and yet understanding of the specific cellular and molecular mechanisms of T3 action in bone and cartilage remains incomplete. Here, the skeleton is considered as a T3-target tissue. The actions of thyroid hormones during skeletal development and in chondrocytes and growth plate cartilage during post-natal linear growth are outlined. The physiological importance of these actions are discussed in relation to patients with autosomal dominant mutations in genes encoding the thyroid hormone receptors TR α 1 and TR β , and in mice harbouring deletions or mutations of the orthologous genes. The role of thyroid hormones and the control of T3 action in bone turnover and maintenance are also outlined, and T3 action in bone-forming osteoblasts and bone-resorbing osteoclasts discussed. The physiological and functional consequences of

T3 action in bone are considered in relation to mutant mouse models and to effects on bone mineral density and fracture susceptibility in humans. Finally, new studies identifying a putative role for thyroid hormone metabolism in articular cartilage maintenance and the pathogenesis of osteoarthritis are considered. The pharmacological context of these new findings is discussed, emphasising the importance of this emerging field of study in thyroid hormone pathophysiology.

Copyright © 2012 European Thyroid Association
Published by S. Karger AG, Basel

Regulation of Thyroid Hormone Action

Hypothalamic-Pituitary-Thyroid Axis

The thyroid gland produces mainly the pro-hormone T4 (3,5,3',5'-L-tetraiodothyronine, thyroxine) but also secretes smaller amounts of the active hormone T3 (3,5,3'-L-triiodothyronine). Most circulating T3 is derived via metabolism of T4, from which an outer-ring iodine atom is removed by activity of the type 1 iodothyronine deiodinase enzyme (Dio1) principally in liver and kidney [1]. The hypothalamic-pituitary-thyroid (HPT) axis senses circulating thyroid hormone concentrations and physiological thyroid status is controlled by negative feedback

regulation of thyrotropin-releasing hormone (TRH) in the hypothalamus and thyroid-stimulating hormone (TSH, thyrotropin) synthesis and secretion in the anterior pituitary [2]. More than 95% of circulating thyroid hormones are bound to plasma proteins and the remaining free hormones (fT4, fT3) represent the biologically available and active fraction.

Cellular T3 Availability

fT4 and fT3 enter peripheral target cells via specific cell membrane transporter proteins with differing patterns of tissue distribution. Several thyroid hormone transporters have been identified, including the monocarboxylate transporters 8 and 10 (MCT8, MCT10), the organic acid transporter protein-1c1 (OATP1c1) and the non-specific L-type amino acid transporters 1 and 2 (LAT1, LAT2) [3]. Once inside the cell, thyroid hormones are metabolized further. The type 2 deiodinase (Dio2) catalyses conversion of T4 to T3 by removal of an outer-ring iodine thus activating the pro-hormone, whereas the type 3 enzyme (Dio3) inactivates T4 and T3 by removal of a 5-iodine atom [4]. The relative tissue activities of these two enzymes determine the intracellular T3 concentration and provide a pre-receptor control mechanism that regulates availability of the active hormone to the cell nucleus.

Thyroid Hormone Receptors

The thyroid hormone receptors TR α 1, TR β 1 and TR β 2, bind ligand with high affinity and act as hormone-inducible transcription factors that regulate expression of target genes in response to T3. The TRs are located in the nucleus even in the absence of hormone and, in the case of genes that are positively regulated by T3, these unoccupied TRs interact with co-repressor proteins to repress target gene transcription [5]. After T3 binding, co-repressor is released and co-activator proteins are recruited so that hormone-dependent activation of gene expression is initiated in T3-responsive cells. The TR α 1 and TR β 1 isoforms are expressed in almost all tissues, but in different relative concentrations that vary at discrete stages of development and with age [6], whereas TR β 2 expression is restricted to the pituitary and hypothalamus [7]. Overall, pre-receptor control of T3 availability, together with the diversity of TR isoform expression, provides a sophisticated mechanism that coordinates and regulates temporal and spatial responses to thyroid hormones in critical target tissues during development and in adulthood [5, 6, 8]. T3 action is also regulated by tissue-specific differences in expression of co-

repressor and co-activator proteins that further modulate TR activity [9–12].

The Skeleton Is a T3-Target Organ

MCT10 appears to be the major thyroid hormone-specific transporter expressed in the growth plate, although MCT8 is expressed widely in cartilage-forming chondrocytes, bone-forming osteoblasts and bone-resorbing osteoclasts, whilst expression of the less specific LAT1 and LAT2 transporters has also been identified in skeletal tissue. By contrast, OATP1c1 mRNA is not expressed in bone or cartilage [13–15]. Thyroid hormone metabolism also occurs in skeletal cells [16]. Although Dio1 is not expressed in cartilage and bone [14, 16], the activating enzyme Dio2 is expressed only in osteoblasts [14, 17], whilst the inactivating Dio3 enzyme is present in all skeletal cell lineages, particularly during development and prior to weaning [14, 18]. The T3 receptors, TR α 1 and TR β 1, are each expressed in the skeleton and quantitative RT-PCR studies indicate that TR α 1 mRNA is expressed at levels at least ten-fold greater than TR β 1 [19, 20]. Both receptor isoforms are present in growth plate chondrocytes, bone marrow stromal cells and bone-forming osteoblasts. However, it is not certain whether TRs are expressed in bone-resorbing osteoclasts or in osteocytes, which communicate environmental, endocrine and paracrine cues and coordinate bone formation and bone resorption [16]. Thus, all the factors required for locally regulated T3 action, including thyroid hormone transporters, metabolizing enzymes and receptors, are present in cartilage and bone indicating the skeleton is a physiological target tissue for thyroid hormone throughout life.

Cartilage

Endochondral Ossification, Bone Formation and Linear Growth

The formation of long bones and linear growth occur by the process of endochondral ossification [21, 22]. During endochondral ossification, condensations of mesenchyme form a cartilage scaffold that supports bone development (fig. 1). Chondrocyte progenitor cells undergo clonal expansion and proliferate before differentiating into hypertrophic chondrocytes and ultimately undergoing programmed cell death. During this process, a mineralizing cartilage matrix is secreted, which forms the scaffold for vascular invasion and bone deposition. Newly laid down bone is continuously remodelled by bone-resorbing osteoclasts and bone-forming osteoblasts in or-

Fig. 1. Endochondral ossification in the epiphyseal growth plate. Chondrocytes are organized into discrete columns, and undergo an ordered process of cell proliferation, differentiation, volume expansion and apoptosis. Differentiating hypertrophic chondrocytes secrete a cartilage matrix that is rich in type X collagen, whilst mature cells secrete proteases and matrix-degrading enzymes including matrix metalloproteinase-13 and aggrecanase. Growth factors and cytokines released during matrix degradation promote new blood vessel formation from the metaphyseal bone marrow in order to facilitate migration of osteoblasts and bone formation. Thyroid hormones act at all levels to regulate the process of ossification thereby controlling the rate of bone maturation and linear growth.

der to shape the developing long bone in response to local mechanical stresses and paracrine signalling cues. Linear growth subsequently occurs via the same orderly mechanism in specialized epiphyseal growth plates positioned at the proximal and distal extremities of long bones. The process of endochondral ossification continues until early adulthood when the growth plates become quiescent and ultimately fuse, resulting in cessation of linear growth [21, 22]. Bone mineral deposition and accrual of bone mass continue after linear growth has stopped such that peak bone mass is reached by the age of approximately 25 years.

Thyroid Hormone Action in Growth Plate Chondrocytes

TR α 1 and TR β 1 are expressed in reserve and proliferating chondrocytes, indicating the epiphyseal growth plate is directly responsive to T3 [23] (fig. 1). T3 stimulates clonal expansion of chondrocyte progenitor cells, but inhibits subsequent cell proliferation whilst promoting hypertrophic chondrocyte differentiation and cell

volume expansion [23–26]. Thus, in primary growth plate chondrocyte cultures, T3 stimulates alkaline phosphatase and collagen X expression, and enhances cartilage matrix mineralization [23]. During endochondral ossification the pace of chondrocyte proliferation and differentiation is regulated by several paracrine factors, including insulin-like growth factor-1 (IGF1), Wnt morphogens, bone morphogenetic proteins (BMPs) and fibroblast growth factors (FGFs), and by a negative feedback loop involving Indian hedgehog (Ihh) and parathyroid hormone-related peptide (PTHrP) [27–29]. The set-point of this feedback loop and the rate of linear growth are sensitive to changes in thyroid status in vivo and regulated by local thyroid hormone metabolism and T3 availability [30, 31]. T3 also regulates expression of growth plate matrix proteoglycans [32] and collagen-degrading enzymes such as aggrecanase-2 (a disintegrin and metalloproteinase with thrombospondin motifs 1, ADAMTS5) and matrix metalloproteinase-13 (MMP13) [33–35]. In summary, thyroid hormone is essential for the coordinated progression of endochondral ossification,

acting to stimulate expression of genes that control chondrocyte maturation and cartilage matrix synthesis, mineralization and degradation.

Thyroid Hormone Effects on Linear Growth and Skeletal Development

During childhood and adolescence, hypothyroidism causes growth arrest resulting in epiphyseal dysgenesis and severely delayed bone age [36, 37]. Thyroxine replacement results in 'catch-up' growth, although full predicted final height may not be reached, especially when the diagnosis and treatment of hypothyroidism is delayed. Thyrotoxicosis in childhood accelerates linear growth and advances bone age, but ultimately results in persistent short stature due to early fusion of the epiphyses. In severe cases, craniosynostosis results from premature closure of the skull sutures and fontanelles and may be associated with intellectual deficit [38, 39]. Furthermore, in congenital hyperthyroidism resulting from constitutive activating mutations of the TSH receptor gene (*TSHR*), advanced bone age has been reported, although early thyroidectomy is effective to prevent long-term consequences of hyperthyroidism [40]. These observations indicate extraordinary sensitivity of the developing skeleton to thyroid hormones, and reveal that euthyroid status is essential for normal skeletal development and post-natal growth.

Syndromes of Resistance to Thyroid Hormone

Resistance to thyroid hormone (RTH) is an autosomal dominant condition caused by mutations of *THRB* that result in expression of dominant-negative TR β proteins. The syndrome is characterized by mildly or moderately increased thyroid hormone concentrations and an inappropriately normal or elevated TSH due to impaired negative feedback control of the HPT axis. Patients with RTH display variable skeletal phenotypes that are confounded by the effects of treatment and the expression of heterogeneous TR β mutations, which have variable functional properties and activities [41]. Two reports recently described the first individuals with an RTH syndrome resulting from heterozygous mutations of *THRA* resulting in expression of dominant-negative TR α 1 proteins [42, 43]. Subjects have normal levels of TSH but free and total T4 levels lie within or just below the normal range and free and total T3 levels are within or just above the normal range, leading to a markedly reduced T4:T3 ratio. Patients display a phenotype reminiscent of the features of hypothyroidism that include delayed growth with persistent short stature, impaired tooth eruption and patent

fontanelles with thickening of the skull vault. These features are consistent with retarded intramembranous and endochondral ossification and demonstrate a critical role for TR α 1 in the human skeleton [42, 43]. Mutations in the selenocysteine insertion sequence binding protein 2 gene (*SECISBP2*) cause a complex multisystem disorder that includes thyroid dysfunction and RTH, which result from abnormal thyroid hormone metabolism due to reduced deiodinase enzyme activity [44, 45]. Affected individuals have growth retardation and delayed bone age that respond to treatment with T3 [46], further demonstrating the requirement for thyroid hormones during growth and skeletal development.

The recent studies in individuals with *THRA* mutations are entirely consistent with conclusions from studies of mice with mutations or deletions affecting the *Thra* and *Thrb* genes [20] (fig. 2). T3 action in bone is mediated principally by TR α 1, which is expressed at much higher levels than TR β in the skeleton [19, 20]. Mice harbouring knockout or dominant-negative point mutations of *Thra* are euthyroid but display a skeletal phenotype characteristic of juvenile hypothyroidism that includes impaired intramembranous and endochondral ossification with reduced bone mineral deposition during skeletal development and delayed growth [20, 47–51]. Mice with knockout or dominant-negative point mutations of *Thrb* have disrupted negative feedback regulation of the HPT axis and RTH, but display a skeletal phenotype consistent with the effects of systemic hyperthyroidism on bone. Thus, juvenile TR β mutant mice have advanced ossification with increased bone mineral deposition but display short stature due to accelerated growth plate maturation [20, 48, 49, 51]. The contrasting phenotypes in mice with *Thra* and *Thrb* mutations demonstrate that TR α 1 is the major mediator of T3 action in the skeleton. In TR α 1 mutant mice skeletal features of hypothyroidism result directly from impaired T3 action in bone and cartilage, whereas the consequences of *Thrb* mutations are indirect because the elevated thyroid hormones result in an increased skeletal response to T3 that is mediated by the wild-type TR α 1 protein expressed in bone [52].

Bone

Bone Remodelling Cycle

The skeleton undergoes continuous remodelling in response to mechanical stress and injury at multiple sites throughout the skeleton in order to maintain structural

Fig. 2. TRα mediates T3 action in bone. Upper panels show the consequences of deletion or mutation of TRα (left) or TRβ (right) on regulation of the hypothalamic-pituitary feedback axis. Mutation of TRα does not influence negative feedback regulation of TRH and TSH by thyroid hormones because TRβ is the predominant receptor expressed in the pituitary. Thus, TRα mutants are euthyroid but display features of impaired T3 action in the skeleton because bone and cartilage predominantly express TRα. By contrast, deletion or mutation of TRβ disrupts the HPT axis leading to systemic hyperthyroidism. Thus, TRβ mutants display features of increased T3 action in the skeleton because the high levels of circulating thyroid hormones overstimulate the intact TRα ex-

pressed in bone. The middle panels show the epiphyseal growth plates in juveniles. In TRα mutants, endochondral ossification is retarded as evidenced by disorganized and wide growth plates and delay in formation of the secondary ossification centre. In TRβ mutants, endochondral ossification is advanced with growth plate narrowing and increased bone deposition evident in the secondary ossification centre and metaphysis. The lower panels show adult trabecular bone in TRα and TRβ mutants. In TRα mutants, increased bone mass (osteosclerosis) results from reduced bone turnover due to impaired T3 action in bone, whereas osteoporosis in TRβ mutants results from accelerated bone turnover due to the effects of systemic hyperthyroidism.

integrity and strength [53, 54] (fig. 3). The cyclical process of bone turnover and repair is initiated by osteocytes. These cells are embedded within calcified bone and communicate via an elaborate network of dendritic processes. Osteocytes respond to changes in mechanical loading or micro-fracture by undergoing apoptosis with release of cytokines and growth factors that attract osteoclasts to

sites of micro-damage. Osteoclasts resorb areas of damaged bone and communicate with osteoblasts, which are then attracted by various growth factors and by degraded matrix proteins released during bone resorption. Osteoblasts subsequently synthesize, secrete and mineralize osteoid to lay down new bone. Completion of the formation phase of the bone remodelling cycle by osteoblasts

Fig. 3. The bone remodelling cycle. Sites of micro-damage or fracture are sensed by osteocytes, which are embedded within bone and respond by undergoing apoptosis. Release of various chemoattractants by these osteocytes results in recruitment of osteoclasts, which resorb damaged bone and communicate with osteoblasts by releasing of degraded bone matrix proteins and growth factors during bone resorption. Osteoblasts respond by secreting and mineralizing osteoid to form new bone that ultimately repairs the site of micro-damage to maintain structural integrity of the region. Typically the period of bone resorption occurs over a period of 50 days in humans and the bone formation phase has a duration of 150 days.

results in the repair of defective bone. Overall, the balanced coupling of bone resorption to bone formation is essential to maintain the architecture, mineralization and strength of bone [53, 54].

Thyroid Hormone Action in Osteoblasts

Studies of T3 action in osteoblasts are contradictory, but overall thyroid hormones stimulate osteoblast activity. T3 stimulates synthesis and post-translational modification of type I collagen [55], induces expression of alkaline phosphatase [56] and regulates synthesis and secretion of the bone matrix proteins osteopontin and osteocalcin [55–57]. T3 also promotes bone matrix remodelling by stimulating expression of matrix metalloproteinases-9 and -13 [35]. Furthermore, thyroid hormones regulate key pathways involved in osteoblast proliferation and differentiation. Thus, T3 induces IGF-I transcription [58] and stimulates expression of its regulatory binding proteins IGF1BP-2 and IGF1BP-4 [59]. Moreover, T3 induces FGFR1 expression and FGF-induced MAPK signalling in osteoblasts [60], and also regulates activity of the Wnt signalling pathway [61]. Together, these studies indicate T3 regulates osteoblast differentiation and function by complex mechanisms involving numerous paracrine and autocrine factors.

In addition, T3 actions in osteoblasts may indirectly influence the activity of osteoclasts by regulating osteo-

protegerin (OPG) [62]. OPG is a decoy receptor that inhibits receptor activator of nuclear factor- κ B ligand (RANKL) mediated activation of osteoclastogenesis [63]. However, the involvement of T3 in regulation of the OPG/RANKL pathway is controversial as other studies suggest effects of T3 on osteoclastogenesis may be independent of RANKL [64, 65]. Thus, although thyrotoxicosis results in increased osteoclast numbers and activity leading to increased bone resorption in vivo, it is not clear whether T3 acts directly in osteoclasts or whether these responses are secondary to direct actions of T3 in osteoblasts.

Thyroid Hormone Action in Osteoclasts

Osteoclasts express both TR α 1 and TR β 1 mRNAs but it is not known whether functional receptors are present because available TR antibodies lack the sensitivity required to detect endogenous proteins reliably. Furthermore, studies using mixed primary cultures containing osteoclast lineage cells and bone marrow stromal cells were inconclusive and contradictory [66–70]. Nevertheless, treatment of immortalized osteoblasts or primary bone marrow stromal cells resulted in increased expression of RANKL, interleukin-6, interleukin-8 and prostaglandin E₂, consistent with a likely indirect effect of thyroid hormones on osteoclast function [68, 70].

Thyroid Hormone Effects on Bone Turnover and Maintenance

The bone remodelling cycle (fig. 3) is regulated by thyroid hormones. In hypothyroidism there is reduced bone turnover with impaired bone resorption and formation phases. The resulting increase in duration of the bone remodelling cycle leads to a prolonged period of secondary mineralization [71, 72]. Conversely, in thyrotoxicosis, high bone turnover osteoporosis is due to shortening of the remodelling cycle with uncoupling of the activities of osteoclast and osteoblasts that results in a loss of about 10% of mineralized bone per cycle [72]. Consistent with histomorphometry data, population studies have demonstrated that hypothyroidism is associated with a two- to three-fold increased risk of fracture [73, 74], whilst thyrotoxicosis is an established cause of osteoporosis and fragility fracture [74, 75]. Additional studies also show that thyroid hormone replacement, sub-clinical hyperthyroidism and even variation of thyroid status within the normal reference range are inversely correlated with BMD and may be associated with an increased risk of fracture, particularly in post-menopausal women [75–82]. Taken together, all these studies, and many other smaller clinical investigations, reveal that the adult skeleton is sensitive to small but prolonged periods of altered thyroid status, especially in post-menopausal women.

Studies in adult mice with deletion or dominant-negative mutations of the *Thra* and *Thrb* genes are consistent with these clinical data and further elucidate understanding of the cellular mechanisms underlying T3 action in bone (fig. 2). Thus, despite systemic euthyroidism, adult TR α knockout and mutant mice display a phenotype of osteosclerosis with increased bone volume and mineralization accompanied by a net reduction in osteoclastic bone resorption [47, 48]. These findings further support the view that impaired T3 action in skeletal cells results from deletion or mutation of TR α 1. By contrast, adult TR β mutant mice exhibit osteoporosis with increased osteoclastic bone resorption due to the effects of thyroid hormone excess in TR α 1 expressing skeletal cells [47, 48, 51, 52]. Studies of mice with deletion of the *Dio2* gene further demonstrate a critical requirement for T3 in osteoblasts [17]. *Dio2* knockout mice have increased bone mineralization density and brittle bones due to impaired osteoblast activity. Histomorphometry data revealed a reduced bone formation rate with prolongation of the bone remodelling cycle and a longer period of secondary mineralization. Thus, the type 2 deiodinase enzyme is essential to maintain intracellular

supplies of T3 in osteoblasts and plays a critical role to maintain optimal bone mineralization and strength [17]. Taken together, all these studies underscore the important role of thyroid hormones in regulating adult bone maintenance.

Articular Cartilage

It is clear that thyroid hormones have essential roles to regulate bone development, growth and maintenance throughout life, but new studies also indicate the potential importance of thyroid hormones in articular cartilage lining the joint surfaces. Genome-wide association and population cohort studies have identified both *DIO2* and *DIO3* as disease susceptibility loci in osteoarthritis [83–85], and a *DIO2* polymorphism has been associated with hip geometry, an independent risk factor for osteoarthritis [86]. In support of a potential role for thyroid hormones in the regulation of articular cartilage maintenance, increased expression of Dio2 protein has been documented in samples of osteoarthritic cartilage [87]. Furthermore, a phase III clinical trial investigating the use of eprotirome, a TR β -selective agonist, for treatment of hypercholesterolaemia [88] was terminated due to pre-clinical toxicology studies that identified dose-related articular cartilage damage in dogs treated for 12 months. As a consequence of this finding, the pharmaceutical company involved is not pursuing further work with eprotirome and the programme has been closed. Together, these studies identify an important and likely role for thyroid hormones in the pathogenesis of osteoarthritis and demonstrate the critical importance of a detailed understating of T3 action in bone and cartilage. Future opportunities for the use of pharmacological modulators of T3 action in the prevention and treatment of chronic disease may be limited in the light of these recent findings and further progress will require full investigation of the long-term consequences to the skeleton and joints.

Conclusions

Studies of the effects of thyroid hormones in cartilage and bone represent a novel and growing field in our understanding of the broader pathophysiological consequences of thyroid hormone action. Although considerable advances have been made in recent years, there are important gaps in our knowledge and a number of specific areas require investigation.

The actions of thyroid hormones in chondrocytes and osteoblasts have been characterized in detail, yet the principal effect of thyroid hormone excess on the adult skeleton is to stimulate osteoclast-mediated bone resorption. It is not known whether T3 exerts direct actions in osteoclasts or whether osteoclast responses are secondary to the actions of T3 in other cells such as osteoblasts, bone marrow stromal cells or macrophages. A major effect of thyroid hormone excess is uncoupling of the resorption and formation phases of the bone remodelling cycle, resulting in a net loss of bone and osteoporosis. This response has been well known for over a century, but the underlying cellular and molecular mechanisms have not been investigated. The osteocyte represents a key candidate T3-target cell as it orchestrates coupling between osteoblasts and osteoclasts to regulate bone turnover, but its possible T3 responsiveness has not yet been investigated. One key area for future study will be to investigate cell lineage-specific actions of thyroid hormones in the skeleton in vivo, perhaps by adopting tissue-specific gene-targeting approaches using cre-lox recombination technology.

The recent and exciting identification of patients with mutations of the *THRA* gene has provided convincing support for the proposed major role for TR α 1 in the skeleton. Yet the patients raise additional questions that will require new studies. Firstly, it is not clear whether treatment of children with supra-physiological doses of thyroid hormones will improve growth or skeletal development in affected individuals or whether other therapeutic approaches such as additional treatment with growth

hormone, IFG-1 or even histone-modifying agents will be beneficial. Studies in mouse models will be invaluable to inform future treatment options in humans. Secondly, there is almost no information about the effect of TR α 1 mutations on the adult skeleton in humans and little detail is known in mutant mouse models. These are obvious areas for further study that will provide translational benefit to patients in the future.

Finally, identification of a potential role for thyroid hormone signalling in the pathogenesis of osteoarthritis has opened a new field that is sure to attract intense study in the future. The unfortunate cartilage toxicity of eprotirome will also require further investigation to determine whether it is due to drug-specific toxicity or whether it represents a class effect for synthetic TR agonists.

These examples clearly demonstrate the importance and diversity of thyroid hormone action in cartilage and bone. It is a field that is certain to expand and provide new and exciting discoveries in the years ahead.

Acknowledgements

This work was supported by grants from the Medical Research Council (G108/502, G0800261 and G0501486), Wellcome Trust (GR076584) and Arthritis Research UK (18292).

Disclosure Statement

G.R.W. has no conflicts of interest to disclose.

References

- 1 Bianco AC, Kim BW: Deiodinases: implications of the local control of thyroid hormone action. *J Clin Invest* 2006;116:2571–2579.
- 2 Kopp P: The TSH receptor and its role in thyroid disease. *Cell Mol Life Sci* 2001;58:1301–1322.
- 3 Van der Deure W, Peeters R, Visser T: Molecular aspects of thyroid hormone transporters, including MCT8, MCT10 and OATPs, and the effects of genetic variation in these transporters. *J Mol Endocrinol* 2010;44:1–11.
- 4 Bianco AC, Larsen PR: Cellular and structural biology of the deiodinases. *Thyroid* 2005;15:777–786.
- 5 Harvey CB, Williams GR: Mechanism of thyroid hormone action. *Thyroid* 2002;12:441–446.
- 6 Forrest D, Sjoberg M, Vennstrom B: Contrasting developmental and tissue-specific expression of α - and β -thyroid hormone receptor genes. *EMBO J* 1990;9:1519–1528.
- 7 Abel ED, Boers ME, Pazos-Moura C, Moura E, Kaulbach H, Zakaria M, Lowell B, Radovick S, Liberman MC, Wondisford F: Divergent roles for thyroid hormone receptor β -isoforms in the endocrine axis and auditory system. *J Clin Invest* 1999;104:291–300.
- 8 Cheng SY, Leonard JL, Davis PJ: Molecular aspects of thyroid hormone actions. *Endocr Rev* 2010;31:139–170.
- 9 Astapova I, Lee LJ, Morales C, Tauber S, Bilban M, Hollenberg AN: The nuclear corepressor, NCoR, regulates thyroid hormone action in vivo. *Proc Natl Acad Sci USA* 2008;105:19544–19549.
- 10 Astapova I, Vella KR, Ramadoss P, Holtz KA, Rodwin BA, Liao XH, Weiss RE, Rosenberg MA, Rosenzweig A, Hollenberg AN: The nuclear receptor corepressor (NCoR) controls thyroid hormone sensitivity and the set point of the hypothalamic-pituitary-thyroid axis. *Mol Endocrinol* 2011;25:212–224.
- 11 Weiss RE, Gehin M, Xu J, Sadow PM, O'Malley BW, Chambon P, Refetoff S: Thyroid function in mice with compound heterozygous and homozygous disruptions of SRC-1 and TIF-2 coactivators: evidence for haploinsufficiency. *Endocrinology* 2002;143:1554–1557.
- 12 Weiss RE, Xu J, Ning G, Pohlenz J, O'Malley BW, Refetoff S: Mice deficient in the steroid receptor co-activator 1 (SRC-1) are resistant to thyroid hormone. *EMBO J* 1999;18:1900–1904.

- 13 Capelo LP, Beber EH, Fonseca TL, Gouveia CH: The monocarboxylate transporter 8 and L-type amino acid transporters 1 and 2 are expressed in mouse skeletons and in osteoblastic MC3T3-E1 cells. *Thyroid* 2009;19:171–180.
- 14 Williams AJ, Robson H, Kester MH, van Leeuwen JP, Shalet SM, Visser TJ, Williams GR: Iodothyronine deiodinase enzyme activities in bone. *Bone* 2008;43:126–134.
- 15 Abe S, Namba N, Abe M, Fujiwara M, Aikawa T, Kogo M, Ozono K: Monocarboxylate transporter 10 functions as a thyroid hormone transporter in chondrocytes. *Endocrinology* 2012;153:4049–4058.
- 16 Waung JA, Bassett JH, Williams GR: Thyroid hormone metabolism in skeletal development and adult bone maintenance. *Trends Endocrinol Metab* 2012;23:155–162.
- 17 Bassett JH, Boyde A, Howell PG, Bassett RH, Galliford TM, Archanco M, Evans H, Lawson MA, Croucher P, St Germain DL, Galton VA, Williams GR: Optimal bone strength and mineralization requires the type 2 iodothyronine deiodinase in osteoblasts. *Proc Natl Acad Sci USA* 2010;107:7604–7609.
- 18 Capelo LP, Beber EH, Huang SA, Zorn TM, Bianco AC, Gouveia CH: Deiodinase-mediated thyroid hormone inactivation minimizes thyroid hormone signaling in the early development of fetal skeleton. *Bone* 2008;43:921–930.
- 19 Bookout AL, Jeong Y, Downes M, Yu RT, Evans RM, Mangelsdorf DJ: Anatomical profiling of nuclear receptor expression reveals a hierarchical transcriptional network. *Cell* 2006;126:789–799.
- 20 O'Shea PJ, Harvey CB, Suzuki H, Kaneshige M, Kaneshige K, Cheng SY, Williams GR: A thyrotoxic skeletal phenotype of advanced bone formation in mice with resistance to thyroid hormone. *Mol Endocrinol* 2003;17:1410–1424.
- 21 Karsenty G, Wagner EF: Reaching a genetic and molecular understanding of skeletal development. *Dev Cell* 2002;2:389–406.
- 22 Kronenberg HM: Developmental regulation of the growth plate. *Nature* 2003;423:332–336.
- 23 Robson H, Siebler T, Stevens DA, Shalet SM, Williams GR: Thyroid hormone acts directly on growth plate chondrocytes to promote hypertrophic differentiation and inhibit clonal expansion and cell proliferation. *Endocrinology* 2000;141:3887–3897.
- 24 Ballock RT, Reddi AH: Thyroxine is the serum factor that regulates morphogenesis of columnar cartilage from isolated chondrocytes in chemically defined medium. *J Cell Biol* 1994;126:1311–1318.
- 25 Ballock RT, Zhou X, Mink LM, Chen DH, Mita BC, Stewart MC: Expression of cyclin-dependent kinase inhibitors in epiphyseal chondrocytes induced to terminally differentiate with thyroid hormone. *Endocrinology* 2000;141:4552–4557.
- 26 Ishikawa Y, Genge BR, Wuthier RE, Wu LN: Thyroid hormone inhibits growth and stimulates terminal differentiation of epiphyseal growth plate chondrocytes. *J Bone Miner Res* 1998;13:1398–1411.
- 27 Vortkamp A, Lee K, Lanske B, Segre GV, Kronenberg HM, Tabin CJ: Regulation of rate of cartilage differentiation by Indian hedgehog and PTH-related protein. *Science* 1996;273:613–622.
- 28 Minina E, Kreschel C, Naski MC, Ornitz DM, Vortkamp A: Interaction of FGF, Ihh/ Pthlh, and BMP signaling integrates chondrocyte proliferation and hypertrophic differentiation. *Dev Cell* 2002;3:439–449.
- 29 St-Jacques B, Hammerschmidt M, McMahon AP: Indian hedgehog signaling regulates proliferation and differentiation of chondrocytes and is essential for bone formation. *Genes Dev* 1999;13:2072–2086.
- 30 Dentice M, Bandyopadhyay A, Gereben B, Callebaut I, Christoffolete MA, Kim BW, Nissim S, Mornon JP, Zavacki AM, Zeold A, Capelo LP, Curcio-Morelli C, Ribeiro R, Harney JW, Tabin CJ, Bianco AC: The hedgehog-inducible ubiquitin ligase subunit WSB-1 modulates thyroid hormone activation and PTHrP secretion in the developing growth plate. *Nat Cell Biol* 2005;7:698–705.
- 31 Stevens DA, Hassserjian RP, Robson H, Siebler T, Shalet SM, Williams GR: Thyroid hormones regulate hypertrophic chondrocyte differentiation and expression of parathyroid hormone-related peptide and its receptor during endochondral bone formation. *J Bone Miner Res* 2000;15:2431–2442.
- 32 Bassett JH, Swinhoe R, Chassande O, Samarut J, Williams GR: Thyroid hormone regulates heparan sulfate proteoglycan expression in the growth plate. *Endocrinology* 2006;147:295–305.
- 33 Himeno M, Enomoto H, Liu W, Ishizeki K, Nomura S, Kitamura Y, Komori T: Impaired vascular invasion of CBFA1-deficient cartilage engrafted in the spleen. *J Bone Miner Res* 2002;17:1297–1305.
- 34 Makihiro S, Yan W, Murakami H, Furukawa M, Kawai T, Nikawa H, Yoshida E, Hamada T, Okada Y, Kato Y: Thyroid hormone enhances aggrecanase-2/ADAM-TS5 expression and proteoglycan degradation in growth plate cartilage. *Endocrinology* 2003;144:2480–2488.
- 35 Pereira RC, Jorgetti V, Canalis E: Triiodothyronine induces collagenase-3 and gelatinase B expression in murine osteoblasts. *Am J Physiol* 1999;277:E496–E504.
- 36 Boersma B, Otten BJ, Stoeltinga GB, Wit JM: Catch-up growth after prolonged hypothyroidism. *Eur J Pediatr* 1996;155:362–367.
- 37 Rivkees SA, Bode HH, Crawford JD: Long-term growth in juvenile acquired hypothyroidism: the failure to achieve normal adult stature. *N Engl J Med* 1988;318:599–602.
- 38 Schlesinger S, MacGillivray MH, Munschauer RW: Acceleration of growth and bone maturation in childhood thyrotoxicosis. *J Pediatr* 1973;83:233–236.
- 39 Segni M, Leonardi E, Mazzoncini B, Pucarelli I, Pasquino AM: Special features of Graves' disease in early childhood. *Thyroid* 1999;9:871–877.
- 40 Supornsilchai V, Sahakitrungruang T, Wongjitrat N, Wacharasindhu S, Supha-phetiporn K, Shotelersuk V: Expanding clinical spectrum of non-autoimmune hyperthyroidism due to an activating germline mutation, p.M453T, in the thyrotropin receptor gene. *Clin Endocrinol* 2009;70:623–628.
- 41 Dumitrescu AM, Refetoff S: The syndromes of reduced sensitivity to thyroid hormone. *Biochim Biophys Acta* 2012, E-pub ahead of print.
- 42 Bochukova E, Schoenmakers N, Agostini M, Schoenmakers E, Rajanayagam O, Keogh JM, Henning E, Reinemund J, Gevers E, Sarri M, Downes K, Offiah A, Albanese A, Halsall D, Schwabe JW, Bain M, Lindley K, Muntoni F, Khadem FV, Dattani M, Farooqi IS, Gurnell M, Chatterjee K: A mutation in the thyroid hormone receptor α gene. *N Engl J Med* 2012;366:243–249.
- 43 Van Mullem A, van Heerebeek R, Chrysis D, Visser E, Medici M, Andrikoula M, Tsatsoulis A, Peeters R, Visser TJ: Clinical phenotype and mutant TR α 1. *N Engl J Med* 2012;366:1451–1453.
- 44 Dumitrescu AM, Liao XH, Abdullah MS, Lado-Abeal J, Majed FA, Moeller LC, Boran G, Schomburg L, Weiss RE, Refetoff S: Mutations in SECISBP2 result in abnormal thyroid hormone metabolism. *Nat Genet* 2005;37:1247–1252.
- 45 Schoenmakers E, Agostini M, Mitchell C, Schoenmakers N, Papp L, Rajanayagam O, Padidela R, Ceron-Gutierrez L, Doffinger R, Prevosto C, Luan J, Montano S, Lu J, Castanet M, Clemons N, Groeneveld M, Castets P, Karbaschi M, Aitken S, Dixon A, Williams J, Campi I, Blount M, Burton H, Muntoni F, O'Donovan D, Dean A, Warren A, Brierley C, Baguley D, Guicheney P, Fitzgerald R, Coles A, Gaston H, Todd P, Holmgren A, Khanna KK, Cooke M, Semple R, Halsall D, Wareham N, Schwabe J, Grasso L, Beck-Peccoz P, Ogunko A, Dattani M, Gurnell M, Chatterjee K: Mutations in the selenocysteine insertion sequence-binding protein 2 gene lead to a multisystem selenoprotein deficiency disorder in humans. *J Clin Invest* 2010;120:4220–4235.
- 46 Hamajima T, Mushimoto Y, Kobayashi H, Saito Y, Onigata K: Novel compound heterozygous mutations in the *sbp2* gene: Characteristic clinical manifestations and the implications of GH and triiodothyronine in longitudinal bone growth and maturation. *Eur J Endocrinol* 2012;166:757–764.

- 47 Bassett JH, Nordstrom K, Boyde A, Howell PG, Kelly S, Vennstrom B, Williams GR: Thyroid status during skeletal development determines adult bone structure and mineralization. *Mol Endocrinol* 2007;21:1893–1904.
- 48 Bassett JH, O'Shea PJ, Sriskantharajah S, Rabier B, Boyde A, Howell PG, Weiss RE, Roux JP, Malaval L, Clement-Lacroix P, Samarut J, Chassande O, Williams GR: Thyroid hormone excess rather than thyrotropin deficiency induces osteoporosis in hyperthyroidism. *Mol Endocrinol* 2007;21:1095–1107.
- 49 Bassett JH, Williams GR: The skeletal phenotypes of TR α and TR β mutant mice. *J Mol Endocrinol* 2009;42:269–282.
- 50 Gauthier K, Plateroti M, Harvey CB, Williams GR, Weiss RE, Refetoff S, Willott JF, Sundin V, Roux JP, Malaval L, Hara M, Samarut J, Chassande O: Genetic analysis reveals different functions for the products of the thyroid hormone receptor α locus. *Mol Cell Biol* 2001;21:4748–4760.
- 51 O'Shea PJ, Bassett JH, Sriskantharajah S, Ying H, Cheng SY, Williams GR: Contrasting skeletal phenotypes in mice with an identical mutation targeted to thyroid hormone receptor α 1 or β . *Mol Endocrinol* 2005;19:3045–3059.
- 52 O'Shea PJ, Bassett JH, Cheng SY, Williams GR: Characterization of skeletal phenotypes of TR α 1 and TR β mutant mice: implications for tissue thyroid status and T3 target gene expression. *Nucl Recept Signal* 2006;4:e011.
- 53 Boyle WJ, Simonet WS, Lacey DL: Osteoclast differentiation and activation. *Nature* 2003;423:337–342.
- 54 Harada S, Rodan GA: Control of osteoblast function and regulation of bone mass. *Nature* 2003;423:349–355.
- 55 Varga F, Rumpler M, Zoehrer R, Turecek C, Spitzer S, Thaler R, Paschalis EP, Klaushofer K: T3 affects expression of collagen I and collagen cross-linking in bone cell cultures. *Biochem Biophys Res Commun* 2010;402:180–185.
- 56 Gouveia CH, Schultz JJ, Bianco AC, Brent GA: Thyroid hormone stimulation of osteocalcin gene expression in ROS 17/2.8 cells is mediated by transcriptional and post-transcriptional mechanisms. *J Endocrinol* 2001;170:667–675.
- 57 Varga F, Rumpler M, Luegmayer E, Fratzel-Zelman N, Glantschnig H, Klaushofer K: Triiodothyronine, a regulator of osteoblastic differentiation: depression of histone H4, attenuation of c-fos/c-jun, and induction of osteocalcin expression. *Calcif Tiss Int* 1997;61:404–411.
- 58 Xing W, Govoni KE, Donahue LR, Kesavan C, Wergedal J, Long C, Bassett JH, Gogakos A, Wojcicka A, Williams GR, Mohan S: Genetic evidence that thyroid hormone is indispensable for prepubertal insulin-like growth factor-I expression and bone acquisition in mice. *J Bone Miner Res* 2012;27:1067–1079.
- 59 Milne M, Quail JM, Rosen CJ, Baran DT: Insulin-like growth factor binding proteins in femoral and vertebral bone marrow stromal cells: expression and regulation by thyroid hormone and dexamethasone. *J Cell Biochem* 2001;81:229–240.
- 60 Stevens DA, Harvey CB, Scott AJ, O'Shea PJ, Barnard JC, Williams AJ, Brady G, Samarut J, Chassande O, Williams GR: Thyroid hormone activates fibroblast growth factor receptor-1 in bone. *Mol Endocrinol* 2003;17:1751–1766.
- 61 O'Shea PJ, Kim DW, Logan JG, Davis S, Walker RL, Meltzer PS, Cheng SY, Williams GR: Advanced bone formation in mice with a dominant-negative mutation in the thyroid hormone receptor β gene due to activation of Wnt/ β -catenin protein signaling. *J Biol Chem* 2012;287:17812–17822.
- 62 Varga F, Spitzer S, Klaushofer K: Triiodothyronine (T₃) and 1,25-dihydroxyvitamin D₃ (1,25-D₃) inversely regulate OPG gene expression in dependence of the osteoblastic phenotype. *Calcif Tiss Int* 2004;74:382–387.
- 63 Khosla S: Minireview: The OPG/RANKL/RANK system. *Endocrinology* 2001;142:5050–5055.
- 64 Saraiva PP, Teixeira SS, Padovani CR, Nogueira CR: Triiodothyronine (T₃) does not induce RANKL expression in rat ROS 17/2.8 cells. *Arq Bras Endocrinol Metabol* 2008;52:109–113.
- 65 Kanatani M, Sugimoto T, Sowa H, Kobayashi T, Kanzawa M, Chihara K: Thyroid hormone stimulates osteoclast differentiation by a mechanism independent of RANKL-RANK interaction. *J Cell Physiol* 2004;201:17–25.
- 66 Allain TJ, Chambers TJ, Flanagan AM, McGregor AM: Tri-iodothyronine stimulates rat osteoclastic bone resorption by an indirect effect. *J Endocrinol* 1992;133:327–331.
- 67 Klaushofer K, Hoffmann O, Gleispach H, Leis HJ, Czerwinka E, Koller K, Peterlik M: Bone-resorbing activity of thyroid hormones is related to prostaglandin production in cultured neonatal mouse calvaria. *J Bone Miner Res* 1989;4:305–312.
- 68 Miura M, Tanaka K, Komatsu Y, Suda M, Yasoda A, Sakuma Y, Ozasa A, Nakao K: A novel interaction between thyroid hormones and 1,25-OH₂D₃ in osteoclast formation. *Biochem Biophys Res Commun* 2002;291:987–994.
- 69 Mundy GR, Shapiro JL, Bandelin JG, Canalis EM, Raisz LG: Direct stimulation of bone resorption by thyroid hormones. *J Clin Invest* 1976;58:529–534.
- 70 Siddiqi A, Parsons MP, Lewis JL, Monson JP, Williams GR, Burrin JM: TR expression and function in human bone marrow stromal and osteoblast-like cells. *J Clin Endocrinol Metab* 2002;87:906–914.
- 71 Eriksen EF, Mosekilde L, Melsen F: Kinetics of trabecular bone resorption and formation in hypothyroidism: evidence for a positive balance per remodeling cycle. *Bone* 1986;7:101–108.
- 72 Mosekilde L, Eriksen EF, Charles P: Effects of thyroid hormones on bone and mineral metabolism. *Endocrinol Metab Clin North Am* 1990;19:35–63.
- 73 Vestergaard P, Mosekilde L: Fractures in patients with hyperthyroidism and hypothyroidism: a nationwide follow-up study in 16,249 patients. *Thyroid* 2002;12:411–419.
- 74 Vestergaard P, Rejnmark L, Mosekilde L: Influence of hyper- and hypothyroidism, and the effects of treatment with antithyroid drugs and levothyroxine on fracture risk. *Calcif Tiss Int* 2005;77:139–144.
- 75 Williams GR: The skeletal system in thyrotoxicosis; in Braverman LE, Cooper DS (eds): *Werner and Ingbar's The Thyroid: a Fundamental and Clinical Text*. Philadelphia, Lippincott Williams & Wilkins, 2012, pp 468–475.
- 76 Bauer DC, Ettinger B, Nevitt MC, Stone KL: Risk for fracture in women with low serum levels of thyroid-stimulating hormone. *Ann Intern Med* 2001;134:561–568.
- 77 Lee JS, Buzkova P, Fink HA, Vu J, Carbone L, Chen Z, Cauley J, Bauer DC, Cappola AR, Robbins J: Subclinical thyroid dysfunction and incident hip fracture in older adults. *Arch Intern Med* 2010;170:1876–1883.
- 78 Leese GP, Flynn RV: Levothyroxine dose and fractures in older adults. *BMJ* 2011;342:d2250.
- 79 Murphy E, Gluer CC, Reid DM, Felsenberg D, Roux C, Eastell R, Williams GR: Thyroid function within the upper normal range is associated with reduced bone mineral density and an increased risk of nonvertebral fractures in healthy euthyroid postmenopausal women. *J Clin Endocrinol Metab* 2010;95:3173–3181.
- 80 Roef G, Lapauw B, Goemaere S, Zmierzczak H, Fiers T, Kaufman JM, Taes Y: Thyroid hormone status within the physiological range affects bone mass and density in healthy men at the age of peak bone mass. *Eur J Endocrinol* 2011;164:1027–1034.
- 81 Van der Deure WM, Uitterlinden AG, Hofman A, Rivadeneira F, Pols HA, Peeters RP, Visser TJ: Effects of serum TSH and fT4 levels and the TSHR-asp727glu polymorphism on bone: The Rotterdam Study. *Clin Endocrinol* 2008;68:175–181.
- 82 Williams GR: The skeletal system in hypothyroidism; in Braverman LE, Cooper DS (eds): *Werner and Ingbar's The Thyroid: a Fundamental and Clinical Text*. Philadelphia, Lippincott Williams & Wilkins, 2012, pp 590–595.
- 83 Meulenbelt I, Bos SD, Chapman K, van der Breggen R, Houwing-Duistermaat JJ, Kremer D, Kloppenburg M, Carr A, Tsezou A, Gonzalez A, Loughlin J, Slagboom PE: Meta-analyses of genes modulating intracellular T₃ bio-availability reveal a possible role for the *Dio3* gene in osteoarthritis susceptibility. *Ann Rheum Dis* 2010;70:164–167.

- 84 Meulenbelt I, Min JL, Bos S, Riyazi N, Houwing-Duistermaat JJ, van der Wijk HJ, Kroon HM, Nakajima M, Ikegawa S, Uitterlinden AG, van Meurs JB, van der Deure WM, Visser TJ, Seymour AB, Lakenberg N, van der Breggen R, Kremer D, van Duijn CM, Kloppenburg M, Loughlin J, Slagboom PE: Identification of Dio2 as a new susceptibility locus for symptomatic osteoarthritis. *Hum Mol Genet* 2008;17:1867–1875.
- 85 Kerkhof HJ, Lories RJ, Meulenbelt I, Jonsdottir I, Valdes AM, Arp P, Ingvarsson T, Jhamai M, Jonsson H, Stolk L, Thorleifsson G, Zhai G, Zhang F, Zhu Y, van der Breggen R, Carr A, Doherty M, Doherty S, Felson DT, Gonzalez A, Halldorsson BV, Hart DJ, Hauksson VB, Hofman A, Ioannidis JP, Kloppenburg M, Lane NE, Loughlin J, Luyten FP, Nevitt MC, Parimi N, Pols HA, Rivadeneira F, Slagboom EP, Stykarsdottir U, Tsezou A, van de Putte T, Zmuda J, Spector TD, Stefansson K, Uitterlinden AG, van Meurs JB: A genome-wide association study identifies an osteoarthritis susceptibility locus on chromosome 7q22. *Arthritis Rheum* 2010;62:499–510.
- 86 Waarsing JH, Kloppenburg M, Slagboom PE, Kroon HM, Houwing-Duistermaat JJ, Weinans H, Meulenbelt I: Osteoarthritis susceptibility genes influence the association between hip morphology and osteoarthritis. *Arthritis Rheum* 2011;63:1349–1354.
- 87 Bos SD, Bovee JV, Duijnisveld BJ, Raine EV, van Dalen WJ, Ramos YF, van der Breggen R, Nelissen RG, Slagboom PE, Loughlin J, Meulenbelt I: Increased type II deiodinase protein in OA-affected cartilage and allelic imbalance of OA risk polymorphism rs225014 at Dio2 in human OA joint tissues. *Ann Rheum Dis* 2012;71:1254–1258.
- 88 Ladenson PW, Kristensen JD, Ridgway EC, Olsson AG, Carlsson B, Klein I, Baxter JD, Angelin B: Use of the thyroid hormone analogue eprotirome in statin-treated dyslipidemia. *N Engl J Med* 2010;362:906–916.