Malaria Diagnostics and Surveillance in the Post-Genomic Era

N.W. Lucchi^a J. Oberstaller^{b, c} J.C. Kissinger^{b–d} V. Udhayakumar^a

^aMalaria Branch, Division of Parasitic Diseases and Malaria, Center for Global Health, Centers for Disease Control and Prevention, Atlanta, Ga., ^bDepartment of Genetics, ^cCenter for Tropical and Emerging Global Diseases, and

^dInstitute of Bioinformatics, University of Georgia, Athens, Ga., USA

Key Words

Bioinformatics · Diagnosis · Genome · Malaria · Molecular tools

Abstract

Genome sequences are available for 3 human-infecting malaria parasites, *Plasmodium falciparum*, *P. vivax* and *P. knowlesi*, and population genomics data are available for many endemic regions. This review summarizes how genomic data have been used to develop new, species-specific molecular targets for better malaria diagnosis. The combination of bioinformatics and genomics has been used to identify new sequence targets suitable for diagnostic applications and assess their viability within the context of global *Plasmodium* sequence variation. The selection criteria maximized the sensitivity and specificity of the novel targets. At least one target from each species was found to be suitable for molecular diagnosis of malaria with some advantages over existing molecular methods. The promise of using genome sequence data to develop sensitive, genus- or species-specific diagnostic methods for other pathogens of public health interest is strong. This undertaking together with what we envision as the future of malaria diagnosis in the ‘omic’ era is discussed.

Copyright © 2013 S. Karger AG, Basel

Introduction

Advances in sequencing technology have led to an explosion of sequence generation, especially for infectious agents. As a result, genome sequences for many eukaryotic and prokaryotic pathogens have been generated and are publicly available. For example, the NCBI Genome Project database (<http://www.ncbi.nlm.nih.gov/genome>) provides access to complete genomes of a myriad of human pathogens. The almost-exponential availability of pathogen genome sequences has been made possible by major technological advances, or ‘next-generation’ DNA sequencing techniques, which have dramatically reduced the cost and time required to generate sequence data [1].

The genome sequence for the most lethal human malaria parasite *Plasmodium falciparum* (~23 Mb) was first published in 2002 [2] based on the laboratory-adapted parasite isolate 3D7. However, thousands of global *Plasmodium* strains and clinical isolates have since been examined for single nucleotide polymorphisms (SNPs) or sequenced to generate a map of global variation and identify and track changes related to major phenotypes such as drug resistance [7, 8]. The nuclear genome sequences for the human-infecting *P. vivax* (~27 Mb) and zoonotic *P. knowlesi* (~24 Mb) are also known [3, 4], and all available genome sequences can be easily accessed at (<http://PlasmoDB.org>) [5] (along with other genome-wide datasets, such as SNPs, expression data, proteomics, etc). Plas-

moDB provides tools to make comparisons between emerging high-throughput data sets for these organisms.

Now, more than ever, the role of genomics in understanding *Plasmodium* parasite biology is evident [6]. It is hoped that the availability of *Plasmodium* genome sequences will revolutionize the discovery of new drug targets, development of new vaccine targets and drug-resistant markers, and lead to improvement in other tools for fighting and controlling malaria such as diagnostics. Recently, genome-based association studies have contributed to identification of genomic regions associated with drug resistance [7, 8]. The significance of genomic data for understanding parasite biology, pathogenesis, drug resistance, and population structure has been extensively reviewed recently [9]. The scope of the current review is to highlight how genomics is contributing to the immediate public health needs of malaria diagnosis.

The molecular biologist's toolbox has improved tremendously since the advent of the polymerase chain reaction (PCR) in the mid-1980s. PCR-based assays are fairly standard in most laboratories and can be used for disease detection, forensics and drug-resistance surveillance testing. While several other disease-detection methods exist, molecular tools are still the most specific and sensitive tools. Molecular diagnostic techniques rely on in-depth knowledge of the target sequence and its global variability. Therefore, molecular diagnostic tools benefit tremendously from the increased availability of pathogen genome sequences. For example, genome sequences can be mined to discover conserved sequence regions suitable for the design of specific primers that do not cross-react with either the host or other species of the same genus. Likewise, highly variable regions can be avoided when designing detection assays, as these would lead to false negative results. Sequences which exist in high copy numbers can be used to design a more sensitive assay compared to assays utilizing single- or low-copy number sequence targets.

Why New Tools for Malaria Diagnosis?

Malaria continues to be a major global public health challenge. There were approximately 225 million cases of malaria in 2009 with an estimated 718,000 deaths reported [10]. Five different *Plasmodium* species with different clinical implications infect humans in different combinations around the world. There is a renewed, concerted, international effort to fight and eliminate malaria through funding from the Global Fund, the U.S. Presidents Malaria Initiative, the Bill and Melinda Gates Foundation and

other private and public sources. Thus, new diagnostic assays and tools that are robust enough to accurately detect the species of infecting parasite(s), for case management, detection of transmission foci of malaria reservoirs (sub-microscopic infections) and for monitoring the success of malaria control and elimination programs are needed.

Existing tools for malaria diagnosis include microscopy, parasite antigen/enzyme detection kits (commonly referred to as rapid diagnostic tests (RDTs)) and molecular tools, which are mostly restricted to reference laboratories (reviewed in [11]). Microscopy remains the gold standard when a suitable infrastructure is available. This is the cheapest method and can differentiate species of malaria parasites and provide quantitative data on the level of parasitemia. One limitation of microscopy is that it can fail to identify mixed infections and/or low levels of parasitemia. Microscopic detection limits range from 50–100 parasites/ μ L, depending on the level of expertise of the microscopist and other factors (reviewed in [11]). As malaria often occurs in communities where even microscopic diagnosis is not easily available, RDTs are an alternative diagnostic tool. Current RDTs capture products such as the histidine-rich protein –2 (Pf HRP-2), which is *P. falciparum*-specific, or the aldolase and lactate dehydrogenase enzyme (LDH) which are genus-specific and are therefore capable of detecting all *Plasmodia*, but are not able to discriminate species. However, highly divergent species such as *P. ovale* can produce negative RDT results due to sequence variation [12]. In addition, the lack of the *hrp-2* gene in some *P. falciparum* parasites in parts of South America and Africa [13, 14] has raised some concerns about HRP-2-based RDT use due to the potential for false negative results. Proteomics data generated from multiple strains can be used to develop new-generation RDTs utilizing multiple protein targets to overcome existing challenges.

Nucleic acid-based techniques such as PCR have revolutionized pathogen detection and identification, by offering high sensitivity and specificity. Molecular diagnostic methods for malaria diagnosis have at least 2 advantages compared to the other methods: they can accurately define the species of malaria parasite(s), and they can detect parasite densities that are well below microscopy limits. Previously, the field of malaria diagnosis was wrapped in debates as to what tool is clinically relevant for diagnosis and treatment of malaria, especially in highly endemic regions; this stemmed from the fact that molecular tools are very sensitive in detecting submicroscopic infections which often do not translate to clinical manifestations (subclinical). Others argued that molecular tests may detect 'lingering' parasite DNA as opposed to active infec-

tions and may therefore not be appropriate assays to use. However, the malaria elimination era makes it paramount that even the 'last parasite' is found. In addition, recent studies have clearly demonstrated that submicroscopic gametocyte carriers (who can only be detected molecularly) are capable of transmitting malaria [15]. Molecular tools are thus appealing for use in both case management and programmatic operations, such as monitoring and evaluation of control programs. Molecular diagnostic tools have helped to identify zoonotic transmission of *P. knowlesi* in parts of Southeast Asia [16]. These infections were previously diagnosed as *P. malariae* or *P. falciparum* by microscopy. Molecular tools are also helping to identify new species of malaria parasites including *P. falciparum*-like parasite species in non-human primates [17].

Evolution of Molecular Tools for Malaria Diagnostics

Several molecular diagnostic tools for malaria are available, the majority of which are PCR-based assays. Recent technologies have led to the development of other amplification tools, such as loop-mediated isothermal amplification assays, that are simpler and more field-adaptable [18, 19]. Molecular diagnosis of malaria parasites began with the use of the 18S ribosomal RNA (18S rRNA) gene as the target about 20 years ago [20], and this method is widely used in many reference laboratories with various modifications. This target was a logical choice in the pre-genomics era. Its regions of conserved sequence allowed cloning from multiple *Plasmodium* species facilitating the subsequent design of species-specific primers. Also, all eukaryotic organisms that had been examined to that date contained multiple, often hundreds of identical copies of 18S rRNA [21], so it seemed likely that this target would lead to a very sensitive assay. *Plasmodium* genome sequences have subsequently revealed that the 18S rRNA target is present in 4–8 divergent copies, depending upon the species [2, 3]. PCR's sensitivity is greatly influenced by the starting target molecule copy number; a low target copy number within the parasite limits the detection capabilities of these assays, especially when the parasitemia is low. Previous multiplex assays for simultaneous detection of malaria parasite species showed decreased sensitivity, particularly in detecting the minor species [22]. The 18S rRNA gene target presents challenges for effective multiplex platforms which would cut back on costs, test time and reduce contamination possibilities resulting from many primers competing for the same target.

Fig. 1. Schematic of diagnostic target screening and development pipeline. Genomic sequences for *P. falciparum* (Pf), *P. vivax* (Pv), and *P. knowlesi* (Pk) were downloaded from PlasmoDB. Data were mined for repeats using the RepeatScout algorithm to construct consensus repeat sequences (CRSs) for each identified repeat family. We identified 418 CRSs for Pf, 428 CRSs for Pv and 423 CRSs for Pk. CRSs were then screened in parallel for tandem repeats, similarity to human sequences and vector sequences. Pf, Pv and Pk CRSs passing these screens and proceeding to the next steps are indicated in parentheses. Any CRSs failing these screens were removed from further consideration. CRSs that were not species-specific or less than 300 bp long were eliminated. Family copy numbers for the remaining candidates were determined via comparison of the CRSs against the appropriate genome sequences. Candidate repeat families containing six or more copies and separated by at least 100 bp were considered for further testing.

Use of Genomics and Bioinformatics to Improve Malaria Diagnostic Assays

The new era of genomics and sequence availability can be harnessed to improve molecular diagnostic tools for malaria. Recently, we developed a bioinformatics approach to identify new sequence targets that are suitable for diagnostic applications (fig. 1) in *P. falciparum*, *P. vivax* and *P. knowlesi*. Ideal targets will be species-specific

Table 1. Summary of the novel candidates identified

Candidate	No. of copies	Location in genome	Molecular test	Limits of detection P/μl	Sensitivity ^a %	Specificity ^a %
Pfr364	41	SB2-subtelomeric; not protein-encoding	PCR assay	10–0.1	100	100
Pvr47	14	Unassigned subtelomeric; not protein-encoding	PCR assay	10–0.1	98.9	100
Pvr64*	6	Mostly unassigned contigs, one chromosome interior; protein-encoding	LAMP assay	200–100	94.6	100
Pkr140	7	Chromosome ends and interiors; not protein-encoding	PCR assay	1	ND	ND

^a Compared to a nested PCR assay [20] as a gold standard. * [33].
P/μl = Parasites/μl; ND = not done.

ic, present in many, well-conserved copies (ensuring high sensitivity) and amenable to PCR or other nucleic acid amplification techniques. Our semi-automated bioinformatics pipeline employed a number of screens to ensure candidate targets met these criteria [23, 24].

Our candidate target selection process began by accessing genome sequence data for *P. falciparum* (strain 3D7), *P. vivax* (strain Sal-1) and *P. knowlesi* (strain H) from PlasmoDB. Each genome was mined for repetitive sequence content, and a consensus repeat sequence (CRS) for each repeat family was generated. We screened and eliminated all CRSs with significant similarity to human sequences, significant similarity to artificial sequences that may have been introduced during genome sequencing and those containing internal tandem repeats that could potentially interfere with PCR amplification and similarity to sequences from other *Plasmodium* spp. Only species-specific CRSs that did not exhibit high levels of polymorphism in existing SNP data sets were considered further. We also eliminated candidates <300 bp to allow for primer design and evaluation of target conservation. Repeat families with at least six copies were considered for further testing, yielding a total of 21 *P. falciparum*, 68 *P. vivax* and 19 *P. knowlesi* candidates.

Six *P. falciparum*, seven *P. vivax* and four *P. knowlesi* putative targets were selected for further validation. Over 64 primer pairs were designed to these targets and empirically tested in conventional PCR amplification assays and multiplex assays (for *P. falciparum* and *P. vivax*). At least one putative target from each species (summarized in table 1) was found to significantly improve existing diagnostic capabilities: candidates Pfr364 (*P. falciparum*), Pvr47 [23] and Pvr64 (*P. vivax*; designed for loop mediated isothermal amplification assays, [33]), and Pkr140 (*P. knowlesi*) [24].

Validation of PCR Primers from Novel Targets

All primers designed to the four novel targets were shown to correctly amplify the gene of interest with high specificity and sensitivity (table 1). In addition, the limits of detection of these assays were improved. Of interest is the novel primer set for detection of *P. knowlesi* which exhibits stage-dependent morphological similarities to *P. malariae* and *P. falciparum* [16]. The commonly used molecular assay for the detection of *P. knowlesi* infection was recently noted to cross-react with *P. vivax*, leading to potential false positive results for a small proportion of human clinical *P. vivax* samples [25] and other simian *Plasmodium* species [24]. Primers designed to the Pkr140 target consistently identified *P. knowlesi* without cross-reacting with other human malaria parasites and five other primate *Plasmodium* parasites (*P. simiovale*, *P. cynomolgi*, *P. inui*, *P. coatneyi*, and *P. hylobati*).

Genomic data mining allowed us to develop sensitive and highly specific PCR tests for three *Plasmodium* species. Similar approaches can be used for the other human-infecting species as data are available. The newly developed tests highlighted here need further validation in the field, and their performance can be improved by modifying reaction parameters as needed.

Limitations of the Described Approach

What we have described above is only an example of how genome data can be exploited for developing new diagnostic methods. The genome search methodology we have developed can be further modified depending upon the specific needs of the tools being considered for development. The use of a single strain to select a candidate sequence as described here is a limitation that can be improved. To ascertain that the designed primers are capable of interspecies detection, one could test the novel primers against different strains from across the world, as

was previously done by Demas et al. [23]. However, as additional SNP and population variation data become freely accessible via PlasmoDB and other resources such as MalariaGen (Genomic Epidemiology Network), these databases can be used to determine sequence variation in much greater detail and aid the design of novel primers capable of detecting field isolates.

Future Directions

New-Generation Molecular Tools

The current 'omic' era greatly complements the growing number of new-generation tests and platforms for molecular diagnostic tests observed today. Recently, a microfluidic technology-based TaqMan low-density array (Life Technologies, Carlsbad, Calif., USA) card for real-time PCR detection of 21 different respiratory pathogens was described [26]. Such array formats can be adapted for detection of multiple malaria parasite species and can be integrated with detection of other pathogens of relevance. Further development of less-expensive, simple, field-usable DNA microarrays for malaria detection will be useful for large-scale surveillance programs. Several isothermal amplification assays have been described, e.g. the SAM-BA HIV-1 test for the rapid visual detection of HIV-1 by a dipstick [27], loop-mediated isothermal amplification [19] and its modification for real-time detection using portable devices [28]. Mens et al. [29] recently developed a direct-blood PCR assay visualized by nucleic acid lateral flow immune-assay capable of detecting *Plasmodium*; this assay attempts to circumvent the DNA isolation step, a major hurdle towards the simplification of molecular tests. The current direction is towards simple, robust and cheap tests that can detect multiple disease agents and hopefully enable their use in point-of-care settings with limited resources.

Use of Proteomics to Improve Malaria Antigen-Based RDTs

The next-generation RDTs will require a better selection of *Plasmodium* antigens that are highly abundant, highly conserved across geographic regions and species-specific. The use of genomics and bioinformatics is not limited to the development of molecular assays; these tools together with proteomics can be used to select for *Plasmodium* antigens that meet these standards. In an effort to improve the current malaria RDTs, Kattenberg et al. [30] recently tested monoclonal antibodies against three *Plasmodium* antigens: glutamate-rich protein, di-

hydrofolate reductase-thymidylate synthase and heme detoxification protein for their potential use in RDTs. This study highlights the feasibility of developing next-generation RDTs with multiple protein targets that could be identified and validated using proteomics data.

Molecular Barcoding for Differentiating Parasite Populations and Their Use in Outbreak and Elimination Programs

In outbreak investigations and during post-elimination monitoring of malaria, it will be important to identify the geographic source of malaria parasites. Recently, a molecular barcode study using 24 SNPs from the *P. falciparum* genome was developed and tested for its utility in differentiating parasite populations [31]. Further validation and improvement of such molecular barcodes and curation of publicly shared databases housing these data will be invaluable for public health investigations and tracking of drug-resistant parasite populations. Using multi-locus genotyping of microsatellite markers, it has been shown that recent Peruvian *P. falciparum* populations have evolved from five clonal lineages in the post-malaria eradication era [32], demonstrating the utility of genomic data in facilitating the public health response to control and eliminate malaria.

Summary

Availability of genome data for human malaria parasites has led to significant advances in understanding the biology of malaria parasites and determination of genetic targets associated with evolving resistance to artemisinin as well as other applications [9]. We have highlighted the value of generating population level genomic data, applying bioinformatics methods and mining these data to identify novel targets for malaria diagnosis and surveillance. We view these initial forays with bioinformatics as only the tip of the iceberg when considering the possible uses of genomic data to address a large variety of public health demands. Novel diagnostic tests with better sensitivity and specificity than current tests can be developed with advancing genomics knowledge of malaria parasites. Such tools will be highly relevant for malaria control programs and efficient public health response.

References

- Chen PE, Willner KM, Butani A, Dorsey S, George M, Stewart A, Lentz SM, Cook CE, Akmal A, Price LB, Keim PS, Mieczyn A, Brahmbhatt TN, Bishop-Lilly KA, Zwick ME, Read TD, Sozhamannan S: Rapid identification of genetic modifications in *Bacillus anthracis* using whole genome draft sequences generated by 454 pyrosequencing. *PLoS One* 2010;5: e12397.
- Gardner MJ, Hall N, Fung E, White O, Berri-man M, Hyman RW, Carlton JM, Pain A, Nelson KE, Bowman S, Paulsen IT, James K, Eisen JA, Rutherford K, Salzberg SL, Craig A, Kyes S, Chan MS, Nene V, Shallom SJ, Suh B, Peterson J, Angiuoli S, Pertea M, Allen J, Selengut J, Haft D, Mather MW, Vaidya AB, Martin DM, Fairlamb AH, Fraunholz MJ, Roos DS, Ralph SA, McFadden GI, Cummings LM, Subramanian GM, Mungall C, Venter JC, Carucci DJ, Hoffman SL, Newbold C, Davis RW, Fraser CM, Barrell B: Genome sequence of the human malaria parasite *Plasmodium falciparum*. *Nature* 2002;419:498–511.
- Carlton JM, Adams JH, Silva JC, Bidwell SL, Lorenzi H, Caler E, Crabtree J, Angiuoli SV, Merino EF, Amedeo P, Cheng Q, Coulson RM, Crabb BS, Del Portillo HA, Essien K, Feldblyum TV, Fernandez-Becerra C, Gilson PR, Gueye AH, Guo X, Kang'a S, Kooij TW, Korsinczyk M, Meyer EV, Nene V, Paulsen I, White O, Ralph SA, Ren Q, Sargeant TJ, Salzberg SL, Stoeckert CJ, Sullivan SA, Yamamoto MM, Hoffman SL, Wortman JR, Gardner MJ, Galinski MR, Barnwell JW, Fraser-Liggett CM: Comparative genomics of the neglected human malaria parasite *Plasmodium vivax*. *Nature* 2008;455:757–763.
- Pain A, Böhme U, Berry AE, Mungall K, Finn RM, Jackson AP, Mourier T, Mistry J, Pasini EM, Aslett MA, Balasubramanian S, Borgwardt K, Brooks K, Carret C, Carver TJ, Che-revach I, Chillingworth T, Clark TG, Galinski MR, Hall N, Harper D, Harris D, Hauser H, Ivens A, Janssen CS, Keane T, Larke N, Lapp S, Marti M, Moule S, Meyer IM, Ormond D, Peters N, Sanders M, Sanders S, Sargeant TJ, Simmonds M, Smith F, Squares R, Thurston S, Tivey AR, Walker D, White B, Zuidewijk E, Churcher C, Quail MA, Cowman AF, Turner CM, Rajandream MA, Kocken CH, Thomas AW, Newbold CI, Barrell BG, Berri-man M: The genome of the simian and human malaria parasite *Plasmodium knowlesi*. *Nature* 2008;455:799–803.
- Aurrecochea C, Brestelli J, Brunk BP, Dommer J, Fischer S, Gajria B, Gao X, Gingle A, Grant G, Harb OS, Heiges M, Innamorato F, Iodice J, Kissinger JC, Kraemer E, Li W, Miller JA, Nayak V, Pennington C, Pinney DF, Roos DS, Ross C, Stoeckert CJ Jr, Treatman C, Wang H: PlasmoDB: a functional genomic database for malaria parasites. *Nucleic Acids Res* 2009;37:D539–D543.
- Volkman SK, Ndiaye D, Diakite M, Koita OA, Nwakanma D, Daniels RF, Park DJ, Neafsey DE, Muskavitch MA, Krogstad DJ, Sabeti PC, Hartl DL, Wirth DF: Application of genomics to field investigations of malaria by the international centers of excellence for malaria research. *Acta Trop* 2012;121:324–332.
- Cheeseman IH, Miller BA, Nair S, Nkhoma S, Tan A, Tan JC, Al Saai S, Phyto AP, Moo CL, Lwin KM, McGready R, Ashley E, Imwong M, Stepniewska K, Yi P, Dondorp AM, Mayxay M, Newton PN, White NJ, Nosten F, Ferdig MT, Anderson TJ: A major genome region underlying artemisinin resistance in malaria. *Science* 2012;336:79–82.
- Park DJ, Lukens AK, Neafsey DE, Schaffner SF, Chang HH, Valim C, Ribacke U, Van Tyne D, Galinsky K, Galligan M, Becker JS, Ndiaye D, Mboup S, Wiegand RC, Hartl DL, Sabeti PC, Wirth DF, Volkman SK: Sequence-based association and selection scans identify drug resistance loci in the *Plasmodium falciparum* malaria parasite. *Proc Natl Acad Sci USA* 2012;109:13052–13057.
- Volkman SK, Neafsey DE, Schaffner SF, Park DJ, Wirth DF: Harnessing genomics and genome biology to understand malaria biology. *Nat Rev Genet* 2012;13:315–328.
- World Health Organization: World Malaria Report 2010. http://www.who.int/malaria/world_malaria_report_2010/en/index.html.
- Erdman LK, Kain KC: Molecular diagnostic and surveillance tools for global malaria control. *Travel Med Infect Dis* 2008;6:82–99.
- Tordrup D, Virenfelt J, Andersen FF, Petersen E: Variant *Plasmodium ovale* isolated from a patient infected in Ghana. *Malar J* 2011;10: 15.
- Koita OA, Doumbo OK, Ouattara A, Tall LK, Konaré A, Diakité M, Diallo M, Sagara I, Masiinde GL, Doumbo SN, Dolo A, Tounkara A, Traoré I, Krogstad DJ: False-negative rapid diagnostic tests for malaria and deletion of the histidine-rich repeat region of the *hrp2* gene. *Am J Trop Med Hyg* 2012;86:194–198.
- Gamboa D, Ho MF, Bendezu J, Torres K, Chiodini PL, Barnwell JW, Incardona S, Perkins M, Bell D, McCarthy J, Cheng Q: A large proportion of *P. falciparum* isolates in the Amazon region of Peru lack *pfrp2* and *pfrp3*: implications for malaria rapid diagnostic tests. *PLoS One* 2010;5:e8091.
- Bousema T, Drakeley C: Epidemiology and infectivity of *Plasmodium falciparum* and *Plasmodium vivax* gametocytes in relation to malaria control and elimination. *Clin Microbiol Rev* 2011;24:377–410.
- Singh B, Kim Sung L, Matusop A, Radhakrishnan A, Shamsul SS, Cox-Singh J, Thomas A, Conway DJ: A large focus of naturally acquired *Plasmodium knowlesi* infections in human beings. *Lancet* 2004;363: 1017–1024.
- Prugnolle F, Durand P, Neel C, Ollomo B, Ayala FJ, Arnathau C, Etienne L, Mpoudi-Ngole E, Nkoghe D, Leroy E, Delaporte E, Peeters M, Renaud F: African great apes are natural hosts of multiple related malaria species, including *Plasmodium falciparum*. *Proc Natl Acad Sci USA* 2010;107:1458–1463.
- Lucchi NW, Demas A, Narayanan J, Sumari D, Kabanyanyi A, Kachur SP, Barnwell JW, Udhayakumar V: Real-time fluorescence loop mediated isothermal amplification for the diagnosis of malaria. *PLoS One* 2010;5:e13733.
- Poon LL, Wong BW, Ma EH, Chan KH, Chow LM, Abeyewickreme W, Tangpukdee N, Yuen KY, Guan Y, Looareesuwan S, Peiris JS: Sensitive and inexpensive molecular test for *falciparum* malaria: detecting *Plasmodium falciparum* DNA directly from heat-treated blood by loop-mediated isothermal amplification. *Clin Chem* 2006;52:303–306.
- Snounou G, Viriyakosol S, Zhu XP, Jarra W, Pinheiro L, do Rosario VE, Thaithong S, Brown KN: High sensitivity of detection of human malaria parasites by the use of nested polymerase chain reaction. *Mol Biochem Parasitol* 1993;61:315–320.
- Mercereau-Puijalon O, Barale JC, Bischoff E: Three multigene families in *Plasmodium* parasites: facts and questions. *Int J Parasitol* 2002; 32:1323–1344.
- Mixson-Hayden T, Lucchi NW, Udhayakumar V: Evaluation of three PCR-based diagnostic assays for detecting mixed *Plasmodium* infection. *BMC Res Notes* 2010;3:88.
- Demas A, Oberstaller J, DeBarry J, Lucchi NW, Srinivasamoorthy G, Sumari D, Kabanyanyi AM, Villegas L, Escalante AA, Kachur SP, Barnwell JW, Peterson DS, Udhayakumar V, Kissinger JC: Applied genomics: data mining reveals species-specific malaria diagnostic targets more sensitive than 18S rRNA. *J Clin Microbiol* 2011;49:2411–2418.
- Lucchi NW, Poorak M, Oberstaller J, DeBarry J, Srinivasamoorthy G, Goldman I, Xayavong M, da Silva AJ, Peterson DS, Barnwell JW, Kissinger J, Udhayakumar V: A new single-step PCR assay for the detection of the zoonotic malaria parasite *Plasmodium knowlesi*. *PLoS One* 2012;7:e31848.
- Imwong M, Tanomsing N, Pukrittayakamee S, Day NP, White NJ, Snounou G: Spurious amplification of a *Plasmodium vivax* small-subunit RNA gene by use of primers currently used to detect *P. knowlesi*. *J Clin Microbiol* 2009;47:4173–4175.
- Kodani M, Yang G, Conklin LM, Travis TC, Whitney CG, Anderson LJ, Schrag SJ, Taylor TH Jr, Beall BW, Breiman RF, Feikin DR, Njenga MK, Mayer LW, Oberste MS, Tondella ML, Winchell JM, Lindstrom SL, Erdman DD, Fields BS: Application of TaqMan low-density arrays for simultaneous detection of multiple respiratory pathogens. *J Clin Microbiol* 2011;49:2175–2182.

- 27 Lee HH, Dineva MA, Chua YL, Ritchie AV, Ushiro-Lumb I, Wisniewski CA: Simple amplification-based assay: a nucleic acid-based point-of-care platform for HIV-1 testing. *J Infect Dis* 2010;201(suppl 1):S65–S72.
- 28 Lucchi NW, Demas A, Narayanan J, Sumari D, Kabanyanyi A, Kachur SP, Barnwell JW, Udhayakumar V: Real-time fluorescence loop mediated isothermal amplification for the diagnosis of malaria. *PLoS One* 2010;5:e13733.
- 29 Mens PF, de Bes HM, Sondo P, Laochan N, Keereecharoen L, van Amerongen A, Flint J, Sak JR, Proux S, Tinto H, Schallig HD: Direct blood PCR in combination with nucleic acid lateral flow immuno-assay for the detection of *Plasmodium* species in malaria endemic settings. *J Clin Microbiol* 2012;50:3520–3525.
- 30 Kattenberg JH, Versteeg I, Migchelsen SJ, González IJ, Perkins MD, Mens PF, Schallig HD: New developments in malaria diagnostics: monoclonal antibodies against *Plasmodium* dihydrofolate reductase-thymidylate synthase, heme detoxification protein and glutamate rich protein. *MAbs* 2012;4:120–126.
- 31 Daniels R, Volkman SK, Milner DA, Mahesh N, Neafsey DE, Park DJ, Rosen D, Angelino E, Sabeti PC, Wirth DF, Wiegand RC: A general SNP-based molecular barcode for *Plasmodium falciparum* identification and tracking. *Malar J* 2008;7:223.
- 32 Griffing SM, Mixson-Hayden T, Sridaran S, Alam MT, McCollum AM, Cabezas C, Marquino Quezada W, Barnwell JW, De Oliveira AM, Lucas C, Arrospide N, Escalante AA, Bacon DJ, Udhayakumar V: South American *Plasmodium falciparum* after the malaria eradication era: clonal population expansion and survival of the fittest hybrids. *PLoS One* 2011;6:e23486.
- 33 Patel JC, Oberstaller J, Xayavong M, Narayanan J, Debarry JD, et al.: Real-time loop-mediated isothermal amplification (Real-Amp) for the species-specific identification of *Plasmodium vivax*. *PLoS One* 2013;8: e54986. doi:10.1371/journal.pone.0054986.