Preface

Eosinophils are mysterious and unique cells with the characteristics of a double-edged sword, i.e. possessing detrimental effects for example in asthma and allergic diseases and beneficial effects such as the elimination of parasitic infections. Eosinophils had been considered end-stage or effector cells involved in such diseases. However, numerous lines of evidence have altered this perspective by showing that eosinophils are pleiotropic multifunctional leukocytes involved in the initiation and propagation of diverse inflammatory responses, as well as modulators of innate and adaptive immunity [1].

In 2000, many investigators stated that eosinophils may play only a limited role in asthma, based on Leckie's report [2] that revealed that humanized anti-interleukin 5 (IL-5) monoclonal antibody treatment was unsuccessful in treating human asthma. However, this report encourages many researchers to reconsider the role of eosinophils in asthma, including airway remodeling. Recent studies demonstrated that administration of anti-IL-5 (mepolizumab) to patients with hypereosinophilic syndrome (HES) and eosinophilic asthma is beneficial [3, 4]. Although these monoclonal antibodies are clinically efficacious, they only partially reduce the eosinophil-mediated effects in asthma. For example, eosinophils are reduced in number, but are not eliminated, and the deposition of granule major basic protein (MBP) is not altered. Long-term studies are required, in which either tissue eosinophils or the deposition of eosinophil-specific granule proteins is effectively and selectively eliminated, in order to establish the exact role of these cells in human allergic inflammation such as asthma.

Again, do eosinophils play only detrimental roles? We have shown that eosinophil infiltration and appreciable degranulation, evaluated using an anti-MBP antibody, consistently occur only in the gastrointestinal tract and not in any other organs [5]. This evidence may suggest that eosinophils play a protective role in the gut. Indeed, MBP has been shown to be cytotoxic against tumor cells. Other evidence suggests that the eosinophil secretory ribonucleases, eosinophil-derived neurotoxin (EDN) and eosinophil cationic protein (ECP), possess antiviral activity against respiratory syncytial virus (RSV) in vitro and the related natural rodent pathogen, pneumonia virus of mice, in vivo [6].

Thus, we must not readily assume that eosinophils have only detrimental effects and play a role in only a single disease. They may play beneficial roles in some situations and may also be involved in other diseases. Further studies are required to delineate the exact role of eosinophils in various diseases as well as in healthy individuals.

The initial workshop on Eosinophils in Allergy and Related Diseases took place in Tokyo in 1991. Since then, this workshop has reported up-to-date research on eosinophils and allergy in Japan. The 24th Annual Workshop was held in Tokyo on June 23, 2012, with 27 research presentations on eosinophils and allergy in the fields of basic and clinical sciences. Notably, Dr. Hirohito Kita from the Mayo Clinic gave a special lecture, which is included as a plenary review in this supplement. This workshop has provided a forum for interested investigators to discuss recent advances in the basic and clinical aspects

of the cell biology of eosinophils and other inflammatory cells and their roles in related diseases. Topics in this workshop include the mechanisms of airway remodeling, activation of eosinophils, intracellular signaling of basophils and mast cells, mediator releases in dendritic and bronchial epithelial cells and potential treatment for eosinophil-associated diseases. These proceedings include 21 peer-reviewed original papers, 1 peer-reviewed case report and 1 review paper. Two abstracts, part of which have been submitted or published elsewhere, are also included. The titles of the remaining 3 presentations are listed at the end of this issue. We hope these studies contribute to a better understanding of the pathophysiology and treatment of allergy and eosinophil-associated diseases.

Finally, we would like to express our sincere appreciation to Novartis Pharma K.K., specifically Mr. Mitsuaki Shozui (Tokyo), for supporting this workshop and to International Medical Publishers (Tokyo) for their excellent editorial assistance in publishing these proceedings.

Masahiko Kato, Gunma, Japan

References

- 1 Gleich GJ: Historical overview and perspective on the role of the eosinophil in health and disease; in Lee JJ, Rosenberg HF (eds): Eosinophils in Health and Disease, ed 1. London, Academic Press, Elsevier Inc., 2013, pp 1–10.
- 2 Leckie MJ, ten Brinke A, Khan J, et al: Effects of an interleukin-5 blocking monoclonal antibody on eosinophils, airway hyper-responsiveness, and the late asthmatic response. *Lancet* 2000;356:2144–2148.
- 3 Rothenberg ME, Klion AD, Roufosse FE, et al, Mepolizumab HES Study Group: Treatment of patients with the hypereosinophilic syndrome with mepolizumab. *N Engl J Med* 2008;358:1215–1228.
- 4 Pavord ID, Korn S, Howarth P, et al: Mepolizumab for severe eosinophilic asthma (DREAM): a multicentre, double-blind, placebo-controlled trial. *Lancet* 2012;380:651–659.
- 5 Kato M, Kephart GM, Talley NJ, et al: Eosinophil infiltration and degranulation in normal human tissue. *Anat Rec* 1998;252:418–425.
- 6 Rosenberg HF, Domachowski JB: Eosinophils, eosinophil ribonucleases, and their role in host defense against respiratory virus pathogens. *J Leukoc Biol* 2001;70:691–698.