Community-Acquired Methicillin-Resistant *Staphylococcus aureus*: The New Face of an Old Foe?

Edet E. Udo

Department of Microbiology, Health Sciences Centre, Faculty of Medicine, Kuwait University, Kuwait

Key Words

Antibiotic resistance · Methicillin-resistant *Staphylococcus aureus*, community-associated · Multilocus sequence typing

Abstract

The burden of infections caused by community-associated methicillin-resistant *Staphylococcus aureus* (CA-MRSA) is increasing among different patient populations globally. As CA-MRSA has become established in healthcare facilities, the range of infections caused by them has also increased. Molecular characterization of CA-MRSA isolates obtained from different centers has revealed significant diversity in their genetic backgrounds. Although many CA-MRSA strains are still susceptible to non- β -lactam antibiotics, multiresistance to non- β -lactam agents has emerged in some clones, posing substantial problems for empirical and directed therapy of infections caused by these strains. Some CA-MRSA clones have acquired the capacity to spread locally and internationally. CA-MRSA belonging to ST80-MRSA-IV and ST30-MRSA-IV appear to be the dominant clones in the countries of the Gulf Cooperation Council (GCC). The emergence of pandemic CA-MRSA clones not only limits therapeutic options but also presents significant challenges for

infection control. Continued monitoring of global epidemiology and emerging drug resistance data is critical for the effective management of these infections.

© 2013 S. Karger AG, Basel

Staphylococcus aureus: The Old Foe

Staphylococcus aureus is an opportunistic human pathogen, which causes a wide range of infections ranging from superficial skin and soft tissue infections to life threatening pneumonia, bloodstream, bone and joint, respiratory, and gastrointestinal and genital tract infections [1, 2].

In the preantibiotic era, infections caused by *S. aureus* were usually fatal [1]. However, the introduction of penicillin to treat infections caused by *S. aureus* greatly improved the prognosis of patients with severe staphylococcal infections [1–3]. Nevertheless, *S. aureus* strains resistant to penicillin soon appeared [3]. Penicillin resistance is due to the production by the bacteria of an enzyme, i.e. penicillinase, which inactivates the antibiotic, rendering it inactive against the bacteria. Consequently, new antibiotics such as streptomycin, tetracycline, erythromycin, and chloramphenicol were developed in the 1950s [4].

However, as new antibiotics were developed and put into clinical use, resistance to them appeared, leading to the emergence of strains that were resistant to multiple antibiotics [5, 6].

Emergence of Methicillin Resistance in *S. aureus*

Methicillin resistance was reported in *S. aureus* in 1961 soon after methicillin was introduced for clinical use against penicillinase-producing *S. aureus* [7, 8]. Methicillin-resistant *S. aureus* (MRSA) was subsequently reported in Australia, Europe, the USA, and Japan [9–12]. MRSA is now an important cause of healthcare-associated infections globally. Methicillin resistance in *S. aureus* emerged via the acquisition of the *mecA* gene located on a mobile genomic island designated staphylococcal chromosome cassette *mec* (SCC*mec*) by methicillin-susceptible *S. aureus* [10, 11]. The *mecA* gene is responsible for the synthesis of a novel penicillin-binding protein known as penicillin-binding protein 2a, which has decreased binding affinity for penicillin and cephalosporins [10, 11]. Therefore, MRSA strains are resistant to all β -lactam antibiotics [10, 11].

MRSA strains that were isolated in the 1960s and 1970s were usually susceptible to non- β -lactam antibiotics although they could carry resistance to tetracycline, erythromycin, and chloramphenicol [12]. However, from the late 1970s onwards, new strains of MRSA appeared that were resistant to multi-non- β -lactam agents including aminoglycosides, with only vancomycin left as an antibiotic of last resort for treating MRSA infections. These strains, described as epidemic MRSA (EMRSA), also possessed the capacity to spread extensively and caused serious infections mostly among hospitalized patients worldwide [12–14]. The risk factors for MRSA infections include old age (>60 years), hospitalization in the past 12 months, previous antibiotic use, admission to intensive care units, and catheterization [15].

Emergence of Community-Associated MRSA

MRSA was initially isolated from patients in large or tertiary care hospitals such as teaching hospitals, specialist hospitals, and nursing homes [16] and was described as healthcare-associated MRSA (HA-MRSA). In the 1990s, a new type of MRSA causing infections in the community among individuals who had no history of hospital admission or medical treatment in the previous

year was reported in Western Australia [17]. These types of MRSA strains were described as community-acquired, community-originated, community-associated, or community-onset MRSA (CA-MRSA) [18]. The initial report from Western Australia was followed by reports from New Zealand [19, 20], the USA [21–23], and Europe [24–26]. CA-MRSA is now a major cause of infections in the community as well as in healthcare facilities worldwide [27].

CA-MRSA have genetic and phenotypic characteristics that are different from those of HA-MRSA. CA-MRSA tend to be susceptible to a variety of non- β -lactam antibiotics, whereas HA-MRSA are typically multiresistant to antibiotics. CA-MRSA are more likely than HA-MRSA isolates to encode a putative virulence factor, i.e. Panton-Valentine leukocidin (PVL), a cytotoxin that has been associated with severe pneumonia [28–30], necrotizing fasciitis, and skin and soft tissue infections [11, 25, 30]. Furthermore, CA-MRSA often express lower levels of resistance to oxacillin (MIC: 8–32 mg/l) and multiply faster than HA-MRSA strains with significantly shorter doubling times, which may help CA-MRSA achieve successful colonization by enabling it to outcompete commensal bacterial flora [11]. In addition, CA-MRSA strains harbor SCC*mec* types IV and V, whereas the HA-MRSA carry SCC*mec* types I, II, and III [31, 32]. SCC*mec* types IV and V are smaller in size than SCC*mec* types I, II, and III [32]. Based on SCC*mec* typing, CA-MRSA strains are typically SCC*mec* IV, V, and VI, while HA-MRSA are usually SCC*mec* I, II, and III [32]. Otter and French [24] suggested that using SCC*mec* typing as a marker for CA-MRSA poses a particular problem because of the presence of successful hospital lineages carrying SCC*mec* IV, such as ST22 EMRSA-15-SCC*mec* IV or ST125-SCC*mec*IV, in some parts of the world. The use of SCC*mec* typing may increase the likelihood that these strains will be misclassified as CA-MRSA. Consequently, genotyping based on a combination of multilocus sequence typing (MLST) or Spa typing and SCC*mec* typing has been recommended as criteria for defining CA-MRSA, especially since CA-MRSA has become a major player in healthcare-associated infections [33].

The genetic characteristics, clinical presentations, virulence characteristics, and recommended treatment regimens for CA-MRSA have been reviewed extensively [27, 34–36]. This review focuses on the recent changes in the range of clinical presentations, the emergence of a multi-drug resistance, and the emergence of epidemic clones among recent CA-MRSA strains.

Epidemiological Typing of *S. aureus*

The diversity of *S. aureus* infections and virulence determinants warrant the institution of typing schemes to identify strains with unusual pathogenic abilities, to investigate the source and route of spread of infection, and for the formulation of effective control measures [37]. Over the years, many typing methods, including phage typing, plasmid analysis, pulsed-field gel electrophoresis (PFGE), and DNA sequencing-based methods have been used to type *S. aureus*. The developments and use of these typing methods have coincided with the emergence and dominance of different epidemic *S. aureus* clones. For simplicity, typing of *S. aureus* for epidemiological purposes is divided into three periods, namely pregenetic, genetic, and genomic eras.

The pregenetic era covers the period from 1950 to 1970. Phage typing was the primary method for typing *S. aureus* during this period. Bacteriophage typing is based on the susceptibility of *S. aureus* strains to lysis by specific bacteriophages. Fisk [38] showed that most coagulase-positive staphylococci carried phages that could be detected by cross-culture of pairs of strains and propagated on the sensitive member of the pair. Lysis by these phages was strain specific and was used to characterize isolates from human sources [38]. Phage typing identified the virulent strain of the penicillinase-producing *S. aureus* strain, i.e. phage 80/81, that was responsible for infections in Australian hospitals in the mid-1950s [4]. The phage 80/81 strains were also important causes of epidemics in hospitals in The Netherlands, the USA, and Canada [4]. Leukocidin was a major virulence factor of the 80/81 strains [37, 39].

The use of phage typing for the study of *S. aureus* outbreaks declined as strains of multiresistant MRSA became nontypeable with the available set of typing phages despite the inclusion of new supplementary phages in the phage typing protocol [40]. This period coincided with the introduction of plasmid analysis for epidemiological typing of *S. aureus*. This signaled the birth of the genetic era. Plasmid contents of bacterial strains and their resistance phenotypes constituted the early genotypic methods used for typing MRSA in the 1970s [40–43]. This was facilitated by the availability of rapid methods for isolating plasmid DNA in *S. aureus* [42, 43]. By combining plasmid content with the location of antibiotic resistance, i.e. on the chromosome or on plasmids in the bacterial strain, Grubb [12] differentiated MRSA that were isolated in the 1960s (classic MRSA) from the epidemic MRSA (EMRSA) strains that were spreading across Australia and Europe in the 1980s.

Epidemiological typing of *S. aureus* was further improved by the introduction of PFGE, MLST, *spa* typing, and SCCmec typing [44–47]. This period, the genomic era, witnessed the global expansion of EMRSA and the introduction of CA-MRSA clones.

PFGE enables the resolution of the whole bacterial chromosome. The procedure involves the application of controlled electric fields that change direction at a predetermined angle to samples of DNA that have been embedded in an agarose gel matrix and digested with an infrequent cutting restriction endonuclease. Adjustment of the electrophoretic conditions enables the separation of DNA fragments with lengths from 10 kilobases up to 10 megabases. The banding patterns are used for epidemiological typing [44]. Because of its high discriminatory capacity, PFGE has been regarded as the gold standard for investigating local outbreaks of bacterial infections [48].

MLST is a nucleotide sequence-based approach to the typing and characterization of bacterial isolates. MLST characterizes bacterial isolates using internal fragment sequences of 6–9 housekeeping genes. The fragments are approximately 450–500 bp in length [45]. For each housekeeping gene, the various sequences present within bacterial sequences are specified as distinct alleles. For each of the isolates, the alleles at all loci define the allelic profile or sequence type. MLST allows sequence data to be easily shared electronically between laboratories [45].

Spa typing is a single locus nucleotide sequence-based method that targets the protein A gene of *S. aureus*. The polymorphism in the variable X region in the *spa* gene encoding the staphylococcal surface protein A (*spa*) can be analyzed and used as a determinant to type *S. aureus* [46, 47]. The *Spa* typing method involves sequencing of the 24-bp variable repeat unit in the 3' region of the protein A gene (*spa*) and comparing the pattern of repeats to a public database [46, 47]. The results of *Spa* typing are comparable between laboratories electronically.

SCCmec typing characterizes MRSA isolates on the basis of the genetic elements known as SCCmec. SCCmec vary in size from 21 to 67 Kb [32]. The diversity of SCCmec types forms the basis of SCCmec typing, which facilitated the genetic differentiation of HA-MRSA from CA-MRSA. Based on SCCmec typing, CA-MRSA strains are typically SCCmec IV, V, and VI, while HA-MRSA ones are usually SCCmec I, II, and III [32]. Collectively, these typing methods are critical for identifying new strains, tracking outbreaks, and monitoring the evolution of *S. aureus*. The continued improvement and expansion

Table 1. Infections associated with CA-MRSA isolates: 2010–2012

Infections	References
Neck abscesses	50
Necrotizing fasciitis	51
Purulent pericardial effusion	52
Bloodstream infections	53, 54
Ocular infections	55–59
Osteomyelitis	60, 61
Epididymitis	62
Infective endocarditis	63
Prostatic abscess	64
Mycotic aneurysm	65
Infections in otolaryngology	66
Scarlet fever	67
Meningitis	68, 69
Liver abscess	70
Epidural abscess	71

of these methods, as well as the development of new diagnostic methods, will ensure that newly emerging *S. aureus* clones are identified quickly [48, 49].

Spectrum of Diseases Caused by CA-MRSA

Although CA-MRSA initially caused infections of the skin and soft tissues in young and healthy individuals [27], they have become more widespread in healthcare facilities. In addition, the spectrum of diseases associated with CA-MRSA has also increased over time. While infections of the skin and soft tissues such as furuncles, carbuncles, abscesses, cellulitis, and pneumonia are still the dominant clinical manifestation of CA-MRSA [14, 16, 27], other less common types of infections associated with CA-MRSA in recent years are presented in table 1.

Involvement of CA-MRSA in Outbreaks

There is a growing number of reports from different countries detailing the establishment of CA-MRSA as the dominant MRSA in healthcare facilities with some of the strains causing outbreaks of infections [72–77]. O'Brien et al. [77] first described an outbreak of infection in a Western Australian metropolitan hospital involving a CA-MRSA clone that was indistinguishable from a clone circulating in a remote community in the state. This was followed by reports of outbreaks caused by CA-MRSA in France [78]. Outbreaks caused by CA-MRSA now occur

worldwide [27], with increasing reports of outbreaks in neonatal units [79–82].

Transmission of CA-MRSA strains within healthcare facilities by some identified epidemic clones has resulted in the displacement of the previously endemic HA-MRSA clones [83]. In Germany, nonmultiresistant MRSA, including CC45-MRSA-IV and CC22-MRSA-IV clones, replaced the previously dominant multiresistant MRSA with the HA-MRSA genotype ST239-MRSA-III [84]. In the United Arab Emirates, MRSA with the HA-MRSA genotypes ST239-MRSA-III and ST22-MRSA-IV, which were the dominant strains in 2003, were replaced by CA-MRSA genotypes ST80-MRSA-IV and ST5-MRSA-IV in 2008 [85]. The ST30-MRSA-IV clone emerged in Singapore in 2004 and by 2005 had become the dominant MRSA clone in their hospitals [76]. Similarly, infections caused by CA-MRSA clone USA300 have surpassed those caused by the traditional HA-MRSA isolates in parts of the USA [27].

Emergence of Multiantibiotic Resistance in CA-MRSA

The earlier community MRSA isolates were generally susceptible to commonly used non- β -lactam antibiotics and were consequently described as nonmultiresistant MRSA [18, 86]. The absence of multiresistance in the early CA-MRSA isolates was likely due to a lack of exposure to many antibiotics rather than an inability to acquire multiantibiotic resistance determinants. In 2004, Munckhof et al. [87] showed that CA-MRSA isolates, when exposed to antibiotics in vitro over time, readily acquired resistance to these antibiotics. Soon after, multiresistance to non- β -lactams began to appear in CA-MRSA strains isolated in different centers. In North America, the USA300 CA-MRSA clone acquired multiantibiotic resistance via the acquisition of the multiresistant conjugative plasmid pUSA03 [88, 89, 90] or pSK41-like plasmids [91]. The conjugative plasmid pUSA03 encodes resistance to erythromycin, clindamycin, and high-level mupirocin. USA300 also express resistance to tetracycline and fluoroquinolones. High-level mupirocin resistance has also been reported in strains belonging to the ST80-SCCmec-IV CA-MRSA clone that was also resistant to tetracycline, fusidic acid, gentamicin, and streptomycin and cadmium acetate [92]. In Brazil, CA-MRSA strains belonging to the lineage ST1-SCCmec IV were reported to be resistant to erythromycin, clindamycin, ciprofloxacin, gentamicin, trimethoprim-sulfamethoxazole, and tetracycline [93]. Some strains belonging to the ST59 clone obtained from patients

Table 2. Epidemic CA-MRSA clones and their geographical distribution obtained from Udo et al. [81] and Mediavilla et al. [99]

Sequence types	SCC <i>mec</i>	Countries where they have been reported
ST1	IV	Australia, Brazil, Canada, China, Denmark, Egypt, Finland, France, Germany, Greece, Ireland, Italy, Japan, Kuwait, Pakistan, Romania, Samoa, Singapore, South Korea, Switzerland, United Arab Emirates, UK, USA
ST5	IV/V/VI	Algeria, Argentina, Australia, Austria, Azores, Cameroon, Canary Islands, China, Egypt, France, Germany, Iceland, Italy, Japan, Kuwait, Morocco, Nigeria, Samoa, Senegal, Spain, Switzerland, UK
ST8	IV/V/VI	Argentina, Australia, Austria, Belgium, Brazil, Bulgaria, Cameroon, Canada, Canary Islands, China, Colombia, Costa Rica, Cuba, Czech Republic, Denmark, Ecuador, Finland, France, French Polynesia, Gabon, Germany, Greece, Hong Kong, Iceland, India, Iraq, Ireland, Italy, Japan, Kuwait, Madagascar, Mexico, The Netherlands, New Zealand, Nigeria, Norway, Pakistan, Peru, Poland, Portugal, Romania, Russia, Samoa, South Korea, Spain, Sweden, Switzerland, Trinidad and Tobago, United Arab Emirates, UK, USA, Uruguay, Venezuela
ST30	IV/V	Australia, Austria, Brazil, Canary Islands, China, Czech Republic, Denmark, Egypt, Finland, France, French Polynesia, Germany, Hong Kong, Ireland, Italy, Japan, Kuwait, Latvia, Madagascar, Malaysia, The Netherlands, New Zealand, Pakistan, Peru, Philippines, Poland, Romania, Russia, Samoa, Singapore, South Korea, Spain, Sweden, Switzerland, Taiwan, Turkey, United Arab Emirates, UK, USA, Uruguay
ST45	IV/V/VI	Australia, Azores, Belgium, Germany, Hong Kong, Switzerland
ST59	IV/V	Australia, China, Denmark, Finland, Germany, Hong Kong, The Netherlands, Singapore, Sweden, Taiwan, UK, USA, Vietnam
ST72	IV/V	Czech Republic, Germany, Portugal, South Korea, Sweden, United Arab Emirates, USA
ST80	IV	Algeria, Australia, Austria, Belgium, Bulgaria, Croatia, Czech Republic, Denmark, Egypt, Finland, France, Germany, Greece, Ireland, Italy, Jordan, Kuwait, Lebanon, Libya, Malta, The Netherlands, Norway, Poland, Portugal, Romania, Saudi Arabia, Singapore, Slovenia, Spain, Sweden, Switzerland, Tunisia, United Arab Emirates, UK
ST88	IV/V	Angola, Cameroon, China, Gabon, Italy, Japan, Madagascar, Mali, The Netherlands, Niger, Nigeria, Portugal, Senegal, Spain, Sweden, United Arab Emirates, UK
ST93	IV	Australia, Czech Republic, Finland, Italy, The Netherlands, Samoa, UK
ST97	IV/V	Brazil, Canary Islands, Denmark, Egypt, Germany, Kuwait, Lebanon, The Netherlands, Sweden, United Arab Emirates, UK, USA
ST152	V	Austria, Canary Islands, Denmark, Germany, Kosovo, Macedonia, Slovenia, Sweden, Switzerland
ST772	IV/V	Australia, Bangladesh, Finland, Germany, Hong Kong, India, Ireland, Italy, Japan, The Netherlands, UK

in Vietnam were resistant to the macrolides, lincosamide, streptogramin, gentamicin, tetracycline, chloramphenicol, and trimethoprim-sulphamethoxazole [94–97].

Emergence of Epidemic CA-MRSA Clones

Several studies have shown that specific clones of CA-MRSA with diverse genetic backgrounds have emerged in different geographical areas, suggesting that these strains acquired SCC*mec* elements independently on multiple occasions [33, 36]. However, an increasing number of CA-MRSA clones have been reported to have attained epidemic status in recent years. These include

USA300, ST80, ST30, ST59, ST93, and ST772 [27, 36, 49, 98, 99]. The widely recognized epidemic CA-MRSA clones and their geographic distributions are summarized in table 2.

The USA300 clone is the dominant CA-MRSA clone causing infections in the USA [90, 99] and Canada [100]. The USA300 (PFGE-based designation) corresponds to MLST sequence type 8 (ST8). Outside of North America, USA300 has been isolated in Western Europe [101], Japan [71, 98], and Australia [102, 103]. The USA300 clones carry the SCC*mec* IV genetic element and produce PVL, and some carry the genes for ACME [30]. Surprisingly, USA300 is rare in the Middle East and Gulf Cooperation Council (GCC) countries.

ST80-MRSA has remained the dominant CA-MRSA clone in European countries [36, 99, 101, 103], North Africa, and the Middle East [85, 104, 105–110]. It has also been reported in Malaysia [111] and Australia [27, 112, 113]. The ST80-MRSA clones carry the *SCCmec* IV genetic element and belong to the accessory gene regulator (*agr*) type 3, and the majority of the strains produce PVL [99, 110]. Curiously, the ST80-MRSA clone does not seem to be successful in North America.

The ST30-MRSA clone, also known as the Southwest Pacific (SWP) clone, emerged in Oceania [20, 113] but has become a pandemic clone reported from many parts of the world, including the USA, Japan, Latin America, Turkey, the Middle East, Egypt, and many countries in Western Europe [36, 96, 109, 112, 113]. ST30 isolates are known as PFGE type USA1100 in the USA [114].

The ST59-MRSA clone emerged in Asia and is the dominant CA-MRSA clone prevalent in Taiwan [94, 115]. However, it has also been isolated in Australia [116, 117], China [95, 118], Hong Kong [119], Japan [120, 121], Europe [103], and the USA, where it is known as PFGE type USA1000 [114]. ST59-MRSA has also been isolated from cats and dogs in Malaysia, suggesting that cats and dogs are potential reservoirs of CA-MRSA [122].

The ST93-MRSA clone, also known as the Queensland MRSA clone, was first identified in Queensland and New South Wales, Australia, in 2000 [123]. It was soon reported in other parts of Australia [124, 125], though it was rarely reported in other countries. ST93-MRSA is highly virulent and has been associated with severe infections, including necrotizing pneumonia [125, 126], although it harbors very few known virulence determinants [126, 127].

The ST772-MRSA clone, also known as the Bengal clone, was first reported in Bangladesh [128]. It has since been reported in India [56, 129, 130], Ireland [82], Australia, Germany, Hong Kong, the United Arab Emirates, and Saudi Arabia [112, 128]. The ST772 clone is rapidly becoming the dominant CA-MRSA clone in Indian hospitals [56, 129, 130]. Interestingly, PVL-producing strains of ST772 isolated in India are relatively susceptible to antimicrobial agents compared to ST772 clones isolated in Western Australia that appear to be more resistant to antibacterial agents [112].

CA-MRSA in the GCC Countries

CA-MRSA strains have emerged as important causes of infections in healthcare facilities in the GCC as in other parts of the world. CA-MRSA have been reported in

Saudi Arabia, the United Arab Emirates, Kuwait, and Bahrain. In Saudi Arabia, CA-MRSA was first reported among patients at the King Fahd Hospital in the Eastern Province of the country in 1998 [131]. Since then, CA-MRSA have been reported in other areas of the kingdom with increasing prevalence, and their antibiotic susceptibility patterns have also changed with increasing levels of resistance to erythromycin, tetracycline, clindamycin, ciprofloxacin, and gentamicin [132]. Genetic analysis of CA-MRSA isolates obtained in a hospital in Riyadh revealed that they belonged to different clones [108].

In the United Arab Emirates, a study conducted at Tawam Hospital showed that the prevalence of CA-MRSA increased from 38.5% of all MRSA isolated at the hospital in 2003 to 73.1% in 2008 [85]. These strains consisted mostly of ST80-MRSA-IV, ST5-MRSA-IV, and ST1-MRSA clones. The study also revealed that only one third of the CA-MRSA infections were actually acquired in the community, indicating that CA-MRSA clones had entered and spread within the hospital [85].

In Bahrain, a study conducted at the Salmaniya Medical Complex in 2005 revealed that 13.3% of their MRSA strains were CA-MRSA. These strains were nonmultiresistant to non- β -lactam antibiotics, belonged to different genetic backgrounds, and were isolated mostly from skin and soft tissue sources [133].

In Kuwait, CA-MRSA was first identified in 2001 [134]. The prevalence of CA-MRSA increased from 1.7% of all MRSA in 2001 to 17% in 2005 [135]. Molecular characterization of CA-MRSA obtained between 2006 and 2007 showed that ST80-*SCCmec*-IV (51.1%), ST30-*SCCmec*-IV (22.2%), and ST5-*SCCmec*-IV (14.1%) were the dominant CA-MRSA clones in the country [110]. The ST80-*SCCmec*-IV strains obtained in Kuwait hospitals belonged to diverse PFGE backgrounds and resistance patterns. Some strains had resistance patterns similar to that of the European clone obtained in Denmark [101].

There are currently no published data on the prevalence of CA-MRSA in Qatar and Oman, although MRSA has been reported in these countries.

Relation of the Current CA-MRSA to the Pandemic *S. aureus* Phage 80/81 Clone

Similarities were observed between the infections and spread of some CA-MRSA clones and the patterns of infections associated with those of the pandemic phage 80/81 MSSA clones. For example, the phage 80/81 clone caused a wide range of syndromes including skin infec-

tions (boils, carbuncles, and pustules) and fatal sepsis or pneumonia [136] similar to the spectrum of infection caused by CA-MRSA [27]. The phage 80/81 clone emerged rapidly in Australia before being reported in the UK, Canada, and the USA in the 1950s, similar to the pattern of emergence of CA-MRSA strains. In addition, outbreaks of infections caused by the phage 80/81 clone occurred initially among newborns, young children, and nursing mothers. This mirrors the initial association of CA-MRSA infections in the young healthy and outbreaks among neonates [137]. These observations, together with the emergence and successful spread of EMRSA-16 in the UK and ST30 globally, prompted a comparison of these clones with strains belonging to the phage 80/81 MSSA to establish their relatedness [114]. To clarify the relationship between phage 80/81 and contemporary CA-MRSA clones, DeLeo et al. [114] sequenced representative isolates belonging to the phage 80/81, SWP CA-MRSA (ST30-MRSA), and EMRSA-16 genomes and compared their virulence determinants. The study revealed that phage 80/81 and EMRSA-16 belonged to the same clonal complex 30 (CC30), suggesting that they may have a common ancestor. The study also suggested that there was a coincident divergence of these clones from a common ancestor but, unlike the suggestion by Robinson et al. [138], the SWP clone was not a descendant of the phage 80/81 clone.

Conclusions

There is significant diversity in MRSA arising in communities worldwide. As CA-MRSA has become established in healthcare facilities, the range of infections caused by them has also increased. Although many CA-MRSA still maintain a nonmultidrug resistant antimicrobial profile, multiresistance to non- β -lactam agents has emerged in some clones, posing substantial problems for empirical and directed therapy of infections caused by these strains. The emergence of pandemic CA-MRSA clones not only limits therapeutic options but also presents significant challenges for infection control. Continued monitoring of global epidemiology and emerging drug resistance data is critical for the effective management of these infections.

Disclosure Statement

The author declares no conflict of interest.

References

- Lowy FD: *Staphylococcus aureus* infections. N Engl J Med 1998;339:520–532.
- Beigelman PM, Rantz LA: Clinical importance of coagulase – positive, penicillin-resistant *Staphylococcus aureus*. N Engl J Med 1950;242:353–358.
- Barber M: Staphylococcal infections due to penicillin-resistant strains. Br Med J 1947;2: 863–865.
- Shanson DC: Antibiotic-resistant *Staphylococcus aureus*. J Hosp Infect 1981;2:11–36.
- Collopy BT, Dalton MF, Wright C, et al: Comparison of the clinical significance of methicillin-resistant and methicillin-sensitive *Staphylococcus aureus* isolates. Med J Aust 1984;140:211–214.
- Smith JT, Amyes SGB: Bacterial resistance to antifolate chemotherapeutic agents mediated by plasmids. Br Med J 1984;40:42–46.
- Jevons PM: Celbenin-resistant staphylococci. Br Med J 1961;1:124.
- Barber M: Methicillin-resistant staphylococci. J Clin Pathol 1961;14:385–393.
- Barrett FF, McGehee RF, Finland M Jr: Methicillin-resistant *Staphylococcus aureus* at Boston City Hospital: bacteriologic and epidemiologic observations. N Engl J Med 1968;279: 441–448.
- Baba T, Takeuchi F, Kuroda M, et al: Genome and virulence determinants of high virulence community-acquired MRSA. Lancet 2002; 359:1819–1827.
- Okuma K, Iwakawa K, Turnidge JD, et al: Dissemination of new methicillin-resistant *Staphylococcus aureus* clones in the community. J Clin Microbiol 2002;40:4289–4294.
- Grubb WB: Genetics of MRSA. Rev Med Microbiol 1998;9:153–162.
- Murchan S, Aucken HM, O'Neill GL, et al: Emergence, spread, and characterization of phage variants of epidemic methicillin-resistant *Staphylococcus aureus* 16 in England and Wales. J Clin Microbiol 2004;42:5154–5160.
- Del Giudice P, Bes M, Hubiche T, et al: Clinical manifestations and outcome of skin infections caused by the community-acquired methicillin-resistant *Staphylococcus aureus* clone ST80-IV. J Eur Acad Dermatol Venerol 2011;25:164–169.
- Wang JT, Wang JL, Fang CT, et al: Risk factors for mortality of nosocomial methicillin-resistant *Staphylococcus aureus* (MRSA) blood infection: with investigation of the potential role of community-associated MRSA strains. J Infect 2010;61:449–459.
- Said-Salim B, Mathema B, Kreiswirth BN: Community-acquired methicillin-resistant *Staphylococcus aureus*: an emerging pathogen. Infect Control Hosp Epidemiol 2003;24: 451–455.
- Udo EE, Pearman JW, Grubb WB: Genetic analysis of community isolates of methicillin-resistant *Staphylococcus aureus* in Western Australia. J Hosp Infect 1993;25:97–108.
- Gosbell IB, Mercer JL, Neville SA, et al: Non-multiresistant and multiresistant methicillin-resistant *Staphylococcus aureus* in community-acquired infections. Med J Aust 2001;174: 627–630.
- Smith JMB: Emerging problems of antibiotic resistant bacteria. NZ Med J 1998;111:441–444.
- Adhikari RP, Cook GM, Lamont I, et al: Phenotypic and molecular characterization of community occurring, Western Samoan phage pattern methicillin-resistant *Staphylococcus aureus*. J Antimicrob Chemother 2002; 50:825–831.

- 21 Herold BC, Immergluck LC, Maranan MC, et al: Community-acquired methicillin-resistant *Staphylococcus aureus* in children with no identified predisposing risk. *JAMA* 1998; 279:593–598.
- 22 Groom AV, Wolsey DH, Naimi TS, et al: Community-acquired methicillin-resistant *Staphylococcus aureus* in a rural American Indian community. *JAMA* 2001;286:1201–1205.
- 23 Centers for Disease Control and Prevention: Four pediatric deaths from community-acquired methicillin-resistant *Staphylococcus aureus* – Minnesota and North Dakota, 1997–1999. *MMWR Morb Mortal Wkly Rep* 1999; 48:707–710.
- 24 Otter JA, French GL: The emergence of community-associated methicillin-resistant *Staphylococcus aureus* at a London teaching hospital, 2000–2006. *Clin Microbiol Infect* 2008;14: 670–676.
- 25 Vandenesch F, Naimi T, Enright MC, et al: Community-acquired methicillin-resistant *Staphylococcus aureus* carrying Panton-Valentine leukocidin genes: worldwide emergence. *Emerg Infect Dis* 2003;9:978–984.
- 26 Wannet WJ, Spalburg E, Heck ME, et al: Emergence of virulent methicillin-resistant *Staphylococcus aureus* strains carrying Panton-Valentine leukocidin genes in The Netherlands. *J Clin Microbiol* 2005;43:3341–3345.
- 27 David MZ, Daum RS: Community-associated methicillin-resistant *Staphylococcus aureus*: epidemiology and clinical consequences of an emerging epidemic. *Clin Microbiol Rev* 2010; 23:616–687.
- 28 Liassine N, Auckenthaler R, Descombes MC, et al: Community-acquired methicillin-resistant *Staphylococcus aureus* isolated in Switzerland contains the Panton-Valentine leukocidin or exfoliative toxin genes. *J Clin Microbiol* 2004;42:825–828.
- 29 Lina G, Piemont Y, Godail-Gamot F, et al: Involvement of Panton-Valentine leukocidin producing *Staphylococcus aureus* in primary skin infections and pneumonia. *Clin Infect Dis* 1999;29:1128–1132.
- 30 Diep BA, Sensabaugh GF, Somboona NS, et al: Widespread skin and soft-tissue infections due to two methicillin-resistant *Staphylococcus aureus* strains harbouring the genes for Panton-Valentine leukocidin. *J Clin Microbiol* 2004;42:2080–2084.
- 31 Ma XX, Ito T, Tiensasitorn C, et al: Novel type of staphylococcal cassette chromosome *mec* identified in community-acquired methicillin-resistant *Staphylococcus aureus* strains. *Antimicrob Agents Chemother* 2002;46: 1147–1152.
- 32 Ito T, Katayama Y, Asada K, et al: Structural comparison of three types of cassette chromosome *mec* integrated in the chromosome in methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* 2001;45: 1323–1336.
- 33 Otter JA, French GL: Community-associated methicillin-resistant *Staphylococcus aureus*: the case for a genotypic definition. *J Hosp Infect* 2012;81:143–148.
- 34 Zetola N, Francis JS, Nuermberger EL, et al: Community-acquired methicillin-resistant *Staphylococcus aureus*: an emerging threat. *Lancet Infect Dis* 2005;5:275–286.
- 35 Pichereau S, Rose WE: Invasive community-associated MRSA infections: epidemiology and antimicrobial management. *Expert Opin Pharmacother* 2010;11:3009–3025.
- 36 Mediavilla JR, Chen L, Mathema B, et al: Global epidemiology of community-associated methicillin resistant *Staphylococcus aureus* (CA-MRSA). *Curr Opin Microbiol* 2012;15: 588–595.
- 37 Parker MT: The significance of phage-typing patterns in *Staphylococcus aureus*; in Easmon CSF, Adlam C (eds): *Staphylococci and Staphylococcal Infections*. London, Academic Press, 1983, vol 1, pp 33–61.
- 38 Fisk RT: Studies on staphylococci. 1. Occurrence of bacteriophage carriers among strains of *Staphylococcus aureus*. *J Infect Dis* 1942;71: 153–160.
- 39 Donahue JA, Baldwin JN: Hemolysin and leukocidin production by 80/81 strains of *Staphylococcus aureus*. *J Infect Dis* 1966;116:324–328.
- 40 Noble WC, Rayman M: Plasmid profiling of epidemic staphylococci from around 1960: a comparison of epidemiological techniques. *J Hyg (Lond)* 1986;97:211–218.
- 41 McGowan JE, Terry PM, Huang T-SR, et al: Nosocomial infections with gentamicin-resistant *Staphylococcus aureus*: plasmid analysis as an epidemiologic tool. *J Infect Dis* 140: 864–972.
- 42 Townsend DE, Ashdown N, Bolton S, et al: The use of cetyltrimethyl ammonium bromide for rapid isolation from *Staphylococcus aureus* of relaxable and non-relaxable plasmid DNA for in vitro manipulation. *Lett Appl Microbiol* 1985;1:87–94.
- 43 Grinstead J, Bennett PM: Isolation and purification of plasmid DNA; in Bennet PM, Grinstead J (eds): *Methods in Microbiology*. London, Academic Press, 1984, vol 17, pp 123–131.
- 44 O'Brien FG, Udo EE, Grubb WB: Contour-clamped homogeneous electric field electrophoresis of *Staphylococcus aureus*. *Nat Protoc* 2006;1:3028–3033.
- 45 Enright MC, Day NPJ, Davies CE, et al: Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. *J Clin Microbiol* 2000;38:1008–1015.
- 46 Shopsis B, Gomez M, Montgomery SO, et al: Evaluation of protein A gene polymorphic region DNA sequencing for typing of *Staphylococcus aureus* strains. *J Clin Microbiol* 1999; 37:3556–3563.
- 47 Harmsen D, Claus H, Witte W, et al: Typing of methicillin-resistant *Staphylococcus aureus* in a University Hospital setting by using novel software for spa repeat determination and database management. *J Clin Microbiol* 2003; 41:5442–5448.
- 48 Deurenberg RH, Stobberingh EE: The evolution of *Staphylococcus aureus*. *Infect Genet Evol* 2008;8:747–763.
- 49 Kennedy AD, DeLeo FR: Epidemiology and virulence of community-associated MRSA. *Clin Microbiol Lett* 2009;31:153–160.
- 50 Duggal P, Naseri I, Sobol SE: The increased risk of community-acquired methicillin *Staphylococcus aureus* neck abscesses in young children. *Laryngoscope* 2011;121:51–55.
- 51 Lulich IJ, Sam-Agudu NA: Community-acquired methicillin-resistant *Staphylococcus aureus* necrotizing fasciitis in a healthy adolescent male. *Minn Med* 2010;93:44–46.
- 52 Arora NP, Kottam A, Mahajan N, et al: Purulent pericardial effusion from community-acquired methicillin-resistant *Staphylococcus aureus*. *Am J Med Sci* 2012;344:160–162.
- 53 Kempker RR, Farley MM, Ladson JL, et al: Association of methicillin-resistant *Staphylococcus aureus* (MRSA) USA300 genotypes with mortality in MRSA bacteremia. *J Infect* 2010; 61:372–381.
- 54 Furuno JP, Johnson JK, Schweizer ML, et al: Community-associated methicillin-resistant *Staphylococcus aureus* bacteremia and endocarditis among HIV patients: a cohort study. *BMC Infect Dis* 2011;11:298.
- 55 Hsiao CH, Chuang CC, Tan HY, et al: Methicillin-resistant *Staphylococcus aureus* ocular infections: a 10-year hospital-based study. *Ophthalmology* 2012;119:522–527.
- 56 Nadig S, Velusamy N, Lalitha P, et al: *Staphylococcus aureus* eye infections in two Indian hospitals: emergence of ST772 as a major clone. *Clin Ophthalmol* 2012;6:165–173.
- 57 Vaska VL, Grimwood K, Gole GA, et al: Community-associated methicillin-resistant *Staphylococcus aureus* causing orbital cellulitis in Australian children. *Pediatr Infect Dis* 2011; 30:1003–1006.
- 58 Pandian DG, Babu RK, Chaitra A, et al: Nine years review of preseptal and orbital cellulitis and emergence of community-acquired methicillin-resistant *Staphylococcus aureus* in a tertiary hospital in India. *Indian J Ophthalmol* 2011;59:431–435.
- 59 Kobayashi D, Givner LB, Yeatts RP, et al: Infantile orbital cellulitis secondary to community-associated methicillin-resistant *Staphylococcus aureus*. *J AAPOS* 2011;15:208–210.
- 60 Tomovic S, Friedel ME, Liu JK, et al: Community-acquired methicillin-resistant *Staphylococcus aureus* skull base osteomyelitis with occipital condylar cerebrospinal fluid leak in an immunocompetent patient. *Laryngoscope* 2012;122:977–981.
- 61 Ulug M, Ayaz C, Celen MK: A case report and literature review: osteomyelitis caused by community-associated methicillin resistant *Staphylococcus aureus*. *J Infect Dev Ctries* 2011;5:896–900.
- 62 Raychaudhuri M, Chew PR: A case of epididymitis associated with Panton-Valentine leukocidin *Staphylococcus aureus*. *Sex Transm Infect* 2012;88:355–356.

- 63 Wu KS, Lee SS, Tsai HC, et al: Non-nosocomial healthcare-associated infective endocarditis in Taiwan: an under recognized disease with poor outcome. *BMC Infect Dis* 2011;11:221.
- 64 Park SC, Lee JW, Rim JS: Prostatic abscess caused by community-acquired methicillin-resistant *Staphylococcus aureus*. *Int J Urol* 2011;18:536–538.
- 65 Lee NY, Kan CD, Chi CY, et al: Community-acquired methicillin-resistant *Staphylococcus aureus* mycotic aneurysm. *Intern Med* 2010; 49:623–625.
- 66 Shaw GY, Montandon SV, Driks MR: Otolaryngologic community-acquired MRSA infections in a suburban private practice. *Ear Nose Throat J* 2010;89:E33–E40.
- 67 Lu YC, Chen SJ, Lo WT: Scarlet fever caused by community-associated methicillin-resistant *Staphylococcus aureus*. *Indian Pediatr* 2011;48:563–565.
- 68 Aguilar J, Urdy-Cornejo V, Donabedian S, et al: *Staphylococcus aureus* meningitis: case series and literature review. *Medicine* 2010;89: 117–125.
- 69 Gattuso G, Palvarini L, Tomasoni D, et al: A case of community-acquired MRSA (CA-MRSA) sepsis complicated by meningoen- cephalitis and cerebral abscess, successfully treated with linezolid. *Infez Med* 2009;17: 244–248.
- 70 Wong VW, Cheung YS, Wong J, et al: A community-acquired methicillin-resistant *Staphylococcus aureus* liver abscess. *Hong Kong Med J* 2010;16:227–229.
- 71 Higashiyama M, Ito T, Han X, et al: Epidural abscess caused by community-associated methicillin-resistant *Staphylococcus aureus* strain USA300 in Japan. *J Infect Chemother* 2010;16:345–349.
- 72 Valsesia G, Rossi M, Bertschy S, et al: Emergence of SCCmec type IV and SCCmec type V methicillin-resistant *Staphylococcus aureus* containing the Panton-Valentine leukocidin genes in a large academic teaching hospital in central Switzerland: external invaders or persisting circulators? *J Clin Microbiol* 2010;48: 720–727.
- 73 Patel M, Hoesley CJ, Moser SA, et al: Dissemination of community-associated methicillin-resistant *Staphylococcus aureus* in a tertiary care hospital. *South Med J* 2008;101:40–45.
- 74 Maree CL, Daum RS, Boyle-Vavra S, et al: Community-associated methicillin-resistant *Staphylococcus aureus* causing healthcare-associated infections. *Emerg Infect Dis* 2007;13: 236–242.
- 75 Gosbell IB: Epidemiology, clinical features and management of infections due to community methicillin-resistant *Staphylococcus aureus* (cMRSA). *Intern Med J* 2005;35(suppl 2):S120–S135.
- 76 Hsu L-Y, Koh Y-L, Chlebiga NL, et al: Establishment of ST30 as the predominant clonal type among community-associated methicillin-resistant *Staphylococcus aureus* isolates in Singapore. *J Clin Microbiol* 2006;44:1090–1093.
- 77 O'Brien FG, Pearman JW, Gracey M, et al: Community strain of methicillin-resistant *Staphylococcus aureus* involved in an outbreak. *J Clin Microbiol* 1999;37:2858–2862.
- 78 Dufour P, Gillet Y, Bes M, et al: Community-acquired methicillin-resistant *Staphylococcus aureus* infections in France: emergence of a single clone that produces Panton-Valentine leukocidin. *Clin Infect Dis* 2002;35:819–824.
- 79 Gould IM, Girvan EK, Browning RA, et al: Report of a hospital neonatal unit outbreak of community-associated methicillin-resistant *Staphylococcus aureus*. *Epidemiol Infect* 2009;137:1242–1248.
- 80 Carey AJ, Long SS: *Staphylococcus aureus*: a continuously evolving and formidable pathogen in the neonatal intensive care unit. *Clin Perinatol* 2010;37:535–546.
- 81 Udo EE, Aly NY, Sarkhoo E, et al: Detection and characterization of an ST97-SCCmec-V community-associated methicillin-resistant *Staphylococcus aureus* clone in a neonatal intensive care unit and special care baby unit. *J Med Microbiol* 2011;60:600–604.
- 82 Brennan GI, Shore AC, Corcoran S, et al: Emergence of hospital- and community-associated Panton-Valentine leukocidin-positive methicillin-resistant *Staphylococcus aureus* genotype ST772-MRSA-V in Ireland and detailed investigation of an ST772-MRSA-V cluster in a neonatal intensive care unit. *J Clin Microbiol* 2012;50:841–847.
- 83 Popovich K, Hota B, Rice T, et al: Phenotypic prediction rule for community-associated methicillin-resistant *Staphylococcus aureus*. *J Clin Microbiol* 2007;45:2293–2295.
- 84 Albrecht N, Jatzwauk L, Slickers P, et al: Clonal replacement of epidemic methicillin-resistant *Staphylococcus aureus* strains in a German university hospital over a period of eleven years. *PLoS One* 2011;6:e28189.
- 85 Sonnevend A, Blair I, Alkaabi M, et al: Change in methicillin-resistant *Staphylococcus aureus* clones at a tertiary care hospital in the United Arab Emirates over a 5-year period. *J Clin Pathol* 2012;65:178–182.
- 86 Nimmo GR, Schoonoveldt J, O'Kane G, et al: Community-acquisition of gentamicin-sensitive methicillin-resistant *Staphylococcus aureus* in southeast Queensland, Australia. *J Clin Microbiol* 2000;38:3926–3931.
- 87 Munkhof WJ, Kleinschmidt SL, Turnidge JD: Resistance development in community-acquired strains of methicillin-resistant *Staphylococcus aureus*: an in vitro study. *Int J Antimicrob Agents* 2004;24:605–608.
- 88 Diep BA, Chambers HF, Graber CJ, et al: Emergence of multidrug-resistant, community-associated, methicillin-resistant *Staphylococcus aureus* clone USA300 in men who have sex with men. *Ann Intern Med* 2008;148: 249–257.
- 89 Simor AE, Stuart TL, Louie L, et al, Canadian Nosocomial Infection Surveillance Program: Mupirocin-resistant, methicillin-resistant *Staphylococcus aureus* strains in Canadian hospitals. *Antimicrob Agents Chemother* 2007;51: 3880–3886.
- 90 Tenover FC, McDougal LK, Goering RV, et al: Characterization of a strain of community-associated methicillin-resistant *Staphylococcus aureus* widely disseminated in the United States. *J Clin Microbiol* 2006;44:108–118.
- 91 McDougal LK, Fosheim GE, Nicholson A, et al: Emergence of resistance among USA300 methicillin-resistant *Staphylococcus aureus* isolates causing invasive disease in the United States. *Antimicrob Agents Chemother* 2010; 54:3804–3811.
- 92 Udo EE, Sarkhoo E: Genetic analysis of high-level mupirocin resistance in the ST80 clone of community-associated methicillin-resistant *Staphylococcus aureus*. *J Med Microbiol* 2010; 59:193–199.
- 93 Silva-Carvalho MC, Bonelli RR, Souza RR, et al: Emergence of multiresistant variants of the community-acquired methicillin-resistant *Staphylococcus aureus* lineage ST1-SCCmecIV in 2 hospitals in Rio de Janeiro, Brazil. *Diagn Microbiol Infect Dis* 2009;65:300–305.
- 94 Boyle-Vavra S, Ereshefsky B, Wang C-C, et al: Successful multiresistant community-associated methicillin-resistant *Staphylococcus aureus* lineage from Taipei, Taiwan, that carries either the novel staphylococcal chromosome cassette mec (SCCmec) type V_T or SCCmec type IV. *J Clin Microbiol* 2005;43: 4719–4730.
- 95 Wang L, Liu Y, Yang Y, et al: Multidrug-resistant clones of community-associated methicillin-resistant *Staphylococcus aureus* isolated from Chinese children and the resistance genes to clindamycin and mupirocin. *J Med Microbiol* 2012;61:1240–1247.
- 96 Takizawa Y, Taneike I, Nakagawa S, et al: A Panton-Valentine leukocidin (PVL)-positive community-acquired methicillin-resistant *Staphylococcus aureus* (MRSA) strains, another such strain carrying a multiple-drug resistance plasmid, and other more typical PVL-negative MRSA strains found in Japan. *J Clin Microbiol* 2005;43:3356–3363.
- 97 Chen C-J, Hsueh P-R, Su L-H, et al: Change in the molecular epidemiology of methicillin-resistant *Staphylococcus aureus* bloodstream infections in Taiwan. *Diagn Microbiol Infect Dis* 2009;65:199–201.
- 98 Shibuya Y, Hara M, Higuchi W, et al: Emergence of the community-acquired methicillin-resistant *Staphylococcus aureus* USA300 clone in Japan. *J Infect Chemother* 2008;14: 439–441.
- 99 Mediavilla JR, Chen L, Mathema B, et al: Global epidemiology of community-associated methicillin resistant *Staphylococcus aureus* (CA-MRSA). *Curr Opin Microbiol* 2012;15: 588–595.

- 100 Simor AE, Gilbert NL, Gravel D, et al, Canadian Nosocomial Infection Surveillance Program: Methicillin-resistant *Staphylococcus aureus* colonization or infection in Canada: National Surveillance and Changing Epidemiology, 1995–2007. *Infect Control Hosp Epidemiol* 2010;31:348–356.
- 101 Larsen AR, Bocher S, Stegger M, et al: Epidemiology of European community-associated methicillin-resistant *Staphylococcus aureus* clonal complex 80 type IV strains isolated in Denmark from 1993 to 2004. *J Clin Microbiol* 2008;46:62–68.
- 102 Nimmo GR: USA300 abroad: global spread of a virulent strain of community-associated methicillin-resistant *Staphylococcus aureus*. *Clin Microbiol Infect* 2012;18:725–734.
- 103 Monecke S, Jatzwauk L, Weber S, et al: DNA microarray-based genotyping of methicillin-resistant *Staphylococcus aureus* strains from eastern Saxony. *Clin Microbiol Infect* 2008;14:534–545.
- 104 Enany S, Yaoita E, Yoshida Y, et al: Molecular characterization of Pantone-Valentine leukocidin-positive community-acquired methicillin-resistant *Staphylococcus aureus* in Egypt. *Microbiol Res* 2010;165:152–162.
- 105 Nejma MB, Mastouri M, Jrad BBH, et al: Characterization of ST80 Pantone-Valentine leukocidin-positive community-acquired methicillin-resistant *Staphylococcus aureus* clone in Tunisia. *Diagn Microbiol Infect Dis* 2008, DOI: 10.1016/j.diagmicrobio.2008.02.010.
- 106 Khalil W, Hashwa F, Shihabi A, et al: Methicillin-resistant *Staphylococcus aureus* ST80-IV clone in children from Jordan. *Diagn Microbiol Infect Dis* 2012;73:228–230.
- 107 Tokajian S, Abou KP, Jabbour D, et al: Molecular characterization of *Staphylococcus aureus* in Lebanon. *Epidemiol Infect* 2010; 138:707–712.
- 108 Monecke S, Skakni L, Hasan R, et al: Characterisation of MRSA strains isolated from patients in a hospital in Riyadh, Kingdom of Saudi Arabia. *BMC Microbiol* 2012;12:146.
- 109 Udo EE, O'Brien FG, Al-Sweih N, et al: Genetic lineages of community-associated methicillin-resistant *Staphylococcus aureus* in Kuwait hospitals. *J Clin Microbiol* 2008; 46:3514–3516.
- 110 Udo EE, Sarkhoo E: The dissemination of ST80-SCCmec-IV community-associated methicillin resistant *Staphylococcus aureus* clone in Kuwait hospitals. *Ann Clin Microbiol Antimicrob* 2010;9:31.
- 111 Ahmad N, Ruzan IN, Abd Ghani MK, et al: Characteristics of community- and hospital-acquired methicillin-resistant *Staphylococcus aureus* strains carrying SCCmec type IV isolated in Malaysia. *J Med Microbiol* 2009;58: 1213–1218.
- 112 Monecke S, Coombs G, Shore AC, et al: A field guide to pandemic, epidemic and sporadic clones of methicillin-resistant *Staphylococcus aureus*. *PLoS One* 2011;6:e17936.
- 113 Coombs GW, Nimmo GR, Bell JM, et al, Australian Group for Antimicrobial Resistance: Genetic diversity among community methicillin-resistant *Staphylococcus aureus* strains causing outpatient infections in Australia. *J Clin Microbiol* 2004;42:4735–4743.
- 114 DeLeo FR, Otto M, Krieswirth BN, et al: Community-associated methicillin-resistant *Staphylococcus aureus*. *Lancet* 2010;375: 1557–1568.
- 115 Takano T, Higuchi W, Otsuka T, et al: Novel characteristics of community-acquired methicillin-resistant *Staphylococcus aureus* strains belonging to multilocus sequence type 59 in Taiwan. *Antimicrob Agents Chemother* 2008;52:837–845.
- 116 Gosbell IB, Barbagiannakos T, Neville SA, et al: Non-multiresistant methicillin-resistant *Staphylococcus aureus* bacteraemia in Sydney, Australia: emergence of EMRSA-15, Oceania, Queensland and Western Australian MRSA strains. *Pathology* 2006;38:239–244.
- 117 Coombs GW, Monecke S, Ehrlich R, et al: Differentiation of clonal complex 59 community-associated methicillin-resistant *Staphylococcus aureus* in Western Australia. *Antimicrob Agents Chemother* 2010;54:1914–1921.
- 118 Li X, Sun J, Wu D, et al: Pantone-Valentine leukocidin gene sequence variation and phage in methicillin-resistant and methicillin-susceptible *Staphylococcus aureus* from children in mainland China. *Microbiol Immunol* 2012;56:155–162.
- 119 Ho PL, Chiu SS, Chan MY, et al: Molecular epidemiology and nasal carriage of *Staphylococcus aureus* and methicillin-resistant *S. aureus* among young children attending day care centers and kindergartens in Hong Kong. *J Infect* 2012;64:500–506.
- 120 Kawaguchiya M, Urushibara N, Kuwahara O, et al: Molecular characteristics of community-acquired methicillin-resistant *Staphylococcus aureus* in Hokkaido, northern main island of Japan: identification of sequence types 6 and 59 Pantone-Valentine leukocidin-positive community-acquired methicillin-resistant *Staphylococcus aureus*. *Microb Drug Resist* 2011;17:241–250.
- 121 Hisata K, Ito T, Matsunaga N, et al: Dissemination of multiple MRSA clones among community-associated methicillin-resistant *Staphylococcus aureus* infections from Japanese children with impetigo. *J Infect Chemother* 2011;17:609–621.
- 122 Aklilu E, Zunita Z, Hassan L, et al: Phenotypic and genotypic characterization of methicillin-resistant *Staphylococcus aureus* (MRSA) isolated from dogs and cats at University Veterinary Hospital, Universiti Putra Malaysia. *Trop Biomed* 2010;27:483–492.
- 123 Munckhof WJ, Schooneveldt J, Coombs GW, et al: Emergence of community-acquired methicillin-resistant *Staphylococcus aureus* (MRSA) infection in Queensland, Australia. *Int J Infect Dis* 2003;7:259–264.
- 124 Coombs GW, Pearson JC, O'Brien FG, et al: Methicillin-resistant *Staphylococcus aureus* clones, Western Australia. *Emerg Infect Dis* 2006;12:241–247.
- 125 Peleg AY, Munckhof WJ, Kleinschmidt SL, et al: Life-threatening community-acquired methicillin-resistant *Staphylococcus aureus* infection in Australia. *Eur J Clin Microbiol Infect Dis* 2005;384–387.
- 126 Coombs GW, Goering RV, Chua KY, et al: The molecular epidemiology of the highly virulent ST93 Australian community *Staphylococcus aureus* strain. *PLoS One* 2012;7:e43037.
- 127 Chua KY, Seemann T, Harrison PF, et al: The dominant Australian community-acquired methicillin-resistant *Staphylococcus aureus* clone ST93-IV[2B] is highly virulent and genetically distinct. *PLoS One* 2012;7:e25887.
- 128 Afroz S, Kobayashi N, Nagashima S, et al: Genetic characterization of *Staphylococcus aureus* isolates carrying Pantone-Valentine leukocidin genes in Bangladesh. *Jpn J Infect Dis* 2008;61:393–396.
- 129 D'Souza N, Rodrigues C, Mehta A: Molecular characterization of methicillin-resistant *Staphylococcus aureus* with emergence of epidemic clones of sequence type (ST) 22 and ST772 in Mumbai, India. *J Clin Microbiol* 2010;48:1806–1811.
- 130 Shambat S, Nadig S, Prabhakara S, et al: Clonal complexes and virulence factors of *Staphylococcus aureus* from several cities in India. *BMC Microbiol* 2012;12:64.
- 131 Bukharie HA, Abdelhadi MS, Saeed IA, et al: Emergence of methicillin-resistant *Staphylococcus aureus* as a community pathogen. *Diagn Microbiol Infect Dis* 2001;40:1–4.
- 132 Bukharie HA: A review of community-acquired methicillin-resistant *Staphylococcus aureus* for primary care physicians. *J Fam Comm Med* 2010;17:117–120.
- 133 Udo EE, Panigrahi D, Jamsheer AE: Molecular typing of methicillin-resistant *Staphylococcus aureus* isolated in a Bahrain hospital. *Med Princ Pract* 2008;17:308–314.
- 134 Udo EE, Al-Sweih N, Noronha B: Characterisation of non-multiresistant methicillin-resistant *Staphylococcus aureus* (including EMRSA-15) in Kuwait hospitals. *Clin Microbiol Infect* 2006;12:262–269.
- 135 Udo EE, Al-Sweih N, Dhar R, et al: Surveillance of antibacterial resistance in *Staphylococcus aureus* in Kuwait hospitals. *Med Princ Pract* 2008;17:71–75.
- 136 Roundtree PM, Beard MA: Hospital strains of *Staphylococcus aureus* with particular reference to methicillin-resistant strains. *Med J Aust* 1968;2:1163–1168.
- 137 Shukla SK, Stemper ME, Ramaswamy SV, et al: Molecular characteristics of nosocomial and Native American community-associated methicillin-resistant *Staphylococcus aureus* clones from rural Wisconsin. *J Clin Microbiol* 2004;42:3752–3757.
- 138 Robinson DA, Kearns AM, Holmes A, et al: Re-emergence of early pandemic *Staphylococcus aureus* as a community-acquired methicillin-resistant clone. *Lancet* 2005;365: 1256–1258.