Original Paper

Age Sensitivity of NF κ B Abundance and Programmed Cell Death in Erythrocytes Induced by NF κ B Inhibitors

Mehrdad Ghashghaeinia^{a,b} Judith C. Cluitmans^c Mahmoud Toulany^d
Mohammad Saki^d Martin Köberle^e Elisabeth Lang^a Peter Dreischer^a
Tilo Biedermann^e Michael Duszenko^f Florian Lang^a Giel J. Bosman^{c,g}
Thomas Wieder^{e,g}

^aDepartment of Physiology, University of Tübingen, Tübingen, ^bDepartment of Dermatology, University Medical Center Kiel, Kiel, Germany; ^cDepartment of Biochemistry, Radboud University Nijmegen Medical Centre, Nijmegen Centre for Molecular Life Sciences, HB Nijmegen, The Netherlands;

^dDivision of Radiobiology and Molecular Environmental Research, Department of Radiation Oncology, University Medical Center Tübingen, ^eDepartment of Dermatology, University Medical Center Tübingen,

^fInterfaculty Institute of Biochemistry, University of Tübingen, Tübingen, Germany; ^gG. J. Bosman and T. Wieder contributed equally to this work and thus share senior authorship

Key Words

Phosphatidylserine • Bay 11-7082 • Parthenolide • Cell volume • Eryptosis

Abstract

Background/Aims: Erythrocytes may enter eryptosis, a suicidal death characterized by cell shrinkage and phosphatidylserine exposure at the erythrocyte outer membrane. Susceptibility to eryptosis is enhanced in aged erythrocytes and stimulated by NF κ B-inhibitors Bay 11-7082 and parthenolide. Here we explored whether expression of NF κ B and susceptibility to inhibitor-induced eryptosis is sensitive to erythrocyte age. **Methods:** Human erythrocytes were separated into five fractions, based on age-associated characteristics cell density and volume. NF κ B compared to β -actin protein abundance was estimated by Western blotting and cell volume from forward scatter. Phosphatidylserine exposure was identified using annexin-V binding. **Results:** NF κ B was most abundant in young erythrocytes but virtually absent in aged erythrocytes. A 24h or 48h exposure to Ringer resulted in spontaneous decrease of forward scatter and increase of annexin V binding, effects more pronounced in aged than in young erythrocytes. Both, Bay 11-7082 (20 μ M) and parthenolide (100 μ M) triggered eryptosis, effects again most pronounced in aged erythrocytes. **Conclusion:** NF κ B protein abundance is lowest and spontaneous eryptosis as well as susceptibility to Bay 11-7082 and parthenolide highest in aged erythrocytes. Thus, inhibition of NF κ B signalling alone is not responsible for the stimulation of eryptosis by parthenolide or Bay 11-7082.

Copyright © 2013 S. Karger AG, Basel

Introduction

Senescence of nucleated cells is defined as an irreversible cellular growth arrest and is induced by internal factors, such as oncogenes [1], or external stimuli, e. g. cytokines [2]. Anucleated, circulating human erythrocytes are unable to proliferate and thus are growth-arrested per se. As a consequence, their life span is limited by a special form of senescence leading to a gradual process of cellular shrinkage and phosphatidylserine exposure, and eventually to clearance of aged erythrocytes after approx. 120 days [3]. Mechanisms accounting for erythrocyte senescence include a conformational change of the band 3 membrane protein domain thus leading to the appearance of an antigen specific for senescent cells. This triggers binding of specific autologous immunoglobulin G and subsequent removal of senescent erythrocytes by macrophages, such as Kupffer cells in the liver [4]. The conformational change presumably results from oxidative damage of membrane lipids, membrane proteins, or hemoglobin [5-9].

Senescent erythrocytes have been shown to be particularly sensitive to triggers of eryptosis [10], the suicidal erythrocyte death characterized by breakdown of cell membrane integrity, phosphatidylserine asymmetry and cell shrinkage [11]. Eryptosis is triggered by increase of the cytosolic Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) [11]. Ca^{2+} may enter erythrocytes through Ca^{2+} -permeable cation channels [12, 13], which are activated by oxidative stress [14]. Increased $[\text{Ca}^{2+}]_i$ activates Ca^{2+} -sensitive K^+ channels [15] and subsequently K^+ exit, hyperpolarization, Cl^- exit and water loss that leads to cell shrinkage [16]. Increased $[\text{Ca}^{2+}]_i$ further triggers cell membrane scrambling with phosphatidylserine exposure at the erythrocyte surface [17]. Ca^{2+} sensitivity of cell membrane scrambling is enhanced by ceramide [18]. Eryptosis is further stimulated by energy depletion [19] and caspase activation [20-24]. Signalling of eryptosis involves several kinases including AMP activated kinase (AMPK) [13], cGMP-dependent protein kinase [25], Janus-activated kinase 3 (JAK3) [26], casein kinase [27, 28], p38 kinase [29], PAK2 kinase [30] as well as sorafenib- [31] and sunifinib- [32] sensitive kinases.

Eryptosis is observed in several clinical conditions [11], such as diabetes [24, 33, 34], renal insufficiency [35], hemolytic uremic syndrome [36], sepsis [37], malaria [38-43], sickle cell disease [44], Wilson's disease [42], iron deficiency [45], malignancy [46], phosphate depletion [47], and metabolic syndrome [48]. Moreover, eryptosis is triggered by a wide variety of xenobiotics [10, 32, 48-67] including Nuclear Factor κ B (NFκB) inhibitors Bay 11-7082 and parthenolide [68]. NFκB is known as a transcription factor fostering cell survival [69-71]. The possibility was therefore considered that NFκB participates in the signalling of eryptosis [72]. Alternatively, the inhibitors are in part effective by mechanisms unrelated to NFκB-inhibition.

The present study explored if the amount of NFκB changes during the erythrocyte's aging and if a correlation between enhanced eryptosis in aged erythrocytes [10] and NFκB abundance exists. To demonstrate the possible functional relationship between age-sensitive NFκB abundance and age-sensitive eryptosis, the effect of Bay 11-7082 and parthenolide have been tested in young and aged erythrocytes. As a result, a correlation was found between erythrocyte age and NFκB abundance on the one hand as well as eryptosis on the other. However, the fact that both NFκB inhibitors showed the highest eryptosis induction in aged erythrocytes, where NFκB abundance is lowest, indicates an NFκB-independent mode of action.

Materials and Methods

Erythrocytes, solutions and chemicals

As described earlier [10], erythrocytes were isolated from the blood of healthy volunteers by following the guidelines of the ethical commission of the Radboud University Nijmegen Medical Centre. Blood was

collected in EDTA, and erythrocytes were fractionated according to cell volume using elutriation followed by a fractionation according to cell density using discontinuous Percoll gradients as described earlier [73]. This yields 24 fractions, that were combined to five fractions (I to V); whereby fraction I comprises the youngest, and fraction V the oldest cells [73, 74]. These fractions have been characterized with respect to cell survival *in vivo*, hemoglobin content, metabolome characteristics and membrane composition. The mean corpuscular volume (MCV) of these fractions was measured using a haematology analyzer (Sysmex XT1800i, Sysmex Corporation, Kobe, Japan).

Fractionated erythrocytes were incubated *in vitro* at a hematocrit of 0.4% in Ringer solution containing (in mM) 125 NaCl, 5 KCl, 1 MgSO₄, 32 N-2-hydroxyethylpiperazine-N-2-ethanesulfonic acid (HEPES), 5 glucose, 1 CaCl₂; pH 7.4 at 37°C for 24 or 48 hours [10]. Bay 11-7082 (Sigma, München, Germany) or parthenolide (Biomol, Hamburg, Germany) were added to the Ringer solution at the indicated concentrations for 24 or 48 hours. For this, stock solutions of Bay 11-7082 or parthenolide in dimethyl sulfoxide (DMSO) were diluted in Ringer solution. DMSO did not exceed final concentrations of 0.1%. Vehicle-treated erythrocytes served as controls.

Phosphatidylserine exposure and forward scatter

Erythrocyte fractions I to V or the corresponding unfractionated population (or whole blood indicated as WB) were incubated in Ringer solution in the absence or presence of Bay 11-7082 or parthenolide. After incubation, erythrocytes were washed in annexin-binding buffer at pH 7.4 containing (in mM): 125 NaCl, 10 HEPES and 5 CaCl₂. Erythrocytes were then stained with Annexin-Fluos from Roche Diagnostics (Mannheim, Germany) at a 1:35 dilution and mixed gently on a vortex mixer. After 20 min incubation in the dark at room temperature, samples were diluted 1:5 with annexin-binding buffer, thoroughly mixed to achieve single cell suspensions, and analysed by flow cytometry on a FACS-Calibur from Becton Dickinson (Heidelberg, Germany). Cell volume differences were estimated by forward scatter (FSC), and annexin-fluorescence intensity was measured in FL-1 with an excitation wavelength of 488 nm and an emission wavelength of 530 nm as described earlier [10].

Western blotting

To determine age-dependent differences in the expression of NFκBs (i.e. p50 and p65 subunits), human erythrocytes were separated into five fractions (fractions I to V). Each fraction (250 µl erythrocyte pellet containing approx. 1.0 × 10⁹ cells) was lysed in 50 ml of 20 mM HEPES/NaOH (pH 7.4) hypotonic shock solution containing 1 "complete protease inhibitor cocktail" tablet from Roche Diagnostics. Ghost membranes were pelleted (15.000 × g, for 30 min at 4°C) and lysed in 250 µl lysis buffer pH 7.4 containing 125 mM NaCl, 25 mM HEPES, 10 mM EDTA, 10 mM NaF, 10 mM sodium pyrophosphate, 0.1% sodium dodecyl sulfate, 0.5% sodium deoxycholate, 1% Triton X-100, 0.4% β-mercaptoethanol and 1 "complete protease inhibitor cocktail" tablet. Lysed ghost membranes were solubilized in Laemmli sample buffer at 95°C for 5 min, resolved by 8% SDS-PAGE and electrophoretically transferred onto a PVDF membrane (Roth, Karlsruhe, Germany) as described earlier [68]. Membranes were then incubated in blocking solution (5% nonfat milk in tris-buffered saline (TBS) containing 0.01% Tween 20 (TBST)) at room temperature for 1 h. For detection of NFκB subunits p65 and p50, the membranes were incubated with a 1:1000 dilution of affinity purified rabbit anti-NFκB p65 or anti-NFκB p50 (Cell Signaling Technology Inc., Danvers, MA, USA) at 4°C overnight. After washing membranes with TBST, immunoreactive proteins were visualized using enhanced chemoluminescence following incubation with a 1:5000 dilution of the secondary donkey-anti-rabbit horse-radish-peroxidase (HRP)-conjugated antibody (GE Healthcare, München, Germany) for 1 h at room temperature. β-actin was used as loading control and its detection was evaluated by an affinity isolated rabbit anti-β-actin antibody (Sigma-Aldrich, Taufkirchen, Germany). Immunoreactive bands were quantified by videodensitometry, and the NFκB p65/β-actin ratio or NFκB p50/β-actin ratio of the samples was calculated.

Statistics

Data are expressed as arithmetic or geometric means ± SEM. Statistical analysis was made using ANOVA with Tukey's test as post hoc test, or using student's t test where appropriate. A p value < 0.05 was considered statistically significant.

Fig. 1. Decreased erythrocyte volume and increased annexin-V binding in old erythrocytes. A. Means erythrocyte age (left) and mean corpuscular volume (MCV, right) in the five fractions isolated by volume and density fractionation. MCV data from one representative fractionation are shown. B. Means $\pm$ SEM ($n = 3$) of the erythrocyte forward scatter (geometric mean) of fractions I to V following incubation for 24 hours at 37°C in control Ringer solution. * significantly different from fraction I ($p < 0.05$; ANOVA). C. Means $\pm$ SEM ($n = 3$) of the erythrocyte forward scatter (geometric mean) of fractions I and V following incubation for 48 hours at 37°C in control Ringer solution. * significantly different from fraction I ($p < 0.05$; student's *t* test). D. Mean percentage $\pm$ SEM ($n = 3$) of annexin V binding of erythrocytes from fractions I to V following incubation for 24 hours at 37°C in control Ringer solution. * significantly different from fraction I ($p < 0.05$; ANOVA). E. Mean percentage $\pm$ SEM ($n = 3$) of annexin V binding of erythrocytes from fractions I and V following incubation for 48 hours at 37°C in control Ringer solution. * significantly different from fraction I ($p < 0.05$; student's *t* test).

Results

According to our previous study, the susceptibility to triggers of eryptosis is a function of erythrocyte age. Thus, erythrocytes have been separated into 5 fractions (I to V) according to age-associated differences in density and volume [10]. As illustrated in Fig. 1A, the MCV steadily declined with increasing age (higher fraction numbers) which served as a quality control for separation. In addition, cell volume decrease was confirmed by measurement of the forward scatter (FSC) which was lower in fraction V as compared with fraction I after incubation for 24 hours in Ringer solution (Fig. 1B). The forward scatter decreased even further following incubation in Ringer for 48 hours (Fig. 1C).

The decrease in forward scatter was paralleled by an increase of annexin V binding. As shown in Fig. 1D, after incubation for 24 hours in Ringer solution the percentage of annexin V binding erythrocytes was higher in fraction V than in fraction I. Incubation of the erythrocytes in Ringer solution for 48 hours resulted in a further increase of the percentage of annexin V binding cells (Fig. 1E).

In theory, the increased eryptosis of aged erythrocytes could be paralleled by changes of NFκB abundance. To explore the impact of erythrocyte age on the amount of NFκB, the levels of the NFκB subunits p65 and p50 have been determined by Western blotting. As shown in Fig. 2A, the abundance of the NFκB subunit p65 was highest in erythrocytes of fraction I. In erythrocytes of fraction V, the NFκB protein subunit p65 was virtually absent (Fig. 2A, C). Similar observations were made with NFκB subunit p50, which was again most abundant in fraction I and almost absent in fraction V (Fig. 2B, D).

Fig. 2. Erythrocyte NFκB subunit abundance as a function of erythrocyte age. A. Western blot demonstrating the expression of NFκB protein subunit p65 (upper panel) and β-actin (lower panel) in fractions I to V of a representative erythrocytes fractionation. WB = whole blood extracts (unfractionated); Co. = positive control for p65 expression. B. Western blot demonstrating the expression of NFκB protein subunit p50 (upper panel) and β-actin (lower panel) in fractions I to V of a representative erythrocytes fractionation. WB = whole blood extracts (unfractionated); Co. = positive control for p50 expression. C. Arithmetic means ± SEM (n = 3) of the NFκB protein subunit p65/β-actin abundance in erythrocytes of fractions I to V. Data are given as relative expression values (% of whole blood). * significantly different from fraction I (p < 0.05). D. Arithmetic means ± SEM (n = 3) of the NFκB protein subunit p50/β-actin abundance in erythrocytes of fractions I to V. Data are given as relative expression values (% of whole blood). * significantly different from fraction I (p < 0.05).

Fig. 3. Positive correlation between erythrocyte NFκB subunit abundance and erythrocyte forward scatter. A. Erythrocyte forward scatter following incubation for 24 hours in control Ringer solution as a function of NFκB protein subunit p65/β-actin abundance. B. Erythrocyte forward scatter following incubation for 24 hours in control Ringer solution as a function of NFκB protein subunit p50/β-actin abundance.

Fig. 3 displays the result of erythrocyte forward scatter after 24 h of incubation as a function NFκB subunit p65 (Fig. 3A) or NFκB subunit p50 (Fig. 3B) abundance. Fig. 4 displays the percentage of annexin V binding erythrocytes after a 48 hours incubation in Ringer solution as a function of NFκB subunit p65 (Fig. 4A) or NFκB subunit p50 (Fig. 4B) abundance.

The positive correlation between the forward scatter, and the negative correlation between the percentage of annexin V-exposing erythrocytes and the amount of NFκB subunits p50 and p65 does not necessarily reflect a causal relationship between the NFκB subunits abundance and the susceptibility to eryptosis. However, if such a causal relationship would exist, then pharmacological inhibition of the NFκB signalling pathway should trigger

Fig. 4. Inverse correlation between erythrocyte NFκB subunit abundance and phosphatidylserine exposure. A. The percentage of Annexin V binding erythrocytes following incubation for 48 hours in control Ringer solution as a function of NFκB protein subunit p65/β-actin abundance. B. The percentage of Annexin V binding erythrocytes following incubation for 48 hours in control Ringer solution as a function of NFκB protein subunit p50/β-actin abundance.

Fig. 5. Effect of Bay 11-7082 on phosphatidylserine exposure of erythrocytes of different age. A. Original histograms of annexin V binding erythrocytes in fraction I (left) or fraction V (right) following exposure for 24 h to Ringer solution in the absence (DMSO) or presence of 21 μM Bay 11-7082. B. Mean percentage ± SEM in fractions I to V (n = 3) of annexin V binding erythrocytes following exposure for 24 h (left) or 48 h (right) to Ringer solution in the absence (DMSO) or presence of 21 μM Bay 11-7082. Note that erythrocytes from fraction V are more susceptible to Bay 11-7082-induced eryptosis as compared with erythrocytes from fraction I. C. Annexin V binding of erythrocytes from fraction I (red squares) or fraction V (black squares) as a function of Bay 11-7082 concentration following exposure for 24 h. Data represent the mean of 3 determinations.

eryptosis. Moreover, pharmacological inhibition of NFκB signalling should dissipate the difference between young and aged erythrocytes, if mainly the lack of NFκB accounted for the enhanced susceptibility of aged erythrocytes to eryptosis. As illustrated in Fig. 5, the opposite was true. Treatment of erythrocytes with 20 μM Bay 11-7082, a concentration which should completely block NFκB, was followed by stimulation of annexin V binding, an effect more pronounced in aged erythrocytes than in young erythrocytes (Fig. 5A, B). Accordingly, titration of Bay 11-7082 did not dissipate but enhanced the difference of annexin V binding between erythrocytes of fraction I and erythrocytes of fraction V (Fig. 5C). Similar observations were made with 100 μM parthenolide (Fig. 6A, B), which again did not dissipate but enhanced the difference of annexin V binding between young and old erythrocytes (Fig.

Fig. 6. Effect of parthenolide on phosphatidylserine exposure of erythrocytes of different age. A. Original histograms of annexin V binding erythrocytes in fraction I (left) or fraction V (right) following exposure for 24 h to Ringer solution in the presence of 100 μM parthenolide. B. Mean percentage ± SEM in fractions I to V (n = 3) of annexin V binding erythrocytes following exposure for 24 h to Ringer solution in the absence (DMSO) or presence of 100 μM parthenolide. Note that erythrocytes from fraction V are more susceptible to parthenolide-induced eryptosis as compared with erythrocytes from fraction I. DMSO controls are the same as shown in Fig. 5B. C. Annexin V binding of erythrocytes from fraction I (red squares) or fraction V (black squares) as a function of parthenolide concentration following exposure for 24 h. Data represent the mean of 3 determinations.

6C). Thus, in aged erythrocytes with the lowest NFκB abundance, the inverse correlation between the higher sensitivity to parthenolide and Bay 11-7082-induced eryptosis and the NFκB expression level has to be based on NFκB-independent modes of action of the inhibitors. As a consequence, young erythrocytes, with their high NFκB abundance and the associated low sensitivity to parthenolide- and Bay 11-7082-induced eryptosis, require a higher concentration of parthenolide or Bay 11-7082 in order to achieve the same rate of eryptosis as aged erythrocytes (Fig. 6C and Fig. 5C, respectively).

Discussion

Our data confirm the previous observation that aged erythrocytes are particularly sensitive to eryptosis [10]. The present study further reveals that the amount of NFκB is constantly decreasing during an erythrocyte's life span and negligible in aged erythrocytes. In view of the earlier observation that pharmacological inhibition of NFκB triggered eryptosis [68], the coincidence of low NFκB protein subunit abundance and high susceptibility to eryptosis could have reflected a causal relationship. If so, pharmacological inhibition of NFκB should be more effective in erythrocytes expressing high NFκB levels than in erythrocytes expressing low levels of NFκB. Accordingly, pharmacological inhibition of NFκB should dissipate the difference between young and aged erythrocytes. As illustrated in Fig. 5 and Fig. 6, the effects

of parthenolide or Bay 11-7082 on eryptosis were more pronounced in aged than in young erythrocytes. Accordingly, the inhibitors augmented the differences between erythrocytes of fraction I and erythrocytes from fraction V. Thus, the difference in susceptibility to inhibitor-induced eryptosis between young and aged erythrocytes was not due to the differences in NFκB abundance. Moreover, even though the substances are expected to inhibit NFκB, their effectivity in erythrocytes of fraction V, i.e. in the virtual absence of NFκB, indicates that the substances trigger eryptosis by mechanisms other than inhibition of NFκB. For example, these mechanisms could be glutathione (GSH) depletion (data not shown) [68, 75-77] and/or caspase activation [78-80]. Parthenolide with its properties to inhibit many components of the canonical NFκB signalling pathway, e.g. IKKs [81-83], and NFκB p65 [84-86] seems to be a much better candidate to investigate the possible role of NFκB signalling in cell death mechanisms in erythrocytes than Bay 11-7082 which in this pathway solely inhibits IKK [87-89]. Bay 11-7082 was originally identified as an inhibitor of the NFκB signalling pathway [90].

We have shown previously that old erythrocytes contain higher Ca^{2+} levels [10]. Thus, age-sensitive eryptosis is partially Ca^{2+} -dependent, e.g. through Ca^{2+} -mediated activation of the scramblase and the Gardos channel. The analysis of the relationship between Redox-sensitive NFκB activity and Ca^{2+} -induced cell death, as demonstrated in nucleated cells [91], will be a challenging task of future eryptosis research.

The enhanced *in vitro* susceptibility of aged erythrocytes towards parthenolide should lead to their elimination, and thus to a drift towards younger erythrocyte populations *in vivo*. In consequence, this could enhance erythropoiesis resulting in increased reticulocyte counts *in vivo*. However, clinical studies using feverfew extracts or parthenolide did not show any significant toxicity [92]. Administration of the NFκB inhibitor ethacrynic acid [93] favourably influences the clinical course of sickle cell anemia by tight covalent binding of this compound to hemoglobin S [94]. Whether or not other NFκB inhibitors, e.g. parthenolide or Bay 11-7082, equally possess an anti-anemia effect or influence the numbers of circulating reticulocytes remains to be shown.

The present results do not address the potential role of NFκB for the regulation of gene expression in erythrocyte progenitor cells. In this context, evidence has been provided that the glucocorticoid receptor is needed for stress erythropoiesis [95]. As glucocorticoid receptors are known to inhibit NFκB transcriptional activity by direct physical interaction in a DNA-independent manner (for review see: [96]), stimulation of eryptosis by NFκB inhibitors might also be related to decreased abundance of glucocorticoid receptors in aged human erythrocytes. In addition, acting as transcription factors in progenitor cells, NFκBs may indeed control the expression of genes relevant for the susceptibility to eryptosis. Mature, circulating erythrocytes are, however, devoid of nuclei and unable to express novel proteins. In those cells, the NFκB protein abundance may be irrelevant for the susceptibility to eryptosis.

Interestingly, we can find certain parallels regarding the inverse correlation between NFκB abundance and eryptosis when we look at the proliferative rate of cancer cells and their NFκB activity. Cancer cells with low NFκB DNA-binding activity also exhibit a significantly higher sensitivity to the anti-proliferative effects of parthenolide and vice versa [97], i.e. cancer cells with a high NFκB DNA-binding activity show a significantly lower sensitivity to the anti-proliferative effects of parthenolide.

In conclusion, our current study clearly indicates an inverse correlation between erythrocyte age and NFκB abundance. Thus, NFκB protein was most abundant in young erythrocytes and virtually absent in aged erythrocytes. Concomitantly, NFκB inhibitor-induced eryptosis was most pronounced in aged erythrocytes, pointing to NFκB-independent mechanisms leading to an enhanced susceptibility of aged erythrocytes to parthenolide- and Bay11-7082-induced eryptosis.

Conflict of Interest

Competing interests: the authors have no competing interests.

Acknowledgements

The authors acknowledge the meticulous preparation of the manuscript by Ali Soleimanpour and Tanja Loch. We thank Ms. Bovee for her help in the cell separations and the department of Quality Control at the Sanquin Bloodbank Southeast Region, Nijmegen, the Netherlands for performing the MCV measurements.

This study was supported by the Deutsche Forschungsgemeinschaft (Nr. La 315/4-3, La 315/6-1 and Wi 1279/3-1), Open Access Publishing Fund of Tübingen University, the Wilhelm Sander-Stiftung (Nr. 2012.056.1), and the Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie (Center for Interdisciplinary Clinical Research).

References

- 1 Wajapeyee N, Serra RW, Zhu X, Mahalingam M, Green MR: Oncogenic BRAF induces senescence and apoptosis through pathways mediated by the secreted protein IGFBP7. *Cell* 2008;132:363-374.
- 2 Braumuller H, Wieder T, Brenner E, Assmann S, Hahn M, Alkhaled M, Schilbach K, Essmann F, Kneilling M, Griessinger C, Ranta F, Ullrich S, Mocikat R, Braungart K, Mehra T, Fehrenbacher B, Berdel J, Niessner H, Meier F, van den Broek M, Haring HU, Handgretinger R, Quintanilla-Martinez L, Fend F, Pesic M, Bauer J, Zender L, Schaller M, Schulze-Osthoff K, Rocken M: T-helper-1-cell cytokines drive cancer into senescence. *Nature* 2013;494:361-365.
- 3 Callender S, Race RR, Paykoc ZV: Hypersensitivity to Transfused Blood. *Br Med J* 1945;2:83-84.
- 4 Kay M: Immunoregulation of cellular life span. *Ann N Y Acad Sci* 2005;1057:85-111.
- 5 Arese P, Turrini F, Schwarzer E: Band 3/complement-mediated recognition and removal of normally senescent and pathological human erythrocytes. *Cell Physiol Biochem* 2005;16:133-146.
- 6 Bosman GJ, Willekens FL, Werre JM: Erythrocyte aging: a more than superficial resemblance to apoptosis? *Cell Physiol Biochem* 2005;16:1-8.
- 7 Low PS, Waugh SM, Zinke K, Drenckhahn D: The role of hemoglobin denaturation and band 3 clustering in red blood cell aging. *Science* 1985;227:531-533.
- 8 Lutz HU, Fasler S, Stammler P, Bussolino F, Arese P: Naturally occurring anti-band 3 antibodies and complement in phagocytosis of oxidatively-stressed and in clearance of senescent red cells. *Blood Cells* 1988;14:175-203.
- 9 Pantaleo A, Giribaldi G, Mannu F, Arese P, Turrini F: Naturally occurring anti-band 3 antibodies and red blood cell removal under physiological and pathological conditions. *Autoimmun Rev* 2008;7:457-462.
- 10 Ghashghaieinia M, Cluitmans JC, Akel A, Dreischer P, Toulany M, Koberle M, Skabytska Y, Saki M, Biedermann T, Duszenko M, Lang F, Wieder T, Bosman GJ: The impact of erythrocyte age on eryptosis. *Br J Haematol* 2012;157:606-614.
- 11 Lang F, Gulbins E, Lerche H, Huber SM, Kempe DS, Föller M: Eryptosis, a window to systemic disease. *Cell Physiol Biochem* 2008;22:373-380.
- 12 Foller M, Kasinathan RS, Koka S, Lang C, Shumilina E, Birnbaumer L, Lang F, Huber SM: TRPC6 contributes to the Ca²⁺ leak of human erythrocytes. *Cell Physiol Biochem* 2008;21:183-192.
- 13 Foller M, Sopjani M, Koka S, Gu S, Mahmud H, Wang K, Floride E, Schleicher E, Schulz E, Munzel T, Lang F: Regulation of erythrocyte survival by AMP-activated protein kinase. *FASEB J* 2009;23:1072-1080.
- 14 Brand VB, Sandu CD, Duranton C, Tanneur V, Lang KS, Huber SM, Lang F: Dependence of Plasmodium falciparum in vitro growth on the cation permeability of the human host erythrocyte. *Cell Physiol Biochem* 2003;13:347-356.
- 15 Brugnara C, de Franceschi L, Alper SL: Inhibition of Ca²⁺-dependent K⁺ transport and cell dehydration in sickle erythrocytes by clotrimazole and other imidazole derivatives. *J Clin Invest* 1993;92:520-526.

- 16 Lang PA, Kaiser S, Myssina S, Wieder T, Lang F, Huber SM: Role of Ca²⁺-activated K⁺ channels in human erythrocyte apoptosis. *Am J Physiol Cell Physiol* 2003;285:C1553-C1560.
- 17 Berg CP, Engels IH, Rothbart A, Lauber K, Renz A, Schlosser SF, Schulze-Osthoff K, Wesselborg S: Human mature red blood cells express caspase-3 and caspase-8, but are devoid of mitochondrial regulators of apoptosis. *Cell Death Differ* 2001;8:1197-1206.
- 18 Lang F, Gulbins E, Lang PA, Zappulla D, Foller M: Ceramide in suicidal death of erythrocytes. *Cell Physiol Biochem* 2010;26:21-28.
- 19 Klari BA, Lang PA, Kempe DS, Niemoeller OM, Akel A, Sobiesiak M, Eisele K, Podolski M, Huber SM, Wieder T, Lang F: Protein kinase C mediates erythrocyte "programmed cell death" following glucose depletion. *Am J Physiol Cell Physiol* 2006;290:C244-C253.
- 20 Bhavsar SK, Bobbala D, Xuan NT, Foller M, Lang F: Stimulation of suicidal erythrocyte death by alpha-lipoic acid. *Cell Physiol Biochem* 2010;26:859-868.
- 21 Foller M, Huber SM, Lang F: Erythrocyte programmed cell death. *IUBMB Life* 2008;60:661-668.
- 22 Foller M, Mahmud H, Gu S, Wang K, Floride E, Kucherenko Y, Luik S, Laufer S, Lang F: Participation of leukotriene C(4) in the regulation of suicidal erythrocyte death. *J Physiol Pharmacol* 2009;60:135-143.
- 23 Lau IP, Chen H, Wang J, Ong HC, Leung KC, Ho HP, Kong SK: In vitro effect of CTAB- and PEG-coated gold nanorods on the induction of eryptosis/erythroptosis in human erythrocytes. *Nanotoxicology* 2012;6:847-856.
- 24 Maellaro E, Leoncini S, Moretti D, Del Bello B, Tanganelli I, De Felice C, Ciccoli L: Erythrocyte caspase-3 activation and oxidative imbalance in erythrocytes and in plasma of type 2 diabetic patients. *Acta Diabetol* DOI: 10.1007/s00592-011-0274-0.
- 25 Foller M, Feil S, Ghoreschi K, Koka S, Gerling A, Thunemann M, Hofmann F, Schuler B, Vogel J, Pichler B, Kasinathan RS, Nicolay JP, Huber SM, Lang F, Feil R: Anemia and splenomegaly in cGKI-deficient mice. *Proc Natl Acad Sci USA* 2008;105:6771-6776.
- 26 Bhavsar SK, Gu S, Bobbala D, Lang F: Janus kinase 3 is expressed in erythrocytes, phosphorylated upon energy depletion and involved in the regulation of suicidal erythrocyte death. *Cell Physiol Biochem* 2011;27:547-556.
- 27 Kucherenko Y, Zelenak C, Eberhard M, Qadri SM, Lang F: Effect of casein kinase 1alpha activator pyrvinium pamoate on erythrocyte ion channels. *Cell Physiol Biochem* 2012;30:407-417.
- 28 Zelenak C, Eberhard M, Jilani K, Qadri SM, Macek B, Lang F: Protein kinase CK1alpha regulates erythrocyte survival. *Cell Physiol Biochem* 2012;29:171-180.
- 29 Gatidis S, Zelenak C, Fajol A, Lang E, Jilani K, Michael D, Qadri SM, Lang F: p38 MAPK activation and function following osmotic shock of erythrocytes. *Cell Physiol Biochem* 2011;28:1279-1286.
- 30 Zelenak C, Foller M, Velic A, Krug K, Qadri SM, Viollet B, Lang F, Macek B: Proteome analysis of erythrocytes lacking AMP-activated protein kinase reveals a role of PAK2 kinase in eryptosis. *J Proteome Res* 2011;10:1690-1697.
- 31 Lupescu A, Shaik N, Jilani K, Zelenak C, Lang E, Pasham V, Zbidah M, Plate A, Bitzer M, Foller M, Qadri SM, Lang F: Enhanced Erythrocyte Membrane Exposure of Phosphatidylserine Following Sorafenib Treatment: An in vivo and in vitro Study. *Cell Physiol Biochem* 2012;30:876-888.
- 32 Shaik N, Lupescu A, Lang F: Sunitinib-sensitive suicidal erythrocyte death. *Cell Physiol Biochem* 2012;30:512-522.
- 33 Calderon-Salinas JV, Munoz-Reyes EG, Guerrero-Romero JF, Rodriguez-Moran M, Bracho-Riquelme RL, Carrera-Gracia MA, Quintanar-Escorza MA: Eryptosis and oxidative damage in type 2 diabetic mellitus patients with chronic kidney disease. *Mol Cell Biochem* 2011;357:171-179.
- 34 Nicolay JP, Schneider J, Niemoeller OM, Artunc F, Portero-Otin M, Haik G Jr, Thornalley PJ, Schleicher E, Wieder T, Lang F: Stimulation of suicidal erythrocyte death by methylglyoxal. *Cell Physiol Biochem* 2006;18:223-232.
- 35 Myssina S, Huber SM, Birka C, Lang PA, Lang KS, Friedrich B, Risler T, Wieder T, Lang F: Inhibition of erythrocyte cation channels by erythropoietin. *J Am Soc Nephrol* 2003;14:2750-2757.
- 36 Lang PA, Beringer O, Nicolay JP, Amon O, Kempe DS, Hermle T, Attanasio P, Akel A, Schafer R, Friedrich B, Risler T, Baur M, Olbricht CJ, Zimmerhackl LB, Zipfel PF, Wieder T, Lang F: Suicidal death of erythrocytes in recurrent hemolytic uremic syndrome. *J Mol Med* 2006;84:378-388.

- 37 Kempe DS, Akel A, Lang PA, Hermle T, Biswas R, Muresanu J, Friedrich B, Dreischer P, Wolz C, Schumacher U, Peschel A, Gotz F, Doring G, Wieder T, Gulbins E, Lang F: Suicidal erythrocyte death in sepsis. *J Mol Med* 2007;85:269-277.
- 38 Ghashghaieinia M, Bobbala D, Wieder T, Koka S, Bruck J, Fehrenbacher B, Rocken M, Schaller M, Lang F, Ghoreschi K: Targeting glutathione by dimethylfumarate protects against experimental malaria by enhancing erythrocyte cell membrane scrambling. *Am J Physiol Cell Physiol* 2010;299:C791-804.
- 39 Bobbala D, Alesutan I, Foller M, Huber SM, Lang F: Effect of anandamide in Plasmodium Berghei-infected mice. *Cell Physiol Biochem* 2010;26:355-362.
- 40 Foller M, Bobbala D, Koka S, Huber SM, Gulbins E, Lang F: Suicide for survival--death of infected erythrocytes as a host mechanism to survive malaria. *Cell Physiol Biochem* 2009;24:133-140.
- 41 Koka S, Bobbala D, Lang C, Boini KM, Huber SM, Lang F: Influence of paclitaxel on parasitemia and survival of Plasmodium berghei infected mice. *Cell Physiol Biochem* 2009;23:191-198.
- 42 Lang PA, Schenck M, Nicolay JP, Becker JU, Kempe DS, Lupescu A, Koka S, Eisele K, Klarl BA, Rubben H, Schmid KW, Mann K, Hildenbrand S, Hefter H, Huber SM, Wieder T, Erhardt A, Haussinger D, Gulbins E, Lang F: Liver cell death and anemia in Wilson disease involve acid sphingomyelinase and ceramide. *Nat Med* 2007;13:164-170.
- 43 Siraskar B, Ballal A, Bobbala D, Foller M, Lang F: Effect of amphotericin B on parasitemia and survival of plasmodium berghei-infected mice. *Cell Physiol Biochem* 2010;26:347-354.
- 44 Lang PA, Kasinathan RS, Brand VB, Duranton C, Lang C, Koka S, Shumilina E, Kempe DS, Tanneur V, Akel A, Lang KS, Foller M, Kun JF, Kremsner PG, Wesselborg S, Laufer S, Clemen CS, Herr C, Noegel AA, Wieder T, Gulbins E, Lang F, Huber SM: Accelerated clearance of Plasmodium-infected erythrocytes in sickle cell trait and annexin-A7 deficiency. *Cell Physiol Biochem* 2009;24:415-428.
- 45 Kempe DS, Lang PA, Duranton C, Akel A, Lang KS, Huber SM, Wieder T, Lang F: Enhanced programmed cell death of iron-deficient erythrocytes. *FASEB J* 2006;20:368-370.
- 46 Qadri SM, Mahmud H, Lang E, Gu S, Bobbala D, Zelenak C, Jilani K, Siegfried A, Foller M, Lang F: Enhanced suicidal erythrocyte death in mice carrying a loss-of-function mutation of the adenomatous polyposis coli gene. *J Cell Mol Med* 2012;16:1085-1093.
- 47 Birka C, Lang PA, Kempe DS, Hoefling L, Tanneur V, Duranton C, Nammi S, Henke G, Myssina S, Krikov M, Huber SM, Wieder T, Lang F: Enhanced susceptibility to erythrocyte "apoptosis" following phosphate depletion. *Pflugers Arch* 2004;448:471-477.
- 48 Zappulla D: Environmental stress, erythrocyte dysfunctions, inflammation, and the metabolic syndrome: adaptations to CO2 increases? *J Cardiometab Syndr* 2008;3:30-34.
- 49 Abed M, Towhid ST, Mia S, Pakladok T, Alesutan I, Borst O, Gawaz M, Gulbins E, Lang F: Sphingomyelinase-induced adhesion of eryptotic erythrocytes to endothelial cells. *Am J Physiol Cell Physiol* 2012;303:C991-999.
- 50 Bottger E, Multhoff G, Kun JF, Esen M: Plasmodium falciparum-infected erythrocytes induce granzyme B by NK cells through expression of host-Hsp70. *PLoS One* 2012;7:e33774.
- 51 Felder KM, Hoelzle K, Ritzmann M, Kilchling T, Schiele D, Heinritzi K, Groebel K, Hoelzle LE: Hemotropic mycoplasmas induce programmed cell death in red blood cells. *Cell Physiol Biochem* 2011;27:557-564.
- 52 Firat U, Kaya S, Cim A, Buyukbayram H, Gokalp O, Dal MS, Tamer MN: Increased caspase-3 immunoreactivity of erythrocytes in STZ diabetic rats. *Exp Diabetes Res* 2012;2012:316384.
- 53 Ganesan S, Chaurasiya ND, Sahu R, Walker LA, Tekwani BL: Understanding the mechanisms for metabolism-linked hemolytic toxicity of primaquine against glucose 6-phosphate dehydrogenase deficient human erythrocytes: evaluation of eryptotic pathway. *Toxicology* 2012;294:54-60.
- 54 Gao M, Cheung KL, Lau IP, Yu WS, Fung KP, Yu B, Loo JF, Kong SK: Polyphyllin D induces apoptosis in human erythrocytes through Ca²⁺ rise and membrane permeabilization. *Arch Toxicol* 2012;86:741-752.
- 55 Jilani K, Lupescu A, Zbidah M, Abed M, Shaik N, Lang F: Enhanced Apoptotic Death of Erythrocytes Induced by the Mycotoxin Ochratoxin A. *Kidney Blood Press Res* 2012;36:107-118.
- 56 Kucherenko YV, Lang F: Inhibitory Effect of Furosemide on Non-Selective Voltage-Independent Cation Channels in Human Erythrocytes. *Cell Physiol Biochem* 2012;30:863-875.
- 57 Lang E, Jilani K, Zelenak C, Pasham V, Bobbala D, Qadri SM, Lang F: Stimulation of suicidal erythrocyte death by benzethonium. *Cell Physiol Biochem* 2011;28:347-354.
- 58 Lang E, Qadri SM, Jilani K, Zelenak C, Lupescu A, Schleicher E, Lang F: Carbon monoxide-sensitive apoptotic death of erythrocytes. *Basic Clin Pharmacol Toxicol* 2012;111:348-355.

- 59 Polak-Jonkisz D, Purzyc L: Ca Influx versus Efflux during Eryptosis in Uremic Erythrocytes. *Blood Purif* 2012;34:209-210.
- 60 Qadri SM, Bauer J, Zelenak C, Mahmud H, Kucherenko Y, Lee SH, Ferlinz K, Lang F: Sphingosine but not sphingosine-1-phosphate stimulates suicidal erythrocyte death. *Cell Physiol Biochem* 2011;28:339-346.
- 61 Qadri SM, Kucherenko Y, Zelenak C, Jilani K, Lang E, Lang F: Dicoumarol activates Ca²⁺-permeable cation channels triggering erythrocyte cell membrane scrambling. *Cell Physiol Biochem* 2011;28:857-864.
- 62 Qian EW, Ge DT, Kong SK: Salidroside protects human erythrocytes against hydrogen peroxide-induced apoptosis. *J Nat Prod* 2012;75:531-537.
- 63 Shaik N, Zbidah M, Lang F: Inhibition of Ca²⁺ entry and suicidal erythrocyte death by naringin. *Cell Physiol Biochem* 2012;30:678-686.
- 64 Vota DM, Maltaner RE, Wenker SD, Nesse AB, Vittori DC: Differential Erythropoietin Action upon Cells Induced to Eryptosis by Different Agents. *Cell Biochem Biophys* 2013;65:145-157.
- 65 Weiss E, Cytlak UM, Rees DC, Osei A, Gibson JS: Deoxygenation-induced and Ca²⁺ dependent phosphatidylserine externalisation in red blood cells from normal individuals and sickle cell patients. *Cell Calcium* 2012;51:51-56.
- 66 Zelenak C, Pasham V, Jilani K, Tripodi PM, Rosacclerio L, Pathare G, Lupescu A, Faggio C, Qadri SM, Lang F: Tanshinone IIA stimulates erythrocyte phosphatidylserine exposure. *Cell Physiol Biochem* 2012;30:282-294.
- 67 Lang E, Qadri SM, Lang F: Killing me softly - suicidal erythrocyte death. *Int J Biochem Cell Biol* 2012;44:1236-1243.
- 68 Ghashghaieina M, Toulany M, Saki M, Bobbala D, Fehrenbacher B, Rupec R, Rodemann HP, Ghoreschi K, Rocken M, Schaller M, Lang F, Wieder T: The NFκB pathway inhibitors Bay 11-7082 and parthenolide induce programmed cell death in anucleated Erythrocytes. *Cell Physiol Biochem* 2011;27:45-54.
- 69 Shen HM, Tergaonkar V: NFκappaB signaling in carcinogenesis and as a potential molecular target for cancer therapy. *Apoptosis* 2009;14:348-363.
- 70 Turco MC, Romano MF, Petrella A, Bisogni R, Tassone P, Venuta S: NF-kappaB/Rel-mediated regulation of apoptosis in hematologic malignancies and normal hematopoietic progenitors. *Leukemia* 2004;18:11-17.
- 71 Zhang X, Jin B, Huang C: The PI3K/Akt pathway and its downstream transcriptional factors as targets for chemoprevention. *Curr Cancer Drug Targets* 2007;7:305-316.
- 72 Ghashghaieina M, Toulany M, Saki M, Rodemann HP, Mrowietz U, Lang F, Wieder T: Potential roles of the NFκappaB and glutathione pathways in mature human erythrocytes. *Cell Mol Biol Lett* 2012;17:11-20.
- 73 Bosch FH, Werre JM, Roerdinkholder-Stoelwinder B, Huls TH, Willekens FL, Halie MR: Characteristics of red blood cell populations fractionated with a combination of counterflow centrifugation and Percoll separation. *Blood* 1992;79:254-260.
- 74 Willekens FL, Werre JM, Groenen-Dopp YA, Roerdinkholder-Stoelwinder B, de Pauw B, Bosman GJ: Erythrocyte vesiculation: a self-protective mechanism? *Br J Haematol* 2008;141:549-556.
- 75 Zhang S, Ong CN, Shen HM: Critical roles of intracellular thiols and calcium in parthenolide-induced apoptosis in human colorectal cancer cells. *Cancer Lett* 2004;208:143-153.
- 76 Rauert-Wunderlich H, Siegmund D, Maier E, Giner T, Bargou RC, Wajant H, Stuhmer T: The IKK Inhibitor Bay 11-7082 Induces Cell Death Independent from Inhibition of Activation of NFκappaB Transcription Factors. *PLoS One* 2013;8:e59292.
- 77 Wen J, You KR, Lee SY, Song CH, Kim DG: Oxidative stress-mediated apoptosis. The anticancer effect of the sesquiterpene lactone parthenolide. *J Biol Chem* 2002;277:38954-38964.
- 78 Pozarowski P, Halicka DH, Darzynkiewicz Z: Cell cycle effects and caspase-dependent and independent death of HL-60 and Jurkat cells treated with the inhibitor of NF-kappaB parthenolide. *Cell Cycle* 2003;2:377-383.
- 79 Zhang S, Ong CN, Shen HM: Involvement of proapoptotic Bcl-2 family members in parthenolide-induced mitochondrial dysfunction and apoptosis. *Cancer Lett* 2004;211:175-188.
- 80 Jin X, Qiu L, Zhang D, Zhang M, Wang Z, Guo Z, Deng C, Guo C: Chemosensitization in non-small cell lung cancer cells by IKK inhibitor occurs via NF-kappaB and mitochondrial cytochrome c cascade. *J Cell Mol Med* 2009;13:4596-4607.
- 81 Kwok BH, Koh B, Ndubuisi MI, Elofsson M, Crews CM: The anti-inflammatory natural product parthenolide from the medicinal herb Feverfew directly binds to and inhibits IkappaB kinase. *Chem Biol* 2001;8:759-766.

- 82 Sinha S, Perdomo G, Brown NF, O'Doherty RM: Fatty acid-induced insulin resistance in L6 myotubes is prevented by inhibition of activation and nuclear localization of nuclear factor kappa B. *J Biol Chem* 2004;279:41294-41301.
- 83 Zhou Y, Eppenberger-Castori S, Marx C, Yau C, Scott GK, Eppenberger U, Benz CC: Activation of nuclear factor-kappaB (NFκappaB) identifies a high-risk subset of hormone-dependent breast cancers. *Int J Biochem Cell Biol* 2005;37:1130-1144.
- 84 Garcia-Pineros AJ, Castro V, Mora G, Schmidt TJ, Strunck E, Pahl HL, Merfort I: Cysteine 38 in p65/NF-kappaB plays a crucial role in DNA binding inhibition by sesquiterpene lactones. *J Biol Chem* 2001;276:39713-39720.
- 85 Garcia-Pineros AJ, Lindenmeyer MT, Merfort I: Role of cysteine residues of p65/NF-kappaB on the inhibition by the sesquiterpene lactone parthenolide and N-ethyl maleimide, and on its transactivating potential. *Life Sci* 2004;75:841-856.
- 86 Skalska J, Brookes PS, Nadtochiy SM, Hilchey SP, Jordan CT, Guzman ML, Maggirwar SB, Briehl MM, Bernstein SH: Modulation of cell surface protein free thiols: a potential novel mechanism of action of the sesquiterpene lactone parthenolide. *PLoS One* 2009;4:e8115.
- 87 Nakanishi C, Toi M: Nuclear factor-kappaB inhibitors as sensitizers to anticancer drugs. *Nat Rev Cancer* 2005;5:297-309.
- 88 Park SA, Na HK, Kim EH, Cha YN, Surh YJ: 4-hydroxyestradiol induces anchorage-independent growth of human mammary epithelial cells via activation of IkappaB kinase: potential role of reactive oxygen species. *Cancer Res* 2009;69:2416-2424.
- 89 Fernandez-Majada V, Aguilera C, Villanueva A, Vilardell F, Robert-Moreno A, Aytes A, Real FX, Capella G, Mayo MW, Espinosa L, Bigas A: Nuclear IKK activity leads to dysregulated notch-dependent gene expression in colorectal cancer. *Proc Natl Acad Sci USA* 2007;104:276-281.
- 90 Pierce JW, Schoenleber R, Jesmok G, Best J, Moore SA, Collins T, Gerritsen ME: Novel inhibitors of cytokine-induced IkappaBalpha phosphorylation and endothelial cell adhesion molecule expression show anti-inflammatory effects in vivo. *J Biol Chem* 1997;272:21096-21103.
- 91 Pahl HL, Baeuerle PA: The ER-overload response: activation of NF-kappa B. *Trends Biochem Sci* 1997;22:63-67.
- 92 Curry EA 3rd, Murry DJ, Yoder C, Fife K, Armstrong V, Nakshatri H, O'Connell M, Sweeney CJ: Phase I dose escalation trial of feverfew with standardized doses of parthenolide in patients with cancer. *Invest New Drugs* 2004;22:299-305.
- 93 Han Y, Englert JA, Delude RL, Fink MP: Ethacrynic acid inhibits multiple steps in the NF-kappaB signaling pathway. *Shock* 2005;23:45-53.
- 94 Kennedy PE, Williams FL, Abraham DJ: Design, synthesis, and testing of potential antisickling agents. 3. Ethacrynic acid. *J Med Chem* 1984;27:103-105.
- 95 Bauer A, Tronche F, Wessely O, Kellendonk C, Reichardt HM, Steinlein P, Schutz G, Beug H: The glucocorticoid receptor is required for stress erythropoiesis. *Genes Dev* 1999;13:2996-3002.
- 96 Smoak KA, Cidlowski JA: Mechanisms of glucocorticoid receptor signaling during inflammation. *Mech Ageing Dev* 2004;125:697-706.
- 97 Nakshatri H, Rice SE, Bhat-Nakshatri P: Antitumor agent parthenolide reverses resistance of breast cancer cells to tumor necrosis factor-related apoptosis-inducing ligand through sustained activation of c-Jun N-terminal kinase. *Oncogene* 2004;23:7330-7344.