A Boy with an LCR3/4-Flanked 10q22.3q23.2 Microdeletion and Uncommon Phenotypic Features

E. Petrova^a C. Neuner^a T. Haaf^a M. Schmid^a J. Wirbelauer^b A. Jurkutat^c
K. Wermke^d I. Nanda^a E. Kunstmann^a

^aInstitute for Human Genetics, ^bUniversity Children's Hospital, ^cDepartment of Special Education, Speech and Language Pathology, and ^dDepartment of Orthodontics, Center for Pre-Speech Development and Developmental Disorders, University of Wuerzburg, Wuerzburg, Germany

Key Words

BMPRI1 · Cleft palate · Language development · LCR3/4-flanked 10q22.3q23.2 deletion

Abstract

The recurrent 10q22.3q23.2 deletion with breakpoints within low copy repeats 3 and 4 is a rare genomic disorder, reported in only 13 patients to date. The phenotype is rather uncharacteristic, which makes a clinical diagnosis difficult. A phenotypic feature described in almost all patients is a delay in speech development, albeit systematic studies are still pending. In this study, we report on a boy with an LCR3/4-flanked 10q22.3q23.2 deletion exhibiting an age-appropriate language development evaluated by a standardized test at an age of 2 years and 3 months. The boy was born with a cleft palate – a feature not present in any of the patients described before. Previously reported cases are reviewed, and the role of the *BMPRI1* gene is discussed. The phenotype of patients with an LCR3/4-flanked 10q22.3q23.2 deletion can be rather variable, so counseling the families regarding the prognosis of an affected child should be done with caution. Long-term studies of affected children are needed to delineate the natural history of this rare disorder.

© 2013 S. Karger AG, Basel

Balciuniene et al. [2007] reported 2 patients with a heterozygous 10q22q23 deletion flanked by low copy repeats (LCRs), designated LCR3 and LCR4. Low copy repeats are known to be hotspots for genomic rearrangements. Since this first report, 13 further cases of such recurrent deletions with breakpoints within LCR3 and LCR4 have been published [Alliman et al., 2010; Reddy et al., 2011; Singh et al., 2011; van Bon et al., 2011]; these are summarized in table 1.

LCR3/4-flanked 10q22.3q23.2 deletions present a recurrent genomic disorder with a well-defined genotype, deleted segments ranging from 7.2 to 7.5 Mbp in size. The common phenotype of the reported patients with such deletions included facial dysmorphic features, such as hypertelorism, up- or downslanting palpebral fissures and flat nasal bridge, developmental delay of varying degrees most notable in language acquisition as well as congenital heart defects (CHD), high-arched palate and club feet in some of the patients. Some of the genes within the deleted region, e.g. *NRG3*, *GRID1*, *BMPRI1*, *GLUD1*, have been discussed as putative candidate genes associated with the phenotype, especially regarding the neuropsychological development [Balciuniene et al., 2007; van Bon et al., 2011]. In this paper, we focus on the role of *BMPRI1* (bone morphogenetic protein receptor, type

Table 1. Summary of published cases with 10q22.3q23.2 deletion

Case	Breakpoints	Deletion size	Inheritance	Gender	Age of evaluation	Language development	CHD	Club foot	Palate	Facial dysmorphism	Other clinical features
Balciuniene et al., 2007, UM10qDel-01	81.62	89,211 (89,140)	maternal	male	3 1/2 y	delayed, evaluation by use of Preschool Language Scale, ed 3				mild dysmorphic features	autism
Balciuniene et al., 2007, JHU10qDel-01	81.63	89,250 (89,140)	de novo	male	1 1/2 y	mild developmental delay, language development not explicitly mentioned				white forelock	retrocerebellar cyst, small cerebellum
Alliman et al., 2010, Case 1	81,682,644	88,931,994	de novo	male	2 7/12 y	delayed receptive language	PDA		high-arched	hypertelorism, upslanting palpebral fissures, micrognathia	autism, arachnodactyly, joint hyperextensibility
Alliman et al., 2010, Case 2	81,682,644	88,931,994	de novo	female	17 1/12 y	at age 28 mo, evaluation revealed speech impairment in articulation and verbal expression with an oral-motor component; at age 7 y 2 mo, additional evaluation showed delays in expression, and receptive and auditory language processing			high-arched	downslanting palpebral fissures, prognathism	ADHD
Alliman et al., 2010, Case 3	81,682,644	88,931,994	de novo	male	8 days	patient was too young to be assessed		bilateral		hypertelorism, mild epicanthal folds, mild micrognathia, low-set and posteriorly rotated ears	
Alliman et al., 2010, Case 4	81,682,644	88,931,994	de novo	male	1 8/12 y	evaluation at the age of 17 mo showed severe receptive and expressive delays			high-arched	downslanting palpebral fissures, posteriorly rotated ears, small mouth, frontal bossing	
van Bon et al., 2011, Patient 1	81.6	88.9–89.1	de novo	female	22 y	spoke her first words at the age of 1 y, attended special school for children with mild cognitive impairment from 6 y of age, learned to read and write	–			hypertelorism, upslanting palpebral fissures, broad nasal base, low-set ears, ptosis	
van Bon et al., 2011, Patient 2	81.6	88.7–89.1	unknown	female	2 1/2 y	spoke only 7 words at 2 y of age	ASD, VSD			hypertelorism, low-set ears, anteverted nares, flat nasal bridge, large mouth, telecanthus	the mother took carbamazepin during the pregnancy
van Bon et al., 2011, Patient 3	81.4	89.1–89.3	de novo	male	3 7/12 y	at 44 mo of age, scored at 28 mo for language development using the Denver Developmental test	–		high-arched	hypertelorism, flat nasal bridge, epicanthal folds, dolichocephaly, maligned teeth	epilepsy, Chiari I malformation
van Bon et al., 2011, Patient 4	81.2–81.6	88.6/89.1	de novo	male	12 y	at 20 mo of age, expressive language consisted of only one 2-syllable word; at the age of 12 y special education necessary	tricuspid and pulmonic regurgitation			hypertelorism, almond-shaped eyes	aggressive behavior, 47,XXY, 9 café au lait spots, pectus excavatum, kyphoscoliosis, radioulnar synostosis

Table 1 (continued)

Case	Breakpoints centromeric telomeric	Deletion size	Inheritance	Gender	Age of evaluation	Language development	CHD	Club foot	Palate	Facial dysmorphism	Other clinical features
van Bon et al., 2011, Patient 5	81.1/81.6 89.14	7.5 Mbp	maternal	male	5 y	first words at the age of 2 y; at 4 1/2 y Schlichting test for language production indicated a clear language/speech delay; at 5 y produced 3-word sentences	-	bilateral		hypertelorism, posteriorly rotated ears, flat nasal bridge, broad nasal base	shawl scrotum
Reddy et al., 2011, Case 2	81,437,039 – 81,541,288	7.7 Mbp	de novo	female	4 y	delay in the area of language acquisition	-	-		broad nose, everted lower lip vermilion border	
Singh et al., 2011, Case Report	81,634,060 89,068,181	7.46 Mbp	de novo	female	5 1/2 y	onset not delayed, lacked fluency; Griffith Mental Developmental assessment performed at 5.5 y scored language skills equivalent to 4–4.5 y	VSD	-		mild epicanthal folds	right-sided exotropia, hyperkeratosis of the 5th toe
Current Case	81,643,451 88,947,473	7.3 Mbp	de novo	male	2 y	age-appropriate development of productive and receptive language, evaluated via SETK	VSD, PFO	right- sided	median cleft palate	hypertelorism, broad nasal bridge, upslanting palpebral fissures, epicanthal folds, retrognathia, glossoptosis	

ADHD = Attention deficit-hyperactivity disorder; ASD = atrial septal defect; CHD = congenital heart disease; mo = months; PDA = patent ductus arteriosus; PFO = persistent foramen ovale; SETK = 'Sprachentwicklungstest für zweijährige Kinder' (Language development test for German children at age of 2 years); VSD = ventricular septal defect; y = years.

1A). *BMPRIA* encodes for a transmembrane serine/threonine kinase which acts as a receptor for ligands of the BMP (bone morphogenetic proteins) family of growth factors. BMP signaling plays an important role in human embryonic development, especially in the formation of bone, cartilage, neural and heart tissues [Chen et al., 2004]. Heterozygous mutations of *BMPRIA* are known to cause juvenile polyposis syndrome (JPS, OMIM 174900), and there is also some evidence that *BMPRIA* mutations may underlie CHD and craniofacial dysmorphism.

Here we report on a boy with an LCR3/4-flanked 10q22.3q23.2 deletion, who did not show any deficits in his language development. However, he is the first reported patient with a cleft palate which seriously affects his speech intelligibility. His speech is characterized by articulation errors frequently heard in the speech of individuals with a cleft palate prior to 3 years of age [Golding-Kushner, 2001].

Clinical Report

This male patient was 2 3/12 years old at the time of genetic evaluation. He was born after 40+1 weeks of gestation via spontaneous vaginal delivery after an uneventful pregnancy. Weight, height and head circumference at birth were 2,990 g, 48.5 cm and 34 cm, respectively. A right-sided club foot, a cleft palate and facial dysmorphic features, including broad nasal bridge, high forehead, upslanting palpebral fissures, epicanthal folds, and retrognathia were noticed after birth. Because of the low O₂ saturation, the boy was transferred to the intensive care unit immediately after birth. Echocardiographic evaluation showed a small, hemodynamically irrelevant ventricular septum defect (VSD) and persistent foramen ovale.

The boy showed recurrent O₂ saturation falls due to retrognathia and glossoptosis; therefore, home-monitoring was provided. Three days after birth, a special palatal plate was inserted (orthodontic plate therapy) [for details, see Kochel et al., 2011]. At the age of 1 2/12 years, surgical closure of the cleft palate was performed. The boy had surgical correction of the club foot at the age of 6 months after an unsuccessful nonsurgical treatment.

By the age of 2 years the VSD and persistent foramen ovale were no longer echocardiographically detectable. The patient showed normal motor development, but at the age of 26 months, his height was 80.5 cm (2 cm below 3rd percentile) and body weight was 10,900 g (3rd percentile). Head circumference was normal. The boy's language outcome was assessed using a standardized test for German children developed to measure both receptive and expressive language skills in 24–35-month-old children (SETK-2) [Grimm, 2000]. It consists of 4 subtests with different items assessing performances in the production and comprehension of words and sentences via direct examination of the children by trained speech and language therapists. He successfully performed all 4 subtests.

The patient's parents (mother 30 years old, father 32 years old at the child's birth) were healthy, nonconsanguineous Caucasians. A pregnancy 7 years earlier ended with spontaneous abortion after 9 weeks of gestation. The further family history was unremarkable.

Fig. 1. FISH results of the proband. **a** Simultaneous hybridization of BAC probes RP11-479O17 (green) and RP11-322M19 (red) to metaphase spread validating a 10q22.310q23.2 deletion on one homolog of chromosome 10 (arrow), as the deleted chromosome displays no hybridization signal for the probe RP11-479O17 (green) representing the deleted region, whereas the hybridization signal of flanking probe RP11-322M19 (red) is marked on both homologs. **b** Hybridization of the same BACs on an interphase nucleus showing a single hybridization signal of RP11-479O17 (green) closely associated to the flanking probe, RP11-322M19 (red). Note

the split signal of the flanking probe, RP11-322M19 (red) is due to the initiation of replication of the region. **c** GTG-bands of chromosome 10 in the proband illustrating no pronounced difference in the banding pattern between the homologs. **d** Human CytoSNP-12 array analysis in the proband shows a 7.3-Mb deletion in 10q22.310q23.2 shown in yellow. The uninterrupted deletion in the region is marked through an absence of heterozygote alleles in B allele frequency (top), a reduced intensity (below normal range) of markers in log R ratio (middle) and the CNV value plot (bottom) indicating a copy number of one.

Genetic Diagnostics

The GTG-bands of the proband's chromosomes showed a normal male karyotype 46,XY. Upon this a SNP array analysis was performed, which revealed a 7,304-kb deletion on the long arm of chromosome 10, at 10q22.3q23.2 [arr(hg19)10q22.3q23.2 (81,643,451–88,947,473)81] with breakpoints within LCR3 and LCR4 [see Balciuniene et al., 2007 for a list of the deleted genes]. The deleted chromosome 10 was verified by FISH analysis with informative BAC probes (fig. 1). More than 100 interphase nuclei and additional 23 metaphases were analyzed in the deleted region. The

signals of the BAC from the deleted region consistently showed a single signal, whereas the flanking BAC had 2 signals in each nuclei. Therefore, a clinically significant mosaicism in the patient could be ruled out. Neither of the parents was carrier of the deletion.

In a recent publication, Nowakowska et al. [2012] showed that approximately 2.1% of the apparently de novo interstitial copy number variations are the result of a submicroscopic insertion in one of the parents. To exclude such parental insertion, which, if present, would change the recurrence risk for further family members, FISH was performed. No parental insertion was detected (not shown).

Methods

Cytogenetics

Chromosomes were prepared from PHA stimulated peripheral blood lymphocytes of the index patient as well as the parents following a standard procedure, and standard karyotyping was performed based on GTG-banding at a level of approximately 450 bands.

Molecular Karyotyping

In order to identify cryptic chromosomal changes, a genome-wide SNP array was performed using the HumanCytoSNP-12v2.1 BeadChip Kit (Illumina Inc., San Diego, Calif., USA). This array contains 300,000 markers distributed with an average interSNP distance of around 10 kb. Briefly, genomic DNA was prepared from peripheral blood following the standard salt extraction method, and 200 ng of genomic DNA from the index patient and parent were hybridized to the BeadChip in an Infinium® HD Ultra Assay according to the manufacturer's protocol. After hybridization, the BeadChip was scanned with the Illumina BeadArray Reader, and the data were analyzed by examining signal intensity (log R ratio) and allelic composition (BAF) with GenomeStudio v2010.1 and cnvPartition v3.1.6 software. A minimum of a 5-probe cut-off value was used to define a copy number change. The call rates of the samples were larger than 99.0%.

FISH Analysis

BAC clones from the 10q22.3q23.2 region were selected from the Ensembl genome browser site (<http://www.ensembl.org/>) and ordered from the Children's Hospital Oakland Research Institute (<http://bacpac.chori.org/>). DNA was extracted by alkaline lysis and labeled by nick translation with Fluorescein-12-dUTP (Roche) or Tetramethyl-Rhodamine-5-dUTP (Roche). After hybridization, washing and counterstaining chromosomes were analyzed with a Zeiss AxioImager microscope. Image acquisition and analysis were performed using a CCD camera and FISHView 2.0 software (Applied Spectral Imaging). At least 20 metaphases and additional 100 interphase nuclei were evaluated per BAC probe.

Discussion

One phenotypic feature present in almost all of the published cases of LCR3/4-flanked 10q22.3q23.2 deletion that have been published is a developmental delay of varying degrees, most prominently in language and speech. However, the latter claim must be put into perspective with the fact that the majority of patients in previous studies were not evaluated by standardized tests. Most available information seems to reflect parental attitudes or occasional clinical evaluation. The current patient was examined by a standardized language test, and in contrast to other patients, he showed a normal development of receptive and expressive language at the age of 2 years. Thus, this is a rare case of a patient with an LCR3/4-flanked 10q22.3q23.2 deletion, who shows no

speech/language impairment upon objective evaluation. In view of the considerable variance in speech/language delay ranging from mild to severe in the majority of published patients (table 1), the normal development of language acquisition and production of the index patient may be the result of an even broader phenotypic expressivity of the underlying chromosomal microdeletion. Otherwise, it is possible that the test was not sensitive enough to detect a mild deficit or that speech/language delay may manifest at a later age. Therefore, further examinations with standardized language tests are planned at the age of 4 and 5 years.

Another phenotypic feature present in all patients is craniofacial dysmorphism, most commonly hypertelorism (8/14, 57%), flat/broad nasal bridge (6/14, 43%), upslanting or downslanting palpebral fissures (5/14, 36%), and high-arched palate (4/14, 29%). Other rare features are micrognathia, low-set or posteriorly-rotated ears, and epicanthal folds; the patient presented here is the first one exhibiting a median cleft palate. In a recent publication, Saito et al. [2012] presented a conditional knockout mouse model, generated by expressing a dominant negative BMPRI1A protein (dnBMPRI1A) in neural-crest-derived cells (dnBMPRI1A lacks the intracellular kinase domain and, thus, inhibits the BMPRI1A-mediated signaling pathway). The mutant mice exhibited either facial fusion defects such as a cleft face and cleft palate, or facial dysmorphology corresponding to hypertelorism and flat nasal bridge in humans. An incomplete expansion of neural-crest-derived mesenchymal cells due to extensive apoptosis was found in the mutant embryos. In addition, 50% of the mutant mice with a facial cleft also showed heart defects. The patient reported here also showed a small, VSD and persistent foramen ovale. Thus, the craniofacial dysmorphic features and the heart septal defects observed in the patients with LCR3/4-flanked 10q22.3q23.2 deletion may be partly caused by a reduction of BMPRI1A-mediated signaling.

The *BMPRI1A* gene is also associated with the JPS that is characterized by the development of hamartomatous polyps in the gastrointestinal tract. Heterozygous point mutations or partial deletions of *BMPRI1A* are found in approximately 20% of patient with JPS [Larsen et al., 2011]. Among patients with 10q22q23 microdeletions, JPS has been observed only in those patients having both genes *BMPRI1A* and *PTEN* deleted. *PTEN* is a gene associated with Cowden syndrome and Bannayan-Riley-Ruvalcaba syndrome involved in the deletion, suggesting that the contiguous deletion of both *BMPRI1A* and *PTEN* is required in order for polyposis to manifest [Delnatte et

al., 2006; Dahdaleh et al., 2012]. *PTEN* is telomeric to *BMPRIA* as well as to LCR4, so it is not involved in the LCR3/4-flanked 10q22.3q23.2 deletion; none of the patients with such deletions were reported to have JPS, supporting the supposition that the contiguous deletion of both *BMPRIA* and *PTEN* is needed for the development of the polyposis phenotype. Unfortunately, the number of described patients is very small, and in rare cases, patients with mutations in *BMPRIA* showed Bannayan-Riley-Ruvalcaba syndrome-like features [Zhou et al., 2001]. Therefore, an accurate genotype-phenotype correlation is not possible. In patients with JPS, the risk of gastrointestinal cancer is increased even though the majority of polyps are benign. Our patient did not show any JPS symptoms at the time of evaluation. Nevertheless, he was considered as a patient of risk for JPS, and baseline screening according to Larsen et al. [2011], including a complete blood count, colonoscopy and upper gastrointestinal endoscopy beginning at 15 years of age or at initial symptoms such as gastrointestinal bleeding was recommended.

Five of the 14 patients (36%), including the one presented here, had CHD, most commonly VSD. Breckpot et al. [2012] published a case report on a boy with VSD, short stature and facial dysmorphism, who had an intragenic *BMPRIA* deletion. Reviewing literature for cases with distal chromosome 10q deletions and using

computed gene prioritization, the authors showed that *BMPRIA* is the best candidate gene for CHD in patients with 10q22q23 deletions. Saito et al. [2012] also evaluated heart morphology of the mutant animals and showed that some of those with a facial cleft exhibited a ventricular septum defect as well. Thus, deletions of *BMPRIA* may contribute to various phenotypes. Detailed phenotypic characterization of patients with *BMPRIA* deletions and other mutations is warranted, to further delineate the role of *BMPRIA* in the clinical presentation of the LCR3/4-flanked 10q22.1q23.2 deletion syndrome and in human embryonic development in general.

In conclusion, the phenotype of the LCR3/4-flanked 10q22.1q23.2 deletion varies significantly, especially in speech and language development, ranging from severe delay to age-appropriate development as in the patient presented here. Thus, the prognosis of infants with an ascertained genotype regarding their language development should be given with caution. Long-term follow-up studies of affected patients are needed to further delineate the natural history of this rare disorder.

Acknowledgement

The authors thank the family described in this clinical report for participating in the study.

References

- Alliman S, Coppinger J, Marcadier J, Thiese H, Brock P, et al: Clinical and molecular characterization of individuals with recurrent genomic disorder at 10q22.3q23.2. *Clin Genet* 78:162–168 (2010).
- Balciuniene J, Feng N, Iyadurai K, Hirsch B, Charnas L, et al: Recurrent 10q22–q23 deletions: a genomic disorder on 10q associated with cognitive and behavioral abnormalities. *Am J Hum Genet* 80:938–947 (2007).
- Breckpot J, Tranchevent LC, Thienpont B, Bauters M, Troost E, et al: *BMPRIA* is a candidate gene for congenital heart defects associated with the recurrent 10q22q23 deletion syndrome. *Eur J Med Genet* 55:12–16 (2012).
- Chen D, Zhao M, Mundy G: Bone morphogenetic proteins. *Growth Factors* 22:233–241 (2004).
- Dahdaleh FS, Carr JC, Calva D, Howe JR: Juvenile polyposis and other intestinal polyposis syndromes with microdeletions of chromosome 10q22–23. *Clin Genet* 81:110–116 (2012).
- Delnatte C, Sanlaville D, Mougenot JF, Vermeesch JR, Houdayer C, et al: Contiguous gene deletion within chromosome arm 10q is associated with juvenile polyposis of infancy, reflecting cooperation between the *BMPRIA* and *PTEN* tumor-suppressor genes. *Am J Hum Genet* 78:1066–1074 (2006).
- Golding-Kushner KJ: *Therapy Techniques for Cleft Palate Speech and Related Disorders* (Singular Publishing, San Diego 2001).
- Grimm H, Aktas M, Frevert S: SETK-2. Sprachentwicklungstest für zweijährige Kinder. Diagnose rezeptiver und produktiver Sprachverarbeitungs-fähigkeiten. Hogrefe-Verlag für Psychologie, Göttingen, 2002.
- Kochel J, Meyer-Marcotty P, Wirbelauer J, Böhm H, Kochel M, et al: Treatment modalities of infants with upper airway obstruction-review of the literature and presentation of novel orthopedic appliances. *Cleft Palate Craniofac J* 48:44–55 (2011).
- Larsen Haidle J, Howe JR: Juvenile Polyposis Syndrome, in Pagon RA, Bird TD, Dolan CR, et al. (eds): *GeneReviews™* (University of Washington, Seattle 1993). <http://www.ncbi.nlm.nih.gov/books/NBK1469/>.
- Nowakowska BA, de Leeuw N, Ruivenkamp CA, Sikkema-Raddatz B, Crolla JA, et al: Parental insertional balanced translocations are an important cause of apparently de novo CNVs in patients with developmental anomalies. *Eur J Hum Genet* 20:166–170 (2012).
- Reddy KS, Mardach R, Bass H: Oligoarray (105K) CGH analysis of chromosome microdeletions within 10q22.1q24.32. *Cytogenet Genome Res* 132:113–120 (2011).
- Saito H, Yamamura KI, Suzuki N: Reduced bone morphogenetic protein receptor type 1A signaling in neural-crest-derived cells causes facial dysmorphism. *Dis Model Mech* 5:948–955 (2012).
- Singh S, Aftimos S, George A, Love DR: Interstitial deletion of 10q23.1 and confirmation of three 10qdel syndromes. *Singapore Med J* 52:e143–146 (2011).
- van Bon BW, Balciuniene J, Fruhman G, Nagamani SC, Broome DL, et al: The phenotype of recurrent 10q22q23 deletions and duplications. *Eur J Hum Genet* 19:400–408 (2011).
- Zhou XP, Woodford-Richens K, Lehtonen R, Kurose K, Aldred M, et al: Germline mutations in *BMPRIA/ALK3* cause a subset of cases of juvenile polyposis syndrome and of Cowden and Bannayan-Riley-Ruvalcaba syndromes. *Am J Hum Genet* 69:704–711 (2001).