Two Japanese Cases of Birt-Hogg-Dubé Syndrome with Pulmonary Cysts, Fibrofolliculomas, and Renal Cell Carcinomas

Yukako Murakami^a Mari Wataya-Kaneda^a Mari Tanaka^a
Aya Takahashi^a Akira Tsujimura^b Koji Inoue^c Norio Nonomura^b
Ichiro Katayama^a

Departments of ^aDermatology, ^bUrology and ^cRespiratory Medicine, Osaka University Graduate School of Medicine, Suita, Japan

Key Words

Birt-Hogg-Dubé syndrome · Fibrofolliculoma · Pulmonary cyst · Renal cell carcinoma

Abstract

Birt-Hogg-Dubé syndrome (BHD) is a rare autosomal dominant inherited disease caused by a germline mutation in the *folliculin* gene mapped in the region of chromosome 17p11.2. BHD predisposes the patient to cutaneous fibrofolliculomas (FFs), pulmonary cysts (PCs), and renal cell carcinoma (RC). Here, we present two cases of BHD in Japanese patients. One patient was a 37-year-old female, and the other a 35-year-old male. Each of the patients was affected by all three symptoms of BHD. Both patients had unremarkable FFs, asymptomatic PCs, and asymptomatic RC. The presence of RC was revealed by abdominal ultrasonic examination. We also summarized the data from 62 Asian cases of BHD from the available literature and found that their FFs were unremarkable. In addition, the proportion of BHD patients with FF is smaller for Asian patients than it is for Caucasian patients. We also found that it is rare for BHD patients in Asia to show all three symptoms of BHD. Careful inspection of the skin as well as skin biopsies are important for the early detection of BHD cases in Asia.

© 2014 S. Karger AG, Basel

Mari Wataya-Kaneda, MD, PhD
Department of Dermatology
Osaka University School of Medicine
2-2 Yamadaoka, Suita, Osaka 565-0871 (Japan)
E-mail mkaneda@derma.med.osaka-u.ac.jp

Case Presentations

Case 1

The first patient was a 37-year-old female who presented with an asymptomatic left renal tumor that was revealed by ultrasound and computed tomography (CT) examinations (fig. 1a). The CT examination also revealed the presence of asymptomatic bilateral multiple PCs (fig. 1e). Since the age of 35, a few firm papules had developed on her face and head (fig. 1g, h). The patient had no family history of pulmonary, renal, or cutaneous disease. A laparoscopic left nephrectomy was performed. On gross examination, the resected kidney was a yellowish, solid tumor with partial hemorrhage (fig. 1b). The tumor was identified as a clear cell renal carcinoma (fig. 1c) positive for vimentin stain (fig. 1d) and negative for CK-7. A skin biopsy of the papule on her head revealed it was a fibrofolliculoma (FF; fig. 1i, j) positive for factor 13a (fig. 1k) and c-kit (fig. 1l), and negative for CD34, α -smooth muscle actin (α -SMA), S100, and CD68. Because the findings for this patient met 1 major criterion and 2 minor criteria [1], she was diagnosed with Birt-Hogg-Dubé syndrome (BHD).

Case 2

The second patient was a 35-year-old male who presented with an asymptomatic right renal tumor. The presence of the tumor was revealed by ultrasound and CT examinations (fig. 2a). A laparoscopic right partial nephrectomy was performed, and the tumor was diagnosed as a chromophobe renal cell carcinoma (RC; fig. 2b). The carcinoma was negative for CD10, vimentin, and c-kit, and was partially positive for CK-7 (fig. 2c) and colloidal iron (fig. 2d). When the patient was 40 years of age, a CT examination revealed the presence of asymptomatic bilateral multiple PCs (fig. 2e). His mother, aunt, uncle, and grandfather all had histories of recurrent pneumothorax (fig. 2f). A few firm papules were also present on the patient's face (fig. 2g). The results of the skin biopsies revealed that the papules were FFs (fig. 2h, i) positive for factor 13a (fig. 2j) and c-kit (fig. 2k), and negative for CD34, α -SMA, S100, and CD68. Because the findings for this patient met 1 major criterion and 2 minor criteria [1], he was diagnosed with BHD.

Discussion

BHD is a rare autosomal dominant inherited disease that is caused by a germline mutation in the *folliculin* gene mapped in the region of chromosome 17p11.2. This gene is involved in the signaling of the mammalian target of rapamycin [1]. In cases from Asia, the mutations are located on exons 5, 6, 9, 11, 12, 13, and 14, and on intron 5 (table 1) [2, 3]. The disease is characterized by the presence of cutaneous FFs, multiple PCs, and RC.

Toro et al. [4] reported that FFs are present in 90% of the families with BHD. The skin lesions usually appear as multiple, dome-shaped, whitish papules on the face after the age of 20 years. They are hamartomas that originate from the hair follicle, and are positive for CD34, factor 13a, c-kit, and CD68, and negative for α -SMA. The FFs in both of our cases were positive for factor 13a and c-kit, and were negative for α -SMA, CD34, and CD68.

Menko et al. [1] reported that PCs are present in more than 80% of the adult patients with BHD. These PCs are most often located as basal lung lesions. The median age at first occurrence is 38 years, and the pathological findings of the lung are not so specific. In case 1, there was a layer of alveolar epithelial cells on the inner wall of the PCs.

Pavlovich et al. [5] reported that renal tumors are present in 27% of the BHD patients at a mean age of 50.4 years. Histologically, they are hybrid oncocytic tumors in 67%, chromo-

phobe RCs in 23%, and clear cell RCs in 7% of the BHD patients. Chromophobe RCs are positive for CK7, c-kit, and colloidal iron, partially positive for CD10, and negative for vimentin. Case 2 was diagnosed with chromophobe RC by hematoxylin-eosin stain, which was positive for CK7, partially positive for colloidal iron and negative for vimentin, CD10, and c-kit. Clear cell RC was positive for vimentin and partially positive for CK-7. Case 1 was diagnosed with clear cell RC by hematoxylin-eosin stain, which was positive for vimentin but negative for CK7.

More than 60 cases of BHD in Asia have been reported in the BHD literature. Kunogi et al. [2] described 30 BHD patients with pneumothorax and/or PCs and an *FCLN* mutation. Six (20%) cases had cutaneous lesions, only 1 (3%) of these was histologically diagnosed as FF, and only 1 patient (3%) had RC (table 2). Furuya and Nakatani [3] described 45 patients from 19 Asian families. Thirteen (29%) had FFs, 40 (89%) had PCs, and 9 (20%) had RCs. PCs, RCs, and FFs were present in only 2 of the patients (4%) (table 2).

We summarized 62 Asian case reports of BHD, not including those by Kunogi et al. [2] and Furuya and Nakatani [3] (table 1) [6–14]. Seventeen (27%) patients had FFs or perifollicular fibromas (PFFs) [6–10], 46 (79%) had PCs [10–13], 11 (18%) had RCs (aged 43–69 years, except for our cases; table 2) [8, 12, 14], 5 (8%) had PCs and FFs or PFFs [10], 5 (8%) had PCs and RC [12], and 2 (3%) had RCs and FFs or PFFs [8]. Only 2 cases had all three symptoms of PCs, RCs, and FFs or PFFs [3]. Therefore, our cases are only the third and fourth reported Asian BHD cases affected by all three symptoms. Compared with the other cases, our patients were younger (37 and 35 years) and had quiet FFs on the face and scalp lesions that were unremarkable. Shin et al. [10] described a Korean case with quiet FFs that appeared to be sebaceous hyperplasia. Kim et al. [12] also described a Korean BHD patient with quiet papular lesions that were 0.5–5 mm in diameter, but they were not diagnosed by biopsy.

Tsai et al. [15] reported that the Chinese scalp contains 0.72 follicular units (FUs) and the Caucasian scalp contains 1 FU/mm² surface area. The average hair density is 1.37 mm² in Chinese and 2 mm² in Caucasians. Total hair count and the number of FUs in Asians are smaller than in Caucasians. This difference may account for the few and unremarkable FFs present on the faces and scalps of Chinese, Japanese, and Korean patients.

Careful inspection of the skin and skin biopsies are important for the early detection of Asian BHD. Early detection will lead to the early diagnosis of RC and reduce BHD mortality.

Acknowledgments

This work was supported by a grant from the Japanese Ministry of Education, Culture, Sports, Science, and Technology, and by a grant from the Japanese Ministry of Health, Labor, and Welfare.

References

- 1 Menko FH, van Steensel MA, Giraud S, Friis-Hansen L, Richard S, Ungari S, Nordenskjold M, Hansen TV, Solly J, Maher ER; European BHDC: Birt-Hogg-Dubé syndrome: diagnosis and management. *Lancet Oncol* 2009;10:1199–1206.
- 2 Kunogi M, Kurihara M, Ikegami TS, Kobayashi T, Shindo N, Kumasaka T, Gunji Y, Kikkawa M, Iwakami S, Hino O, Takahashi K, Seyama K: Clinical and genetic spectrum of Birt-Hogg-Dubé syndrome patients in whom pneumothorax and/or multiple lung cysts are the presenting feature. *J Med Genet* 2010;47:281–287.
- 3 Furuya M, Nakatani Y: Birt-Hogg-Dubé syndrome: clinicopathological features of the lung. *J Clin Pathol* 2013;66:178–186.

Murakami et al.: Two Japanese Cases of Birt-Hogg-Dubé Syndrome with Pulmonary Cysts, Fibrofolliculomas, and Renal Cell Carcinomas

- 4 Toro JR, Wei MH, Glenn GM, Weinreich M, Toure O, Vocke C, Turner M, Choyke P, Merino MJ, Pinto PA, Steinberg SM, Schmidt LS, Linehan WM: BHD mutations, clinical and molecular genetic investigations of Birt-Hogg-Dubé syndrome: a new series of 50 families and a review of published reports. *J Med Genet* 2008;45:321–331.
- 5 Pavlovich CP, Grubb RL 3rd, Hurley K, Glenn GM, Toro J, Schmidt LS, Torres-Cabala C, Merino MJ, Zbar B, Choyke P, Walther MM, Linehan WM: Evaluation and management of renal tumors in the Birt-Hogg-Dubé syndrome. *J Urol* 2005;173:1482–1486.
- 6 Aoki M, Kawana S: Guess what! Diagnosis and comments: Birt-Hogg-Dubé syndrome. *Eur J Dermatol* 2000;10:407–409.
- 7 Kawasaki H, Sawamura D, Nakazawa H, Hattori N, Goto M, Sato-Matsumura KC, Akiyama M, Shimizu H: Detection of 1733insC mutations in an Asian family with Birt-Hogg-Dubé syndrome. *Br J Dermatol* 2005;152:142–145.
- 8 Imada K, Dainichi T, Yokomizo A, Tsunoda T, Song YH, Nagasaki A, Sawamura D, Nishie W, Shimizu H, Fukagawa S, Urabe K, Furue M, Hashimoto T, Naito S: Birt-Hogg-Dubé syndrome with clear-cell and oncocytic renal tumour and trichoblastoma associated with a novel FLCN mutation. *Br J Dermatol* 2009;160:1350–1353.
- 9 Park G, Kim HR, Na CH, Choi KC, Shin BS: Genetic study in a case of Birt-Hogg-Dubé syndrome. *Ann Dermatol* 2011;23:S188–S192.
- 10 Shin WW, Baek YS, Oh TS, Heo YS, Son SB, Oh CH, Song HJ: Birt-Hogg-Dubé syndrome, a rare case in Korea confirmed by genetic analysis. *Ann Dermatol* 2011;23:S193–S196.
- 11 Gunji Y, Akiyoshi T, Sato T, Kurihara M, Tominaga S, Takahashi K, Seyama K: Mutations of the Birt Hogg Dube gene in patients with multiple lung cysts and recurrent pneumothorax. *J Med Genet* 2007;44:588–593.
- 12 Kim J, Yoo JH, Kang DY, Cho NJ, Lee KA: Novel in-frame deletion mutation in FLCN gene in a Korean family with recurrent primary spontaneous pneumothorax. *Gene* 2012;499:339–342.
- 13 Nishii T, Tanabe M, Tanaka R, Matsuzawa T, Okudela K, Nozawa A, Nakatani Y, Furuya M: Unique mutation, accelerated mTOR signaling and angiogenesis in the pulmonary cysts of Birt-Hogg-Dubé syndrome. *Pathol Int* 2013;63:45–55.
- 14 Murakami T, Sano F, Huang Y, Komiya A, Baba M, Osada Y, Nagashima Y, Kondo K, Nakaigawa N, Miura T, Kubota Y, Yao M, Kishida T: Identification and characterization of Birt-Hogg-Dubé associated renal carcinoma. *J Pathol* 2007;211:524–531.
- 15 Tsai RY, Lee SH, Chan HL: The distribution of follicular units in the Chinese scalp: implications for reconstruction of natural-appearing hairlines in Orientals. *Dermatol Surg* 2002;28:500–503.
- 16 Ishiko A, Konohana I, Ikutomi M, Araki Y: A case of multiple fibrofolliculomas (Birt-Hogg-Dubé syndrome) (in Japanese). *Rinshohifuka* 1990;44:1237–1243.
- 17 Takahashi A, Hayashi T, Yoshida O, Ueda K, Furukawa F, Shuin T: Renal cell carcinoma in the Birt-Hogg-Dubé syndrome: report of a case (in Japanese). *Hinyokika Kyou* 2001;47:719–721.
- 18 Ishihara S, Kimoto M, Konohana A, Yamaguchi J, Nagasaka T, Ishiko A: Birt-Hogg-Dubé syndrome (in Japanese). *Hifubyoushinryou* 2003;25:1115–1118.
- 19 Nagashima Y, Mitsuya T, Shioi K I, Noguchi S, Kishida T, Hamano A, Ohgo Y, Tsuura Y, Ogawa T, Aoki I, Yao M: Renal oncocytosis. *Pathol Int* 2005;55:210–215.
- 20 Saito R, Sawada M, Ito H, Ishizaki S, Harada T, Nakayama S: A case of Multiple perifollicular fibroma (in Japanese). *Hifu Rinshou* 2007;49:205–209.
- 21 Sadamasa H, Ikegami N, Yasui M, Yaguchi H, Hiruma M, Ohara H, Kimura T: A case of Birt-Hogg-Dubé syndrome (in Japanese). *Skin Research* 2008;7:189–193.
- 22 Misago N, Joh K, Yatsuki H, Soejima H, Narisawa Y: A BHD germline mutation identified in an Asian family with Birt-Hogg-Dubé syndrome. *Acta Derm Venereol* 2008;88:423–425.
- 23 Kim EH, Jeong SY, Kim HJ, Kim YC: A case of Birt-Hogg-Dubé syndrome. *J Korean Med Sci* 2008;23:332–335.
- 24 Miyasaka Y, Sakuraba M, Oh S, Takamochi K, Miyamoto H, Suzuki K: A case of Birt-Hogg-Dubé syndrome with intrathoracic heterotopic endometriosis (in Japanese). *Nihon Kokyuki Gakkai Zasshi* 2009;23:641–646.
- 25 Ando K, U T, Omori T, Tajiri M, Ogura T: A case of Birt-Hogg-Dubé syndrome (in Japanese). *Nihon Kokyuki Gakkai Zasshi* 2009;23:807–811.
- 26 Ishii H, Oka H, Amemiya Y, Iwata A, Otani S, Kishi K, Shirai R, Tokimatsu I, Kawahara K, Kadota J: A Japanese family with multiple lung cysts and recurrent pneumothorax: a possibility of Birt-Hogg-Dubé syndrome. *Intern Med* 2009;48:1413–1417.
- 27 Koga S, Furuya M, Takahashi Y, Tanaka R, Yamaguchi A, Yasufuku K, Hiroshima K, Kurihara M, Yoshino I, Aoki I, Nakatani Y: Lung cysts in Birt-Hogg-Dubé syndrome: histopathological characteristics and aberrant sequence repeats. *Pathol Int* 2009;59:720–728.
- 28 So SY: Spontaneous pneumothorax due to Birt-Hogg-Dubé syndrome in a Chinese family. *Respirology* 2009;14:775–776.
- 29 Hayashi M, Takayanagi N, Ishiguro T, Sugita Y, Kawabata Y, Fukuda Y: Birt-Hogg-Dubé syndrome with multiple cysts and recurrent pneumothorax: pathological findings. *Intern Med* 2010;49:2137–2142.
- 30 Nakagawa K, Kitadate S, Saito M, Fujimoto Y, Kojima K, Oikawa T, Tsuchihara K, Iguchi M, Kou T, Osanai K, Toga H, Niida Y: Two cases of Birt-Hogg-Dubé syndrome in a family (in Japanese). *Rynsho hoshasen* 2011;56:133–137.

Murakami et al.: Two Japanese Cases of Birt-Hogg-Dubé Syndrome with Pulmonary Cysts, Fibrofolliculomas, and Renal Cell Carcinomas

- 31 Sakairi Y, Yoshino I, Okamoto T, Hoshino H, Yoshida S, Tanaka R, Koga S, Takahashi Y, Nakatani Y: Two surgical cases of pneumothorax with Birt-Hogg-Dubé syndrome; the hypothesis of cyst formation (in Japanese). *JJSPCLD* 2011;11:25–28.
- 32 Nagashima Y, Furuya M, Gotohda H, Takagi S, Hes O, Michal M, Grossmann P, Tanaka R, Nakatani Y, Kuroda N: FLCN gene-mutated renal cell neoplasms: mother and daughter cases with a novel germline mutation. *Int J Urol* 2012;19:468–470.
- 33 Ishizuka M, Miyazaki Y, Okamoto T, Komazaki Y, Tamaoka M, Seyama K, Inase N: A case of Birt-Hogg-Dubé syndrome with family history of pneumothorax and typical CT findings (in Japanese). *Ochanomizu Igaku Zasshi* 2012;60:113–117.
- 34 Kashiwada T, Shimizu H, Tamura K, Seyama K, Horie Y, Mizoo A: Birt-Hogg-Dubé syndrome and familial adenomatous polyposis: an association or a coincidence? *Intern Med* 2012;51:1789–1792.
- 35 Tobino K, Seyama K: Birt-Hogg-Dubé syndrome with renal angiomyolipoma. *Intern Med* 2012;51:1279–1280.
- 36 Ema R, Morioka S, Sakurai A, Miki Y, Tomita K, Nakamura H: A case of Birt-Hogg-Dubé syndrome with bilateral renal carcinoma and pneumothorax has a novel mutation in exon 6 of the folliculin gene (in Japanese). *Nihon Kokyuki Gakkai Zasshi* 2013;2:13–17.
- 37 Haga T, Fukuoka M, Morita M, Cho K, Kataoka H, Kurihara M: Birt-Hogg-Dubé syndrome cases of a mother and her son (in Japanese). *Nihon Kyobu Rinsyo* 2013;72(6):678–682.

Murakami et al.: Two Japanese Cases of Birt-Hogg-Dubé Syndrome with Pulmonary Cysts, Fibrofolliculomas, and Renal Cell Carcinomas

Table 1. Asian cases of BHD except Kunogi et al. [2] and Furuya and Nakatani [3]

Case	Year	First author	Age, years	Gender	Skin histology	PC	Family history	RC	Mutation
1	1990	Ishiko [16]	46	M	ff, td, ac	–	–	–	UK
2	2000	Aoki [6]	65	M	ff, td	–	+s	–	UK
3	2001	Takahashi [17]	53	F	ff	–	+ff	+	UK
4	2003	Ishihara [18]	52	F	ff, td, ac	–	+s	–	UK
5	2004	Kawasaki [7]	26	M	ff	–	+s	–	E11
6			UK	M	NT	–	fath. of 5	–	E11
7	2005	Nagashima [19]	51	F	NF	–	–	+	NF
8	2007	Murakami T [14]	55	UK	NF	–	–	+	E11
9	2007	Gunji [11]	30	F	NF	+	+p	–	E13
10			38	F	NF	+	+p	–	E11
11			40	F	NF	+	+p, r	–	E6
12			37	F	NF	+	+p, r	–	I5
13			38	F	NF	+	+p, r	–	E12
14	2007	Saito [20]	35 (Ch)	M	pff	–	–	–	UK
15	2008	Sadamasa [21]	35	M	ff, td	+	+s	–	UK
16	2008	Misago [22]	67	F	ff, td	–	+s	–	E13
17			38	F	not ff, td	+	dau. of 16	–	E13
18			35	F	not ff, td	+	dau. of 16	–	E13
19	2008	Kim [23]	31 (Ko)	F	pff, td	+	+p	–	E14
20	2009	Imada [8]	68	M	ff	–	+s, r	+	E12
21	2009	Miyasaka [24]	34	F	trunk, NT	+	+p	–	E12
22	2009	Ando [25]	58	F	ff, ac?	+	+p	–	E12
23	2009	Ishii [26]	29	F	NF	+	+p	–	UK
24			40	M	NF	+	bro. of 23	–	E6
25			72	M	NF	+	fath. of 23	–	UK
26	2009	Koga [27]	41	F	NF	+	+ff	–	E12
27			UK	M	ff	–	fath. of 26	–	E12
28	2009	So [28]	57 (Ch)	F	ff	+	+p	–	E13?
29			31	F	NF	+	dau. of 28	–	E13?
30			28	F	NF	+	dau. of 28	–	E13?
31	2010	Hayashi [29]	39	F	NF	+	–	–	E13
32	2011	Nakagawa [30]	30'	F	NF	+	+p, r	–	E12
33			30'	M	NF	+	bro. of 32	–	UK
34	2011	Sakairi [31]	41	F	NF	+	+ff	–	E12
35			55	F	NF	+	+p	–	E12
36	2011	Park [9]	43 (Ko)	M	ff, td	–	+s, p	–	E11
37	2011	Shin [10]	46 (Ko)	F	NF	+	dau. of 37	+	E5
38			69	F	NF	+	+p, r	+	E5
39	2012	Nagashima [32]	69	F	NF	+	–	+	E5
40	2012	Ishizuka [33]	45	F	NT	+	+p	–	E11
41	2012	Kim [12]	40 (Ko)	F	not ff	+	+p, r	–	E6
42			UK	M	UK	+	fath. of 41	+	UK
43			UK	F	NF	+	sis. of 41	–	E6
44			UK	F	NF	+	sis. of 41	–	E6
45	2012	Kashiwada [34]	60	F	NT?	+	+s, p	–	E11

Murakami et al.: Two Japanese Cases of Birt-Hogg-Dubé Syndrome with Pulmonary Cysts, Fibrofolliculomas, and Renal Cell Carcinomas

46			32	M	NT?	+	son of 39	–	E11?
47	2012	Tobino [35]	43	F	NF	+	+p	+*	E12
48			UK	M	NF	+	son of 47	–	UK
49	2013	Ema [36]	53	F	not ff	+	–	+	E6
50	2013	Nishii [13]	33	F	not ff	+	+s, p	–	I5
51			UK	M	extrm., NT	+	fath. of 50	–	I5?
52			UK	M	UK	+	uncle of 50	–	UK
53			UK	F	UK	+	G.M. of 50	–	UK
54	2013	Haga [37]	61	F	NF	+	+p	–	E11
55			29	M	NF	+	son of 54	–	E11
56			UK	M	NF	+	bro. of 54	–	UK
57	This report Murakami Y		37	F	ff	+	NF	+	UK
58			35	M	ff	+	+p	+	UK
59			UK	F	UK	+	moth. of 58	–	UK
60			UK	F	UK	+	aunt of 58	–	UK
61			UK	M	UK	+	uncle of 58	–	UK
62			UK	M	UK	+	G.F. of 58	–	UK

Ch = Chinese; Ko = Korean; UK = unknown; ff = fibrofolliculoma; td = trichodyscoma; ac = acrochordon; NF = no cutaneous change could be found; NT = not tested; extrm. = extremity; s = skin lesion; p = pulmonary lesion; r = renal lesion; dau. = daughter; sis. = sister; bro. = brother; fath. = father; G.M. = grandmother; moth. = mother; G.F. = grandfather; E = exon; I = intron.

Diagnosed as renal cyst by abdominal sonography without pathological examination.

* Diagnosed as angiomyolipoma by CT images without pathological examination.

Table 2. Clinical findings of Asian cases compared with those in the USA or Europe

	Toro [4], Menko [1], Pavlovich [5]	Kunogi [2]	Furuya [3]	Murakami Y, this report
Fibrofolliculoma	90%	20%*	29%	27%
Pulmonary cyst	80%	100%	89%	79%
Renal cell carcinoma	27%	3%	20%	18%

* Only 3% of the patients had a pathological diagnosis.

Murakami et al.: Two Japanese Cases of Birt-Hogg-Dubé Syndrome with Pulmonary Cysts, Fibrofolliculomas, and Renal Cell Carcinomas

Fig. 1. The abdominal CT revealed a low-density area on the left kidney of a 37-year-old female with BHD (a), which was histopathologically diagnosed as clear cell carcinoma. On gross examination, the resected kidney contained a partially hemorrhagic, yellowish, solid tumor (b). The tumor was composed of cells with clear to acidophilic cytoplasm (hematoxylin-eosin stain; c), which were positive for vimentin stain (d; 200× magnification). The thoracic CT revealed bilateral, multiple PCs in the basilar and mediastinal regions (e). Microscopic findings for the lung revealed that the septum of the cyst wall contained capillaries and was lined by pneumocytes on both surfaces (f; 200× magnification). Quiet, skin-colored papules were present on the patient's face (g). There were few pinkish-colored, verrucous papules on the scalp (h). Histopathological images (hematoxylin-eosin) of the FF. The stoma was rich in fibroblasts and was oriented in parallel bundles of root sheath-like fibers and neighboring enlarged hair follicles [20× magnification (i) and 200× magnification (j)] positive for factor 13a (k) and c-kit (l; 400× magnification).

Murakami et al.: Two Japanese Cases of Birt-Hogg-Dubé Syndrome with Pulmonary Cysts, Fibrofolliculomas, and Renal Cell Carcinomas

Fig. 2. The abdominal CT revealed a low-density area on the right kidney of a 35-year-old male with BHD (a). The tumor was composed of cells with eosinophilic cytoplasm (hematoxylin-eosin; b), was partially positive for CK7 (c; 200× magnification) and colloidal iron (d; 400× magnification), and histopathologically diagnosed as chromophobe cell carcinoma. A thoracic CT revealed the presence of bilateral, basally located, multiple PCs (e). Pedigrees of the families affected with recurrent pneumothorax (f). Skin-colored papules were present on the nose (g). Histopathological images (hematoxylin-eosin) of the FFs [40× magnification (h) and 200× magnification (i)] positive for factor 13a (j) and c-kit (k; 400× magnification).