Malignant Melanoma with Concurrent BRAF E586K and NRAS Q81K Mutations

Rodney Shackelford^a Maressa Pollen^a Moise Vora^a Tamara T. Jusion^b
James Cotelingam^a Binu Nair^c

^aDepartment of Pathology, ^bClinical Pathology Molecular Pathology Consultative Services, and ^cHematology/Oncology, LSU Health Shreveport, Shreveport, La., USA

Key Words

Cutaneous melanoma · Concurrent BRAF E586K and NRAS Q81K mutations · Skin cancer

Abstract

Cutaneous melanoma is an aggressive malignant tumor of melanocytes which accounts for 80% of skin cancer-related deaths. A number of driver mutations have been identified in melanoma, with the mutually exclusive BRAF V600E and NRAS Q61A mutations together accounting for roughly 70% of mutations. Simultaneous BRAF V600E and NRAS Q61A mutations in melanoma are rare, with evidence suggesting that up to 2.9% (2/69) of primary cutaneous melanomas carry both mutations. Here we describe a 42-year-old man with concurrent BRAF E586K and NRAS Q81K driver mutations. To our knowledge, this is the first description of these driver mutations occurring simultaneously in primary cutaneous melanoma.

© 2014 S. Karger AG, Basel

Introduction

Primary cutaneous melanoma is an aggressive malignancy of melanocytes which accounts for only 4% of dermatologic malignancies, but is responsible for 80% of skin cancer-related deaths and, with metastatic disease, has a 5-year survival rate of only 14% [1]. The risk factors for melanoma are immunosuppression, ultraviolet radiation exposure, a family history of melanoma, and multiple benign or atypical nevi [2]. Histologically, melanomas are characterized by different growth patterns, such as superficial spreading, nodular, acral lentiginous, and lentigo maligna melanomas [3]. Many different melanoma driver mutations have been identified, with approximately 50–60% being mutated *BRAF* and 15–20%

Rodney E. Shackelford
Department of Pathology, LSU Health Shreveport
1501 Kings Highway
Shreveport, LA 71130 (USA)
E-Mail rdnyshac@aol.com

mutated *NRAS*. Interestingly, the V600E and Q16R mutations account for roughly 90% of *BRAF* and *NRAS* mutations, respectively [4, 5]. Less common driver mutations include other *BRAF* and *NRAS* mutations as well as mutations in other genes such as *c-Kit* (2–6%), *CTNNB1* (2–3%), *GNA11* (2%), *GNAQ* (1%), and *MEK1* (2–6%) [4–6]. In general, melanomas with concurrent driver mutations are rare; Goel et al. [6] examined 69 cases of primary cutaneous melanoma and found that only 2.9% (2/69) carried concurrent *BRAF V600E* and *NRAS Q16R* mutations.

The activating *BRAF E586K* mutation, causing constitutive enzyme heterodimerization and activation, is rare in melanomas [7, 8]. Similarly, the *NRAS Q81K* mutation accounts for a small percentage of melanoma *NRAS* driver mutations [5]. Here we describe the first reported case of concurrent *BRAF E586K* and *NRAS Q16R* driver mutations in the presence of primary cutaneous melanoma.

Case Report

A 42-year-old male Caucasian with a history of chronic sun exposure presented with a mole on his left shoulder that had recently become red, itchy, and inflamed. The lesion was excised and HE sections were prepared. Histologic examination revealed sheets of pleomorphic epithelial cells and a histology consistent with cutaneous melanoma (fig. 1a). Immunohistochemical (IHC) S100 staining showed diffuse membranous and cytoplasmic staining in most malignant cells (fig. 1b). Dual Mart-1 and Ki-67 IHC staining showed strong membranous Mart-1 staining (red stain) and strong Ki-67 nuclear staining (brown stain). Dual S100 and Mart-1 IHC positivity is typical of melanomas (fig. 1b, c) [2]. The nuclear Ki-67 positivity showed a brisk mitotic index of approximately 20% (fig. 1c). IHC staining was performed at Reliopath (Opelousas, La., USA) according to regulations of the College of American Pathologists.

Several weeks later, a PET/CT scan of the patient revealed an abnormal fluorodeoxyglucose uptake by soft tissue nodules in the anterolateral chest wall which were suggestive of metastatic disease. Additionally, the resection of a jejunal mass and 4 local lymph nodes showed a metastatic melanoma. Next-generation sequencing was done on the metastatic jejunal melanoma to analyze the tumor for possible druggable driver mutations. A sequencing analysis of the *BRAF* exons 11 and 15 and the *NRAS* exons 2 and 3 revealed concurrent *BRAF E586K* and *NRAS Q16R* driver mutations. The analysis of the *c-Kit* exons 9, 11, 13, 17, and 18 revealed no mutations.

Discussion

NRAS mutations in melanoma commonly compromise the ability of the enzyme to hydrolyze guanosine triphosphate, allowing the enzyme to be constitutively active and activate downstream signaling proteins such as *RAS* and *BRAF* [9]. *RAS* and *BRAF* activities contribute to the malignant proliferation by inducing the activities of proteins including mTOR, cyclin D1, and NF- κ B [9]. Additionally, *RAS* also activates the phosphatidylinositol 3-kinase/PTEN/Akt pathway, leading to apoptosis inhibition and malignant cell survival [9]. Concurrent *NRAS* and *BRAF* activating driver mutations are not expected to be common events, as they lie on the same molecular signal transduction pathway, and once one activating driver mutation occurs there would be little selective pressure for a second mutation to occur [6, 9].

To our knowledge, concurrent *BRAF E586K* and *NRAS Q81K* driver mutations have not been previously identified in melanomas. The clinical treatment of this rare combination of concurrent driver mutations in this case may be difficult, as it is unknown whether the *BRAF E586K* mutation responds to *BRAF* inhibitors. Additionally, there is in vitro evidence that *NRAS* mutations may confer resistance to *BRAF* inhibitors [5, 10].

References

- 1 Cancer Facts and Figures, 2003. Atlanta, American Cancer Society, 2003.
- 2 Miller AJ, Mihm MC Jr: Melanoma. *N Engl J Med* 2006;355:51–65.
- 3 Smoller BR: Histologic criteria for diagnosing primary cutaneous malignant melanoma. *Mod Pathol* 2006;19(suppl 2):S34–S40.
- 4 My Cancer Genome: molecular profiling of melanoma.
<http://www.mycancergenome.org/content/disease/melanoma>.
- 5 Kunz M: Oncogenes in melanoma: an update. *Eur J Cell Biol* 2014;93:1–10.
- 6 Goel VK, Lazar AJ, Warneke CL, Redston MS, Haluska FG: Examination of mutations in BRAF, NRAS, and PTEN in primary cutaneous melanoma. *J Invest Dermatol* 2006;126:154–160.
- 7 Wan PT, Garnett MJ, Roe SM, Lee S, Niculescu-Duvaz D, Good VM, Jones CM, Marshall CJ, Springer CJ, Barford D, Marais R; Cancer Genome Project: Mechanism of activation of the RAF-ERK signaling pathway by oncogenic mutations of B-RAF. *Cell* 2004;116:855–867.
- 8 Rajakulendran T, Sahmi M, Lefrançois M, Sicheri F, Therrien M: A dimerization-dependent mechanism drives RAF catalytic activation. *Nature* 2009;461:542–545.
- 9 Sullivan RJ, Flaherty K: MAP kinase signaling and inhibition in melanoma. *Oncogene* 2013;32:2373–2379.
- 10 Nazarian R, Shi H, Wang Q, Kong X, Koya RC, Lee H, Chen Z, Lee MK, Attar N, Sazegar H, Chodon T, Nelson SF, McArthur G, Sosman JA, Ribas A, Lo RS: Melanomas acquire resistance to B-RAF(V600E) inhibition by RTK or N-RAS upregulation. *Nature* 2010;468:973–977.

Fig. 1. Representative HE and IHC staining results of the cutaneous melanoma removed from the patient's back. **a** High-power staining of the melanoma showing epithelioid pleomorphic malignant cells. **b** IHC staining showing S100 positivity typical of melanomas. **c** IHC staining showing dual Mart-1 (membranous) and Ki-67 (nuclear) positivity, also consistent with melanomas.