Treatment of Dry Age-Related Macular Degeneration

Giuseppe Querques^{a, b} Philip J. Rosenfeld^c Edoardo Cavallero^b
Enrico Borrelli^b Federico Corvi^b Lea Querques^b Francesco M. Bandello^b
Marco A. Zarbin^d

^aDepartment of Ophthalmology, University of Paris Est Créteil, Centre Hospitalier Intercommunal de Créteil, Créteil, France; ^bDepartment of Ophthalmology, University Scientific Institute San Raffaele, Milan, Italy; ^cBascom Palmer Eye Institute, University of Miami Miller School of Medicine, Miami, Fla., and ^dInstitute of Ophthalmology and Visual Science, Rutgers-New Jersey Medical School, Newark, N.J., USA

Key Words

Drusen · Geographic atrophy · Age-related macular degeneration · Treatment · Pathway · Drugs · Stem cells · Neurotrophic factors · Visual cycle · Oxidative damage

Abstract

A number of different approaches are under development for treating nonexudative manifestations of age-related macular degeneration (AMD). Some interventions target specific pathways that are believed to play a role in AMD pathogenesis, e.g. oxidative damage, lipofuscin accumulation, chronic inflammation (including complement activation), extracellular matrix changes (e.g. β -amyloid accumulation), impaired choroidal blood flow, and apoptosis. In principle, these therapies can be combined ('combination therapy'), which may lead to synergistic effects that include better visual outcome, less likelihood for 'escape' (i.e. drug resistance), and less frequent treatment.

© 2014 S. Karger AG, Basel

Introduction

A number of treatments for nonexudative age-related macular degeneration (AMD) are under study. Several processes have been implicated in the pathogenesis of AMD, including oxidative damage, accumulation of toxic visual cycle products, chronic inflammation (including complement activation), and choroidal vascular insufficiency [1]. In this chapter we discuss interventions that target these processes to prevent the progression from early- to late-stage nonexudative AMD.

Antioxidants

The Age-Related Eye Disease Study (AREDS, NCT00594672) showed that the use of a daily dose of 80 mg zinc oxide, 2 mg cupric oxide, 15 mg β -carotene, 500 mg vitamin C, and 400 IU vitamin E in patients with extensive intermediate drusen, at least one large ($\geq 125 \mu\text{m}$) druse, noncentral geographic atrophy (GA) in 1 or both eyes, choroidal neovascularization in 1 eye, or visual acu-

ity $\leq 20/40$ in 1 eye due to AMD reduces the risk of moderate visual loss by 19% (during a 5-year follow-up period) [2]. Because the use of β -carotene increases the risk of lung cancer in smokers (or among persons who have stopped smoking within the previous 8 years) [3, 4], patients who smoke should not use this formulation unless they have stopped smoking at least 8 years previously. In general, it may be wise to ask the patient's internist to approve the use of these supplements due to these and other concerns (e.g. use of high-dose zinc in patients with a history of prostate cancer or copper in patients with a history of dementia). The AREDS did not show a statistically significant benefit of the formulation for either the development of new GA or for involvement of the fovea in eyes with preexisting GA. In part, this result may be due to the paucity of patients with GA who were enrolled in the study.

Cysteine is an important antioxidant involved in the regulation of apoptosis and immune function. Moriarty-Craige et al. [5] conducted an ancillary study in which a subset of AREDS subjects at two sites were studied at two time points, an average of 1.7 and 6.7 years after enrollment, to determine whether antioxidant supplements alter the plasma glutathione and/or cysteine redox potential in AMD patients. The AREDS antioxidant supplements reduced oxidation of cysteine to cystine but had no effect on glutathione. The authors suggested that because cysteine is important for cell growth, apoptosis, and immune function, the beneficial effect of antioxidant supplementation on progression to advanced AMD may be explained, in part, by its effect on the ratio of cysteine/cystine and/or its effect on cysteine availability. Brantley et al. [6] carried out a study to determine whether short-term AREDS antioxidant and zinc supplementation affects biomarkers of oxidative stress, possibly serving as a predictor of their efficacy. Short-term AREDS supplementation significantly lowered mean plasma levels of cystine in participants on a regulated diet. The authors concluded that a 5-day course of antioxidant and zinc supplements can modify plasma levels of cystine, suggesting that this oxidative stress biomarker could help predict how likely an individual is to benefit from the AREDS supplementation.

AREDS2 (NCT00345176), a randomized, double-masked, multicenter phase III clinical trial, was designed to determine whether adding lutein and zeaxanthin, docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA), or both to the AREDS formulation decreases the risk of developing advanced AMD and to evaluate the effect of eliminating β -carotene, lowering the zinc dose (to

25 mg), or both in the AREDS formulation [7]. Participants were randomized to receive lutein (10 mg) + zeaxanthin (2 mg), DHA (350 mg) + EPA (650 mg), lutein + zeaxanthin, and DHA + EPA, or placebo. All participants were also asked to either take the original AREDS formulation or accept a secondary randomization to four variations of the AREDS formulation, including elimination of β -carotene, lowering of zinc dose, or both [7]. A comparison with placebo in the primary analyses demonstrated no statistically significant reduction in the progression to advanced AMD: lutein + zeaxanthin (hazard ratio, HR, 0.90; 98.7% CI, 0.76–1.07; $p = 0.12$), DHA + EPA (HR, 0.97; 98.7% CI, 0.82–1.16; $p = 0.70$), and lutein + zeaxanthin and DHA + EPA (HR, 0.89; 98.7% CI, 0.75–1.06; $p = 0.10$). There was no apparent effect of β -carotene elimination or lower-dose zinc on the progression to advanced AMD [7]. More lung cancers were noted in the β -carotene versus the no β -carotene group: 23 (2.0%) versus 11 (0.9%; nominal $p = 0.04$), mostly in former smokers [7]. The authors concluded that the addition of lutein + zeaxanthin, DHA + EPA, or both to the AREDS formulation in primary analyses did not further reduce the risk of progression to advanced AMD. However, because of the potential of increased incidence of lung cancer in former smokers, lutein + zeaxanthin could be an appropriate carotenoid substitute in the AREDS formulation [7]. In an exploratory analysis of lutein/zeaxanthin versus no lutein/zeaxanthin [8], the HR of the development of late AMD was 0.90 (95% CI, 0.82–0.99; $p = 0.04$). An exploratory analyses of a direct comparison of lutein/zeaxanthin with β -carotene showed HR of 0.82 (95% CI, 0.69–0.96; $p = 0.02$) for the development of late AMD, 0.78 (95% CI, 0.64–0.94; $p = 0.01$) for the development of neovascular AMD and 0.94 (95% CI, 0.70–1.26; $p = 0.67$) for the development of central GA [8]. In analyses restricted to eyes with bilateral large drusen at baseline, the direct comparison of lutein/zeaxanthin with β -carotene showed HR of 0.76 (95% CI, 0.61–0.96; $p = 0.02$) for the progression to late AMD, 0.65 (95% CI, 0.49–0.85; $p = 0.002$) for neovascular AMD and 0.98 (95% CI, 0.69–1.39; $p = 0.91$) for central GA [8]. Thus, replacing β -carotene with lutein and zeaxanthin may not only be safer but might also provide greater protection against the development of late AMD, specifically neovascular complications and not GA.

Carotenoids, especially lutein and zeaxanthin, comprise the macular pigment. The primary direct antioxidant function of carotenoids is to scavenge singlet oxygen, but they also quench the triplet state of photosensitizers and delay the peroxidation of membrane

phospholipids [9, 10]. Factors associated with increased risk for AMD and increased risk for low macular pigment density include age, cigarette smoking, female gender, light iris color, and increasing lens density [11–14], but not all clinical studies confirm the association between low macular pigment density and increased risk for AMD [15]. Two postmortem studies revealed decreased retinal lutein and zeaxanthin in AMD eyes versus controls [16, 17]. Increasing age and advanced AMD in the fellow eye have been associated with a relative absence of macular pigment [18]. The Intervention Trial in Early Age-related Macular Degeneration (I-TEAM; NCT01694680) is an ongoing, randomized, placebo-controlled multicenter study aimed to assess the effect of 1 year of daily consumption of a lutein-enriched egg beverage on the maintenance of visual function in subjects with early signs of AMD (i.e. AREDS categories 2 or 3).

Visual Cycle Inhibitors

Excessive lipofuscin accumulation in the retinal pigment epithelium (RPE) may play an important role in AMD pathogenesis [19]. In RPE cells, the main source of lipofuscin is probably partially degraded components of phagocytized outer segments [20]. Components of lipofuscin arise from the visual cycle.

The visual cycle involves the following sequence of events. In vertebrate photoreceptors, light causes isomerization of 11-*cis*-retinylidene to all-*trans*-retinylidene followed by the release of all-*trans*-retinal from the opsin binding pocket and its reduction to all-*trans*-retinol [21]. ABCA4, an ATP-binding cassette transporter present in the outer segment of rods and cones, transports N-retinylidene-phosphatidylethanolamine from the outer segment discs to the photoreceptor cytoplasm [22, 23]. Retinol dehydrogenases in the inner and outer segments reduce all-*trans*-retinal to all-*trans*-retinol [24, 25]. Vitamin A (all-*trans*-retinol) diffuses to RPE where it is esterified by lecithin/retinol acyltransferase to all-*trans*-retinyl esters and is stored in retinosomes [26, 27]. All-*trans*-retinyl esters are isomerized to 11-*cis*-retinol in a reaction involving RPE65 [28–30]. Next, 11-*cis*-retinol is reduced to 11-*cis*-retinal [31, 32] which then diffuses across the extracellular space to photoreceptors and recombines with rod and cone opsin proteins to regenerate visual pigments.

Within the outer segment discs, ethanolamine can combine with two retinaldehyde molecules to form N-retinylidene-N-retinylethanolamine (A2E); A2E is a ma-

ior fluorophore in RPE lipofuscin [33]. Inhibition of the visual cycle, by blocking all-*trans*-retinol RPE uptake or by blocking the formation of 11-*cis*-retinal, should reduce the production of lipofuscin and A2E [34, 35].

Retinol-binding protein (RBP) has a high-affinity binding site for all-*trans*-retinol. Retinol binding to RBP creates a high-affinity binding site for transthyretin. N-(4-hydroxyphenyl) retinamide (fenretinide, Sirion Therapeutics, Tampa, Fla., USA) displaces all-*trans*-retinol from RBP and prevents the interaction of RBP with transthyretin. In the absence of transthyretin binding, the RBP-fenretinide complex is excreted in urine (due to its relatively small size). Thus, fenretinide treatment causes a dose-dependent, reversible reduction in circulating RBP and retinol. Due to the unique requirement of the eye for retinol delivered by RBP, during chronic fenretinide administration levels of retinol within the eye will be reduced while other extrahepatic tissues obtain retinol from alternative sources. Fenretinide reduces lipofuscin and A2E accumulation in the RPE of ABCA4^{-/-} mice and causes modest delays in dark adaptation [34]. However, data from preclinical models indicate that other mechanisms may be involved in all-*trans*-retinol delivery to the eye, and restriction of dietary vitamin A may be important in achieving a reduction of ocular all-*trans*-retinol levels even when using fenretinide [36, 37].

The efficacy of fenretinide (100 and 300 mg daily, orally) to slow lesion growth in GA patients was examined in a 2-year, placebo-controlled, double-masked trial (NCT00429936) [38]. Fenretinide treatment produced dose-dependent reversible reductions in serum RBP-retinol that were associated with trends in reduced GA growth rates. Patients in the 300-mg group who achieved serum retinol levels of $\leq 1 \mu\text{M}$ ($\leq 2 \text{ mg/dl}$ RBP), for example, showed a reduction in the yearly lesion growth rate compared with subjects in the placebo group (1.70 vs. 2.03 mm/year, respectively; $p = 0.1848$). Fenretinide treatment also reduced, in a nondose-dependent manner, the incidence of choroidal neovascularization (about 45% reduction in incidence rate in the combined fenretinide groups vs. placebo; $p = 0.0606$) [38].

Isotretinoin or 13-*cis*-retinoic acid (Accutane; Hoffmann-La Roche, Inc., Nutley, N.J., USA) is used for the treatment of acne. This visual cycle modulator is associated with a high incidence of nyctalopia because it inhibits the conversion of all-*trans*-retinyl esters in retinosomes to 11-*cis*-retinol, as well as the conversion of 11-*cis*-retinol to 11-*cis*-retinal by retinol dehydrogenase. As expected, isotretinoin inhibits lipofuscin formation in ABCA4^{-/-} mice [39, 40].

Emixustat (ACU-4429; Acucela, Seattle, Wash., USA) is a nonretinoid that inhibits the conversion of all-*trans*-retinyl ester to 11-*cis*-retinol via the blockade of RPE-specific 65-kDa protein (RPE65) or another protein needed for isomerization of all-*trans*-retinol [41]. ACU-4429 also reduces RPE lipofuscin and A2E accumulation in mice and reduces retinal neovascularization in a model of retinopathy of prematurity [42]. A phase I study (NCT00942240) demonstrated that oral administration of ACU-4429 produced a dose-dependent inhibition of the electroretinogram b-wave, was well tolerated up to 75 mg, and demonstrated linear pharmacokinetics across doses [43]. Adverse events were dose dependent, mild and visual in nature (dyschromatopsia and alteration in dark adaptation), transient, and resolved within a few days after cessation of therapy [43]. A phase II study (NCT01002950) to evaluate the safety, tolerability, pharmacokinetics, and pharmacodynamics of ACU-4429 in subjects with GA has been completed, but no results have been published as yet. A phase IIb/III (NCT01802866) multicenter, randomized, double-masked study to determine whether ACU-4429 reduces the rate of progression of GA compared with placebo in subjects with dry AMD is in progress.

Anti-Inflammatory Agents

AMD is associated with chronic inflammation in the region of the RPE, Bruch's membrane, and the choroid [44–46]. Corticosteroids have antiangiogenic and anti-inflammatory effects. Iluvien (Alimera Sciences, Alpharetta, Ga., USA) is an injectable, nonerodible, intravitreal implant containing 190 µg of the corticosteroid fluocinolone acetonide. A phase II study (NCT00695318) was recently completed after the recruitment of 40 patients with bilateral GA who were randomized to a high (0.5 µg/day) or low (0.2 µg/day) dose of Iluvien, with the fellow eye used as a control. The primary outcome of the study was the rate of GA enlargement in treated versus untreated eyes. The results of the study are unpublished at this time.

Drusen and GA are associated with mutations in components of the complement pathway, which is part of the innate immune system [45, 47]. Mutations in components of the complement pathway modulate the risk of developing AMD and include the following loci: complement factor H (CFH), complement component 2 (C2), complement factor B (CFB), complement component 3 (C3), and complement factor I (CFI) [48–61]. The alter-

native complement pathway is continuously activated in the fluid phase, and tissue surfaces require continuous complement inhibition to prevent spontaneous autologous cell injury [62, 63]. Oxidative stress can compromise the regulation of the complement system by RPE cells [64–66].

C3 inhibition might be effective at blocking complement activation that arises from many different mutations in the complement cascade (thus targeting a relatively large population of AMD patients). This degree of complement inhibition, however, may create risks such as an increased risk of injection-associated endophthalmitis [67–69]. POT-4 (Potentia Pharmaceuticals, Louisville, Ky., USA, and Alcon, Hünenberg, Switzerland) is a C3 inhibitor that binds C3 and prevents its proteolysis to C3a and C3b. It is administered by intravitreal injection, and gel-like deposits form in the vitreous when POT-4 is injected at high concentrations, thus providing a sustained-release delivery system that lasts for approximately 6 months. A phase I study of POT-4 in dry AMD eyes was completed successfully without safety concerns (NCT00473928).

With the inhibition of C5, terminal complement activity is blocked, but proximal complement functions (e.g. C3a anaphylatoxin production, C3b opsonization, immune complex and apoptotic body clearance) remain intact. ARC1905 (Ophthotech Corp., Princeton, N.J., USA) is an anti-C5 aptamer administered by intravitreal injection. A phase I clinical trial (NCT00950638) in patients with nonexudative (as well as exudative) complications of AMD has been completed, but the results are unpublished. Eculizumab (Soliris; Alexon Pharmaceuticals, Cheshire, Conn., USA) is a humanized monoclonal antibody that blocks C5, is administered intravenously and is approved by the US Food and Drug Administration (FDA) for the treatment of paroxysmal nocturnal hemoglobinuria. A phase II placebo-controlled study (NCT00935883) to assess the effect on the rate of GA progression and change in drusen volume showed that intravenous eculizumab was well tolerated in AMD patients but did not decrease the growth rate of GA significantly [70].

LFG316 (Novartis Pharmaceuticals, Whippany, N.J., USA) is an antibody directed against C5 and is administered intravitreally. The phase I study (NCT01255462) to assess the safety and tolerability of intravitreal LFG316 in patients with advanced AMD, GA or choroidal new vessels was completed, and a phase II study to evaluate the efficacy of 6 successive monthly doses on the growth of GA is underway (NCT01527500). Factor D is the rate-

limiting enzyme in the activation of the alternative complement pathway. Genetic polymorphisms as well as hyperactivity of the alternative complement pathway have been implicated in the development of AMD, including GA [71]. Factor D inhibitors, e.g. BCX1470 (Alcon, Fort Worth, Tex., USA) and lampalizumab (FCFD4514S; Genentech/Roche, San Francisco, Calif., USA) block the activation of the alternative complement pathway while preserving the activation of the classical and lectin pathways, thus possibly reducing the risk of infectious complications associated with treatment. Lampalizumab is a monoclonal F_{ab} fragment directed against factor D. A phase II clinical trial (NCT01602120, MAHALO) involving 143 patients with bilateral GA assessed the effects of monthly versus bimonthly intravitreal lampalizumab injections of 5 mg, 10 mg, or sham on the change in the GA area during an 18-month period, using fundus autofluorescence images to identify areas of GA. Preliminary results indicate a 20.4% reduction in the rate of GA enlargement at the 18-month follow-up ($p < 0.1179$, statistically significant per prespecified protocol criteria) with efficacy observed beginning at month 6 [72]. In GA patients with CFI mutations treated monthly with lampalizumab, the GA progression rate was decreased by 44% ($p < 0.005$) at 18 months. In this study, 57% of the patients were positive for CFI mutations. If these results are confirmed in a phase III trial, they may mean that complement inhibition is useful even at a late stage of AMD. In addition, patients will require genetic testing to assess their likelihood to respond to the therapy.

Sirolimus (rapamycin; MacuSight/Santen, Union City, Calif., USA) is a macrolide fungicide that blocks the mammalian target of rapamycin, a protein kinase that regulates proliferation, motility, survival, and protein synthesis. Sirolimus has anti-inflammatory, antiangiogenic and antifibrotic effects. A single-center, open-label phase II trial (NCT00766649) enrolled 11 participants with bilateral GA; 8 participants completed 24 months of follow-up [73]. Sirolimus (440 µg) was administered every 3 months as a subconjunctival injection in only 1 randomly assigned eye/participant for 24 months. Fellow eyes served as untreated controls. The primary efficacy outcome measure was the change in the total GA area at 24 months. At month 24, the mean GA area increased by 54.5 and 39.7% in the study and fellow eyes, respectively ($p = 0.41$), whereas mean visual acuity decreased by 21.0 and 3.0 letters in the study and fellow eyes, respectively ($p = 0.03$). Substantial differences in mean change in the drusen area, central retinal thickness and macular sensitivity were not detected for all analysis time points up to 24 months.

Glatiramer acetate (Copaxone; TEVA, Petah Tikva, Israel) induces glatiramer acetate-specific suppressor T cells and downregulates inflammatory cytokines. It can be administered subcutaneously and has been studied in a phase II/III clinical trial in patients with drusen (NCT00466076). A small, controlled study demonstrated efficacy after 12 weeks of subcutaneous injections: 4 patients were treated with weekly subcutaneous injections of Copaxone (20 mg), and all 8 study eyes showed a decrease in the total drusen area [74].

Drusen vesicles contain fibrillar amyloid composed, in part, of amyloid- β [75, 76]. Amyloid- β induces the production of interleukin-1 β and tumor necrosis factor- α by macrophages and microglia, which can cause an increased expression of CFB in RPE and may contribute to AMD progression [77]. RN6G (PF-4382923; Pfizer, New York, N.Y., USA) is a humanized monoclonal antibody that targets the C-termini of amyloid- β_{40} and amyloid- β_{42} . Treatment with intravenous RN6G is intended to prevent the accumulation and cytotoxic effects of amyloid- β_{40} and amyloid- β_{42} by removing amyloid from the tissue compartment through mass action kinetics. Beneficial effects were documented in a mouse model of AMD [78]. Phase I and II clinical trials have been completed successfully to evaluate the safety of the treatment in patients with advanced dry AMD (NCT01003691, NCT01577381). We are not aware of plans to carry out phase III studies of RN6G. GlaxoSmithKline has successfully completed a phase I study of intravenous GSK933776, an intravenously administered humanized monoclonal antibody directed against amyloid- β . This antibody reduces the accumulation of amyloid- β and C3a deposition in Bruch's membrane of *cfh*^{-/-} mice [79]. A phase II study (NCT01342926) is underway.

Drugs That Increase Choroidal Blood Flow

Choroidal blood flow decreases with age, probably due to a decrease in choriocapillaris diameter and density [80]. In addition, decreased choroidal blood flow is correlated positively with fundus findings associated with an increased risk of choroidal neovascularization (i.e. drusen, pigmentary changes) [81]. The administration of vasodilators may improve choroidal blood flow, possibly delaying the progression of nonexudative as well as exudative manifestations of late AMD [82]. A pilot study (NCT01013376) investigating the safety of MC-1101, an eye drop developed to increase choroidal blood flow by the generation of nitric oxide to induce vasodilation, has

demonstrated that it is safe and well tolerated in humans [83]. Increased choroidal blood volume and velocity were demonstrated in treated AMD eyes. MC-1101 is also anti-inflammatory and an antioxidant. The active ingredient of MC-1101 is already used as an approved oral antihypertensive agent. A randomized, double-masked phase II/III trial (NCT01601483) investigating patients with mild-to-moderate nonexudative AMD is ongoing to evaluate the efficacy and safety (visual function is the primary outcome after 2 years) of topical 1% MC-1101 administered 3 times a day.

Neuroprotective Therapy

Apoptosis, programmed cell death, has an important homeostatic function; however, apoptosis has been implicated in the pathogenesis of AMD [84]. Neuroprotective drugs are intended to preserve macular function by preventing apoptosis of viable RPE cells and photoreceptors.

Ciliary neurotrophic factor (CNTF), a member of the IL-6 family of neurotrophic cytokines, delays the loss of photoreceptors in animal models of retina degenerative disease [85]. CNTF can be delivered using encapsulated cell technology (ECT) [86]. Using ECT, genetically modified RPE that overexpresses CNTF is housed in a semi-permeable capsule with small pores that permit CNTF to escape into the vitreous cavity and protect the allogeneic RPE cells from immune rejection. The ECT device is implanted in the vitreous cavity and comprises a sustained CNTF delivery system.

Zhang et al. [87] reported the results of a 1-year, phase II randomized, double-masked, controlled dose-ranging study involving the use of CNTF-ECT (NT-501 implant; Neurotech, Lincoln, R.I., USA) to treat GA. A total of 51 individuals were randomly assigned to receive high-dose ($n = 27$) or low-dose ($n = 12$) implants or sham surgery ($n = 12$). At the 12-month follow-up, 96.3% of patients in the high-dose NT-501-treated cohort lost <15 letters (ETDRS) versus 75% of patients in the sham group ($p = 0.078$). No increase in vision occurred, and no serious adverse events were reported. The trend in visual stabilization at 12 months was preceded (at 4 months) by a dose-dependent, statistically significant increase in retinal thickness ($p < 0.001$ and $p = 0.013$ for the high- and low-dose cohorts, respectively) by optical coherence tomography. CNTF-induced increased retinal thickness has been reported in laboratory animals with retinitis pigmentosa-like conditions [88, 89]. In mice, this thickness

change reflects, in part, increased photoreceptor nuclear size and increased amounts of euchromatin, and in *rd*-1 dogs it reflects increased photoreceptor nuclear size as well as swelling of photoreceptors and/or Müller cell processes, with expansion of the outer limiting membrane towards the RPE.

Brimonidine is an α -2 adrenergic receptor agonist that is approved by the US FDA for the treatment of glaucoma. Brimonidine protects retinal ganglion cells, bipolar cells, and photoreceptors from degeneration in several models of experimental injury, including retinal ischemia, partial optic nerve crush, ocular hypertension, and retinal phototoxicity [90–92]. The proposed mechanisms of neuroprotection include the following: increased expression of basic fibroblast growth factor, a cytokine that delays apoptosis; upregulation of endogenous production of trophic factors such as brain-derived neurotrophic factor in retinal ganglion cells; stabilization of mitochondrial transmembrane potential under conditions of oxidative stress, and suppression of the accumulation of glutamate causing neuronal cell death [92, 93]. Brimonidine is delivered by a sustained-release biodegradable implant (Allergan, Irvine, Calif., USA) that is injected into the vitreous through a 22-gauge needle with the same applicator system used to deliver dexamethasone (Ozurdex; Allergan). A randomized, double-masked, dose-response, sham-controlled phase II study of the safety and efficacy of brimonidine intravitreal implants in patients with GA has been completed (NCT00658619). In this 2-year follow-up study, 119 patients with bilateral GA and visual acuity between 20/40 and 20/320 were randomized to receive the brimonidine delivery system containing either 200 or 400 μ g of drug or sham treatment. The patients received the drug or sham treatment in 1 eye and sham treatment in the fellow eye on day 1 and a repeat of the assigned treatment and sham treatment at month 6. The primary outcome measure was the change of GA lesion area in the study eye from baseline to month 12. The results of this trial showed equivocal results. A second study is planned.

Stem Cell Therapy

Stem cell-derived RPE and photoreceptors have been shown to rescue the retina, replace lost retinal neurons, and restore vision in various animal models of retinal degenerative disease (reviewed in Zarbin [94]). Stem cells also provide a unique opportunity to create *in vitro* models of retinal diseases such as AMD [95]. The pathology

and pathophysiology of AMD indicate that RPE transplants may prevent disease progression through the replacement of dead and dying RPE [96–98]. Because of their unlimited proliferative potential, stem cells may be an ideal source of tissue from which large numbers of therapeutic cells may be derived for transplantation purposes [99]. Induced pluripotent stem cell-derived human RPE has been shown to rescue the retina in animal models of retinal degeneration involving RPE mutations that preclude normal phagocytic function [100]. In addition, stem cell-derived RPE produces neurotrophic factors that should support photoreceptor survival [101]. Schwartz et al. [102] reported the results of embryonic stem cell-derived RPE transplantation in patients with AMD and Stargardt disease. This report is preliminary and involves a small number of patients with a short follow-up, but promising visual results were reported in 1 patient with Stargardt disease whose vision improved from hand motions to 20/800 in the treated eye during the first 3 months after subretinal transplantation of the stem cell-derived RPE. It is not clear if the basis of the improved vision was due to the replacement of native RPE or simply to a rescue

effect mediated by neurotrophic factor production from the transplanted cells.

Other stem cell studies are in early phases of development. A phase I/II study of subretinal transplantation of human central nervous system stem cells (NCT01632527) in eyes with GA, for example, is currently underway and designed to evaluate the safety and efficacy of this treatment. At this time it is clear that, with certain precautions, stem cells can be produced en masse safely and that they can be induced to differentiate into ocular cells with the potential for replacement and rescue therapy. Challenges to successful stem cell therapy for degenerative retinal disease include the following: the abnormal microenvironment of the diseased eye, which can compromise cell survival and differentiation [103]; synaptic rewiring that accompanies retinal degeneration and which may compromise replacement strategies [104]; maintenance of functional phenotype over time; immune rejection, particularly for embryonic stem cell-derived RPE [105], and transplantation of ‘healthy’ cells, particularly for induced pluripotent stem cell-derived RPE from AMD donors.

References

- Zarbin MA, Rosenfeld PJ: Pathway-based therapies for age-related macular degeneration: an integrated survey of emerging treatment alternatives. *Retina* 2010;30:1350–1367.
- Age-Related Eye Disease Study Research Group: A randomized, placebo-controlled, clinical trial of high-dose supplementation with vitamins C and E, beta carotene, and zinc for age-related macular degeneration and vision loss: AREDS report No 8. *Arch Ophthalmol* 2001;119:1417–1436.
- Omenn GS, Goodman GE, Thornquist MD, et al: Risk factors for lung cancer and for intervention effects in CARET, the Beta-Carotene and Retinol Efficacy Trial. *J Nat Cancer Inst* 1996;88:1550–1559.
- The effect of vitamin E and beta carotene on the incidence of lung cancer and other cancers in male smokers. The Alpha-Tocopherol, Beta Carotene Cancer Prevention Study Group. *N Engl J Med* 1994;330:1029–1035.
- Moriarty-Craige SE, Adkison J, Lynn M, et al: Antioxidant supplements prevent oxidation of cysteine/cystine redox in patients with age-related macular degeneration. *Am J Ophthalmol* 2005;140:1020–1026.
- Brantley MA Jr, Osborn MP, Sanders BJ, et al: The short-term effects of antioxidant and zinc supplements on oxidative stress biomarker levels in plasma: a pilot investigation. *Am J Ophthalmol* 2012;153:1104–1109.e2.
- Age-Related Eye Disease Study 2 Research Group: Lutein + zeaxanthin and omega-3 fatty acids for age-related macular degeneration: the Age-Related Eye Disease Study 2 (AREDS2) randomized clinical trial. *JAMA* 2013;309:2005–2015.
- Age-Related Eye Disease Study 2 Research Group; Chew EY, Clemons TE, et al: Secondary analyses of the effects of lutein/zeaxanthin on age-related macular degeneration progression: AREDS2 report No 3. *JAMA Ophthalmol* 2014;132:142–149.
- Packer L: Antioxidant action of carotenoids in vitro and in vivo and protection against oxidation of human low-density lipoproteins. *Ann NY Acad Sci* 1993;691:48–60.
- Bohm F, Tinkler JH, Truscott TG: Carotenoids protect against cell membrane damage by the nitrogen dioxide radical. *Nat Med* 1995;1:98–99.
- Hammond BR Jr, Curran-Celentano J, Judd S, et al: Sex differences in macular pigment optical density: relation to plasma carotenoid concentrations and dietary patterns. *Vision Res* 1996;36:2001–2012.
- Hammond BR Jr, Wooten BR, Snodderly DM: Cigarette smoking and retinal carotenoids: implications for age-related macular degeneration. *Vision Res* 1996;36:3003–3009.
- Hammond BR Jr, Fuld K, Snodderly DM: Iris color and macular pigment optical density. *Exp Eye Res* 1996;62:293–297.
- Hammond BR Jr, Wooten BR, Snodderly DM: Density of the human crystalline lens is related to the macular pigment carotenoids, lutein and zeaxanthin. *Optom Vis Sci* 1997;74:499–504.
- Berendschot TT, Willemse-Assink JJ, Bastiaanse M, et al: Macular pigment and melanin in age-related maculopathy in a general population. *Invest Ophthalmol Vis Sci* 2002;43:1928–1932.
- Landrum JT, Bone RA, Kilburn MD: The macular pigment: a possible role in protection from age-related macular degeneration. *Adv Pharmacol* 1997;38:537–556.
- Bone RA, Landrum JT, Mayne ST, et al: Macular pigment in donor eyes with and without AMD: a case-control study. *Invest Ophthalmol Vis Sci* 2001;42:235–240.
- Beatty S, Murray IJ, Henson DB, et al: Macular pigment and risk for age-related macular degeneration in subjects from a Northern European population. *Invest Ophthalmol Vis Sci* 2001;42:439–446.
- Holz FG, Bellman C, Staudt S, et al: Fundus autofluorescence and development of geographic atrophy in age-related macular degeneration. *Invest Ophthalmol Vis Sci* 2001;42:1051–1056.
- Kennedy CJ, Rakoczy PE, Constable IJ: Lipofuscin of the retinal pigment epithelium: a review. *Eye* 1995;9:763–771.

- 21 Lamb TD, Pugh EN Jr: Dark adaptation and the retinoid cycle of vision. *Prog Retin Eye Res* 2004;23:307–380.
- 22 Weng J, Mata NL, Azarian SM, et al: Insights into the function of Rim protein in photoreceptors and etiology of Stargardt's disease from the phenotype in abcr knockout mice. *Cell* 1999;98:13–23.
- 23 Beharry S, Zhong M, Molday RS: N-retinylidene-phosphatidylethanolamine is the preferred retinoid substrate for the photoreceptor-specific ABC transporter ABCA4 (ABCR). *J Biol Chem* 2004;279:53972–53979.
- 24 Rattner A, Smallwood PM, Nathans J: Identification and characterization of all-*trans*-retinol dehydrogenase from photoreceptor outer segments, the visual cycle enzyme that reduces all-*trans*-retinal to all-*trans*-retinol. *J Biol Chem* 2000;275:11034–11043.
- 25 Maeda A, Maeda T, Imanishi Y, et al: Retinol dehydrogenase (RDH12) protects photoreceptors from light-induced degeneration in mice. *J Biol Chem* 2006;281:37697–37704.
- 26 Imanishi Y, Batten ML, Piston DW, et al: Noninvasive two-photon imaging reveals retinyl ester storage structures in the eye. *J Cell Biol* 2004;164:373–383.
- 27 Imanishi Y, Gerke V, Palczewski K: Retinoids: new insights into intracellular managing of hydrophobic substances in lipid bodies. *J Cell Biol* 2004;166:447–453.
- 28 Redmond TM, Poliakov E, Yu S, et al: Mutation of key residues of RPE65 abolishes its enzymatic role as isomerohydrolase in the visual cycle. *Proc Natl Acad Sci USA* 2005;102:13658–13663.
- 29 Jin M, Li S, Moghrabi WN, et al: RPE65 is the retinoid isomerase in bovine retinal pigment epithelium. *Cell* 2005;122:449–459.
- 30 Moiseyev G, Chen Y, Takahashi Y, et al: RPE65 is the isomerohydrolase in the retinoid visual cycle. *Proc Natl Acad Sci USA* 2005;102:12413–12418.
- 31 Jang GF, Van Hooser JP, Kuksa V, et al: Characterization of a dehydrogenase activity responsible for oxidation of 11-*cis*-retinol in the retinal pigment epithelium of mice with a disrupted RDH5 gene. A model for the human hereditary disease fundus albipunctatus. *J Biol Chem* 2001;276:32456–465.
- 32 Farjo KM, Moiseyev G, Takahashi Y, et al: The 11-*cis*-retinol dehydrogenase activity of RDH10 and its interaction with visual cycle proteins. *Invest Ophthalmol Vis Sci* 2009;50:5089–5097.
- 33 Sparrow JR, Fishkin N, Zhou J, et al: A2E, a byproduct of the visual cycle. *Vision Res* 2003;43:2983–2990.
- 34 Radu RA, Han Y, Bui TV, et al: Reductions in serum vitamin A arrest accumulation of toxic retinal fluorophores: a potential therapy for treatment of lipofuscin-based retinal diseases. *Invest Ophthalmol Vis Sci* 2005;46:4393–4401.
- 35 Kuksa V, Imanishi Y, Batten M, et al: Retinoid cycle in the vertebrate retina: experimental approaches and mechanisms of isomerization. *Vision Res* 2003;43:2959–2981.
- 36 Quadro L, Blamer WS, Salchow DJ, et al: Impaired retinal function and vitamin A availability in mice lacking retinol-binding protein. *EMBO J* 1999;18:4633–4644.
- 37 Vogel S, Piantadosi R, O'Byrne SM, et al: Retinol-binding protein-deficient mice: biochemical basis for impaired vision. *Biochemistry* 2002;41:15360–15368.
- 38 Mata NL, Lichter JB, Vogel R, et al: Investigation of oral fenretinide for treatment of geographic atrophy in age-related macular degeneration. *Retina* 2013;33:498–507.
- 39 Sieving PA, Chaudhry P, Kondo M, et al: Inhibition of the visual cycle in vivo by 13-*cis* retinoic acid protects from light damage and provides a mechanism for night blindness in isotretinoin therapy. *Proc Natl Acad Sci USA* 2001;98:1835–1840.
- 40 Radu RA, Mata NL, Nusinowitz S, et al: Treatment with isotretinoin inhibits lipofuscin accumulation in a mouse model of recessive Stargardt's macular degeneration. *Proc Natl Acad Sci USA* 2003;100:4742–4747.
- 41 Maeda A, Maeda T, Golczak M, et al: Effects of potent inhibitors of the retinoid cycle on visual function and photoreceptor protection from light damage in mice. *Mol Pharmacol* 2006;70:1220–1229.
- 42 Akula JD, Hansen RM, Tzekov R, et al: Visual cycle modulation in neurovascular retinopathy. *Exp Eye Res* 2010;91:153–161.
- 43 Kubota R, Boman NL, David R, et al: Safety and effect on rod function of ACU-4429, a novel small-molecule visual cycle modulator. *Retina* 2012;32:183–188.
- 44 Anderson DH, Mullins RF, Hageman GS, et al: A role for local inflammation in the formation of drusen in the aging eye. *Am J Ophthalmol* 2002;134:411–431.
- 45 Gehrs KM, Jackson JR, Brown EN, et al: Complement, age-related macular degeneration and a vision of the future. *Arch Ophthalmol* 2010;128:349–358.
- 46 Cherepanoff S, McMenamin P, Gillies MC, et al: Bruch's membrane and choroidal macrophages in early and advanced age-related macular degeneration. *Br J Ophthalmol* 2010;94:918–925.
- 47 Johnson LV, Leitner WP, Staples MK, et al: Complement activation and inflammatory processes in Drusen formation and age related macular degeneration. *Exp Eye Res* 2001;73:887–896.
- 48 Hageman GS, Anderson DH, Johnson LV, et al: A common haplotype in the complement regulatory gene factor H (HF1/CFH) predisposes individuals to age-related macular degeneration. *Proc Natl Acad Sci USA* 2005;102:7227–7232.
- 49 Klein RJ, Zeiss C, Chew EY, et al: Complement factor H polymorphism in age-related macular degeneration. *Science* 2005;308:385–389.
- 50 Edwards AO, Ritter R 3rd, Abel KJ, et al: Complement factor H polymorphism and age-related macular degeneration. *Science* 2005;308:421–424.
- 51 Haines JL, Hauser MA, Schmidt S, et al: Complement factor H variant increases the risk of age-related macular degeneration. *Science* 2005;308:419–421.
- 52 Hughes AE, Orr N, Esfandiary H, et al: A common CFH haplotype, with deletion of CFHR1 and CFHR3, is associated with lower risk of age-related macular degeneration. *Nat Genet* 2006;38:1173–1177.
- 53 Skerka C, Lauer N, Weinberger AA, et al: Defective complement control of factor H (Y402H) and FHL-1 in age-related macular degeneration. *Mol Immunol* 2007;44:3398–3406.
- 54 Magnusson KP, Duan S, Sigurdsson H, et al: CFH Y402H confers similar risk of soft drusen and both forms of advanced AMD. *PLoS Med* 2006;3:e5.
- 55 Li M, Atmaca-Sonmez P, Othman M, et al: CFH haplotypes without the Y402H coding variant show strong association with susceptibility to age-related macular degeneration. *Nat Genet* 2006;38:1049–1054.
- 56 Gold B, Merriam JE, Zernant J, et al: Variation in factor B (BF) and complement component 2 (C2) genes is associated with age-related macular degeneration. *Nat Genet* 2006;38:458–462.
- 57 Spencer KL, Hauser MA, Olson LM, et al: Protective effect of complement factor B and complement component 2 variants in age-related macular degeneration. *Hum Mol Genet* 2007;16:1986–1992.
- 58 Yates JR, Sepp T, Matharu BK, et al: Complement C3 variant and the risk of age-related macular degeneration. *N Engl J Med* 2007;357:553–561.
- 59 Ennis S, Jomary C, Mullins R, et al: Association between the SERPING1 gene and age-related macular degeneration: a two-stage case-control study. *Lancet* 2008;372:1828–1834.
- 60 Maller JB, Fagerness JA, Reynolds RC, et al: Variation in complement factor 3 is associated with risk of age-related macular degeneration. *Nat Genet* 2007;39:1200–1201.
- 61 Fagerness JA, Maller JB, Neale BM, et al: Variation near complement factor I is associated with risk of advanced AMD. *Eur J Hum Genet* 2009;17:100–104.
- 62 Thurman JM, Holers VM: The central role of the alternative complement pathway in human disease. *J Immunol* 2006;176:1305–1310.
- 63 Sohn JH, Kaplan HJ, Suk HJ, et al: Chronic low level complement activation within the eye is controlled by intraocular complement regulatory proteins. *Invest Ophthalmol Vis Sci* 2000;41:3492–3502.
- 64 Thurman JM, Renner B, Kunchithapatham K, et al: Oxidative stress renders retinal pigment epithelial cells susceptible to complement-mediated injury. *J Biol Chem* 2009;284:16939–16947.

- 65 Wu Z, Lauer TW, Sick A, et al: Oxidative stress modulates complement factor H expression in retinal pigmented epithelial cells by acetylation of FOXO3. *J Biol Chem* 2007; 282:22414–22425.
- 66 Zhou J, Jang YP, Kim SR, et al: Complement activation by photooxidation products of A2E, a lipofuscin constituent of the retinal pigment epithelium. *Proc Natl Acad Sci USA* 2006;103:16182–16187.
- 67 Engelbert M, Gilmore MS: Fas ligand but not complement is critical for control of experimental *Staphylococcus aureus* endophthalmitis. *Invest Ophthalmol Vis Sci* 2005;46:2479–2486.
- 68 Giese MJ, Mondino BJ, Glasgow BJ, et al: Complement system and host defense against staphylococcal endophthalmitis. *Invest Ophthalmol Vis Sci* 1994;35:1026–1032.
- 69 Aizuss DH, Mondino BJ, Sumner HL, et al: The complement system and host defense against *Pseudomonas* endophthalmitis. *Invest Ophthalmol Vis Sci* 1985;26:1262–1266.
- 70 Yehoshua Z, Alexandre de Amorim Garcia Filho C, Nunes RP, et al: Systemic complement inhibition with eculizumab for geographic atrophy in age-related macular degeneration: the COMPLETE study. *Ophthalmology* 2014;121:693–701.
- 71 Seddon JM, Yu Y, Miller EC, et al: Rare variants in CFI, C3 and C9 are associated with high risk of advanced age-related macular degeneration. *Nat Genet* 2013;45:1366–1370.
- 72 Regillo CD: Lampalizumab (anti-factor D) in patients with geography atrophy: the MAHALO phase II results. Presented at the 2013 Ann Meet Am Acad Ophthalmol, New Orleans, November 2013.
- 73 Wong WT, Dresner S, Forooghian F, et al: Treatment of geographic atrophy with subconjunctival sirolimus: results of a phase I/II clinical trial. *Invest Ophthalmol Vis Sci* 2013; 54:2941–2950.
- 74 Landa G, Butovsky O, Shoshani J, et al: Weekly vaccination with Copaxone (glatiramer acetate) as a potential therapy for dry age-related macular degeneration. *Curr Eye Res* 2008; 33:1011–1013.
- 75 Johnson LV, Leitner WP, Rivest AJ, et al: The Alzheimer's A β -peptide is deposited at sites of complement activation in pathologic deposits associated with aging and age-related macular degeneration. *Proc Natl Acad Sci USA* 2002; 99:11830–11835.
- 76 Isas JM, Luibl V, Johnson LV, et al: Soluble and mature amyloid fibrils in drusen deposits. *Invest Ophthalmol Vis Sci* 2010;51:1304–1310.
- 77 Wang J, Ohno-Matsui K, Yoshida T, et al: Amyloid- β up-regulates complement factor B in retinal pigment epithelial cells through cytokines released from recruited macrophages/microglia: another mechanism of complement activation in age-related macular degeneration. *J Cell Physiol* 2009;220:119–128.
- 78 Ding JD, Johnson LV, Herrmann R, et al: Anti-amyloid therapy protects against retinal pigmented epithelium damage and vision loss in a model of age-related macular degeneration. *Proc Natl Acad Sci USA* 2011;108:E279–E287.
- 79 Catchpole I, Germaschewski V, Hoh Kam J, et al: Systemic administration of A-beta mAb reduces retinal deposition of A-beta and activated complement C3 in age-related macular degeneration mouse model. *PLoS One* 2013; 8:e65518.
- 80 Grunwald JE, Hariprasad SM, DuPont J: Effect of aging on foveolar choroidal circulation. *Arch Ophthalmol* 1998;116:150–154.
- 81 Grunwald JE, Metelitsina TI, Dupont JC, et al: Reduced foveolar choroidal blood flow in eyes with increasing AMD severity. *Invest Ophthalmol Vis Sci* 2005;46:1033–1008.
- 82 Boltz A, Luksch A, Wimpfing B, et al: Choroidal blood flow and progression of age-related macular degeneration in the fellow eye in patients with unilateral choroidal neovascularization. *Invest Ophthalmol Vis Sci* 2010; 51:4220–4225.
- 83 Ralston PG, Sloan D, Waters-Honcu D, Saigal S, Torkildsen G: A pilot, open-label study of the safety of MC-1101 in both normal volunteers and patients with early nonexudative age-related macular degeneration. *Invest Ophthalmol Vis Sci* 2010;51:E-Abstract 913.
- 84 Dunaief JL, Dentichev T, Ying GS, et al: The role of apoptosis in age-related macular degeneration. *Arch Ophthalmol* 2002;120: 1435–1442.
- 85 Li Y, Tao W, Luo L, et al: CNTF induces regeneration of cone outer segments in a rat model of retinal degeneration. *PLoS One* 2010;5:e9495.
- 86 Tao W, Wen R, Goddard MB, et al: Encapsulated cell-based delivery of CNTF reduces photoreceptor degeneration in animal models of retinitis pigmentosa. *Invest Ophthalmol Vis Sci* 2002;43:3292–3298.
- 87 Zhang K, Hopkins JJ, Heier JS, et al: Ciliary neurotrophic factor delivered by encapsulated cell intraocular implants for treatment of geographic atrophy in age-related macular degeneration. *Proc Natl Acad Sci USA* 2011; 108:6241–6245.
- 88 Bok D, Yasumura D, Matthes MT, et al: Effects of adeno-associated virus-vectored ciliary neurotrophic factor on retinal structure and function in mice with a P216L rds/peripherin mutation. *Exp Eye Res* 2002;74:719–735.
- 89 Zeiss CJ, Allore HG, Towle V, et al: CNTF induces dose-dependent alterations in retinal morphology in normal and rcd-1 canine retina. *Exp Eye Res* 2006;82:395–404.
- 90 Lai RK, Chun T, Hasson D, et al: Alpha-2 adrenoceptor agonist protects retinal function after acute retinal ischemic injury in the rat. *Vis Neurosci* 2002;19:175–185.
- 91 WoldeMussie E, Ruiz G, Wijono M, et al: Neuroprotection of retinal ganglion cells by brimonidine in rats with laser-induced chronic ocular hypertension. *Invest Ophthalmol Vis Sci* 2001;42:2849–2855.
- 92 Wheeler L, WoldeMussie E, Lai R: Role of alpha-2 agonists in neuroprotection. *Surv Ophthalmol* 2003;48(suppl 1):S47–S51.
- 93 Tatton W, Chen D, Chalmers-Redman R, et al: Hypothesis for a common basis for neuroprotection in glaucoma and Alzheimer's disease: anti-apoptosis by alpha-2-adrenergic receptor activation. *Surv Ophthalmol* 2003;48(suppl 1):S25–S37.
- 94 Zarbin M: The promise of stem cells for age-related macular degeneration and other retinal degenerative diseases. *Drug Discov Today Ther Strateg* 2012.
- 95 Melville H, Carpinello M, Hollis K, et al: Stem cells: a new paradigm for disease modeling and developing therapies for age-related macular degeneration. *J Transl Med* 2013; 11:53.
- 96 Sarks JP, Sarks SH, Killingsworth MC: Evolution of geographic atrophy of the retinal pigment epithelium. *Eye (Lond)* 1988;2: 552–577.
- 97 Zhu D, Wu J, Spee C, et al: BMP4 mediates oxidative stress-induced retinal pigment epithelial cell senescence and is overexpressed in age-related macular degeneration. *J Biol Chem* 2009;284:9529–9539.
- 98 Shen JK, Dong A, Hackett SF, et al: Oxidative damage in age-related macular degeneration. *Histol Histopathol* 2007;22:1301–1308.
- 99 Casaroli-Marano RP, Zarbin MA: Cell-Based Therapy for Retinal Degenerative Disease. Basel, Karger, 2014.
- 100 Carr AJ, Vugler AA, Hikita ST, et al: Protective effects of human iPS-derived retinal pigment epithelium cell transplantation in the retinal dystrophic rat. *PLoS One* 2009; 4:e8152.
- 101 Sugino IK, Sun Q, Wang J, et al: Comparison of FRPE and human embryonic stem cell-derived RPE behavior on aged human Bruch's membrane. *Invest Ophthalmol Vis Sci* 2011;52:4979–4997.
- 102 Schwartz SD, Hubschman JP, Heilwell G, et al: Embryonic stem cell trials for macular degeneration: a preliminary report. *Lancet* 2012;379:713–720.
- 103 Sugino IK, Rapista A, Sun Q, et al: A method to enhance cell survival on Bruch's membrane in eyes affected by age and age-related macular degeneration. *Invest Ophthalmol Vis Sci* 2011;52:9598–9609.
- 104 Jones BW, Watt CB, Frederick JM, et al: Retinal remodeling triggered by photoreceptor degenerations. *J Comp Neurol* 2003;464:1–16.
- 105 Fairchild PJ: The challenge of immunogenicity in the quest for induced pluripotency. *Nat Rev Immunol* 2010;10:868–875.