Temporal Gene Expression in the Hippocampus and Peripheral Organs to Endotoxin-Induced Systemic Inflammatory Response in Caspase-1-Deficient Mice

Claudio Alberto Mastronardi^a Gilberto Paz-Filho^a Martina Zanoni^c
Nicolas Molano-González^d Mauricio Arcos-Burgos^a Julio Licinio^b Ma-Li Wong^b

^aGenomics and Predictive Medicine Group, Genome Biology Department, John Curtin School of Medical Research, Australian National University, Canberra, A.C.T., and ^bMind and Brain Theme, South Australian Health and Medical Research Institute and Flinders, University of South Australia, Adelaide, S.A., Australia; ^cDepartment of Psychiatry, University of Verona, Verona, Italy; ^dCenter for Autoimmune Diseases Research (CREA), School of Medicine and Health Sciences, Universidad del Rosario, Bogotá, Colombia

Key Words

Inflammation · Lipopolysaccharide · Transcription · *Nos2* · *Cox2* · Chemokines · *Cxcl1* · *Cxcl10* · *Adamts1* · Coexpression

Abstract

Objectives: Caspase-1 (casp1), a key protease involved in the systemic inflammatory response syndrome (SIRS), controls the brain expression of a set of eight genes: *Nos2* and *Ptgs2* (nitric oxide synthase 2 and prostaglandin-endoperoxide synthase 2, two inducible enzymes), *Cxcl1* and *Cxcl10* (C-X-C motif chemokine ligand 1 and ligand 10), *Tgtp* and *Gbp2* (T cell-specific GTPase 1 and guanylate-binding protein 2, two GTPases), *Adamts1* (a disintegrin-like and metalloprotease with thrombospondin type 1 motif, 1, a metalloprotease) and *Il1rn* (interleukin-1 receptor antagonist). Our objective was to ascertain whether casp1 also controlled the peripheral expression of these genes and, if so, to compare their central versus peripheral patterns of gene expression in immune and endocrine tissues during SIRS. **Methods:** Wild-type (wt) and casp1 knockout (casp1^{-/-}) mice were injected

with either saline or a high dose of endotoxin/lipopolysaccharide (LPS; 800 µg/mice i.p.). Saline-injected mice were immediately euthanized after injection, whereas LPS-injected mice were sacrificed 6 and 12 h after LPS administration. Hippocampal, splenic and adrenal gene expressions were determined by real-time PCR. **Results:** Overall, casp1^{-/-} mice showed a lower inflammatory response than wt mice. The expression levels of powerful proinflammatory factors such as *Nos2* and *Ptgs2* was reduced in casp1^{-/-} mice. Moreover, a hierarchical clustering analysis aimed at studying patterns of gene coexpression revealed large alterations in the hippocampal pattern of casp1^{-/-} mice. Surprisingly, the expression of *Adamts1* was increased in the hippocampus and adrenals of casp1^{-/-} mice. **Conclusions:** The resilience of casp1^{-/-} mice to SIRS lethality is associated with a lower inflammatory response, loss of hippocampal gene coexpression patterns, and increased hippocampal *Adamts1* gene expression. The latter might be beneficial for casp1^{-/-} mice, since ADAMTS1 is likely to play a role in neuronal plasticity. The mechanisms described here may help the development of either novel biomarkers or therapeutic targets against SIRS/sepsis.

© 2015 S. Karger AG, Basel

Introduction

Systemic inflammatory response syndrome (SIRS) defines a serious medical condition of extreme and severe generalized inflammation in the absence of infection, whereas infection is the underlying process in sepsis [1]. Both SIRS and sepsis are leading causes of death in intensive care units [2].

The molecular mechanisms underlying SIRS remain elusive and, consequently, effective therapeutic strategies have not yet been developed. Animal models have been developed to aid the elucidation of the pivotal role of the innate immune system on SIRS [3–10], such as the administration of endotoxin/lipopolysaccharide (LPS) from Gram-negative bacteria to genetically engineered mouse strains in order to mimic the generalized inflammation present in SIRS [11, 12]. One of these strains is the caspase-1 ($\text{casp1}^{-/-}$) knockout mouse that is resilient to LPS-induced lethality [11]. Casp1 is a cysteine protease that may be a crucial element in SIRS lethality. It cleaves pro-interleukin (IL)-1 β and pro-IL-18 into mature biologically active cytokines. By submitting $\text{casp1}^{-/-}$ mice to LPS-induced SIRS, we previously observed that casp1 orchestrates the differential expression of eight genes in the central nervous system (CNS): two inducible enzymes [nitric oxide synthase 2 (*Nos2*) and prostaglandin-endoperoxide synthase 2/cyclooxygenase 2 (*Ptgs2*)], two chemokines [C-X-C motif ligand 1 (*Cxcl1*) and 10 (*Cxcl10*)], two GTPases [T cell-specific GTPase (*Tgtp*) and guanylate nucleotide-binding protein 2 (*Gbp2*)], one metalloprotease [a disintegrin-like and metallopeptidase with thrombospondin type 1 (*Adamts1*)] and the IL-1 receptor antagonist gene (*Il1rn*), an endogenous antagonist of IL-1 [12].

Because SIRS is characterized by neuro-immuno-endocrine derangements leading to death [13–16], we focused our studies on neural (hippocampus), immune (spleen) and endocrine (adrenals) tissues of wild-type (wt; control) and $\text{casp1}^{-/-}$ mice. The hippocampus was chosen because it is a brain area that plays a decisive role in the regulation of the HPA (hypothalamic-pituitary-adrenal) axis during different types of stress [17], including immunological challenges [18, 19]. The adrenals are the effector organs of the HPA-axis and were studied because of their crucial role in the secretion of corticoids during SIRS, whereas the spleen was chosen for being a prominent immune organ.

Given the fact that many of the proinflammatory factors that are triggered during LPS-induced SIRS require de novo synthesis, we measured their mRNA expression before (0 h) and after LPS injection (6 and 12 h later). We

hypothesized that their involvement in SIRS-induced lethality would be reflected by differential mRNA expression levels between wt (control) and $\text{casp1}^{-/-}$ mice (resilient to SIRS-induced lethality).

We also assessed gene-gene interactions in wt and $\text{casp1}^{-/-}$ mice and compared them to the patterns of gene expression in the CNS (hippocampal). Herein, we present evidence that a lower inflammatory response, pattern disruption of hippocampal gene coexpression of the aforementioned set of eight genes, and activation of hippocampal *Adamts1* may be key mechanisms underpinning resilience to SIRS lethality in $\text{casp1}^{-/-}$ mice.

Materials and Methods

Animals

We used virus- and antibody-free 8 to 10-week-old male wt and $\text{casp1}^{-/-}$ mice that were bred in the same genetic background (C57BL/6). $\text{Casp1}^{-/-}$ mice were created, characterized and generously provided by BASF BioResearch (Cambridge, Mass., USA) [11]. These mice are fertile, develop normally, and have no histopathological abnormalities. It is noteworthy that $\text{casp1}^{-/-}$ mice are resistant to the lethal effects of LPS [11] as well as to the anorectic effects exerted by the central action of LPS-induced cytokines [20]. Mice were kept in a light- (12 h on/12 h off) and temperature-controlled environment, with food and water ad libitum. All procedures were performed under established guidelines for the humane care and use of animals; they were approved by the University of Miami Institutional Animal Care and Use, and the Australian National University Animal Experimentation Ethics Committee. Experiments were designed to avoid confounding variables such as infection and stress, which could alter mRNA levels.

Experimental Protocol

Wt and $\text{casp1}^{-/-}$ mice ($n = 23$ per genotype) were injected with either 0.9% NaCl (saline) or a lethal dose of LPS (800 $\mu\text{g}/\text{mice}$ i.p. from *Escherichia coli* serotype O111:B4; Merk Millipore, Bayswater, Vic., Australia) [11]. Control animals were immediately euthanized after saline injection i.p. (0 h; wt mice, $n = 7$, and $\text{casp1}^{-/-}$ mice, $n = 8$), whereas LPS-injected mice were euthanized at 6 h (wt, $n = 13$, and $\text{casp1}^{-/-}$, $n = 12$), and 12 h (wt and $\text{casp1}^{-/-}$, $n = 3$ per genotype) after LPS injection. We chose to use the specific LPS serotype, dose and administration route because this regimen was previously shown to cause 100% lethality in wt mice during the first 36 h after injection [11]. Injections were given between 7.00 and 9.00 a.m.; mice were sacrificed by decapitation one at a time in a room distant from the animal holding area in a manner to avoid stress triggered by blood scent. Organs were extracted, snap-frozen and stored at -80°C until processed.

RNA Isolation and Reverse Transcription

We chose one brain area (hippocampus) and two peripheral tissues (spleen and adrenals), to determine variations in central and peripheral mRNA expression levels. Total RNA was extracted from individual tissues using RNeasy Lipid Mini Kit including DNase digestion (Qiagen, Germantown, Md., USA) and quantified using

Table 1. List of real-time PCR primers used to determine the gene expression levels of the eight selected genes

Gene	Forward	Reverse	NCBI Ref. Seq.
<i>β-actin</i>	5-CCTAGGCACCAGGGTGTGAT-3	5-CATGTCGTCGCCAGTTGGTAACA-3	NM_007393
<i>Ptgs2</i>	5-TGGGCCATGGAGTGGACTT-3	5-GGATGTGAGGAGGGTAGATCATCT-3	NM_011198.3
<i>Nos2</i>	5-AAGGCCACATCGGATTTTAC-3	5-TGTTCTCTATTTTTGCCTCTTTAAAG-3	NM_010927.3
<i>Cxcl1</i>	5-ACCGAAGTCATAGCCACACTCA-3	5-CACCTTTTAGCATCTTTTGGACAA-3	NM_008176.3
<i>Cxcl10</i>	5-CTGGGTCTGAGTGGGACTCAA-3	5-TCATTGCCACGATGAAAAAGAAT-3	NM_021274.1
<i>Tgtp</i>	5-CACCACTGCTGAGCTTCTACCATA-3	5-TCGATGTCTCTCAGTACCTTTTCAA-3	NM_011579.3
<i>Gbp2</i>	5-TTGAGAAGGGTGACAACCAGAA-3	5-TGGTTCCTATGCTGTTGTAGATGAA-3	NM_010260.1
<i>Il-1ra</i>	5-CCAGCTGGAGGAAGTTAACATCA-3	5-ACGGTCAGCCTCTAGTGTGTG-3	NM_031167.5
<i>Adamts1</i>	5-GACAATAACGGAAAAACGTTTCA-3	5-GTGAGCTTGACCTGTCCTTT-3	NM_009621.4

a Nanodrop ND-1000 spectrophotometer (Thermo-Fisher Scientific, Waltham, Mass., USA); 1 µg of RNA was reverse transcribed to cDNA with the Omniscript RT Kit (Qiagen) using random hexamer primers (Invitrogen, Carlsbad, Calif., USA).

PCR Primers

The genes of interest were selected from our previous microarray study [12]. To avoid amplification of genomic DNA traces, forward and reverse primers were designed in different exons with Primer Express Software (Applied Biosystems, Foster City, Calif., USA). We used negative and positive controls (respectively, reverse transcription with genomic DNA and cDNA from mouse brain treated with LPS). Table 1 lists our primer pair sequences, which amplified cDNA only.

Quantitative Real-Time PCR

We performed SYBR green-based real-time PCR to determine variations in mRNA expression profiles. Standard curves of pooled, serially diluted cDNAs were constructed for each primer pair. Experiments were performed using the 7900HT Fast Real-Time PCR System (Applied Biosystems). cDNA samples were run in triplicates; SYBR green cycle threshold values were averaged for each sample, and the RNA input for each target gene was calculated using standard curves. Fold changes were expressed as a ratio to β -actin expression for each sample. Each reaction included cDNA from 20 ng of RNA, 4.5 pmol of each primer, and 7.5 µl of SYBR green PCR Mastermix (Applied Biosystems) in a final volume of 15 µl. The PCR parameters were: 50°C for 2 min, 95°C for 10 min, 40 cycles of 95°C for 15 s, and 60°C for 1 min.

Statistical Analysis

Variance Analyses

Two-way ANOVA tests were performed to assess the effects of genotype (wt or $\text{caspl}^{-/-}$) and time (0, 6 or 12 h) both individually and in combination on gene expression levels in adrenal, hippocampus and spleen tissues. Tukey's honest significant difference test was used within each gene/tissue combination to perform pair-wise comparisons maintaining the family-wise probability of coverage at 95%. We present 95% confidence intervals (CI) for the differences and family-wise-corrected p values to assess statistical significance. All analyses were performed in R package version 3.0.2 (R Foundation for Statistical Computing, 2013).

Correlation Analysis

We calculated pair-wise Pearson's correlation coefficients to examine the relationships between gene expression in wt and $\text{caspl}^{-/-}$ mice. As a complementary strategy, nonparametric local regressions (LOESS) were fitted to the data to assess the relationship between pair-wise gene expression levels. Statistical comparison of the linear correlation coefficient was performed based on a normal distribution, and the resulting p values were corrected using the false discovery rate [21].

Hierarchical Clustering

We used hierarchical clustering to analyze gene coexpression in our samples. Hierarchical clustering uses an agglomerative algorithm that joins the most similar component features, and then joins the next most similar using the first aggregation as a single combined unit. In hierarchical clustering, the clusters were generated via complete linkage. To assess the uncertainty of the analysis, approximately unbiased (AU) and bootstrap probability (BP) p values were calculated by pvclust [22], R package version 3.0.2. AU p values were computed by multiscale bootstrap resampling, and are a better approximation to unbiased p values than BP-based p values (computed by normal bootstrap resampling).

Results

LPS-Induced Differential Temporal Gene Expression in Peripheral and Central Tissues of wt and $\text{caspl}^{-/-}$ Mice

Our study was conducted up to 12 h after LPS injection. The health status of mice of both genotypes gradually worsened with the passage of time. Towards the last collection point, at 12 h after LPS injection, wt mice were quite immobile but without signs of being moribund (e.g. hard-labored breathing), whereas $\text{caspl}^{-/-}$ mice were less active but still responsive to external stimuli such as a gentle tapping and/or shaking of the cage. We corroborated the resilience of $\text{caspl}^{-/-}$ mice to

Table 2. Survival rate of wt and casp1^{-/-} mice after receiving an LD₁₀₀ injection of LPS

	n	Survival rate after LPS injection, %	
		12 h	24 h
wt	4	100	0
casp1 ^{-/-}	5	100	100

At 24 h all wt mice had died, whereas all casp1^{-/-} mice were still alive.

the lethal effects of LPS in a separate experiment. We confirmed that the same LPS serotype and dose used in the present studies caused 100% lethality in wt mice (n = 4) within 12–24 h after injection, whereas all casp1^{-/-} mice (n = 5) were alive at 24 h, which confirmed previous results showing that 100% of casp1^{-/-} survived for at least 30 h after receiving the same i.p. dose and serotype of LPS injection [11] (table 2). Figure 1 illustrates gene expression level changes in the hippocampus, spleen and adrenals.

In wt mice, most of the studied genes displayed the highest expression levels (23 out of 24 measurements) at 6 h after injection (χ^2 , $p < 0.0001$), except for adrenal *Adamts1* which peaked at 12 h. In casp1^{-/-} mice, the highest mRNA levels (19 out of 24 measurements) were also seen at 6 h after injection (χ^2 , $p < 0.01$), except for splenic *Il1rn*, *Ptgs2*, *Nos2* and *Adamts1*, and adrenal *Adamts1*, which peaked at 12 h (fig. 1). We found that LPS induced significant increases (above baseline levels observed at 0 h) in the expression of most of the eight studied genes in the three tissues of both genotypes, except for the splenic expression of *Il1rn*, *Nos2*, *Ptgs2* and *Adamts1* genes in casp1^{-/-} mice, and the splenic *Il1rn* gene in wt mice (fig. 1; table 3); therefore, four out of the eight genes were not activated in the spleen of casp1-deficient mice. Thus, it appears that the temporal expression of the studied genes mostly followed an early and sharp increase pattern similar to that described for proinflammatory elements during the initial stage of the inflammatory response. A summary of the two-way

ANOVA significance probabilities for the differential expression of each gene within each organ is given in table 3. The following sections provide a brief summary of the gene expression changes for each gene.

Nos2

Nos2 gene expression was significantly lower at 6 h in all three casp1^{-/-} mice tissues when compared to those of wt mice; it was 30% lower in the hippocampus ($p < 0.01$), 75% lower in the spleen ($p < 0.001$), and 40% lower in the adrenals ($p < 0.001$; fig. 1a). The LPS-induced expression of *Nos2* followed a similar pattern in both genotypes, but of decreased magnitude in casp1^{-/-} mice. Both genotypes showed activation of gene expression at 6 h, and a decrease towards baseline at 12 h (except for *Nos2* expression in wt adrenal, which remained elevated at 12 h; $p < 0.01$ vs. 0 h).

Ptgs2

The peak *Ptgs2* mRNA level at 6 h was significantly lower by 25 and 75% in casp1^{-/-} mice in the hippocampus ($p < 0.05$) and the spleen ($p < 0.001$), respectively, whereas no significant changes were observed within the adrenals of both genotypes (fig. 1b). *Ptgs2* transcriptional patterns were similar to those of *Nos2*, i.e. *Ptgs2* gene expression was increased at 6 h and returned to baseline values at 12 h.

Cxcl1, *Cxcl10* and *Tgtp*

In the two peripheral organs, casp1^{-/-} mice showed a significantly lower and variable magnitude of increases in these three genes compared to wt mice, and casp1^{-/-} mice displayed 30–80% lower mRNA levels ($p < 0.05$ to < 0.001 ; fig. 1c–e). These genes showed similar transcriptional patterns in the three tissues/organs studied from both genotypes. At the CNS level, their hippocampal expression showed no significant differences between the two genotypes.

Gbp2

Adrenal mRNA expression of *Gbp2* was significantly lower in casp1^{-/-} mice than wt mice, by 35% at 6 h ($p < 0.001$; fig. 1f). Thereafter, *Gbp2* expression had decreased

Fig. 1. mRNA expression pattern of the eight selected genes in the hippocampus, spleen and adrenals of wt and casp1^{-/-} mice. The time-dependent pattern of expression of each gene is shown: *Nos2* (a), *Ptgs2* (b), *Cxcl1* (c), *Cxcl10* (d), *Tgtp* (e), *Gbp2* (f), *Il1rn* (g) and *Adamts1* (h). The mean expression levels at 0, 6 and 12 h after LPS

injection are represented by white and black columns, respectively, for wt and casp1^{-/-} mice; the line above represents one standard error of the mean. Statistically significant differences between wt and casp1^{-/-} mice at the same time point are shown: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$. (For figure see next page.)

Table 3. Two-way ANOVA summary

Organ	Gene	Component			wt			casp1 ^{-/-}		
		genotype	time	interaction	6 vs. 0 h	12 vs. 0 h	12 vs. 6 h	6 vs. 0 h	12 vs. 0 h	12 vs. 6 h
Hippocampus	<i>Nos2</i>	***	***	n.s.	***	n.s.	***	***	n.s.	**
	<i>Ptgs2</i>	**	***	*	***	n.s.	*	***	n.s.	*
	<i>Cxcl1</i>	**	***	n.s.	***	***	n.s.	***	n.s.	**
	<i>Cxcl10</i>	n.s.	***	n.s.	***	***	n.s.	***	*	n.s.
	<i>Tgtp</i>	n.s.	***	n.s.	***	**	n.s.	***	n.s.	***
	<i>Gbp2</i>	n.s.	***	n.s.	***	***	n.s.	***	***	n.s.
	<i>Il1rn</i>	***	***	*	***	***	n.s.	***	n.s.	*
	<i>Adamts1</i>	***	***	***	***	***	n.s.	***	***	n.s.
Spleen	<i>Nos2</i>	***	***	**	***	n.s.	**	n.s.	n.s.	n.s.
	<i>Ptgs2</i>	***	***	**	***	n.s.	**	n.s.	n.s.	n.s.
	<i>Cxcl1</i>	***	***	**	***	n.s.	**	**	n.s.	n.s.
	<i>Cxcl10</i>	**	***	n.s.	***	n.s.	***	***	n.s.	**
	<i>Tgtp</i>	**	***	*	***	n.s.	***	**	n.s.	n.s.
	<i>Gbp2</i>	n.s.	***	n.s.	***	n.s.	n.s.	***	n.s.	n.s.
	<i>Il1rn</i>	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
	<i>Adamts1</i>	***	*	*	*	n.s.	n.s.	n.s.	n.s.	n.s.
Adrenals	<i>Nos2</i>	***	***	*	***	**	n.s.	***	n.s.	*
	<i>Ptgs2</i>	n.s.	***	*	**	**	n.s.	***	n.s.	n.s.
	<i>Cxcl1</i>	***	***	**	***	***	n.s.	***	n.s.	n.s.
	<i>Cxcl10</i>	***	***	*	***	*	n.s.	***	n.s.	*
	<i>Tgtp</i>	***	***	*	***	n.s.	n.s.	*	n.s.	n.s.
	<i>Gbp2</i>	***	***	**	***	***	n.s.	***	n.s.	**
	<i>Il1rn</i>	***	***	n.s.	***	*	n.s.	***	n.s.	n.s.
	<i>Adamts1</i>	***	***	***	n.s.	*	n.s.	***	***	***

The significance of the three major components (genotype, time and interaction) is shown for the eight genes in each of the tissues studied. p values are family-wise corrected. The comparison among time points within genotypes is also shown for wt and casp1^{-/-} mice. Statistically significant differences between time points are shown: * p < 0.05; ** p < 0.01; *** p < 0.001. n.s. = Not significant. Bold n.s. indicates no increased expression when compared with 0 h. For example, splenic expression of *Nos2*, *Ptgs2* and *Adamts1* remained significantly unchanged in casp1^{-/-} mice, whereas the splenic expression of *Il1rn* remained significantly unchanged in both genotypes.

in both genotypes at 12 h to similar levels. Hippocampal and splenic *Gbp2* expression levels showed no genotype differences.

Il1rn

In the hippocampus of both genotypes, LPS elicited similarly large *Il1rn* expression increases at 6 h. At 12 h, *Il1rn* mRNA remained elevated in wt mice and declined by 70% in casp1^{-/-} mice (p ≤ 0.05; fig. 1g). Adrenal *Il1rn* activation for both genotypes was similar to that of the hippocampus, reaching significantly larger increases of 40% at 6 h (p < 0.05) and of 115% at 12 h (p < 0.001) in wt mice. Splenic *Il1rn* mRNA expression was unchanged in both genotypes.

Adamts1

Surprisingly, hippocampal *Adamts1* mRNA expression was of a higher magnitude in casp1^{-/-} mice, increased by 23% at 6 h (p ≤ 0.01) and 45% at 12 h (p < 0.001; fig. 1h), and followed a similar temporal pattern in wt and casp1^{-/-} mice. Adrenal *Adamts1* expression was also larger in casp1^{-/-} mice: they were 50% higher at 6 h (p < 0.05) and 2.3-fold higher at 12 h (p < 0.001). Splenic *Adamts1* mRNA expression increased significantly at 6 h and returned to baseline levels at 12 h in wt mice, whereas no changes above baseline were observed in casp1^{-/-} mice. The genotype comparison showed that splenic *Adamts1* mRNA levels were 70% lower in casp1^{-/-} mice at 6 h (p < 0.001).

Correlation Analysis

Hippocampus

All correlations between hippocampal mRNA expression levels of the genes studied were significant within each genotype; however, correlation coefficients comparing genotypes (wt vs. *casp1*^{-/-} mice) showed significant differences only between *Tgtp/Gbp2* ($p_{\text{FDR}} < 0.05$) and *Tgtp/Adamts1* ($p_{\text{FDR}} < 0.01$; online suppl. fig. 1S; for all online suppl. material, see www.karger.com/doi/10.1159/000368310).

Spleen

In wt mice, correlation analyses of splenic mRNA levels were significant, except for all *Il1rn* correlations (online suppl. fig. 2S). In *casp1*^{-/-} mice, most splenic mRNA level correlations were significant, and most *Il1rn* and *Adamts1* correlations were not significant, except for the correlation between *Il1rn* and *Adamts1*, which was significant (online suppl. fig. 2S). Splenic correlation coefficients comparing genotype showed that only three coefficient pairs, all related to *Adamts1*, were significantly different: *Ptgs2* and *Adamts1* ($p_{\text{FDR}} < 0.001$), *Nos2* and *Adamts1* ($p_{\text{FDR}} = 0.006$), and *Gbp2* and *Adamts1* ($p_{\text{FDR}} = 0.018$; online suppl. fig. 2S).

Adrenals

Correlation analysis of adrenal mRNA levels showed that all correlations were significant in wt mice, except for *Nos2* and *Adamts1*, whereas six pairs of correlations related to *Adamts1* were not significant in *casp1*^{-/-} mice: *Ptgs2* and *Adamts1*, *Nos2* and *Adamts1*, *Cxcl1* and *Adamts1*, *Cxcl10* and *Adamts1*, *Tgtp* and *Adamts1*, and *Gbp2* and *Adamts1*. Adrenal correlation coefficients comparing genotypes did not show any significant differences (online suppl. fig. 3S). Overall, there was a loss of correlation for the *Adamts1* gene in peripheral tissues in *casp1*-deficient mice.

Hierarchical Clustering

This analysis was done to ascertain patterns of gene coexpression using clustering strategies by comparing: (i) the same tissue between wt and *casp1*^{-/-} mice, and (ii) the three tissues within the same genotype (fig. 2).

Clustering Comparison between wt and *casp1*^{-/-} Mice

In the hippocampus (fig. 2a, b), two clusters were supported at the 95% CI level in wt animals. The first cluster was comprised of the *Cxcl10* and *Il1rn* genes, whereas the second was defined by the *Gbp2*, *Tgtp* and *Adamts1* genes

(fig. 2a, b). Interestingly, no clusters were supported at the 95% level in *casp1*^{-/-} mice (fig. 2a, b). There were qualitative similarities between splenic data of both genotypes (fig. 2c, d). In the spleen of wt mice, all genes except the *Il1rn* gene formed one cluster (fig. 2c, d), whereas in *casp1*^{-/-} mice, two clusters were supported at the 95% level (the *Cxcl1* and *Cxcl10* cluster, and the *Ptgs2*, *Gbp2*, *Nos2* and *Tgtp* cluster; fig. 2d). High similarities were found between adrenal data from wt and *casp1*^{-/-} mice: they had one cluster consisting of six and seven genes, respectively, at a 95% significance level. The *Ptgs2* (in wt mice) and the *Adamts1* (in wt and *casp1*^{-/-} mice) genes were not included in the clusters (fig. 2f).

Clustering Comparison among Tissues

More similarities were found between the two peripheral organs (spleen and adrenals) than between those tissues and the hippocampus. In wt mice, five genes were coexpressed both within the spleen and adrenals (*Gbp2*, *Tgtp*, *Cxcl1*, *Cxcl10* and *Nos2*). In *casp1*^{-/-} mice, six out of the seven genes that clustered together within the adrenal gland were divided into two clusters (one with two genes, *Cxcl1* and *Cxcl10*, and another with four genes *Ptgs2*, *Gbp2*, *Nos2* and *Tgtp*) in the spleen. There were two clusters within the hippocampus of wt mice. One of them showed a coexpression of two genes (*Cxcl10* and *Il1rn*) that were also shown to be coexpressed in the adrenals of wt mice (fig. 2a, b, e, f). The other cluster within the brain of wt mice displayed three genes that were also shown to be coexpressed within the spleen (fig. 2a–d). Overall, at the 95% level, *Adamts1* did not coexpress with any gene in *casp1*^{-/-} mice tissues, while in wt it coexpressed in 2 of the 3 tissues.

Discussion

We used LPS-induced SIRS in *casp1*-deficient mice to detail patterns of gene expression that may contribute to their resilience to LPS-induced lethality, which could lead to the development of novel therapeutic targets for SIRS. In this study, we described central and peripheral changes in the expression of a set of eight genes in the context of *casp1* deficiency and a lethal dose of LPS, and showed that endotoxin elicited an inflammatory cascade of much lower magnitude in *casp1*^{-/-} mice. Overall, these changes occurred in a time-dependent manner, with peak mRNA expression of most genes occurring 6 h after LPS injection, and declining to baseline levels by 12 h. In previous studies we have shown evidence suggesting that activated transcripts during inflammation show two

Fig. 2. Hierarchical cluster analysis of gene expression levels with AU- and BP-based p values (in red and green, respectively; colors refer to the online version only). The vertical distance between the gene expression levels indicate how dissimilar they are. Results for

hippocampus (**a, b**), spleen (**c, d**) and adrenal tissues (**e, f**) are presented in wt (**a, c, d**) and *casp1*^{-/-} (**b, d, f**) mice. The genes shown inside the red boxes clustered together according to an AU probability set at ≥95% level, after bootstrap resampling.

types of temporal patterns: transient and sustained [23]. With exception of *Il1rn*, it appears that within the spleen most of the genes followed a transient pattern of increase by augmenting at 6 h and returning to baseline at 12 h, whereas most of the genes seemed to follow a more sustained pattern of activation in the hippocampus and adrenals, which lasted for at least 12 h.

The most contrasting changes in gene expression between both genotypes occurred in the spleen. After LPS injection, the splenic expression of three genes, namely *Nos2*, *Ptgs2* and *Adamts1*, was largely increased in wt mice, whereas it was not significantly altered in *casp1*^{-/-} mice.

There are a couple of exceptions to the generalized idea that the inflammatory cascade elicited by LPS was of a lower magnitude in *casp1*^{-/-} mice. For instance, we found that the expression of *Adamts1* remained significantly increased in the hippocampus and adrenals of *casp1*-deficient mice.

Our previous work suggested that the expression of these eight genes was controlled by *casp1* [12]. In order to study gene-gene interactions, we evaluated their relationship by performing correlation and hierarchical clustering analyses. There was an overall loss of correlation for the *Adamts1* gene in peripheral tissues of *casp1*^{-/-} mice. Differences observed in the hierarchical clustering

analyses seem to support that casp1 deficiency altered the coexpression of these genes in a more significant manner within the brain than in the peripheral organs. Indeed, these analyses revealed similar clusters within the adrenals and the spleen of wt and casp1^{-/-} mice, whereas no similarities were evident in the hippocampus of both genotypes. In addition, coexpression analyses showed more similarities between peripheral organs (spleen and adrenals) than between these peripheral organs and the hippocampus; there was also a tendency of coexpression loss for *Adamts1* in casp1^{-/-} mice.

Products of three out of the eight genes reported here, specifically the *Nos2*, *Ptgs2* and *Il1rn* genes, have been pharmacologically targeted in preclinical and clinical trials of SIRS/sepsis. The inhibition of nitric oxide (NO) synthesis, a free radical largely synthesized by NOS2 during SIRS/sepsis, improved hemodynamic parameters in immunologically challenged rodents and septic patients [24–30]. However, multicenter randomized placebo-controlled studies showed that the mortality rate of septic shock patients treated with NOS inhibitors was not improved [29, 30]. Similarly, preclinical studies largely support the concept that pharmacological inhibition of cyclooxygenase and/or genetic ablation of PTGS2 resulted in beneficial effects on mortality [31]. Unfortunately, the beneficial effects of cyclooxygenase inhibition observed in preclinical studies have so far not been convincingly and consistently translated into the clinical setting [32–34]. Only a sub-study from one clinical trial showed a significant reduction in the 30-day mortality rate of sepsis [35]. Studies involving IL-1ra also yielded promising outcomes in preclinical trials [36]. Likewise casp1^{-/-} mice, overexpressing IL-1ra have also shown resilience to LPS-induced lethality [36]. In both casp1^{-/-} and IL-1ra-overexpressing mice, the resilience to LPS-induced lethality was observed for at least 6 and 7 days, respectively, indicating that in these two strains of mice the improvement in survival was long lasting [11, 36]. However, these results could not be effectively translated into the clinical setting as rhIL1ra failed to reduce sepsis mortality [37, 38].

The contribution of the products of the other five genes that were also differentially expressed between wt and casp1^{-/-} mice has not been clearly ascertained in preclinical/clinical studies of SIRS/sepsis. Two of them are the chemokines *Cxcl1* and *Cxcl10*. Chemokines and their receptors are expressed in immune and endothelial cells, and play a fundamental role in recruiting immune cells into the inflammatory site [39, 40]. Therefore, the reduced expression of *Cxcl1* and *Cxcl10* observed in casp1^{-/-} mice could have contributed to a better outcome against

LPS-induced lethality due to decreased immune cell recruitment.

We also investigated the expression of the GTPases *Tgtp* and *Gbp2*, another two genes that were differentially expressed between casp1^{-/-} and wt mice. These two interferon-induced GTPases are expressed in immune cells, but their roles in SIRS/sepsis have not been clearly ascertained [41–43]. Overall, guanylate binding proteins have been characterized as downstream effectors of the interferon signaling pathway [44]. They play essential roles in fighting intracellular pathogens [44]. A recent study showed that intracellular LPS from Gram-negative bacteria activated guanylate-binding proteins, which in turn participated in a programmed cell death pathway termed pyroptosis [45]. Since we employed a large dose of LPS, it is likely that increased cytosolic accumulation of endotoxin could activate guanylate-binding protein-induced pyroptosis. Thus, their decreased expression in casp1^{-/-} mice could result in decreased inflammation and, consequently, in improved survival after being exposed to a lethal dose of endotoxin. However, further studies need to be performed to elucidate the roles of GBP2 and TGTP during SIRS.

Finally, ADAMTS1, an enzyme that belongs to the family of the metalloproteases, was also differentially expressed between the two genotypes. ADAMTS1 is necessary for normal growth, female fertility and kidney, adrenal and female reproductive organ structure and function [46]. Recently, it was shown that LPS dramatically increased the synthesis of ADAMTS1 within the splenic lymph and interstitial fluid to a concentration that was much higher than that of plasma, suggesting that the spleen is a major source of ADAMTS1, and that this metalloprotease is involved in the endotoxin-induced disease process [47]. Within the brain, metalloproteases play a key role in controlling blood-brain barrier permeability during LPS-induced inflammation, and ADAMTS1 may have a role in regulating neural plasticity, as null female mice have a selective decline of synaptic protein levels in the frontal cortex [48].

There are apparent discrepancies between our current and previous studies. In our previous study, the transcriptional differences between both genotypes appeared to occur specifically within the brain, and not in the periphery, since both wt and casp1^{-/-} mice showed similar LPS-induced increases in their expression level within the spleen [12]. In this study, we found that only the hippocampal transcriptional expression of two out of the eight genes, *Nos2* and *Ptgs2*, showed similar differences to those previously reported in the whole brain of both genotypes [12]. These differences could be due to the fact that we previ-

ously employed a lower dose and less inflammatory serotype of LPS (e.g. *E. coli* 055:B5, 25 mg/kg) than the one employed in this study [12]. The previous dose and serotype of LPS was shown to cause a slower progression towards death in wt mice (100% mortality after 5 days of injection) [49] than the LPS dose and serotype reported here (100% mortality within approximately 1.25 days) [11]. Furthermore, in our previous studies we employed whole brains [12], whereas here we used hippocampi. Moreover, we studied a small group of mice at 12 h after LPS injection, and this could have lowered the statistical power to detect differences at that particular time point.

It was recently shown that casp1-targeted mice carry a 129 ES cell-derived mutation in the *casp11* gene, which appeared to be involved, at least in part, in their resilience to LPS-induced lethality [50]. The synergistic key role of two of the downstream effectors controlled by casp1, namely IL-18 and IL-1 β , during LPS-induced lethality was recently shown [51]. The survival rate of casp1/11^{-/-} was very similar to that of IL-1 β /IL-18-deficient mice after receiving an LD₁₀₀ of LPS of the same serotype as the one used in the present studies [51]. The latter results clearly highlight the relevance of the casp1-orchestrated pathway during LPS-induced SIRS. However, when casp1/11^{-/-} and IL-1 β /IL-18-deficient mice were subjected to other immunological challenges, such as the cecal ligation and puncture procedure or the administration of a lethal dose of tumor necrosis factor- α , casp1/11^{-/-} displayed a lower survival rate than IL-1 β /IL-18-deficient

mice, indicating that targeting downstream effectors could be more efficacious than inhibiting the upstream activators CASP1 or CASP11 [51].

In summary, we described tissue and time-dependent expression profiles of a set of genes regulated by casp-1 after LPS-induced SIRS. Our data suggest that three elements may contribute to casp1^{-/-} mice resilience to SIRS-induced lethality: (1) a decreased inflammatory cascade due to a lower expression of casp1-orchestrated genes and a faster resolution of the proinflammatory process, (2) increased *Adamts1* activation in the hippocampus, which could potentially increase neural plasticity, and (3) loss of gene-gene interaction in the hippocampus. Future studies need to confirm whether changes in CNS gene-gene interactions or ADAMTS1 biological activity are associated with the increased survival of casp1^{-/-} mice after a lethal dose of LPS. Additionally, we will examine whether inhibitors of casp1 activity promote similar changes in genetic expression. The elucidation of these transcription pattern changes could provide a foundation for the development of molecular biomarkers and therapeutic targets against SIRS/sepsis.

Acknowledgments

This research was supported in part by a National Institutes of Health grant K24RR017365 to M.L.W., and by institutional funds from the University of Miami and the Australian National University. The funders had no role in the study design, data collection and analysis, decision to publish or preparation of the manuscript.

References

- 1 Takala A, Nupponen I, Kylanpaa-Back ML, Repo H: Markers of inflammation in sepsis. *Ann Med* 2002;34:614–623.
- 2 Hotchkiss RS, Karl IE: The pathophysiology and treatment of sepsis. *N Engl J Med* 2003; 348:138–150.
- 3 Wong ML, Xie B, Beatini N, Phu P, Marathe S, Johns A, Gold PW, Hirsch E, Williams KJ, Licinio J, Tabas I: Acute systemic inflammation up-regulates secretory sphingomyelinase in vivo: a possible link between inflammatory cytokines and atherogenesis. *Proc Natl Acad Sci U S A* 2000;97: 8681–8686.
- 4 Mastronardi CA, Srivastava V, Yu WH, Les Dees W, McCann SM: Lipopolysaccharide-induced leptin synthesis and release are differentially controlled by α -melanocyte-stimulating hormone. *Neuroimmunomodulation* 2005;12:182–188.
- 5 Mastronardi CA, Yu WH, McCann S: Lipopolysaccharide-induced tumor necrosis factor- α release is controlled by the central nervous system. *Neuroimmunomodulation* 2001;9:148–156.
- 6 Mastronardi CA, Yu WH, Rettori V, McCann S: Lipopolysaccharide-induced leptin release is not mediated by nitric oxide, but is blocked by dexamethasone. *Neuroimmunomodulation* 2000;8:91–97.
- 7 Wong ML, Bongiorno PB, Rettori V, McCann SM, Licinio J: Interleukin (IL) 1 β , IL-1 receptor antagonist, IL-10, and IL-13 gene expression in the central nervous system and anterior pituitary during systemic inflammation: pathophysiological implications. *Proc Natl Acad Sci U S A* 1997;94:227–232.
- 8 Wong M-L, Rettori V, Al-Shekhlee A, Bongiorno PB, Canteros G, McCann SM, Gold PW, Licinio J: Inducible nitric oxide synthase gene expression in the brain during systemic inflammation. *Nature Med* 1996;2:581–584.
- 9 Mastronardi CA, Yu WH, Srivastava VK, Dees WL, McCann SM: Lipopolysaccharide-induced leptin release is neurally controlled. *Proc Natl Acad Sci U S A* 2001;98:14720–14725.
- 10 Wong ML, O’Kirwan F, Khan N, Hannestad J, Wu KH, Elashoff D, Lawson G, Gold PW, McCann SM, Licinio J: Identification, characterization, and gene expression profiling of endotoxin-induced myocarditis. *Proc Natl Acad Sci U S A* 2003;100:14241–14246.
- 11 Li P, Allen H, Banerjee S, Franklin S, Herzog L, Johnston C, McDowell J, Paskind M, Rodman L, Salfeldt J, et al: Mice deficient in IL-1 β -converting enzyme are defective in production of mature IL-1 β and resistant to endotoxic shock. *Cell* 1995;80:401–411.
- 12 Mastronardi C, Whelan F, Yildiz OA, Hannestad J, Elashoff D, McCann SM, Licinio J, Wong ML: Caspase 1 deficiency reduces inflammation-induced brain transcription. *Proc Natl Acad Sci U S A* 2007;104: 7205–7210.
- 13 Chesnokova V, Melmed S: Minireview: neuro-immuno-endocrine modulation of the hypothalamic-pituitary-adrenal (HPA) axis by gp130 signaling molecules. *Endocrinology* 2002;143:1571–1574.

- 14 Carnio EC, Moreto V, Giusti-Paiva A, Antunes-Rodrigues J: Neuro-immune-endocrine mechanisms during septic shock: role for nitric oxide in vasopressin and oxytocin release. *Endocr Metab Immune Disord Drug Targets* 2006;6:137–142.
- 15 Chrousos GP: The hypothalamic-pituitary-adrenal axis and immune-mediated inflammation. *New Engl J Med* 1995;332:1351–1362.
- 16 Tracey KJ: Physiology and immunology of the cholinergic antiinflammatory pathway. *J Clin Invest* 2007;117:289–296.
- 17 De Kloet ER, Vreugdenhil E, Oitzl MS, Joels M: Brain corticosteroid receptor balance in health and disease. *Endocr Rev* 1998;19:269–301.
- 18 Schobitz B, Sutanto W, Carey MP, Holsboer F, de Kloet ER: Endotoxin and interleukin 1 decrease the affinity of hippocampal mineralocorticoid (type I) receptor in parallel to activation of the hypothalamic-pituitary-adrenal axis. *Neuroendocrinology* 1994;60:124–133.
- 19 Bener D, Wohlman A, Itzik A, Yirmiya R, Ben-Hur T, Weidenfeld J: Glucocorticoid resistance following herpes simplex-1 infection: role of hippocampal glucocorticoid receptors. *Neuroendocrinology* 2007;85:207–215.
- 20 Yao JH, Ye SM, Burgess W, Zachary JF, Kelley KW, Johnson RW: Mice deficient in interleukin-1 β converting enzyme resist anorexia induced by central lipopolysaccharide. *Am J Physiol* 1999;277:R1435–R1443.
- 21 Benjamini Y, Hochberg Y: Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Ser B* 1995;57:289–300.
- 22 Suzuki R, Shimodaira H: Pvcust: an R package for assessing the uncertainty in hierarchical clustering. *Bioinformatics* 2006;22:1540–1542.
- 23 Mastronardi CA, Licinio J, Wong ML: Candidate biomarkers for systemic inflammatory response syndrome and inflammation: a pathway for novel translational therapeutics. *Neuroimmunomodulation* 2010;17:359–368.
- 24 Thiemeermann C, Vane J: Inhibition of nitric oxide synthesis reduces the hypotension induced by bacterial lipopolysaccharides in the rat in vivo. *Eur J Pharmacol* 1990;182:591–595.
- 25 Kilbourn RG, Gross SS, Jubran A, Adams J, Griffith OW, Levi R, Lodato RF: NG-methyl-L-arginine inhibits tumor necrosis factor-induced hypotension: implications for the involvement of nitric oxide. *Proc Natl Acad Sci U S A* 1990;87:3629–3632.
- 26 Petros A, Bennett D, Vallance P: Effect of nitric oxide synthase inhibitors on hypotension in patients with septic shock. *Lancet* 1991;338:1557–1558.
- 27 Petros A, Lamb G, Leone A, Moncada S, Bennett D, Vallance P: Effects of a nitric oxide synthase inhibitor in humans with septic shock. *Cardiovasc Res* 1994;28:34–39.
- 28 Avontuur JA, Tutein Nolthenius RP, van Bodegom JW, Bruining HA: Prolonged inhibition of nitric oxide synthesis in severe septic shock: a clinical study. *Crit Care Med* 1998;26:660–667.
- 29 Lopez A, Lorente JA, Steingrub J, Bakker J, McLuckie A, Willatts S, Brockway M, Anzueto A, Holzapfel L, Breen D, Silverman MS, Takala J, Donaldson J, Arneson C, Grove G, Grossman S, Grover R: Multiple-center, randomized, placebo-controlled, double-blind study of the nitric oxide synthase inhibitor 546C88: effect on survival in patients with septic shock. *Crit Care Med* 2004;32:21–30.
- 30 Kinasewitz GT, Privalle CT, Imm A, Steingrub JS, Malczynski JT, Balk RA, DeAngelo J: Multicenter, randomized, placebo-controlled study of the nitric oxide scavenger pyridoxalated hemoglobin polyoxyethylene in distributive shock. *Crit Care Med* 2008;36:1999–2007.
- 31 Aronoff DM: Cyclooxygenase inhibition in sepsis: is there life after death? *Mediators Inflamm* 2012;2012:696897.
- 32 Bernard GR, Wheeler AP, Russell JA, Schein R, Summer WR, Steinberg KP, Fulkerson WJ, Wright PE, Christman BW, Dupont WD, Higgins SB, Swindell BB: The effects of ibuprofen on the physiology and survival of patients with sepsis. *New Engl J Med* 1997;336:912–918.
- 33 Haupt MT, Jastremski MS, Clemmer TP, Metz CA, Goris GB: Effect of ibuprofen in patients with severe sepsis: a randomized, double-blind, multicenter study. The Ibuprofen Study Group. *Crit Care Med* 1991;19:1339–1347.
- 34 Memis D, Karamanlioglu B, Turan A, Koyuncu O, Pamukcu Z: Effects of lornoxicam on the physiology of severe sepsis. *Crit Care* 2004;8:R474–R482.
- 35 Arons MM, Wheeler AP, Bernard GR, Christman BW, Russell JA, Schein R, Summer WR, Steinberg KP, Fulkerson W, Wright P, Dupont WD, Swindell BB: Effects of ibuprofen on the physiology and survival of hypothermic sepsis. Ibuprofen in Sepsis Study Group. *Crit Care Med* 1999;27:699–707.
- 36 Hirsch E, Irikura VM, Paul SM, Hirsh D: Functions of interleukin 1 receptor antagonist in gene knockout and overproducing mice. *Proc Natl Acad Sci U S A* 1996;93:11008–11013.
- 37 Opal SM, Fisher CJ Jr, Dhainaut JF, Vincent JL, Brase R, Lowry SF, Sadoff JC, Slotman GJ, Levy H, Balk RA, Shelly MP, Pribble JP, LaBrecque JF, Lookabaugh J, Donovan H, Dubin H, Baughman R, Norman J, DeMaria E, Matzel K, Abraham E, Seneff M: Confirmatory interleukin-1 receptor antagonist trial in severe sepsis: a phase III, randomized, double-blind, placebo-controlled, multicenter trial. The Interleukin-1 Receptor Antagonist Sepsis Investigator Group. *Crit Care Med* 1997;25:1115–1124.
- 38 Fisher CJ Jr, Opal SM, Lowry SF, Sadoff JC, LaBrecque JF, Donovan HC, Lookabaugh JL, Lemke J, Pribble JP, Stromattet SC, et al: Role of interleukin-1 and the therapeutic potential of interleukin-1 receptor antagonist in sepsis. *Circ Shock* 1994;44:1–8.
- 39 Puneet P, Mochhala S, Bhatia M: Chemokines in acute respiratory distress syndrome. *Am J Physiol Lung Cell Mol Physiol* 2005;288:L3–L15.
- 40 Oynebraten I, Bakke O, Brandtzaeg P, Johansen FE, Haraldsen G: Rapid chemokine secretion from endothelial cells originates from 2 distinct compartments. *Blood* 2004;104:314–320.
- 41 Carlow DA, Teh SJ, Teh HS: Specific antiviral activity demonstrated by TGTP, a member of a new family of interferon-induced GTPases. *J Immunol* 1998;161:2348–2355.
- 42 Carlow DA, Marth J, Clark-Lewis I, Teh HS: Isolation of a gene encoding a developmentally regulated T cell-specific protein with a guanine nucleotide triphosphate-binding motif. *J Immunol* 1995;154:1724–1734.
- 43 Lafuse WP, Brown D, Castle L, Zwilling BS: Cloning and characterization of a novel cDNA that is IFN- γ -induced in mouse peritoneal macrophages and encodes a putative GTP-binding protein. *J Leukoc Biol* 1995;57:477–483.
- 44 MacMicking JD: IFN-inducible GTPases and immunity to intracellular pathogens. *Trends Immunol* 2004;25:601–609.
- 45 Pilla DM, Hagar JA, Haldar AK, Mason AK, Degrandi D, Pfeffer K, Ernst RK, Yamamoto M, Miao EA, Coers J: Guanylate binding proteins promote caspase-11-dependent pyroptosis in response to cytoplasmic LPS. *Proc Natl Acad Sci U S A* 2014;111:6046–6051.
- 46 Shindo T, Kurihara H, Kuno K, Yokoyama H, Wada T, Kurihara Y, Imai T, Wang Y, Ogata M, Nishimatsu H, Moriyama N, Oh-hashu Y, Morita H, Ishikawa T, Nagai R, Yazaki Y, Matsushima K: ADAMTS-1: a metalloproteinase-disintegrin essential for normal growth, fertility, and organ morphology and function. *J Clin Invest* 2000;105:1345–1352.
- 47 Oveland E, Karlsen TV, Haslene-Hox H, Semaeva E, Janaczky B, Tenstad O, Wiig H: Proteomic evaluation of inflammatory proteins in rat spleen interstitial fluid and lymph during LPS-induced systemic inflammation reveals increased levels of ADAMTS1. *J Proteome Res* 2012;11:5338–5349.
- 48 Howell MD, Torres-Collado AX, Iruela-Arispe ML, Gottschall PE: Selective decline of synaptic protein levels in the frontal cortex of female mice deficient in the extracellular metalloproteinase ADAMTS1. *PLoS One* 2012;7:e47226.
- 49 Lee JH, Jung HS, Giang PM, Jin X, Lee S, Son PT, Lee D, Hong YS, Lee K, Lee JJ: Blockade of nuclear factor- κ B signaling pathway and anti-inflammatory activity of cardamomin, a chalcone analog from *Alpinia conchigera*. *J Pharmacol Exp Ther* 2006;316:271–278.
- 50 Kayagaki N, Warming S, Lamkanfi M, Vande Walle L, Louie S, Dong J, Newton K, Qu Y, Liu J, Heldens S, Zhang J, Lee WP, Roose-Girma M, Dixit VM: Non-canonical inflammasome activation targets caspase-11. *Nature* 2011;479:117–121.
- 51 Vanden Bergh T, Demon D, Bogaert P, Vandendriessche B, Goethals A, Depuydt B, Vuylsteke M, Roelandt R, Van Wouterghem E, Vandembroeck J, Choi SM, Meyer E, Krautwald S, Declercq W, Takahashi N, Cauwels A, Vandenabeele P: Simultaneous targeting of IL-1 and IL-18 is required for protection against inflammatory and septic shock. *Am J Respir Crit Care Med* 2014;189:282–291.