Review

Some Aspects of the Renin-Angiotensin-System in Hemodialysis Patients

Umar Malik Veena Raizada

University of New Mexico School of Medicine, Albuquerque, New Mexico, USA

Key Words

RAS • Renin-angiotensin system • Angiotensin II • ACE • ACE2 • Hemodialysis • Hypertension • ESRD • Cardiovascular disease

Abstract

Understanding of the renin-angiotensin system (RAS) has changed remarkably over the past decade. Renin, angiotensin converting enzyme (ACE), angiotensin II (Ang II), and Ang II receptors are the main components of the RAS. Recent studies identified the ACE2/Ang 1-7/Mas receptor axis, which counter-regulates the classical RAS. Many studies have examined the effects of the RAS on the progression of cardiovascular disease and chronic kidney disease (CKD). In addition, many studies have documented increased levels of ACE in hemodialysis (HD) patients, raising concerns about the negative effects of RAS activation on the progression of renal disease. Elevated ACE increases the level of Ang II, leading to vasoconstriction and cell proliferation. Ang II stimulation of the sympathetic system leads to renal and cardiovascular complications that are secondary to uncontrolled hypertension. This review provides an overview of the RAS, evaluates new research on the role of ACE2 in dialysis, and reviews the evidence for potentially better treatments for patients undergoing HD. Further understanding of the role of ACE and ACE2 in HD patients may aid the development of targeted therapies that slow the progression of CKD and cardiovascular disease.

© 2015 The Author(s)
Published by S. Karger AG, Basel

Introduction

In 1898, Robert Tigerstedt and his student Per Bergman at the Karolinska Institute in Sweden stumbled across a monumental discovery that forever changed our understanding of renal hemodynamic control. They identified a molecule from the renal tissue of a rabbit that increased blood pressure in nephrectomized animals [1] and named this substance “renin”, after the organ from which it was derived. This discovery led to a widespread search

for additional compounds with similar characteristics, and eventually to a full elucidation of the now classic renin-angiotensin system (RAS). Our understanding of this system has grown remarkably since these early discoveries. We now know that local RAS pathways exist in diverse tissues, as well as the kidneys, and this further complicates treatment regimens for patients with chronic kidney disease (CKD) [2]. Moreover, renin can activate local RAS systems by binding to its receptor in selected tissues [3]. In addition, research of the various angiotensin II (Ang II) receptors indicated that Ang II can have positive and negative downstream effects on circulation [4]. Most recently, the discovery of angiotensin converting enzyme 2 (ACE2) led to investigations of its counter-regulatory role compared to ACE [5, 6].

These advances in basic research have been matched by the development of therapies that inhibit RAS medications that can improve patient outcomes due to their inhibition of several disease processes. For example, a multicenter, prospective, observational study by Yamamoto et al. showed that patients on hemodialysis who used ACE inhibitors (ACEIs) and aldosterone receptor blockers (ARBs) had a lower rate of fracture-related hospitalization [7]. Medications that inhibit RAS have a large impact on cardiovascular disease (CVD). Thus, Ikeda et al. showed that patients with Stage 5 CKD who used ARBs had a lower prevalence of congestive heart failure (CHF) [8] and Tang et al. showed that long-term hemodialysis (HD) patients with heart failure had reduced all-cause and cardiovascular mortality when they were given RAS blockade with ACEI and/or ARB therapy [9]. Interestingly, a retrospective cohort study of patients with left ventricular ejection fraction (LVEF) of 45% or less (n = 45) showed that only 31% received an ACEI and only 9% received an ARB [10]. A double-blind, placebo-controlled, multicenter trial by Cice et al. showed that the addition of telmisartan (an ARB) with an ACEI reduced all-cause mortality, cardiovascular death, and hospitalization for heart failure in patients with LVEF of 40% or less who were on HD [11]. In addition, a retrospective cohort study by Shireman et al. showed that use of antihypertensive medications -- including ACEIs or ARBs -- reduced the adjusted hazard ratios for mortality in HD patients [12].

Although therapies that inhibit RAS provide important benefits to patients on HD, they also have negative side effects. For example, hyperkalemia is associated with ACEI and ARB therapy due to downstream effects on aldosterone, which normally increases uptake of sodium at the expense of potassium. Interestingly, Han et al. showed that use of an ACEI and/or ARB did not increase the risk of hyperkalemia in patients on maintenance HD [13]. The RAS also plays an important role in the secretion of erythropoietin. Vlahakos et al. demonstrated that the RAS stimulated erythropoietin secretion in HD patients [14]. In addition, a systematic review of 29,061 patients by Cheungpasitporn et al. showed an association between development of anemia and use of medications that inhibit RAS, including ACEIs and ARBs [15].

Perhaps of most interest, the alterations in RAS activity can affect the progression of CKD and CVD. Luciano et al. highlight the importance of vitamin D in the progression of CKD, citing evidence for vitamin D analogs reducing proteinuria by suppressing RAS activity [16]. In addition, numerous renin and ACE gene polymorphisms increase the risk for coronary artery disease (CAD) [17, 18]. Lei et al. showed that gene polymorphisms of ACE increased the risk of CAD in individuals with Type II diabetes mellitus (DM) [19] and Kato et al. demonstrated that gene polymorphisms of ACE increased the risk of CVD in individuals with hypertension [20].

In the early 1980's many studies evaluated the discovery of varying ACE levels in HD patients. These studies (see below and Table 1) had conflicting results, and the explanations for these findings were incomplete at that time. Data on ACE activity in HD patients remains limited, but our increasing understanding of the complicated nature of the RAS and its effects in CKD and CVD, as described above, justify exploration of possible mechanisms for these interesting findings.

The RAS has significant effects on CVD and residual renal function. Elevated ACE is important because it increases the level of Ang II, which promotes the progression of CKD and CVD through

vasoconstriction and cell proliferation. In addition, sympathetic activation *via* Ang II leads to uncontrolled hypertension that has negative effects on the renal and cardiovascular systems. Moreover, elevated ACE and increased Ang II lead to secretion of aldosterone from the adrenal cortex. However, the function and importance of aldosterone are beyond the scope of this review, and we will focus on the renin-angiotensin system (RAS) and its importance to patients on HD.

Table 1. Initial studies that measured plasma or serum levels of angiotensin converting enzyme in different patient populations

Authors	Plasma or Serum ACE	Patient population
Patel et al. 1979	Serum	CKD on HD
Muira et al. 1984	Serum	CKD on HD
Silverstein et al. 1984	Serum	CKD on HD
Nielsen et al. 1985	Serum	CKD on HD
Letizia et al. 1995	Serum	CKD on HD
Docci et al. 1988	Serum	CKD on HD
Schweisfurth et al. 1979	Serum	Hepatitis/liver cirrhosis
Lieberman et al. 1980	Serum	Diabetes Mellitus
Koo et al. 1987	Plasma	CKD on HD
Anguiano et al. 2015	Plasma	CKD Stage 3-5

Abbreviations: ACE: angiotensin converting enzyme; CKD: chronic kidney disease; HD: hemodialysis.

Elevated ACE in Hemodialysis Patients

In 1979, Patel et al. performed one of the first and best known evaluations of the levels of ACE in HD patients [21]. They studied 19 patients with CKD on long-term HD and 19 controls, and found that ACE activity was increased in 58% of the patients [21]. They proposed that these increased serum ACE levels might be secondary to increased pulmonary vascular area in patients with renal failure, because ACE was thought to be produced only in the lungs. In 1982, Muira et al. confirmed these results but proposed a different mechanism. They noted that the ACE activity of patients with CKD on regular HD was higher than that of age-matched controls [22], and they also found higher ACE levels in patients on HD for longer than 3 years and in patients with diabetic nephropathy. It is important to note that elevated ACE also occurs in DM [23]. Muira et al. found no relationships of enzyme activity with age and sex [22]. Based on studies reporting accelerated atherosclerosis in patients on maintenance HD, they speculated that advanced endothelial damage could lead to elevated serum ACE in HD patients, because vascular endothelial cells of the lungs and kidneys produce most of the ACE.

Additional studies have found similar results, but only select articles postulated possible mechanisms. Silverstein et al. showed elevated serum ACE in 16 of 48 non-HD patients ($p < 0.001$) who had various types of CKD [24]. They also found increased serum ACE in 17 of 52 patients on maintenance HD ($p < 0.001$) in a comparison with healthy adult controls. Nielsen et al. showed that serum ACE levels increased by 14.3% ($p < 0.001$) in patients after a 240 min session of HD [25]. They suggested that increased serum ACE was due to a complement-mediated sequestration of leukocytes within the pulmonary vasculature, as previously described by Craddock et al. [26]. This accumulation of leukocytes leads to injury of the vascular endothelium with interstitial edema and subsequently to pulmonary abnormalities, as also postulated by Muira et al. [22]. Koo et al. showed that serum ACE was significantly elevated in patients with CKD at 5 h after an HD session ($p < 0.001$) [27]. They suggested that elevated ACE in HD patients is more closely related to the HD procedure than to the renal disease itself. In support, Craddock et al. have shown complement-mediated pulmonary dysfunction in patients after HD sessions [28].

Letizia et al. showed that patients with CKD on HD and patients who received renal transplantation had significantly higher serum ACE activity than healthy controls [29]. Plasma renin activity (PRA) and plasma aldosterone (PA) also increased after an HD session ($p < 0.026$ and $p < 0.044$, respectively). Iitake et al. also documented elevations of PRA in

HD patients [30]. These elevations in PRA raise additional concerns regarding the negative effects of the RAS on HD patients. As such, research into renin inhibitors has produced some promising results regarding the control of these effects in HD patients. Ishimitsu et al. showed that PRA decreased following administration of aliskiren (a direct renin inhibitor) and this was accompanied by a modest decrease in blood pressure [31]. Morishita et al. also demonstrated that use of aliskiren was associated with decreased BP and cardio-protective effects in hypertensive patients on long-term HD [32]. Ito et al. showed that aliskiren and an ARB had similar effects on BP control, but there was a significant decrease in PRA with aliskiren but not with an ARB [33]. In addition, Moriya et al. demonstrated that aliskiren improved vascular endothelial function in patients on hemodialysis [34]. These findings were independent of antihypertensive effect from the medication. Advances by Xiong et al. have demonstrated the importance of the CD38 gene in mice on the intracellular regulation of renin production [35]. This study provides some insight into this poorly understood mechanism and may highlight future studies aimed at developing more targeted therapies for suppressing PRA. Overall, these studies illustrate the importance of appropriate RAS blockade for patients on long-term HD.

In contrast to Koo et al., Letizia's group found no significant difference in serum ACE activity before and after an HD session. Ulrich et al. examined ACE expression in monocytes of HD patients with CVD and controls with normal renal function and found that HD patients had greater ACE expression [36]. Docci et al. compared the effect of different dialyzer membranes on serum ACE, and found that all tested membranes caused similar increases in ACE. They proposed that the elevation of ACE was less a cause of the HD procedure itself, and instead was secondary to hemoconcentration [37]. Table 1 summarizes the above studies that examined serum and plasma levels of ACE in various patient populations.

Clinical Implications

The increased ACE in HD patients and the consequent elevation of Ang II can promote the progression of CVD and renal disease. Patients on maintenance HD often suffer from hypertension at baseline, and ultrafiltration during the HD procedure typically normalizes blood pressure by reducing volume overload. However, some patients on HD have the opposite response, and suffer from hypertension during or after the HD procedure. Previous studies have postulated several possible mechanisms for this intradialytic hypertension, including electrolyte imbalance, removal of antihypertensive medications during HD, and activation of the RAS [38-41]. RAS activation can occur due to the decreased intravascular volume during the ultrafiltration process of HD [42, 43]. Activation of RAS leads to subsequent elevation of Ang II, and this enzyme can cause vasoconstriction, salt and water retention mediated *via* aldosterone, and cell proliferation. In addition, sympathetic activation occurs because Ang II acts on Ang II receptor type 1 (AT1 receptor) in the central nervous system, leading to increased systemic vascular resistance [44]. As such, Ang II promotes hypertension through water retention, vasoconstriction, and sympathetic activation, underlining the important role of ACE inhibitors in the treatment of hypertension in HD patients. Thus, treatment of high blood pressure in HD patients with elevated ACE may be particularly challenging because increased activation of local-RAS systems -- independent of the primary RAS -- may contribute to hypertension that is sustained and difficult to treat.

An excess of Ang II may also be deleterious in other patient groups. For example, elevated Ang II increases the morbidity and mortality of patients with CHF despite ACE inhibitor therapy [45]. This indicates the growing need to understand RAS activation in renal diseases and CVDs. ACE is upregulated in the proximal tubules of rats with proteinuria, likely resulting in further renal damage and progression of renal disease [46]. Sustained progression of renal and CVD in HD patients may also be linked to increased proteinuria, so therapies that decrease proteinuria may benefit such patients.

Local-RAS pathways

Our review of the role of the RAS in HD patients will conclude with a discussion of the local RAS, one of the most recent discoveries in this field. It is now widely accepted that the kidneys themselves have a local RAS. Although the liver produces most of the angiotensinogen (Agt), epithelial cells of the proximal tubule also synthesize this hormone, and it can even be detected in urine [52]. The collecting tubule is another source of renin, and renin expression is linked with stimulation of Ang II [53, 54]. ACE occurs in the brush border membrane of the proximal tubule [52]. Thus, the presence of these important compounds may lead to intrarenal production of Ang II. This local system may affect serum ACE levels (*i.e.* at the level of the whole body) in patients with renal disease. Interestingly, studies have shown that patients given ACE inhibitors still have elevated intrarenal Ang II production. This indicates a role of other pathways in the conversion of Ang I to Ang II [55]. The intrarenal-RAS is not yet fully understood and the role of this system in patients undergoing HD is unknown. A more complete understanding of the intrarenal-RAS is needed because it can alter ACE levels in patients with CKD who are on maintenance HD.

Conclusion

In this article, we reviewed numerous studies that examined the effect of elevated ACE in patients with CKD who were undergoing HD. There are several possible explanations for the elevated serum ACE in these patients. For example, existing underlying diseases, such as DM and liver disease, may contribute to the elevated ACE in these patients [23, 56]. Another possibility is that the diseased kidney secretes ACE into the serum, although the evidence for this is weak. In fact, Patel et al. found that plasma ACE was elevated in patients with bilateral nephrectomies [21]. It is possible that elevated ACE in HD patients is due to decreased degradation of the enzyme or decreased excretion by the kidney. Several of the studies reviewed here suggest that elevated ACE may be due to the effects of HD on the pulmonary vasculature *via* complement-mediated pathways [26, 28].

Another possibility is that ACE is elevated in patients with CKD due to disruption of its negative feedback loop by Ang II [57]. It is possible that dysregulation of the negative feedback mechanism of Ang II upon renin in the presence of CKD may increase RAS activity and the levels of ACE [58]. As discussed above, prorenin itself can activate local-RAS pathways through the pro-renin receptor (PRR) [3]. This activation may further increase serum ACE levels, secondary to the dysregulation of the negative feedback mechanism of the RAS. The intrarenal-RAS may affect whole body levels of ACE through diffusion abnormalities within the diseased kidney, although this requires further study. Immuno-staining experiments showed increased ACE expression within the glomerular and renal vasculature of rats with DM and albuminuria [59]. Soler et al. also showed increased ACE expression in the renal vessels of diabetic mice [60] and immunohistochemical studies in patients with non-insulin dependent DM indicated increased ACE in those with nephropathy [61].

It is important to know whether the elevation of ACE in HD patients is due to renal disease itself, the HD procedure, or both. However, the mechanisms underlying the elevation of ACE are likely multifaceted and related to multiple interactions. It is important to understand these mechanisms in order to develop better treatments, slow disease progression, and reduce the severity of CKD and CVD in HD patients. Although the RAS is associated with development of hypertension and DM, ACE/ACE2 levels are not directly related to the progression of CKD. Hypertension and DM develop as consequences of many different factors and have many potential etiologies. Future studies are needed to provide answers to these questions and to develop new questions in our quest to better understand, treat, and prevent these diseases. Further evaluation and determination of these mechanisms is imperative in our ever-changing understanding of the once simple RAS, originally discovered by Tigerstedt and Bergman.

Disclosure Statement

The authors of this review have no financial or nonfinancial relationships to disclose.

References

- 1 Tigerstedt R, Bergman PG: Niere und Kreislauf. *Skand Arch Physiol* 1898;8:223-271.
- 2 Kobori H, Nangaku M, Navar LG, Nishiyama A: The intrarenal renin-angiotensin system: from physiology to the pathobiology of hypertension and kidney disease. *Pharmacol Rev* 2007;59:251-287.
- 3 Nguyen G, Delarue F, Burcklé C, Bouzahir L, Giller T, Sraer J-D: Pivotal role of the renin/prorenin receptor in angiotensin II production and cellular responses to renin. *J Clin Invest* 2002;109:1417-1427.
- 4 Yayama K, Okamoto H: Angiotensin II-induced vasodilation via type 2 receptor: role of bradykinin and nitric oxide. *Int Immunopharmacol* 2008;8:312-318.
- 5 Wysocki J, Battle D: Reduced plasma ACE2 activity in dialysis patients: another piece in the conundrum of factors involved in hypertension and cardiovascular morbidity? *Nephrol Dial Transplant* 2013;28:2200-2202.
- 6 Roberts MA, Velkoska E, Ierino FL, Burrell LM: Angiotensin-converting enzyme 2 activity in patients with chronic kidney disease. *Nephrol Dial Transplant* 2013;28:2287-2294.
- 7 Yamamoto S, Kido R, Onishi Y, Fukuma S, Akizawa T, Fukagawa M, Kazama JJ, Narita I, Fukuhara S: Use of renin-angiotensin system inhibitors is associated with reduction of fracture risk in hemodialysis patients. *PLoS ONE* 2015;10:e0122691.
- 8 Ikeda M, Nakao M, Hirano K, Yokoyama K, Yokoo T, Joki N, Ando R, Shinoda R, Inaguma D, Yamaka T, Komatsu Y, Koiwa F, Sakaguchi T, Negi S, Shigematsu T: Possible prevention of dialysis-requiring congestive heart failure by angiotensin-II receptor blockers in non-dialysis Japanese patients with Stage 5 chronic kidney disease. *J Renin Angiotensin Aldosterone Syst* 2015, in press.
- 9 Tang C-H, Chen T-H, Wang C-C, Hong C-Y, Huang K-C, Sue Y-M: Renin-angiotensin system blockade in heart failure patients on long-term haemodialysis in Taiwan. *Eur J Heart Fail* 2013;15:1194-1202.
- 10 Derthoo D, Belmans A, Claes K, Bammens B, Ciarka A, Droogné W, Vanhaecke J, Van Cleemput J, Janssens S: Survival and heart failure therapy in chronic dialysis patients with heart failure and reduced left ventricular ejection fraction: an observational retrospective study. *Acta Cardiol* 2013;68:51-57.
- 11 Cice G, Di Benedetto A, D'Isa S, D'Andrea A, Marcelli D, Gatti E, Calabro R: Effects of telmisartan added to Angiotensin-converting enzyme inhibitors on mortality and morbidity in hemodialysis patients with chronic heart failure a double-blind, placebo-controlled trial. *J Am Coll Cardiol* 2010;56:1701-1708.
- 12 Shireman TI, Phadnis MA, Wetmore JB, Zhou X, Rigler SK, Spertus JA, Ellerbeck EF, Mahnken JD: Antihypertensive medication exposure and cardiovascular outcomes in hemodialysis patients. *Am J Nephrol* 2014;40:113-122.
- 13 Han S-W, Won Y-W, Yi J-H, Kim H-J: No impact of hyperkalemia with renin-angiotensin system blockades in maintenance haemodialysis patients. *Nephrol Dial Transplant* 2007;22:1150-1155.
- 14 Vlahakos DV, Balodimos C, Papachristopoulos V, Vassilakos P, Hinari E, Vlachojannis JG: Renin-angiotensin system stimulates erythropoietin secretion in chronic hemodialysis patients. *Clin Nephrol* 1995;43:53-59.
- 15 Cheungpasitporn W, Thongprayoon C, Chiasakul T, Korpaisarn S, Erickson SB: Renin-angiotensin system inhibitors linked to anemia: a systematic review and meta-analysis. *QJM* 2015;108:879-884.
- 16 Lucisano S, Buemi M, Passantino A, Aloisi C, Cernaro V, Santoro D: New insights on the role of vitamin D in the progression of renal damage. *Kidney Blood Press Res* 2013;37:667-678.
- 17 Fragoso JM, Alvarez-León E, Delgadillo-Rodríguez H, Arellano-González M, López-Pacheco FC, Cruz-Robles D, Peña-Duque MA, Pérez-Méndez O, Martínez-Ríos MA, Vargas-Alarcón G: The C4280A (rs5705) gene polymorphism of the renin (REN) gene is associated with risk of developing coronary artery disease, but not with restenosis after coronary stenting. *Exp Mol Pathol* 2015;99:128-132.
- 18 Martínez-Ríos MA, Alvarez-León E, Totomoch A, Angeles J, Peña-Duque MA, Delgadillo-Rodríguez H, Martínez-Rodríguez N, Ramírez-Fuentes S, Fragoso JM, Vargas-Alarcón G: Haplotypes of the angiotensin-converting enzyme (ACE) gene are associated with coronary artery disease but not with restenosis after coronary stenting. *Exp Mol Pathol* 2014;97:166-170.

- 19 Lei H-P, Chen H-M, Zhong S-L, Yao Q-Z, Tan H-H, Yang M, Lin QX, Shan ZX, Zheng ZW, Zhu JN, Zhou ZL, Lin SG, Yu XY: Association between polymorphisms of the renin-angiotensin system and coronary artery disease in Chinese patients with type 2 diabetes. *J Renin Angiotensin Aldosterone Syst* 2012;13:305-313.
- 20 Kato N, Tatara Y, Ohishi M, Takeya Y, Onishi M, Maekawa Y, Rakugi H: Angiotensin-converting enzyme single nucleotide polymorphism is a genetic risk factor for cardiovascular disease: a cohort study of hypertensive patients. *Hypertens Res* 2011;34:728-734.
- 21 Patel R, Ansari A: Serum angiotensin converting enzyme activity in patients with chronic renal failure on long term hemodialysis. *Clin Chim Acta* 1979;92:491-495.
- 22 Miura H, Nakayama M, Sato T: Serum angiotensin converting enzyme (S-ACE) activity in patients with chronic renal failure on regular hemodialysis. *Jpn Heart J* 1984;25:87-92.
- 23 Lieberman J, Sastre A: Serum angiotensin-converting enzyme: elevations in diabetes mellitus. *Ann Intern Med* 1980;93:825-826.
- 24 Silverstein E, Brunswick J, Rao TK, Friedland J: Increased serum angiotensin-converting enzyme in chronic renal disease. *Nephron* 1984;37:206-210.
- 25 Nielsen AH, Knudsen F, Kristensen SD: Serum angiotensin-converting enzyme increases during hemodialysis. Evidence for injury of the pulmonary vascular endothelium? *Nephron* 1985;40:100-103.
- 26 Craddock PR, Fehr J, Brigham KL, Kronenberg RS, Jacob HS: Complement and leukocyte-mediated pulmonary dysfunction in hemodialysis. *N Engl J Med* 1977;296:769-774.
- 27 Koo WS, Lee YJ, Kim HS, Kim SY, Choi EJ, Chang YS, Yoon YS, Bang BK: Changes of plasma angiotensin-converting enzyme activity during hemodialysis. *Korean J Intern Med* 1987;2:62-65.
- 28 Craddock PR, Fehr J, Dalmaso AP, Brigham KL, Jacob HS: Hemodialysis leukopenia. Pulmonary vascular leukostasis resulting from complement activation by dialyzer cellophane membranes. *J Clin Invest* 1977;59:879-888.
- 29 Letizia C, Mazzaferro S, Morabito S, De Ciocchis A, Cerci S, D'Ambrosio C, Cinotti GA, Scavo D: Response of serum angiotensin converting enzyme, plasma renin activity and plasma aldosterone to conventional dialysis in patients on chronic haemodialysis. *Int Urol Nephrol* 1995;27:465-470.
- 30 Iitake K, Kimura T, Matsui K, Ota K, Shoji M, Inoue M, Yoshinaga K: Effect of haemodialysis on plasma ADH levels, plasma renin activity and plasma aldosterone levels in patients with end-stage renal disease. *Acta Endocrinol* 1985;110:207-213.
- 31 Ishimitsu T, Ohta S, Ohno E, Takahashi T, Numabe A, Okamura A, Ohba S, Hashimoto A, Matsuoka H: Long-term antihypertensive effects of aliskiren, a direct renin inhibitor, in chronic hemodialysis patients. *Ther Apher Dial* 2013;17:524-531.
- 32 Morishita Y, Hanawa S, Chinda J, Iimura O, Tsunematsu S, Kusano E: Effects of aliskiren on blood pressure and the predictive biomarkers for cardiovascular disease in hemodialysis-dependent chronic kidney disease patients with hypertension. *Hypertens Res* 2011;34:308-313.
- 33 Ito T, Ishikawa E, Fujimoto N, Okubo S, Ito G, Ichikawa T, Nomura S, Ito M: Effects of aliskiren on blood pressure and humoral factors in hypertensive hemodialysis patients previously on angiotensin II receptor antagonists. *Clin Exp Hypertens* 2014;36:497-502.
- 34 Moriya H, Kobayashi S, Ohtake T, Tutumi D, Mochida Y, Ishioka K, Oka M, Maesato K, Hidaka S, Nomura S: Aliskiren, a direct renin inhibitor, improves vascular endothelial function in patients on hemodialysis independent of antihypertensive effect ~ a pilot study~. *Kidney Blood Press Res* 2013;37:190-198.
- 35 Xiong J, Xia M, Yi F, Abais JM, Li N, Boini KM, Li PL: Regulation of renin release via cyclic ADP-ribose-mediated signaling: evidence from mice lacking CD38 gene. *Cell Physiol Biochem* 2013;31:44-55.
- 36 Ulrich C, Heine GH, Garcia P, Reichart B, Georg T, Krause M, Köhler H, Girndt M: Increased expression of monocytic angiotensin-converting enzyme in dialysis patients with cardiovascular disease. *Nephrol Dial Transplant* 2006;21:1596-1602.
- 37 Docci D, Delvecchio C, Turci F, Baldrati L, Gollini C: Effect of different dialyzer membranes on serum angiotensin-converting enzyme during hemodialysis. *Int J Artif Organs* 1988;11:28-32.
- 38 Dousdampanis P, Trigka K, Giannopoulos A, Fourtounas C: Pathogenesis and management of intradialytic hypertension. *Curr Hypertens Rev* 2014;10:171-176.
- 39 Dolson GM, Ellis KJ, Bernardo MV, Prakash R, Adrogué HJ: Acute decreases in serum potassium augment blood pressure. *Am J Kidney Dis* 1995;26:321-326.
- 40 Matzke GR, Aronoff GR, Atkinson AJ, Bennett WM, Decker BS, Eckardt KU, Golper T, Grabe DW, Kasiske B, Keller F, Kielstein JT, Mehta R, Mueller BA, Pasko DA, Schaefer F, Sica DA, Inker LA, Umans JG, Murray P:

- Drug dosing consideration in patients with acute and chronic kidney disease-a clinical update from Kidney Disease: Improving Global Outcomes (KDIGO). *Kidney Int* 2011;80:1122-1137.
- 41 Hörl MP, Hörl WH: Hemodialysis-associated hypertension: pathophysiology and therapy. *Am J Kidney Dis* 2002;39:227-244.
- 42 Locatelli F, Cavalli A, Tucci B: The growing problem of intradialytic hypertension. *Nat Rev Nephrol* 2010;6:41-48.
- 43 Inrig JK: Intradialytic hypertension: a less-recognized cardiovascular complication of hemodialysis. *Am J Kidney Dis* 2010;55:580-589.
- 44 Zucker IH, Gao L: The regulation of sympathetic nerve activity by angiotensin II involves reactive oxygen species and MAPK. *Circ Res* 2005;97:737-739.
- 45 Roig E, Perez-Villa F, Morales M, Jiménez W, Orús J, Heras M, Sanz G: Clinical implications of increased plasma angiotensin II despite ACE inhibitor therapy in patients with congestive heart failure. *Eur Heart J* 2000;21:53-57.
- 46 Largo R, Gómez-Garre D, Soto K, Marrón B, Blanco J, Gazapo RM, Plaza JJ, Egido J: Angiotensin-converting enzyme is upregulated in the proximal tubules of rats with intense proteinuria. *Hypertension* 1999;33:732-739.
- 47 Zhang Y, Liu J, Luo J-Y, Tian XY, Cheang WS, Xu J, Lau CW, Wang L, Wong WT, Wong CM, Lan HY, Yao X, Raizada MK, Huang Y: Upregulation of Angiotensin (1-7)-Mediated Signaling Preserves Endothelial Function Through Reducing Oxidative Stress in Diabetes. *Antioxid Redox Signal* 2015;23:880-892.
- 48 Jin H-Y, Song B, Oudit GY, Davidge ST, Yu H-M, Jiang Y-Y, Gao PJ, Zhu DL, Ning G, Kassiri Z, Penninger JM, Zhong JC: ACE2 deficiency enhances angiotensin II-mediated aortic profilin-1 expression, inflammation and peroxynitrite production. *PLoS ONE* 2012;7:e38502.
- 49 Mizuiri S, Hemmi H, Arita M, Ohashi Y, Tanaka Y, Miyagi M, Sakai K, Ishikawa Y, Shibuya K, Hase H, Aikawa A: Expression of ACE and ACE2 in individuals with diabetic kidney disease and healthy controls. *Am J Kidney Dis* 2008;51:613-623.
- 50 Anguiano L, Riera M, Pascual J, Valdivielso JM, Barrios C, Betriu A, Mojal S, Fernández E, Soler MJ: Circulating angiotensin-converting enzyme 2 activity in patients with chronic kidney disease without previous history of cardiovascular disease. *Nephrol Dial Transplant* 2015;30:1176-1185.
- 51 Da Silveira KD, Pompermayer Bosco KS, Diniz LRL, Carmona AK, Cassali GD, Bruna-Romero O, de Sousa LP, Teixeira MM, Santos RA, Simões e Silva AC, Ribeiro Vieira MA: ACE2-angiotensin-(1-7)-Mas axis in renal ischaemia/reperfusion injury in rats. *Clin Sci* 2010;119:385-394.
- 52 Kobori H, Ozawa Y, Suzaki Y, Prieto-Carrasquero MC, Nishiyama A, Shoji T, Cohen EP, Navar LG: Young Scholars Award Lecture: Intratubular angiotensinogen in hypertension and kidney diseases. *Am J Hypertens* 2006;19:541-550.
- 53 Kang JJ, Toma I, Sipos A, Meer EJ, Vargas SL, Peti-Peterdi J: The collecting duct is the major source of prorenin in diabetes. *Hypertension* 2008;51:1597-1604.
- 54 Prieto-Carrasquero MC, Kobori H, Ozawa Y, Gutiérrez A, Seth D, Navar LG: AT1 receptor-mediated enhancement of collecting duct renin in angiotensin II-dependent hypertensive rats. *Am J Physiol Renal Physiol* 2005;289:F632-F637.
- 55 Nishiyama A, Seth DM, Navar LG: Renal interstitial fluid angiotensin I and angiotensin II concentrations during local angiotensin-converting enzyme inhibition. *J Am Soc Nephrol* 2002;13:2207-2212.
- 56 Schweisfurth H, Wernze H: Changes of serum angiotensin I converting enzyme in patients with viral hepatitis and liver cirrhosis. *Acta Hepatogastroenterol (Stuttg)* 1979;26:207-210.
- 57 Schunkert H, Ingelfinger JR, Hirsch AT, Pinto Y, Remme WJ, Jacob H, Dzau VJ: Feedback regulation of angiotensin converting enzyme activity and mRNA levels by angiotensin II. *Circ Res* 1993;72:312-318.
- 58 Atlas SA: The renin-angiotensin aldosterone system: pathophysiological role and pharmacologic inhibition. *J Manag Care Pharm* 2007;13:9-20.
- 59 Anderson S, Jung FF, Ingelfinger JR: Renal renin-angiotensin system in diabetes: functional, immunohistochemical, and molecular biological correlations. *Am J Physiol* 1993;265:F477-F486.
- 60 Soler MJ, Wysocki J, Ye M, Lloveras J, Kanwar Y, Battle D: ACE2 inhibition worsens glomerular injury in association with increased ACE expression in streptozotocin-induced diabetic mice. *Kidney Int* 2007;72:614-623.
- 61 Mizuiri S, Yoshikawa H, Tanegashima M, Miyagi M, Kobayashi M, Sakai K, Hayashi I, Aikawa A, Ohara T, Hasegawa A: Renal ACE immunohistochemical localization in NIDDM patients with nephropathy. *Am J Kidney Dis* 1998;31:301-307.