Chromosome Segregation in *Vibrio cholerae*

Revathy Ramachandran Jyoti Jha Dhruva K. Chattoraj

Laboratory of Biochemistry and Molecular Biology, Center for Cancer Research, NCI, National Institutes of Health, Bethesda, Md., USA

Key Words

Bacterial mitosis · Divided genomes · Coordination of replication and segregation · Par-independent polar segregation

Abstract

The study of chromosome segregation is currently one of the most exciting research frontiers in cell biology. In this review, we discuss our current knowledge of the chromosome segregation process in *Vibrio cholerae*, based primarily on findings from fluorescence microscopy experiments. This bacterium is of special interest because of its eukaryotic feature of having a divided genome, a feature shared with 10% of known bacteria. We also discuss how the segregation mechanisms of *V. cholerae* compare with those in other bacteria, and highlight some of the remaining questions regarding the process of bacterial chromosome segregation.

© 2015 S. Karger AG, Basel

Introduction

The three main events of the cell cycle are chromosome replication, chromosome segregation, and cell division. Coordination of these processes is essential for efficient maintenance of the genome during cell prolifera-

tion. Because these processes are functionally integrated, an effective understanding of any one of them is only possible in the context of the other events of the cell cycle. Therefore, in our discussion of chromosome segregation we try to make connections to chromosome replication and cell division.

In bacteria, chromosome segregation is not as well understood as chromosome replication and cell division. In eukaryotes, chromosomes can be visualized using a simple microscope, and mitosis was described more than a century ago. In contrast, visualization of chromosome dynamics in bacteria was realized only a decade or two ago with the introduction of fluorescence microscopy, which allowed visualization of smaller objects such as specific proteins and DNA loci [Webb et al., 1997]. It is now clear that even in apparently compartment-less organisms such as bacteria, there is considerable spatiotemporal order in the organization and segregation of chromosomes [Niki et al., 2000; Possoz et al., 2012; Wang et al., 2013; Wiggins et al., 2010]. These cytological studies are also contributing to the understanding of how the segregation process is coordinated with chromosome replication and cell division. Just as great strides in molecular biology and biochemistry owe a great deal to X-ray crystallography for the discovery of DNA structure, the illumination of the inner workings of the bacterial cell in space and time, and ‘bacterial mitosis’ in particular, owe a great deal to fluorescence microscopy.

Segregation of *Vibrio cholerae* Chromosomes

V. cholerae is closely related to *Escherichia coli*, except that the *V. cholerae* genome is divided into two chromosomes, chr1 and chr2. The close relationship has contributed both conceptually and experimentally to studies in *V. cholerae* and has made it the leading system for studying chromosome dynamics in bacteria with divided genomes.

The two chromosomes of *V. cholerae* are circular, ~3 (chr1) and 1.1 (chr2) Mbp long. The replication systems of chr1 and chr2 are distinct from each other and regulated by replicon-specific initiators, similarly to the situation for the *E. coli* chromosome and one of its plasmid, P1 [Egan and Waldor, 2003]. Both chromosomes encode the three-component *parABS* segregation system. This system was first discovered and studied in plasmids, and is found in the chromosomes of about 70% of sequenced bacteria [Gerdes et al., 2000; Livny et al., 2007]. The *parABS* system allows active mobilization of centromeric sites, which comprise the *parS* component. ParB is a *parS*-specific DNA-binding protein, and ParA is an ATPase that activates segregation. Although the *par* systems in general are highly homologous, including the ones in *V. cholerae* (*parABS1* and *parABS2*) [Gerdes et al., 2000], they function in a chromosome-specific fashion [Dubarry et al., 2006; Kadoya et al., 2011; Saint-Dic et al., 2006]. Underlying this specificity are differences in *parS* sequences that ensure binding to cognate ParB proteins [Yamaichi et al., 2007a]. The ParB-*parS* complexes in turn also recognize their cognate ParA partner through specific protein-protein interactions [Radnedge et al., 1998].

The choreography of chromosome dynamics in *V. cholerae* has been studied in live cells by tagging chromosomal loci with either operator arrays or the plasmid ParB-*parS* systems [David et al., 2014; Fiebig et al., 2006; Fogel and Waldor, 2006; Srivastava and Chattoraj, 2007]. The localization patterns of *parS* sites (and the nearby origin region) are different for the two chromosomes. In newborn cells that are obtained under slow growth conditions, the chr1 origin *ori1* is found near the old pole. After duplication, *ori1* copies segregate asymmetrically; one *ori1* stays at the pole of duplication and the other moves to the opposite pole in a directed fashion, a process requiring hydrolysis of ParA1-bound ATP (fig. 1) [Fogel and Waldor, 2006]. In contrast, the chr2 origin *ori2* is found at mid-cell. After duplication, *ori2* copies segregate symmetrically to cell quarter positions in a *parAB2*-dependent fashion [Yamaichi et al., 2007b]. Under rapid

growth conditions, cells are born with two copies of *ori1*, one at each pole, but with one *ori2* at mid-cell [Srivastava and Chattoraj, 2007]. Initiation from *ori1* occurs from both poles, and segregation of the duplicated copies appears similar in the two halves of the cell. The old and the new pole therefore appear equally efficient in initiating replication and segregating the sisters. This is noteworthy because in *Caulobacter crescentus*, whose chromosomal origin also migrates from one pole to the other pole, only the origin at the old pole fires in the next rounds [Sliushenko et al., 2011]. This is rationalized from the fact that two cell halves of *C. crescentus* have distinct behavioral characteristics, which is not the case in *V. cholerae*. The higher number of *ori1* over *ori2* (*ori1:ori2* = 2:1) in *V. cholerae* at fast growth rates is expected because the time to replicate the larger chr1 but not the smaller chr2 exceeds the cell generation time [Skarstad et al., 1985; Srivastava and Chattoraj, 2007; Stokke et al., 2011].

Chromosome segregation studies have revealed that the origin is the first region that segregates, even in bacteria that do not have *parABS* system, and this is believed to set the course of separation for the remainder of the chromosome [Wang et al., 2013]. The region where replication terminates (*ter*) has also drawn special attention because its segregation makes room for cell division. Complete segregation of sister chromosomes clears the mid-cell area and allows septum closure without steric hindrance from the nucleoid [Bernard et al., 2010; Cambridge et al., 2014]. In newborn *V. cholerae* cells, the chr1 *ter* (*ter1*) is found at the new pole opposite to the pole where *ori1* is found (fig. 1) [Srivastava et al., 2006]. The *ter1* region then gradually migrates to the mid-cell in older cells, where following replication the copies remain together until the time of septum closure. In contrast, the duplicated copies of chr2 *ter* (*ter2*), which are formed at about the same time as *ter1* [Rasmussen et al., 2007], can segregate away from the mid-cell significantly before septum formation. However, in a significant fraction of cells, *ter2* copies also remain together at the mid-cell until the time of septum closure, similar to the *ter1* copies.

The reason for the heterogeneous behavior of *ter2* and the basis for the different timing of separation of *ter1* and *ter2* appear to be due to the differential role that the MatP/*matS* system plays at the two termini [Demarre et al., 2014]. The MatP/*matS* system was discovered in *E. coli* and is specifically involved in organizing the *ter* region [Mercier et al., 2008]. The *E. coli* chromosome appears to be divided into a few large domains (macrodomains) [see the review by Messerschmidt and Wald-

Fig. 1. Model of *parABS*-mediated segregation of *V. cholerae* chr1. The chromosome is represented by an ellipse, whose major axis has the ParB1-bound centromere (B) at one end and the replication terminus region (capped) at the other. In newborn cells, the centromere is found anchored to the old pole by the HubP (P) transmembrane polar protein via ParB1, ParA1 and HubP interactions. ParA1 is shown as a cloud whose density is maximal near the pole. After replication initiation and duplication of the nearby centromeric region, one of the daughter centromeres migrates towards the new pole where ParA1 is present at higher concentration, shown by a larger cloud. The cloud meets the ParB1/*parS1* centromeric complex when replication elongation has progressed further. The complex is then rapidly pulled to the pole by retraction of the ParA1 cloud. In predivisional (mother) cells, the terminus region is found together at mid-cell. Adapted from Fogel and Waldor [2006], Srivastava et al. [2006], and Yamaichi et al. [2012].

minghaus, this issue, pp. 301–315], one of which encompasses the *ter* region [Mercier et al., 2008; Valens et al., 2004]. Special sites (*matS*) in the *ter* macrodomain bind MatP, which causes compaction of the region, and in *E. coli*, tethers the domain to the closing division septum [Bailey et al., 2014]. *V. cholerae* has MatP, and both *ter1* and *ter2* regions have *matS* sites [Demarre et al., 2014]. Without MatP, both *ter* regions show segregation earlier than the onset of septal closing. The significance of the differential behavior of *ter1* and *ter2* remains to be understood. It is an intriguing issue for future investigation, given the timing and locations of replication termination are so similar for chr1 and chr2.

The segregation of the bulk of the chromosome has also been followed in some bacterial species, including *V. cholerae* [David et al., 2014]. The location of markers between *ori* and *ter* determines the overall topology of the chromosomes both during and between replication phases. In both chr1 and chr2, markers on the two replichores (the halves of the chromosome between *ori* and *ter* loci) are arranged along the longitudinal axis of the cell (fig. 2) [David et al., 2014]. The order of location of the markers follows the order in which they are present on the chromosome, and the markers segregate sequentially as they are replicated. Orientation of chr1 is dictated by *parS1*.

When *parS1* is displaced to an ectopic locus, it still migrates to the pole, causing rotation of the bulk of the chromosome. An exception is the *ter1* region that retains its normal mid-cell location when *parS1* is misplaced. As mentioned earlier, the *ter* region is specifically controlled by the MatP/*matS* system, which probably confines it to the mid-cell irrespective of the pull from *parS1*. Displacing *parS* also rotates the *C. crescentus* chromosome, including the *ter* region [Umbarger et al., 2011]. The rotation of *ter* was possibly because there is no MatP/*matS* system in *C. crescentus*.

The location of chromosome-wide markers also indicated the relative positions of the two *V. cholerae* chromosomes. In newborn cells, chr1 extends over the entire cell length, whereas chr2 occupies only the cell half containing the new pole (fig. 2). Chr2 thus overlaps the terminal half of chr1. Naked DNA occupies <1% of cell space, although the nucleoid could occupy ~75% of the cell space [Fisher et al., 2013]. Whether the overlap between the chromosomes causes any steric hindrance for their independent segregation remains to be addressed. Demixing of the two chromosomes is expected from entropic reasons [Jun and Wright, 2010]. Other forces (e.g. anchoring of *parS* and *ter*) might be at work as well.

Fig. 2. Spatial arrangement of chromosomes in newborn bacterial cells. Note that the locations of the origin/*parS* region (B) and the terminus region (capped) can vary in different bacteria. In *E. coli*, the terminus region is compacted by the MatP/*matS* system, but the region also must extend to join the two arms (replichores) of the chromosome, which do not overlap under slow growth conditions. The junctions are indicated by thinner lines. In other examples, the replichores overlap over their entire length. Some other noteworthy features are as follows. The second (smaller) chromosomes in *V. cholerae* and in *Brucella abortus* are located differently [Deghelt et al., 2014]. The extra lines in *B. abortus* next to the centromere and the terminus indicate that their locations are not that fixed and move around their mean positions a bit. In *P. aeruginosa*, the origin and terminus regions are more compacted than in other bacteria [Vallet-Gely and Boccard, 2013]. Both regions are also more removed from the poles. In *M. xanthus*, the regions are also removed from the pole. It is worth noting that centromeres are poleward in all examples (an exception is *E. coli*, which may not have a centromere).

Generality in Chromosome Segregation Systems

Chromosome segregation has been studied in many bacterial species to varying depth: *Bacillus subtilis* [Wang et al., 2014a], *C. crescentus* [Viollier et al., 2004], *E. coli*

[Nielsen et al., 2006; Wang et al., 2006], *Helicobacter pylori* [Sharma et al., 2014], *Pseudomonas aeruginosa* [Vallet-Gely and Boccard, 2013], *Streptomyces coelicolor* (only *ori* region) [Jakimowicz et al., 2007] and *Myxococcus xanthus* (only *ori* region) [Harms et al., 2013]. Although these bacteria belong to different phyla, their segregation systems appear to have a lot in common. First, wherever tested, the chromosomal loci are found to have fixed addresses, and the overall loci organization changes during replication/segregation in a reproducible manner. Second, the chromosomal loci segregate sequentially as they are replicated [David et al., 2014; Vallet-Gely and Boccard, 2013; Viollier et al., 2004; Wang et al., 2006]. (An exception is found in *E. coli*, where some loci stay together longer than others [Bates and Kleckner, 2005; Joshi et al., 2013].) Third, the two chromosomal arms flanking *ori* (the replichores) overlap over their entire length. An exception is seen in *E. coli*, where the replichores stay away from each other in separate cell halves under slow growth conditions (fig. 2). However, in fast growth conditions, the replichores overlap even in *E. coli* [Youngren et al., 2014]. Recently, the two patterns of replichore orientations, overlapping versus nonoverlapping, were found to occur at different stages of the same replication cycle in *B. subtilis* [Wang et al., 2014a]. When overlapping, the daughter chromosomes show mirror symmetry in the mother cell, whereas when nonoverlapping, they are directly repeated [Wang et al., 2006]. Fourth, the *parAB* genes show a remarkable degree of conservation [Gerdes et al., 2000]. The *parS* sites are also conserved or have undergone minor changes [Lin and Grossman, 1998; Livny et al., 2007; Passot et al., 2012]. When present, the *parABS* system makes at least some contribution to the segregation process. Fifth, the *parAB* genes as well as multiple *parS* sites (up to about 10) are found close to the replication origin. The origin region is the only one that is actively segregated in a directed manner, both in *parABS*-dependent and -independent systems. In dependent systems, the *parS* sites migrate ahead of the origin [Fogel and Waldor, 2006; Yamaichi et al., 2012]. Finally, the segregation process is interfaced with other cell cycle events, particularly cell division, discussed below under Chromosome Segregation and the Cell Cycle.

The ‘Spindle’ Pole

It is to be expected that the segregation process should not only move the chromosomal loci to their designated addresses but also retain them after the segregation com-

pletes. Although no specific host factors have been found in plasmid segregation and positioning, in the case of chromosomes there are several examples of proteins that appear to tether segregated *parS* regions to cell poles, including the *parS* region of chr1 [Yamaichi et al., 2012]. A transmembrane polar protein, HubP, directly interacts with ParA1. Since ParA1 interacts with the ParB1-*parS* complex, the HubP-ParA1 interaction helps to retain the *parS* region at the pole. A similar mechanism operates in *C. crescentus*, where a polar protein, PopZ, directly interacts with the ParB-*parS* complex and anchors *parS* to the pole [Bowman et al., 2008; Ebersbach et al., 2008]. An anchoring mechanism was described even earlier in sporulating *B. subtilis*, in which there are several centromeric sites in and around the origin where the ParB analog RacA binds. RacA interacts with the polar protein DivIVA, which localizes itself by recognizing the negative curvature of the pole [Ben-Yehuda et al., 2005; Lenarcic et al., 2009; Ramamurthi and Losick, 2009]. DivIVA also participates in chromosome segregation in another Gram-positive bacterium, *Corynebacterium glutamicum* [Donovan et al., 2012]. In *S. coelicolor*, ParA interacts with a polar protein, Scy, to coordinate segregation with growth [Ditkowski et al., 2013].

Models for Segregation Mechanism

Several models for active segregation of chromosomes have been proposed [Wang et al., 2013]. Here, we discuss the ones that are germane to centromere movement. The centromeres of *V. cholerae* chr1 appear to segregate by a mitosis-like pulling mechanism [Fogel and Waldor, 2006]. A similar mechanism seems to operate in *C. crescentus*, where chromosome segregation has been studied in the most detail [Lim et al., 2014; Ptacin et al., 2014; Shebelut et al., 2010]. In both organisms, the origins travel from one pole to the other with the help of *parABS* systems and are anchored by polar proteins, HubP in *V. cholerae* [Yamaichi et al., 2012], and PopZ and TipN in *C. crescentus* (fig. 1) [Bowman et al., 2008; Ebersbach et al., 2008; Schofield et al., 2010]. The poleward movement requires hydrolysis of ParA-bound ATP [Fogel and Waldor, 2006; Ptacin et al., 2010; Schofield et al., 2010]. A model invoking pulling by ParA has also been described for *M. xanthus*, although in this system the origins stay a bit removed from the pole (fig. 2) [Harms et al., 2013].

The pulling mechanism appears to involve retraction of the ParA ‘cloud’ towards the new pole. ParA-ATP concentrates (forms a ‘cloud’) at the pole, particularly the

new pole, due to its attraction for proteins, such as PopZ and HubP, which are concentrated more at the new pole (fig. 1). Interaction of the ParB bound centromeres with ParA-ATP hydrolyzes ATP, and this is believed to dissolve the cloud frontier, and the retracting cloud causes a directional movement of the centromeres towards the pole. The directional retraction mechanism includes features of the diffusion-ratchet model proposed for plasmid movement, where the ParA-ATP is attracted to the nucleoid (rather than to polar proteins) due to nonspecific DNA binding activity of ParA-ATP [Hwang et al., 2013; Ptacin et al., 2014; Szardenings et al., 2011; Vecchiarelli et al., 2013]. The ParA-ATP cloud serves as a matrix for ParB-bound plasmid movement because interaction of plasmid-bound ParB with ParA-ATP stimulates the release of ParA-ATP from DNA, which is likely to be coupled to ParB-stimulated ATP hydrolysis by ParA. Re-binding of ParA to DNA requires not only ATP binding but also a conformational change of ParA-ATP, which is a slow step. The time delay prevents ParA from rebinding the nucleoid in the vicinity of the ParB-bound plasmid. The resulting ParA gradient that surrounds the ParB-bound plasmid allows it to drift with bias towards locations on the nucleoid where ParA-ATP concentrations are higher. (Note that the time delay is directly related to forming the ParA gradient, not force generation. It is the ParA gradient that is directly related to the force generation mechanism.) The model adequately explains maximal separation of plasmids away from each other.

In principle, ParB-bound chromosomal centromeres can also surf on a nucleoid-associated ParA gradient, depleting ParA from the old to the new pole direction as segregation proceeds. For this mechanism to be effective, the released ParA should not readily rebind (back track) to the depleted regions of the nucleoid towards the old pole. It appears that in *C. crescentus*, the polar proteins PopZ and TipN sequester released ParA molecules and thereby favor directional movement of the centromeres [Ptacin et al., 2014; Schofield et al., 2010]. In *V. cholerae*, it is not known whether HubP has a similar sequestering function. Backtracking would also be discouraged if the reactivation rate of released ParA to a DNA-binding competent state is slow, the key feature of the diffusion-ratchet model. Prevention of backtracking must be efficient because the ParB1/*parS* complex is found at the leading edge of the nucleoid, significantly ahead of the nearby chromosomal region, and the poleward retraction of ParA1 coincides with ParB1/*parS* movement [Fogel and Waldor, 2006]. These facts were initially accommodated in a model wherein ParA1 polymerizes to

form filaments (rather than a cloud), which are bound to HubP at one end and depolymerize from the ParB1/*parS1* bound end [Fogel and Waldor, 2006]. At present, the evidence for ParA polymerization is generally clear only in vitro, not in vivo [Vecchiarelli et al., 2012]. Also, filament formation may not be obligatory since a concentration gradient of ParA dimers or small oligomers with maximal density at the new pole could also explain the results. The latter scenario is more likely because ParA appears as a cloud in most cases, and a recent high-resolution study in *C. crescentus* has argued convincingly against the existence of filaments [Lim et al., 2014]. It is noteworthy that the movement of chromosomal *parS* sites in the polar region is unlikely to be due to cloud formation over the nucleoid because the polar regions are usually devoid of DNA due to exclusion from polysomes [Bakshi et al., 2012] and because *parS1* moves significantly ahead of the nearby chromosomal regions in *V. cholerae* [Fogel and Waldor, 2006]. The ParA1 cloud formation in the DNA-free polar region most likely owes to polar proteins (fig. 1).

The recent *C. crescentus* study also found the cloud-based diffusion-ratchet model to be inadequate to account for the speed of ParB-*parS* movement, and proposed that the movement is aided by the chromosomal DNA itself because of its elastic nature. The idea here is that ParA bound to the chromosome pushes the ParB-*parS* complex from one point to another ('DNA-relay') when the stretched (elastic) chromosomal regions relax. Thus, a combination of diffusion-ratchet and intrinsic chromosome dynamics seems to account for the speed, while the burden of providing the direction of movement still rests with the ParA concentration gradient.

Which aspects of these mechanisms are general and which are particular to certain systems need to be established. The DNA-relay model was proposed on the basis that ParA is static on DNA and the ParA concentration is particularly low (90 dimers per cell) in *C. crescentus* [Lim et al., 2014]. If these are not generally found, other mechanisms might exist. *V. cholerae* chr1 is less dependent on *par* than the *C. crescentus* chromosome. The *Caulobacter* origin travels no more than 40% of the cell length without *par* [Shebelut et al., 2010; Toro et al., 2008]. In contrast, *V. cholerae* chr1 origin travels about 80% of cell length without *par1* [Fogel and Waldor, 2006; Kadoya et al., 2011]. In *V. cholerae*, the role of *par1* appears to be only in anchoring the centromere to the pole at the final step of segregation. *V. cholerae* and *C. crescentus* also differ in another respect: whereas *C. crescentus* depends on ParB to sequester MipZ, a potent cell division inhibitor [Than-

bichler and Shapiro, 2006], *par1* is not essential for *V. cholerae*. This bacterium does not have MipZ and therefore may not need ParB1 to allow cell division.

Unlike *par1*, *par2* (*parABS2*) is essential for segregation of *V. cholerae* chr2 [Yamaichi et al., 2007b]. Although the overall organization and segregation program of chr2 are known, the mechanism of segregation is not [Demarre et al., 2014; Fiebig et al., 2006; Fogel and Waldor, 2006; Srivastava and Chattoraj, 2007]. In particular, the mechanisms underlying the directionality of centromere movement and retention of the centromeres once they reach their destination in cell quarter positions remain to be determined (fig. 2). Since the location of markers in chr2 follows their order on the chromosome, it should be actively organized as in chr1. It is to be noted that although chr2 is believed to have originated from a plasmid, the maintenance mode of chr2, particularly its replication, is typical of chromosomes rather than plasmids. Whereas both the timing of replication initiation and centromere movement are random for plasmids [Leonard and Helmstetter, 1988; Sengupta et al., 2010], chr2 replication starts at a fixed time in the cell cycle that might well depend on the timing of chr1 replication, and the sister origins progressively separate away from each other [Baek and Chattoraj, 2014; Fiebig et al., 2006; Fogel and Waldor, 2005; Rasmussen et al., 2007; Srivastava and Chattoraj, 2007]. Because the plasmid diffusion-ratchet mechanism accounts for directional movement over a short range only, it may not be adequate to explain the ordered movement of chr2 over a quarter of cell length.

Competition (Incompatibility) between chr1 and chr2 for Segregation

Mis-segregation of chromosomes in eukaryotes is avoided through sister chromatid cohesion, and sensing of tension from microtubules pulling the sister centromeres towards the opposite spindle poles. There is no competition among the chromosomes because the cell cycle-specific factors that participate in segregation, such as cohesins and microtubules, are apparently not limiting. From studies in a few bacteria with a multipartite genome (*V. cholerae*, *Burkholderia cenocepacia* [Dubarry et al., 2006], *B. cereus* [Deghelt et al., 2014]), it is becoming clear that the factors that control chromosome replication and segregation are chromosome specific, whereas in eukaryotes they are not. It has been well established that by encoding specific control factors, plasmids avoid competition (incompatibility) with the host as well as

with other plasmids if they happen to be present in the same bacterium. This principle of coexistence of different plasmids with each other and the host chromosome has been retained in bacteria with a multipartite genome. This is to be expected since in these bacteria the largest chromosome is equivalent to the single chromosome of bacteria with undivided genomes, and the other (smaller) chromosomes appear to have originated from plasmids [Egan et al., 2005]. In the plasmid to chromosome transition, the plasmid-like strategy for avoiding incompatibility has apparently been retained. It is worth noting that overlapping binding specificities have been discerned among the ParBs and centromeres of the Burkholderiales multipartite genomes [Passot et al., 2012]. The binding affinities, however, differ widely, and this might suffice to avoid incompatibility in this system.

par-Independent Segregation of chr1

About 30% of sequenced bacteria do not have *par* genes, including *E. coli*. How chromosomes move in these bacteria is not clear. Even the bacteria that do have *par* genes (e.g. *B. subtilis*) may not require them for chromosome segregation. The same can be said for *V. cholerae* chr1. In the absence of *par1*, chr1 segregation appears normal except that the mean position of centromeres becomes ~10% further removed from the pole [Fogel and Waldor, 2006; Kadoya et al., 2011]. These results indicate that the separation of the duplicated *ori1* and their poleward migration can occur independently of the *par1* system. This initial *par*-independent and subsequent *par*-dependent migration is also true for the *C. crescentus* chromosome except that the *par*-dependent step is more extensive, as discussed above [Kadoya et al., 2011; Shebelut et al., 2010]. A recent study indicates that the origin region could provide the centromeric function in the absence of *par1* in *V. cholerae* [David et al., 2014]. This was evident when an extra *ori1* was placed in chr1 distal to the native *ori1*. Either one of the origins could go to the pole indicating that the origin activity suffices for polar segregation. Notably, *parS1* is dominant over *ori1*: when both are present, *parS1* occupies the polar position.

Chromosome Segregation and the Cell Cycle

Accumulating evidence supports a connection between chromosome replication, segregation and cell division in many bacteria, including in *V. cholerae*.

Connection of replication to segregation is suggested by the proximity of *parS* sites to *ori* [Livny et al., 2007] and co-replicative segregation of replicated regions [David et al., 2014]. The free energy of DNA polymerization could be a source of motive force for bidirectional segregation of newly replicated sister DNA [Lemon and Grossman, 2001]. Similarly, the ‘snap’-mediated build-up of tension due to ongoing replication could be another source of force to push sisters apart when the tension is released [Joshi et al., 2013]. As discussed above, the best evidence for replication contributing to segregation has come from recent evidence that *V. cholerae* *ori1* can fulfill centromeric function in the absence of *par1* [David et al., 2014]. We note that the role of the origin as centromere was initially tested in *E. coli* and was considered unlikely [Gordon et al., 2002]. Another link between replication and segregation is the fact that the *V. cholerae* chr2 replication initiator RctB regulates the chr2 *par* operon [Yamaichi et al., 2011].

The evidence for segregation proteins influencing replication is more direct. In *B. subtilis* and *Streptococcus pneumoniae*, ParB organizes the replication origin by loading a condensin protein in the vicinity of the origin [Gruber and Errington, 2009; Minnen et al., 2011; Sullivan et al., 2009; Wang et al., 2014b]. ParA controls the activity of DnaA and thereby controls replication initiation in *B. subtilis* and in *V. cholerae* for chr1 [Kadoya et al., 2011; Scholefield et al., 2012]. *V. cholerae* ParB2 also controls replication of chr2 using two separate mechanisms [Venkova-Canova et al., 2013; Yamaichi et al., 2011]. It is noteworthy that Par proteins control replication by targeting the most important regulators of replication, the DnaA initiator for chr1 and the *B. subtilis* chromosome, and the RctB initiator for chr2. The proximity of centromeres advances replication timing in several eukaryotes [Tanaka et al., 2013]. Recently, one of the replication initiator proteins (Orc1) was shown to control centriole duplication [Hossain and Stillman, 2012]. It is only a matter of time before more connections like these will be evident.

There is also considerable evidence for coordination between segregation and cell division. Nucleoid occlusion directly prevents cytokinesis in *B. subtilis* and *E. coli* [Bernard et al., 2010; Cambridge et al., 2014]. The *ter* macrodomain at the cell center may also serve as a landmark for the formation of the septal ring [Bailey et al., 2014]. The best evidence so far has come from studies in *C. crescentus*, where sequestration of the cell division inhibitor MipZ by ParB is required for cell viability [Thanbichler and Shapiro, 2006]. This has been one of the most

insightful findings in understanding how the sequence of the three basic cell cycle processes can be coordinated. Only upon replication initiation and segregation of the origins to the poles can MipZ be sequestered away from the mid-cell, allowing FtsZ polymerization at the right place (mid-cell) and at the right time (only after ensuring that replication and segregation have initiated). Clear evidence has also come from the finding that the *ter* macrodomain organizer MatP interacts with ZapB, a component of the cell division apparatus of *E. coli* [Espéti et al., 2012]. Since *V. cholerae* has MatP and both *ter1* and *ter2* regions contain *matS* sites, both chromosomes might interface with the cell division apparatus [Demarre et al., 2014].

Future

Since the discovery in plasmids of ParA and ParB proteins and the finding that they serve motor function and kinetochore formation, respectively, the search has been on for the missing components of the mitotic apparatus, the spindle. It was expected that the host might provide an equivalent of the spindle to be used as tracks for plasmid segregation. However, no convincing evidence of host factor participation was obtained. The plasmid segregation models can now explain how segregation can be achieved without requiring additional proteins, although some plasmid segregation systems do require the help from the nucleoid [Hester and Lutkenhaus, 2007; Szardenings et al., 2011]. However, there is more order in chromosome segregation than is the case for plasmid segregation, and there is evidence that additional factors (MipZ, TipN, PopZ, HubP) play crucial roles in ordered segregation of the chromosome. Initial studies using bacterial and yeast two-hybrid systems in *V. cholerae* have indicated that there are proteins other than HubP that interact with both ParA and ParB [Baek et al., 2014]. The interactions need to be authenticated and their physiological significance determined. These studies are likely to help understand how the different cell cycle processes are coordinated.

Segregation falls in the general category of pattern formation, and theories on biological pattern formation have been considered to explain plasmid segregation. The most recent among them is the diffusion-ratchet mechanism [Meinhardt and Gierer, 2000; Vecchiarelli et al., 2010]. As mentioned earlier, the diffusion-ratchet mechanism has also been invoked to explain chromosome segregation [Lim et al., 2014; Ptacin et al., 2014]. In this con-

text, it would be of interest to know the reactivation rate of inactive ParA in vitro, which is the essence of the diffusion-ratchet model, and how the chromosomal Par proteins compare with their plasmid counterparts in making ParA-free zones on the nucleoid. In other words, can the nucleoid be used as scaffold for its own segregation?

Chromosome segregation is a multistep process that includes not only motion but condensation (keeping the chromosome bulk together), anchoring of centromeres, clearing the nucleoids from mid-cell and coordination with the cell cycle. Understanding how chromosomal markers move in a directional fashion is the most pressing problem confronting the chromosome segregation field. Tracking the mover ParA in space and time and at super-resolution is crucial for a deeper understanding of the dynamics. Use of superresolution microscopy to measure the diffusion rate of ParA molecules (free and nucleoid bound, as has been done for RNA polymerase [Bakshi et al., 2013]) is in order. Efforts have already begun in *C. crescentus* [Lim et al., 2014; Ptacin et al., 2014], and it remains to be seen how closely the *V. cholerae* segregation system resembles that of *C. crescentus*.

Understanding the basis of *par*-independent segregation is more challenging. From the examples in figure 2, it appears that movement towards the pole is an inherent feature of the origin region. Determining the role of replication in segregation appears to be the first order of business. The exciting possibility that a completely new mechanism may await discovery is also there.

The levels of Par proteins are crucial in thinking about segregation models [Lim et al., 2014], especially as overexpression of *par* genes is often detrimental [Kadoya et al., 2011; Mohl and Gober, 1997; Schofield et al., 2010]. The regulation of chromosomal *par* genes are generally not known except in some isolated cases [Casart et al., 2008]. In plasmids, the *parAB* genes form an autoregulated operon, but this was not the case when chromosomal *par* genes were tested in *P. aeruginosa* and in *V. cholerae* for chr1 [Baek et al., 2014; Bartosik et al., 2014]. Whether and how *par* genes of chr1 are regulated remains to be determined. Studies in both *P. aeruginosa* and *V. cholerae* also showed that Par proteins directly or indirectly control several other genes apparently unrelated to segregation. The mechanism of this regulation remains totally unknown, and its understanding may reveal how Par proteins play their wider role in bacteria.

Cohesins play major and variable roles in mitosis and meiosis [Tanaka et al., 2013]. Although cohesins per se

have not been found in bacteria, a similar role of lesser duration could be played by SeqA [Joshi et al., 2013; Stokke et al., 2011]. SeqA has been found to affect segregation and cell division over and above its role in replication in *V. cholerae* [Demarre and Chattoraj, 2010; Saint-Dic et al., 2008]. The condensins on the other hand (e.g. MukB in *E. coli* and SMC in *B. subtilis*) could play a significant role in cell cycle events [see the reviews by Kleine Borgmann and Graumann, this issue, pp. 384–395, and Rybenkov et al., this issue, pp. 371–383; Danilova et al., 2007; Gruber and Errington, 2009; Sullivan et al., 2009]. Unexpectedly, the role of MukB does not appear critical for *V. cholerae*, although this remains to be further studied [David et al., 2014].

Cell division requires clearing the mid-cell region of chromosomal DNA [Bernard et al., 2010; Cambridge et al., 2014]. This is a rather intricate process, requiring not only completion of replication but also decatenation of the sisters, resolution of dimers that occasionally form when sisters recombine, and organization of the *ter* macromodomain. These processes are fundamental prerequisites for cytokinesis [see the review by Crozat et al., this issue, pp. 396–408] and, although not yet demonstrated, must be operating on both chromosomes of *V. cholerae*.

V. cholerae provides the opportunity to examine the behavior of two distinct chromosomal maintenance sys-

tems operating in the same environment. It is not known whether the presence of one chromosome influences the segregation of the other. We know only that preventing chr2 replication does not influence chr1 replication or segregation [Kadoya and Chattoraj, 2012]. *V. cholerae* strains are available with large-scale rearrangement of the genome [Val et al., 2012]. Segregation studies in these cells could reveal whether the chromosomal bulk causes any steric hindrance to segregation. These cells are also of interest for investigating whether a divided genome has helped in chromosome segregation, as has been argued [Srivastava and Chattoraj, 2007].

Acknowledgements

The authors are indebted to Christophe Possoz [David et al., 2014], François-Xavier Barre [Demarre et al., 2014] and Xavier De Bolle [Deghelt et al., 2014] for communicating results prior to publication. The authors are also indebted to Abhishek Goel for help in drawing the figures, and to David Lane, Christophe Possoz, Anthony Vecchiarelli, Zemer Gitai, Michael Lichten and Michael Yarmolinsky for thoughtful comments, and Jemima Barrowman for editing the manuscript. This work was supported by the Intramural Research Program, Center for Cancer Research, The National Cancer Institute.

References

- Baek JH, Chattoraj DK: Chromosome I controls chromosome II replication in *Vibrio cholerae*. PLoS Genet 2014;10:e1004184.
- Baek JH, Rajagopala SV, Chattoraj DK: Chromosome segregation proteins of *Vibrio cholerae* as transcription regulators. MBio 2014; 5:e01061–01014.
- Bailey MW, Bisicchia P, Warren BT, et al: Evidence for divisome localization mechanisms independent of the Min system and SlmA in *Escherichia coli*. PLoS Genet 2014; 10:e1004504.
- Bakshi S, Dalrymple RM, Li W, et al: Partitioning of RNA polymerase activity in live *Escherichia coli* from analysis of single-molecule diffusive trajectories. Biophys J 2013;105:2676–2686.
- Bakshi S, Siryaporn A, Goulian M, et al: Super-resolution imaging of ribosomes and RNA polymerase in live *Escherichia coli* cells. Mol Microbiol 2012;85:21–38.
- Bartosik AA, Glabski K, Jecz P, et al: Transcriptional profiling of *parA* and *parB* mutants in actively dividing cells of an opportunistic human pathogen *Pseudomonas aeruginosa*. PLoS One 2014;9:e87276.
- Bates D, Kleckner N: Chromosome and replisome dynamics in *Escherichia coli*: loss of sister cohesion triggers global chromosome movement and mediates chromosome segregation. Cell 2005;121:899–911.
- Ben-Yehuda S, Fujita M, Liu XS, et al: Defining a centromere-like element in *Bacillus subtilis* by identifying the binding sites for the chromosome-anchoring protein RacA. Mol Cell 2005;17:773–782.
- Bernard R, Marquis KA, Rudner DZ: Nucleoid occlusion prevents cell division during replication fork arrest in *Bacillus subtilis*. Mol Microbiol 2010;78:866–882.
- Bowman GR, Comolli LR, Zhu J, et al: A polymeric protein anchors the chromosomal origin/ParB complex at a bacterial cell pole. Cell 2008;134:945–955.
- Cambridge J, Blinkova A, Magnan D, et al: A replication-inhibited unsegregated nucleoid at mid-cell blocks Z-ring formation and cell division independently of SOS and the SlmA nucleoid occlusion protein in *Escherichia coli*. J Bacteriol 2014;196:36–49.
- Casart Y, Gamero E, Rivera-Gutierrez S, et al: Par genes in *Mycobacterium bovis* and *Mycobacterium smegmatis* are arranged in an operon transcribed from ‘sigge’ promoters. BMC Microbiol 2008;8:51.
- Danilova O, Reyes-Lamothe R, Pinskaya M, et al: MukB colocalizes with the *oriC* region and is required for organization of the two *Escherichia coli* chromosome arms into separate cell halves. Mol Microbiol 2007;65:1485–1492.
- David A, Demarre G, Muresan L, et al: The two *cis*-acting sites, *parS1* and *oriC1*, contribute to the longitudinal organisation of *Vibrio cholerae* chromosome I. PLoS Genet 2014; 10:e1004448.
- Deghelt M, Mullier C, Sternon J-F, et al: G1-arrested newborn cells are the predominant infectious form of the pathogen *Brucella abortus*. Nat Commun 2014;5:4366.
- Demarre G, Chattoraj DK: DNA adenine methylation is required to replicate both *Vibrio cholerae* chromosomes once per cell cycle. PLoS Genet 2010;6:e1000939.

- Demarre G, Galli E, Muresan L, et al: Differential management of the replication terminus regions of the two *Vibrio cholerae* chromosomes during cell division. *PLoS Genet* 2014; 10ED:e1004557.
- Ditkowski B, Holmes N, Rydzak J, et al: Dynamic interplay of ParA with the polarity protein, Scy, coordinates the growth with chromosome segregation in *Streptomyces coelicolor*. *Open Biol* 2013;3:130006.
- Donovan C, Sieger B, Kramer R, et al: A synthetic *Escherichia coli* system identifies a conserved origin tethering factor in actinobacteria. *Mol Microbiol* 2012;84:105–116.
- Dubarry N, Pasta F, Lane D: Parabs systems of the four replicons of *Burkholderia cenocepacia*: new chromosome centromeres confer partition specificity. *J Bacteriol* 2006;188:1489–1496.
- Ebersbach G, Briegel A, Jensen GJ, et al: A self-associating protein critical for chromosome attachment, division, and polar organization in *Caulobacter*. *Cell* 2008;134:956–968.
- Egan ES, Fogel MA, Waldor MK: Divided genomes: negotiating the cell cycle in prokaryotes with multiple chromosomes. *Mol Microbiol* 2005;56:1129–1138.
- Egan ES, Waldor MK: Distinct replication requirements for the two *Vibrio cholerae* chromosomes. *Cell* 2003;114:521–530.
- Espéli O, Borne R, Dupaigne P, et al: A MatP-divisome interaction coordinates chromosome segregation with cell division in *Escherichia coli*. *EMBO J* 2012;31:3198–3211.
- Fiebig A, Keren K, Theriot JA: Fine-scale time-lapse analysis of the biphasic, dynamic behaviour of the two *Vibrio cholerae* chromosomes. *Mol Microbiol* 2006;60:1164–1178.
- Fisher JK, Bourniquel A, Witz G, et al: Four-dimensional imaging of *Escherichia coli* nucleoid organization and dynamics in living cells. *Cell* 2013;153:882–895.
- Fogel MA, Waldor MK: Distinct segregation dynamics of the two *Vibrio cholerae* chromosomes. *Mol Microbiol* 2005;55:125–136.
- Fogel MA, Waldor MK: A dynamic, mitotic-like mechanism for bacterial chromosome segregation. *Genes Dev* 2006;20:3269–3282.
- Gerdes K, Moller-Jensen J, Bugge Jensen R: Plasmid and chromosome partitioning: surprises from phylogeny. *Mol Microbiol* 2000;37:455–466.
- Gordon GS, Shivers RP, Wright A: Polar localization of the *Escherichia coli* *oriC* region is independent of the site of replication initiation. *Mol Microbiol* 2002;44:501–507.
- Gruber S, Errington J: Recruitment of condensin to replication origin regions by ParB/SpoOJ promotes chromosome segregation in *B. subtilis*. *Cell* 2009;137:685–696.
- Harms A, Treuner-Lange A, Schumacher D, et al: Tracking of chromosome and replisome dynamics in *Myxococcus xanthus* reveals a novel chromosome arrangement. *PLoS Genet* 2013; 9:e1003802.
- Hester CM, Lutkenhaus J: Soj (ParA) DNA binding is mediated by conserved arginines and is essential for plasmid segregation. *Proc Natl Acad Sci USA* 2007;104:20326–20331.
- Hossain M, Stillman B: Meier-gorlin syndrome mutations disrupt an Orc1 Cdk inhibitory domain and cause centrosome reduplication. *Genes Dev* 2012;26:1797–1810.
- Hwang LC, Vecchiarelli AG, Han YW, et al: ParA-mediated plasmid partition driven by protein pattern self-organization. *EMBO J* 2013;32:1238–1249.
- Jakimowicz D, Zydek P, Kois A, et al: Alignment of multiple chromosomes along helical ParA scaffolding in sporulating *Streptomyces* hyphae. *Mol Microbiol* 2007;65:625–641.
- Joshi MC, Magnan D, Montminy TP, et al: Regulation of sister chromosome cohesion by the replication fork tracking protein SeqA. *PLoS Genet* 2013;9:e1003673.
- Jun S, Wright A: Entropy as the driver of chromosome segregation. *Nat Rev Microbiol* 2010;8: 600–607.
- Kadoya R, Baek JH, Sarker A, et al: Participation of chromosome segregation protein ParA1 of *Vibrio cholerae* in chromosome replication. *J Bacteriol* 2011;193:1504–1514.
- Kadoya R, Chattoraj DK: Insensitivity of chromosome I and the cell cycle to blockage of replication and segregation of *Vibrio cholerae* chromosome II. *MBio* 2012;3:00067–00012.
- Lemon KP, Grossman AD: The extrusion-capture model for chromosome partitioning in bacteria. *Genes Dev* 2001;15:e2031–e2041.
- Lenarcic R, Halbedel S, Visser L, et al: Localisation of DivIVA by targeting to negatively curved membranes. *EMBO J* 2009;28:2272–2282.
- Leonard AC, Helmstetter CE: Replication patterns of multiple plasmids coexisting in *Escherichia coli*. *J Bacteriol* 1988;170:1380–1383.
- Lim HC, Surovtsev IV, Beltran BG, et al: Evidence for a DNA-relay mechanism in *parABS*-mediated chromosome segregation. *Elife* 2014;3:e02758.
- Lin DC, Grossman AD: Identification and characterization of a bacterial chromosome partitioning site. *Cell* 1998;92:675–685.
- Livny J, Yamaichi Y, Waldor MK: Distribution of centromere-like *parS* sites in bacteria: insights from comparative genomics. *J Bacteriol* 2007;189:8693–8703.
- Meinhardt H, Gierer A: Pattern formation by local self-activation and lateral inhibition. *Bioessays* 2000;22:753–760.
- Mercier R, Petit MA, Schbath S, et al: The MatP/matS site-specific system organizes the terminus region of the *Escherichia coli* chromosome into a macrodomain. *Cell* 2008;135: 475–485.
- Minnen A, Attaiach L, Thon M, et al: SMC is recruited to *oriC* by ParB and promotes chromosome segregation in *Streptococcus pneumoniae*. *Mol Microbiol* 2011;81:676–688.
- Mohl DA, Guber JW: Cell cycle-dependent polar localization of chromosome partitioning proteins in *Caulobacter crescentus*. *Cell* 1997;88: 675–684.
- Nielsen HJ, Ottesen JR, Youngren B, et al: The *Escherichia coli* chromosome is organized with the left and right chromosome arms in separate cell halves. *Mol Microbiol* 2006;62: 331–338.
- Niki H, Yamaichi Y, Hiraga S: Dynamic organization of chromosomal DNA in *Escherichia coli*. *Genes Dev* 2000;14:212–223.
- Passot FM, Calderon V, Fichant G, et al: Centromere binding and evolution of chromosomal partition systems in the *Burkholderiales*. *J Bacteriol* 2012;194:3426–3436.
- Possoz C, Junier I, Espeli O: Bacterial chromosome segregation. *Front Biosci (Landmark Ed)* 2012;17:1020–1034.
- Ptacin JL, Gahlmann A, Bowman GR, et al: Bacterial scaffold directs pole-specific centromere segregation. *Proc Natl Acad Sci USA* 2014; 111:E2046–E2055.
- Ptacin JL, Lee SF, Garner EC, et al: A spindle-like apparatus guides bacterial chromosome segregation. *Nat Cell Biol* 2010;12:791–798.
- Radnedge L, Youngren B, Davis M, et al: Probing the structure of complex macromolecular interactions by homolog specificity scanning: the P1 and P7 plasmid partition systems. *EMBO J* 1998;17:6076–6085.
- Ramamurthi KS, Losick R: Negative membrane curvature as a cue for subcellular localization of a bacterial protein. *Proc Natl Acad Sci USA* 2009;106:13541–13545.
- Rasmussen T, Jensen RB, Skovgaard O: The two chromosomes of *Vibrio cholerae* are initiated at different time points in the cell cycle. *EMBO J* 2007;26:3124–3131.
- Saint-Dic D, Frushour BP, Kehrl JH, et al: A ParA homolog selectively influences positioning of the large chromosome origin in *Vibrio cholerae*. *J Bacteriol* 2006;188:5626–5631.
- Saint-Dic D, Kehrl J, Frushour B, et al: Excess SeqA leads to replication arrest and a cell division defect in *Vibrio cholerae*. *J Bacteriol* 2008;190:5870–5878.
- Schofield WB, Lim HC, Jacobs-Wagner C: Cell cycle coordination and regulation of bacterial chromosome segregation dynamics by polarly localized proteins. *EMBO J* 2010;29:3068–3081.
- Schofield G, Errington J, Murray H: Soj/ParA stalls DNA replication by inhibiting helix formation of the initiator protein DnaA. *EMBO J* 2012;31:1542–1555.
- Sengupta M, Nielsen HJ, Youngren B, et al: P1 plasmid segregation: accurate redistribution by dynamic plasmid pairing and separation. *J Bacteriol* 2010;192:1175–1183.
- Sharma A, Kamran M, Verma V, et al: Intracellular locations of replication proteins and the origin of replication during chromosome duplication in the slowly growing human pathogen *Helicobacter pylori*. *J Bacteriol* 2014;196: 999–1011.
- Shebelut CW, Guberman JM, van Teeffelen S, et al: *Caulobacter* chromosome segregation is an ordered multistep process. *Proc Natl Acad Sci USA* 2010;107:14194–14198.

- Skarstad K, Steen HB, Boye E: *Escherichia coli* DNA distributions measured by flow cytometry and compared with theoretical computer simulations. *J Bacteriol* 1985;163:661–668.
- Sliusarenko O, Heinritz J, Emonet T, et al: High-throughput, subpixel precision analysis of bacterial morphogenesis and intracellular spatio-temporal dynamics. *Mol Microbiol* 2011;80:612–627.
- Srivastava P, Chattoraj DK: Selective chromosome amplification in *Vibrio cholerae*. *Mol Microbiol* 2007;66:1016–1028.
- Srivastava P, Fekete RA, Chattoraj DK: Segregation of the replication terminus of the two *Vibrio cholerae* chromosomes. *J Bacteriol* 2006;188:1060–1070.
- Stokke C, Waldminghaus T, Skarstad K: Replication patterns and organization of replication forks in *Vibrio cholerae*. *Microbiology* 2011;157:695–708.
- Sullivan NL, Marquis KA, Rudner DZ: Recruitment of SMC by ParP-*parS* organizes the origin region and promotes efficient chromosome segregation. *Cell* 2009;137:697–707.
- Szardenings F, Guymer D, Gerdes K: ParA ATPases can move and position DNA and subcellular structures. *Curr Opin Microbiol* 2011;14:712–718.
- Tanaka TU, Clayton L, Natsume T: Three wise centromere functions: see no error, hear no break, speak no delay. *EMBO Rep* 2013;14:1073–1083.
- Thanbichler M, Shapiro L: MipZ, a spatial regulator coordinating chromosome segregation with cell division in *Caulobacter*. *Cell* 2006;126:147–162.
- Toro E, Hong SH, McAdams HH, et al: *Caulobacter* requires a dedicated mechanism to initiate chromosome segregation. *Proc Natl Acad Sci USA* 2008;105:15435–15440.
- Umbarger MA, Toro E, Wright MA, et al: The three-dimensional architecture of a bacterial genome and its alteration by genetic perturbation. *Mol Cell* 2011;44:252–264.
- Val ME, Skovgaard O, Ducos-Galand M, et al: Genome engineering in *Vibrio cholerae*: a feasible approach to address biological issues. *PLoS Genet* 2012;8:e1002472.
- Valens M, Penaud S, Rossignol M, et al: Macrodomain organization of the *Escherichia coli* chromosome. *EMBO J* 2004;23:4330–4341.
- Vallet-Gely I, Boccard F: Chromosomal organization and segregation in *Pseudomonas aeruginosa*. *PLoS Genet* 2013;9:e1003492.
- Vecchiarelli AG, Han YW, Tan X, et al: ATP control of dynamic P1 ParA-DNA interactions: a key role for the nucleoid in plasmid partition. *Mol Microbiol* 2010;78:78–91.
- Vecchiarelli AG, Hwang LC, Mizuuchi K: Cell-free study of F plasmid partition provides evidence for cargo transport by a diffusion-ratchet mechanism. *Proc Natl Acad Sci USA* 2013;110:E1390–E1397.
- Vecchiarelli AG, Mizuuchi K, Funnell BE: Surfing biological surfaces: exploiting the nucleoid for partition and transport in bacteria. *Mol Microbiol* 2012;86:513–523.
- Venkova-Canova T, Baek JH, Fitzgerald PC, et al: Evidence for two different regulatory mechanisms linking replication and segregation of *Vibrio cholerae* chromosome II. *PLoS Genet* 2013;9:e1003579.
- Viollier PH, Thanbichler M, McGrath PT, et al: Rapid and sequential movement of individual chromosomal loci to specific subcellular locations during bacterial DNA replication. *Proc Natl Acad Sci USA* 2004;101:9257–9262.
- Wang X, Liu X, Possoz C, et al: The two *Escherichia coli* chromosome arms locate to separate cell halves. *Genes Dev* 2006;20:1727–1731.
- Wang X, Montero Llopis P, Rudner DZ: Organization and segregation of bacterial chromosomes. *Nat Rev Genet* 2013;14:191–203.
- Wang X, Montero Llopis P, Rudner DZ: *Bacillus subtilis* chromosome organization oscillates between two distinct patterns. *Proc Natl Acad Sci USA* 2014a;111:12877–12882.
- Wang X, Tang OW, Riley EP, et al: The SMC condensin complex is required for origin segregation in *Bacillus subtilis*. *Curr Biol* 2014b;24:287–292.
- Webb CD, Teleman A, Gordon S, et al: Bipolar localization of the replication origin regions of chromosomes in vegetative and sporulating cells of *B. subtilis*. *Cell* 1997;88:667–674.
- Wiggins PA, Cheveralls KC, Martin JS, et al: Strong intranucleoid interactions organize the *Escherichia coli* chromosome into a nucleoid filament. *Proc Natl Acad Sci USA* 2010;107:4991–4995.
- Yamaichi Y, Bruckner R, Ringgaard S, et al: A multidomain hub anchors the chromosome segregation and chemotactic machinery to the bacterial pole. *Genes Dev* 2012;26:2348–2360.
- Yamaichi Y, Fogel MA, McLeod SM, et al: Distinct centromere-like *parS* sites on the two chromosomes of *Vibrio* spp. *J Bacteriol* 2007a;189:5314–5324.
- Yamaichi Y, Fogel MA, Waldor MK: Par genes and the pathology of chromosome loss in *Vibrio cholerae*. *Proc Natl Acad Sci USA* 2007b;104:630–635.
- Yamaichi Y, Gerding MA, Davis BM, et al: Regulatory cross-talk links *Vibrio cholerae* chromosome II replication and segregation. *PLoS Genet* 2011;7:e1002189.
- Youngren B, Nielsen HJ, Jun S, et al: The multi-fork *Escherichia coli* chromosome is a self-duplicating and self-segregating thermodynamic ring polymer. *Genes Dev* 2014;28:71–84.