Examination of the Influence of Leptin and Acute Metabolic Challenge on RFRP-3 Neurons of Mice in Development and Adulthood

Matthew C. Poling^{a,b} Morris P. Shieh^a Nagambika Munaganuru^a Elena Luo^a
Alexander S. Kauffman^a

^aDepartment of Reproductive Medicine and ^bBiomedical Sciences Graduate Program, University of California, San Diego, La Jolla, Calif., USA

Key Words

RFRP-3 · GnIH · Leptin · Reproduction · Development

Abstract

Background: The neuropeptide RFamide-related peptide-3 (RFRP-3; mammalian ortholog to gonadotropin-inhibiting hormone) can inhibit luteinizing hormone (LH) release and increases feeding, but the regulation and development of RFRP-3 neurons remains poorly characterized, especially in mice. **Methods and Results:** We first confirmed that peripheral injections of murine RFRP-3 peptide could markedly suppress LH secretion in adult mice, as in other species. Second, given RFRP-3's reported orexigenic properties, we performed double-label in situ hybridization for metabolic genes in *Rfrp* neurons of mice. While *Rfrp* neurons did not readily coexpress neuropeptide Y, thyrotropin-releasing hormone, or MC4R, a small subset of *Rfrp* neurons did express the leptin receptor in both sexes. Surprisingly, we identified no changes in *Rfrp* expression or neuronal activation in adult mice after acute fasting. However, we determined that *Rfrp* mRNA levels in the dorsal-medial nucleus were significantly reduced in adult obese (Ob) mice of both sexes. Given the lower *Rfrp* levels observed in adult Ob mice, we asked whether leptin might also regulate RFRP-3 neuron development. *Rfrp* gene expression changed markedly over juvenile development, correlating with the timing of the juvenile 'leptin surge' known to govern hypothalamic feeding

circuit development. However, the dramatic developmental changes in juvenile *Rfrp* expression did not appear to be leptin driven, as the pattern and timing of *Rfrp* neuron development were unaltered in Ob juveniles. **Conclusion:** Leptin status modulates RFRP-3 expression in adulthood, but is not required for normal development of the RFRP-3 system. Leptin's regulation of adult RFRP-3 neurons likely occurs primarily via indirect signaling, and may be secondary to obesity, as only a small subset of RFRP-3 neurons express the long form of the leptin receptor (LepRb).

© 2014 S. Karger AG, Basel

Introduction

Neuronal networks of the hypothalamus integrate peripheral endocrine and metabolic signals to regulate gonadotropin-releasing hormone (GnRH) neurons, which in turn stimulate the reproductive axis. In most cases, hormonal and metabolic information are transmitted to GnRH neurons indirectly via upstream reproductive neural circuitry. One possible intermediary neuropeptide is RFamide-related peptide 3 (RFRP-3), the mammalian ortholog of avian gonadotropin-inhibiting hormone [1, 2]. In rodents, RFRP-3 treatment inhibits both GnRH neuronal activity and luteinizing hormone (LH) secretion through central GnRH-dependent mechanisms [3–6]. In mice and rats, some GnRH neurons are apposed by

RFRP-3 axonal fibers [2, 7, 8], and a subset of GnRH neurons express Gpr147 [8, 9], a high-affinity receptor for RFRP-3 [10, 11]. In rodents, RFRP-3-producing neurons reside in a scattered pattern solely within and just ventral to the dorsal-medial nucleus of the hypothalamus (collectively abbreviated as DMN) [2, 12]. Likewise, the rodent *Rfrp* gene, encoding RFRP-3, is expressed in a scattered pattern exclusively in the DMN and, to a much lesser extent, ventrally in the neighboring area just dorsal of the VMN [2, 9, 10]. The DMN is known to regulate aspects of energy balance, feeding behavior, and thermoregulation [13–15]. However, the precise role(s) of RFRP-3 neurons in the DMN is unclear, in part due to a lack of knowledge of both the neuropeptides and receptors coexpressed in RFRP-3 neurons and the identity of regulatory factors that govern RFRP-3 synthesis and secretion.

Leptin, a hormone secreted from adipocytes, has strong effects on hypothalamic regulation of satiety, energy expenditure, and body weight, not to mention a stimulatory (permissive) role in reproductive function. Obese mice (Ob) have a nonfunctional leptin gene and are overweight, hyperphagic, have low LH levels, and are infertile [16, 17], illustrating the importance of leptin in maintenance of both energy homeostasis and reproduction. Leptin does not directly regulate GnRH neurons, as the long form of the leptin receptor (LepRb), which is responsible for signal transduction, is not expressed in GnRH neurons [18, 19]. Rather, leptin acts on GnRH neurons indirectly, through upstream intermediates which have yet to be fully identified. RFRP-3 neurons may be one potential relay system through which leptin signals are mediated, as the DMN (where RFRP-3 neurons reside) is a leptin-responsive brain region [20–22]. Central injections of RFRP-3 not only inhibit LH secretion, but also stimulate feeding behavior [23, 24]. Moreover, RFRP-3 neurons are activated by chronic mild food restriction in hamsters [25], and RFRP-3's receptor, Gpr147, is required in mice to suppress LH secretion after acute food deprivation (FD) [26], suggesting that RFRP-3 plays a role in both energy balance and reproduction, as does leptin [27]. Thus, leptin might inhibit the production and/or secretion of RFRP-3 in order to facilitate reproductive function and/or suppress feeding behavior.

In addition to its roles in adulthood, leptin has important developmental effects on the hypothalamus. During the second week of postnatal life, serum leptin levels increase drastically and transiently in a postnatal leptin 'surge' [28]. This temporary increase in juvenile leptin levels regulates the development of axonal projections from the arcuate nucleus to the DMN, as well as other

brain regions [29, 30]. We previously demonstrated that DMN *Rfrp* expression, as measured by cell number and *Rfrp* mRNA levels per cell, is dramatically higher in juveniles on postnatal day (PND) 10 than at birth [9], but whether this developmental difference is caused, fully or in part, by the juvenile leptin surge is unknown.

In this study, we addressed whether neural *Rfrp* expression is regulated by metabolic manipulations, such as leptin deficiency seen in Ob animals or short-term FD. We also ascertained whether the developmental maturation of the neural *Rfrp* system is leptin dependent, owing to the developmental surge in leptin secretion during the juvenile period. Specifically, we determined whether (1) the mouse RFRP-3 peptide suppresses LH secretion, as reported for the rat RFRP-3 peptide variant in other species, (2) *Rfrp* neurons coexpress important metabolic genes, including *LepRb*, that are known to be expressed in the DMN, (3) *Rfrp* expression and/or neuronal activation is enhanced during FD or altered in adult Ob mice, and (4) the developmental pattern of *Rfrp* expression in postnatal mice parallels developmental leptin secretion (the juvenile leptin surge), and if so, whether normal *Rfrp* development is dependent on leptin signaling.

Materials and Methods

Animals, Gonadectomies, and Tissue Collection

Experiments utilized either C57BL6 mice or mice from the Ob strain [wild-type (WT) and Ob]. Mice from the Ob line were purchased from Harlan Laboratories and maintained on a C57BL6 background in the lab. All animals were housed on a 12:12-hour light-dark cycle, with food and water available ad libitum (ad lib), except where indicated in experiment 3. All experiments were conducted in accordance with the NIH Animal Care and Use Guidelines and with approval of the Animal Care and Use Committee of the University of California, San Diego, Calif., USA.

For adult C57BL6 mice, 7- to 9-week-old female (diestrous) or male mice were anesthetized with isoflurane, weighed, blood was collected via retro-orbital bleeding, and then they were rapidly decapitated. Prior to RFRP-3 injections and for the FD experiment, all C57BL6 mice were bilaterally gonadectomized (GDX), as previously described [9]. Gonadectomy was implemented to promote high levels of circulating LH for RFRP-3 or food restriction to inhibit, and to simultaneously control for differences in sex steroids between individuals since estradiol and testosterone can both mildly suppress *Rfrp* levels in mice of both sexes [9, 31]. For adult Ob mice, 7- to 8-week-old female or male WT or Ob mice were GDX (again, to control for sex steroids between genotypes and allow for maximal *Rfrp* expression levels in the absence of steroid inhibition) and then sacrificed 7 days later for blood and tissue collection.

For developmental experiments, infantile and juvenile pups were generated by C57BL6 breeder pairs or heterozygous Ob

breeder pairs (producing WT, Ob and heterozygous littermates). The date of birth was designated as PND 1. Newborn and juvenile pups of various ages were anesthetized with isoflurane, weighed, and rapidly decapitated. Trunk blood was collected from each animal to measure serum leptin levels. For mice of the Ob strain, tail samples were taken postmortem to determine the Ob genotype. Note that the obesity phenotype in Ob mice is not present until 4 weeks of age or later [16, 17].

Brains from all animals were collected at sacrifice, frozen on dry ice, stored at -80°C , and then sectioned on a cryostat into five coronal series of 20- μm brain sections which were thaw-mounted onto Superfrost-plus slides. Slides were stored at -80°C until use for in situ hybridization (ISH).

Hormone Assays

Blood from adult animals was collected by retro-orbital bleed. Blood from juvenile animals was collected via trunk blood. Serum LH was measured by the UVA Ligand Core (range: 0.04–37.4 ng/ml). Serum leptin was measured using the Quantikine Mouse Leptin ELISA Kit (R&D Systems) following the manufacturer's protocol (range: 1.25–80 ng/ml).

Single-Label and Double-Label ISH

The *Rfrp*, *c-fos*, *Trh*, and *LepRb* cRNA ISH riboprobes have been described and validated previously [9, 32–34]. Neuropeptide Y (*Npy*) and melanocortin receptor 4 (*Mc4r*) were cloned from mouse hypothalamic RNA into pBluescript SK-plasmid and transcribed using T7 polymerase. The probes correspond to the following bases of their respective murine mRNAs: *Npy*, bases 7–468 (NM_023456); *Trh*, bases 308–572 (NM_009426); *LepRb*, bases 3,379–3,912 (NM_146146), and *Mc4r*, bases 589–1,442 (NM_016977).

Single-label ISH was performed as previously described [9, 35]. Briefly, for each assay, slide-mounted brain sections encompassing the entire DMN from 1 of the 5 sets of serial brain sections were fixed in 4% paraformaldehyde, pretreated with acetic anhydride, rinsed in 2X SSC (sodium citrate, sodium chloride), delipidated in chloroform, dehydrated in ethanol, and air-dried. Radiolabeled (^{33}P) *Rfrp* antisense riboprobe (0.04 pmol/ml) was combined with tRNA, heat-denatured, added to hybridization buffer, and applied to each slide (100 μl /slide). Slides were cover-slipped and placed in a 55°C humidity chamber overnight. The slides were then washed in 4X SSC and placed into RNase A treatment for 30 min at 37°C , then in RNase buffer without RNase at 37°C for 30 min. After washing in 2X SSC at room temperature, slides were washed in 0.1X SSC at 62°C for 1 h, dehydrated in ethanols, and air-dried. Slides were then dipped in Kodak NTB emulsion, air-dried, and stored at 4°C for 3–4 days (depending on the assay) before being developed and cover-slipped.

Double-label ISH was performed as previously described [35, 36]. Briefly, one set of slide-mounted brain sections encompassing the entire DMN was fixed in 4% paraformaldehyde, pretreated with acetic anhydride, rinsed in 2X SSC (sodium citrate, sodium chloride), delipidated in chloroform, dehydrated in ethanols, and air-dried. Radio-labeled (^{33}P) antisense *Npy*, *Trh*, *LepRb*, or *Mc4r* (0.05 pmol/ml) were combined with digoxigenin (DIG)-labeled *Rfrp* riboprobes (Roche DIG labeling kit, 1:500) along with tRNA, denatured by boiling, and dissolved together in hybridization buffer. The probe mix was then applied to slides and hybridized at 55°C overnight. The slides were then washed

in 4X SSC and placed into RNase A treatment for 30 min at 37°C , then in RNase buffer without RNase at 37°C for 30 min. After washing in 2X SSC at room temperature, slides were washed in 0.1X SSC at 62°C for 1 h. Slides were then incubated in 2X SSC with 0.05% Triton X-100 and 3% sheep serum (NSS) for 75 min at room temperature, and then incubated overnight at room temperature with anti-DIG antibody conjugated to alkaline phosphatase (Roche; diluted 1:500 in buffer 1 containing 1% NSS and 0.3% Triton X-100). The next day, slides were washed with buffer 1 and then incubated with Vector Red alkaline phosphatase substrate (Vector Labs) for 1 h at room temperature. The slides were then air-dried, dipped in Kodak NTB emulsion, stored at 4°C , and developed and cover-slipped 3–9 days later, depending on the assay.

ISH slides were analyzed with an automated image processing system (Dr. Don Clifton, University of Washington) by a person blinded to the treatment group [37]. For single-label *Rfrp* experiments, the software counted the number of ISH silver grain clusters representing *Rfrp* cells as well as the number of silver grains over each cell (which provides a semiquantitative index of *Rfrp* mRNA expressed per cell) [38–40]. Cells were considered *Rfrp* positive when the number of silver grains in a cluster exceeded that of background by 3-fold. We have previously described two obvious *Rfrp* cell populations interspersed within the DMN that dramatically differ in silver grain (i.e. mRNA) expression. Therefore, as in previous studies, we divided *Rfrp* cells into two categories based on their *Rfrp* mRNA expression level; 'high expressing' (HE) cells have a silver grain cluster diameter greater than $>20\text{ }\mu\text{m}$, whereas 'low expressing' (LE) cell have a silver grain cluster less than $<20\text{ }\mu\text{m}$. Previous studies have determined that silver grain number and silver grain cell area are highly dependent, representative of mRNA differences, and not due to morphological cell size differences [9]. All *Rfrp* cells from one set of serial sections were quantified in each animal. For double-label ISH assays, all red fluorescent DIG-containing cells (*Rfrp* cells) from one set of serial brain sections were identified under fluorescence microscopy and the grain-counting software was used to quantify silver grains (representing either *LepRb*, *Npy*, *Trh*, or *Mc4r* mRNA) overlying each DIG cell. Signal-to-background ratios for individual cells were calculated, and a cell was considered double-labeled if its ratio was >3 [35, 36, 41, 42].

Statistical Analysis

All data are expressed as means $\pm$ SEM for each group. In all experiments, differences were analyzed by Student's t test, ANOVA, or 2-way ANOVA (for age and genotype), followed by post hoc comparisons for individual age/genotype groups via Fisher's (protected) LSD. Statistical significance was set at $p < 0.05$. All analyses were performed in StatView 5.0.1 (SAS Institute, Cary, N.C., USA).

Experiment 1: Does RFRP-3 Inhibit LH Secretion in Mice?

The conserved role of RFRP-3 in the inhibition of LH secretion is assumed to be true in mice, but, surprisingly, data demonstrating this has not yet been reported. In order to test if the mouse RFRP-3 peptide is functionally able to suppress LH secretion in mice, as occurs with other RFRP-3 variants in other species, adult ovariectomized mice were subjected to either 100 ng or 500 ng of the murine RFRP-3 peptide (VNMEAGTRSHFPLPQRF-NH₂, Genscript USA Inc.) dissolved in 100 μl of saline or saline vehicle.

Twenty minutes after intraperitoneal injection, blood was collected by retro-orbital bleed and the serum was assayed for LH (n = 7–8 per treatment group).

*Experiment 2: What Metabolic Neuropeptides and Receptors Are Expressed in *Rfrp* Neurons?*

The phenotypic identity of *Rfrp* neurons is virtually unknown. Besides estrogen and glucocorticoid receptors, other receptors or secreted cofactors, such as neurotransmitters or neuropeptides, have not been identified in *Rfrp* neurons. Any number of metabolic genes expressed in the DMN region may colocalize with *Rfrp*, and if so, would give insight to the regulation and potential metabolic functions of RFRP-3 neurons. We therefore examined whether transcripts of important metabolic genes already known to be expressed in the DMN specifically colocalize with *Rfrp*. Adult C57BL6 mice of both sexes (females in diestrous) were sacrificed and their brains collected for double-label ISH analyses. Using alternate series of coronal brain sections (encompassing the entire rostral to caudal span of the DMN) from each mouse, we examined in 4 separate assays the coexpression of neuropeptide Y (*Npy*) [43–45], *Thr* [34, 46], *LepRb* [47], and melanocortin receptor 4 (*Mcr4r*) [48] in *Rfrp* neurons (n = 4–6 per sex).

*Experiment 3: Is *Rfrp* Expression or Neuron Activity Responsive to Short-Term Metabolic Challenge?*

Given that RFRP-3 treatment increases food intake [7, 49] and that a subset of *Rfrp* neurons express the leptin receptor (experiment 2), it is possible that transiently diminished serum leptin levels achieved via short-term FD may alter *Rfrp* expression and/or *Rfrp* neuronal activation. Furthermore, a recent report demonstrated that LH secretion is not suppressed after 12 h of FD in *Gpr147* knockout mice, suggesting RFRP-3 is essential during acute FD to suppress LH [26]. To further examine this, adult C57BL6 female mice were first GDX (to equate sex steroids between groups since E₂ can inhibit *Rfrp* levels) and were then later subjected to short-term (12-hour) FD or kept on ad lib feeding (controls) (n = 7 per group). All food was removed from FD animals at lights off (18:00 h) and all animals were sacrificed 12 h later at lights on (06:00 h) the following morning. Cages of ad lib control mice were opened, but food was not removed. Brains and blood were collected immediately at the end of the 12-hour FD or ad lib feeding period. *Rfrp* expression in the DMN was analyzed by single-label ISH and, in an alternate set of brain sections from the same animals, double-label ISH was performed for *Rfrp* and *c-fos*, a marker of neuronal activation. Blood serum collected at the time of sacrifice was assayed for circulating LH and leptin.

*Experiment 4: Is *Rfrp* Expression Altered in Adult Ob Animals?*

Since experiment 2 found that a subset of *Rfrp* neurons express *LepRb*, we hypothesized that *Rfrp* expression may be leptin regulated. To determine if *Rfrp* levels are altered with chronic leptin deprivation, we measured *Rfrp* neuron numbers and cellular *Rfrp* expression levels using single-label ISH in adult male and female GDX Ob mice (leptin deficient) and their WT GDX littermates (n = 5–7 animals per genotype per sex).

*Experiment 5: Does *Rfrp* Expression during Postnatal Development Coincide with Serum Leptin Levels?*

Experiment 4 demonstrated that adult Ob mice display altered *Rfrp* expression, suggesting that leptin may regulate RFRP-

3 neurons. In development, there is a large transient rise in serum leptin that occurs during the second week of life in mice. Around this same age, *Rfrp* expression is significantly higher than on the day of birth [9]. In order to see if the juvenile leptin surge coincides with developmental changes in *Rfrp* neurons, C57BL6 pups were sacrificed by rapid decapitation on PND 1, 3, 6, 8, 10, 12, 14, and 16 (n = 5–9 animals per age, per sex). Only litters of 4–8 pups were used. Trunk blood was collected for serum hormone analysis, and brains were collected and frozen on dry ice. For each sex, brain slices containing the entire DMN were assayed and analyzed for *Rfrp* expression by single-label ISH, using the HE and LE *Rfrp* cell criteria described previously (see the section ‘Single-label and Double-Label ISH’ above) [36]; each sex was assayed independently. Serum leptin was measured by ELISA. Along with the postnatal mice, cohorts of control adult diestrous female or male mice were simultaneously assayed for *Rfrp* expression and serum leptin levels.

In a complementary experiment, to determine if the long-form leptin receptor is expressed in juvenile *Rfrp* neurons at different developmental stages, an alternate set of brain slides from female PND 1, 10, and 12 mice (along with adult diestrous female controls) was assayed for *LepRb* coexpression in *Rfrp* neurons using double-label ISH (n = 6–7 animals per age group).

*Experiment 6: Are the Juvenile Changes in *Rfrp* Expression during Development Dependent on Leptin?*

Given the synchronous developmental changes in juvenile *Rfrp* expression and postnatal leptin secretion observed in experiment 5, we hypothesized that the developmental changes in *Rfrp* neurons may be leptin dependent. To test this, we repeated experiment 5 using heterozygous Ob breeding pairs to produce WT, Ob, and heterozygous pups. Brains, trunk blood, and tail snips for genotyping were collected on PND 1, 6, 10, 12, and 16. Only litters of 4–8 pups were used. Brains from WT and Ob males of these various postnatal ages were analyzed for *Rfrp* mRNA expression levels via single-label ISH (n = 5–9 animals per age per genotype). Serum leptin for each animal was measured via ELISA to confirm the presence or absence of circulating leptin.

Results

Experiment 1: The Murine RFRP-3 Peptide Inhibits LH Secretion in Adult Female Mice

To determine if peripherally administered murine RFRP-3 peptide can suppress LH secretion in mice, as occurs with other RFRP-3 variants in other species, 100 ng or 500 ng of RFRP-3 or saline was injected i.p. in adult GDX female mice. Both the 100-ng and 500-ng doses of murine RFRP-3 were able to significantly suppress LH secretion, by nearly 70% (fig. 1; p < 0.05). There was no difference in the efficacy of the two doses of murine RFRP-3 on LH suppression.

Fig. 1. Effect of peripheral murine RFRP-3 injection on LH secretion in mice. Adult GDX female mice were injected i.p. with 100 ng or 500 ng of murine RFRP-3 dissolved in 100 μ l of saline, or 100 μ l of saline alone, and bled 20 min later. RFRP-3 significantly suppressed LH secretion in GDX mice at both doses ($p < 0.05$).

Fig. 2. Expression of hypothalamic energy balance genes in the dorsal-medial hypothalamus. Representative photomicrographs of double-label ISH of neuropeptide Y (*NPY*), *TRH*, *LepRb*, or *Mc4r* (silver grains) in *Rfrp* neurons (red fluorescence). Single-labeled *Rfrp* neurons are identified with blue arrows and single-labeled *NPY*, *TRH*, *LepRb*, and *Mc4r* neurons are marked with yellow arrows. Double-labeled *LepRb/Rfrp* and double-labeled *Mc4r/Rfrp* neurons are marked with green arrowheads.

Experiment 2: A Subset of *Rfrp* Neurons Express *LepRb* mRNA

Double-label ISH was performed for energy balance-related genes known to be expressed in the DMN [34, 43–48] to determine if they are coexpressed in *Rfrp* neu-

rons of adult male or female mice. We found that virtually no *Rfrp* neurons coexpressed *Npy* or *Trh*, despite notable *Npy* and *Trh* expression in the same DMN region, often near *Rfrp* neurons (fig. 2; table 1). Next, to assess the possibility that metabolic hormones/neuropeptides

Table 1. Summary of four double-label ISH assays for metabolic-related genes in the DMN coexpressed with *Rfrp* neurons

	Intact males	DE females
NPY/ <i>Rfrp</i>	1.5±0.7%	3.0±1.5%
TRH/ <i>Rfrp</i>	1.1±0.5%	1.6±0.8%
LepRb/ <i>Rfrp</i>	16.9±1.6%	13.0±1.6%
Mc4r/ <i>Rfrp</i>	6.9±1.0%	8.7±0.7%

Adult diestrous (DE) females and intact male mice were assayed for neuropeptide Y (NPY), TRH, LepRb, or Mc4r mRNA in *Rfrp* cells in the DMN. Values are average percent coexpression ± SEM. n = 4–6 animals per sex.

might directly regulate the RFRP-3 system, we looked for coexpression of several metabolic signaling factor receptors in *Rfrp* neurons. *LepRb* mRNA was found to be notably coexpressed in a subset of *Rfrp* neurons, suggesting that a proportion of adult *Rfrp* neurons may be leptin responsive. The degree of *LepRb-Rfrp* coexpression was similar in both sexes (~15%). An even smaller subset of *Rfrp* neurons (~8%) in both sexes was found to coexpress the primary melanocortin receptor, *Mc4r*. A summary of the coexpression levels for each gene with *Rfrp* is in table 1.

Experiment 3: *Rfrp* Neurons in Mice Appear Unresponsive to 12-Hour FD

To test if energetic challenge induced by short-term FD modifies *Rfrp* mRNA expression levels or the activity of RFRP-3 neurons, adult female mice were subjected to short-term (12-hour) FD. Demonstrating the effectiveness of the energetic challenge, 12-hour FD mice had a significant decrease in body weight (5.3%; $p < 0.05$) and significant decreases in serum LH and leptin relative to ad lib control females (fig. 3a–c; $p < 0.05$). However, despite this negative impact of 12-hour FD on body weight and endocrine physiology, there were no significant differences in either the total number of *Rfrp* neurons or the total *Rfrp* mRNA between ad lib and FD animals (fig. 3d–f). Moreover, when *Rfrp* neurons were categorized as either HE or LE, there were no statistical differences between ad lib and FD animals (data not shown).

In addition to mRNA levels, we also used double-label ISH to examine alterations in neuronal activation of RFRP-3 neurons in 12-hour FD mice. As with the single-label assay, the total number of *Rfrp* cells in this double-label assay did not differ between 12-hour FD and ad lib mice (117 ± 8 cells vs. 121 ± 8 cells, respectively). More-

over, there was no group difference in the percent of *Rfrp* neurons demonstrating recent neuronal activation, as measured by *c-fos* mRNA coexpression in *Rfrp* neurons (fig. 3g, h).

Experiment 4: *Rfrp* Expression Is Decreased in Adult Ob Mice of Both Sexes

This experiment assessed whether there are notable differences in *Rfrp* expression in adult Ob (leptin deficient) and WT mice that were all GDX beforehand to remove any masking effects of circulating E_2 (which moderately decreases *Rfrp* expression). As expected, Ob animals of both sexes were significantly heavier, weighing almost twice as much than their WT counterparts ($p < 0.05$, data not shown). In the brain, the total number of detectable *Rfrp* neurons was not significantly different between Ob and WT mice in either sex (fig. 4a, b). However, using the HE and LE *Rfrp* cell criteria, there was a significant decrease in the number of HE *Rfrp* neurons in female Ob mice compared to female WT mice (fig. 4a; $p < 0.05$), with no significant differences in the number of LE *Rfrp* cells between genotypes. Likewise, a similar reduction in the number of HE *Rfrp* neurons was detected in male Ob mice compared to WT male littermates (fig. 4b; $p < 0.05$). In addition to these changes in cell number, the total level of *Rfrp* mRNA in the DMN was significantly lower in both male and female Ob mice compared to their WT littermate controls (fig. 4a, b; $p < 0.05$).

Experiment 5: Changes in *Rfrp* Expression over Postnatal Development Correlate with the Leptin Surge

We previously reported that *Rfrp* levels in the brain are dramatically higher in juvenile mice (on PND 10) compared to birth. Since experiment 4 demonstrated that *Rfrp* expression is significantly altered in adult Ob animals of both sexes, we hypothesized that the juvenile leptin surge [28] might govern the developmental changes in *Rfrp* expression. We first determined whether there was a correlation between juvenile leptin levels and postnatal developmental changes in *Rfrp* expression. C57BL6 mice of both sexes were sacrificed on PND 1, 3, 6, 8, 10, 12, 14, and 16, and their brains were assayed for *Rfrp* expression (fig. 5) and compared to serum leptin levels at each age. The total number of detectable *Rfrp* neurons was highest at birth and was significantly lower at later postnatal ages, with the lowest number of *Rfrp* neurons in adulthood (diestrous stage for females; fig. 6a). However, the amount of *Rfrp* expressed per cell was lowest at birth and increased substantially with age. Indeed, the

Fig. 3. Effects of short-term food deprivation on *Rfrp* expression and *Rfrp* neuron activation in GDX female mice. **a** 12-hour FD had a significant effect on the change in body weight ($p < 0.05$) with deprived animals losing almost 6% of body weight on average. **b** Serum LH is significantly less in 12-hour FD GDX animals than ad lib animals ($p < 0.05$). **c** Serum leptin is significantly less in 12-hour FD GDX animals than ad lib animals ($p < 0.05$). **d** Representative photomicrographs of single-label ISH for *Rfrp* mRNA in ad lib and 12-hour FD female GDX mice. 3V = Third ventricle. **e** The total number of *Rfrp* neurons is not significantly different between

ad lib and 12-hour FD animals. **f** The total number of *Rfrp* mRNA is not significantly different between ad lib and 12-hour FD animals. **g** Representative photomicrographs of double-label ISH for *c-fos* and *Rfrp* mRNAs in ad lib and 12-hour FD female GDX mice. Double-labeled *c-fos* and *Rfrp* (green arrowhead), single-labeled *c-fos* cells (yellow arrow), and single-labeled *Rfrp* neurons (blue arrow) are shown. **h** Quantification of percent *c-fos* and *Rfrp* coexpression in ad lib and 12-hour FD animals showed no significant difference between the two groups.

Fig. 4. *Rfrp* expression in adult Ob male and female GDX mice. **a** Representative photomicrographs of single-label ISH for *Rfrp* mRNA in WT and Ob female GDX mice. 3V = Third ventricle. **b** Quantification of total *Rfrp* neurons, HE *Rfrp* neurons, LE *Rfrp* neurons, and total *Rfrp* mRNA from single-label ISH in female GDX mice. The number of HE cells and the total *Rfrp* mRNA was significantly less in Ob females than WT littermates ($p < 0.05$).

There were no significant differences between the total number of *Rfrp* neurons or number of LE cells. **c** Quantification of total *Rfrp* neurons, HE *Rfrp* neurons, LE *Rfrp* neurons, and total *Rfrp* mRNA from single-label ISH in male GDX mice. The number of HE cells and the total *Rfrp* mRNA was significantly less in Ob males than WT littermates ($p < 0.05$). There were no significant differences between the total number of *Rfrp* neurons or number of LE cells.

Fig. 5. *Rfrp* expression over the first 16 days of postnatal life. Representative photomicrographs of single-label ISH for *Rfrp* mRNA on PND 1, 3, 6, 8, 10, 12, 14, and 16 in female mice. Day of birth is designated PND 1. 3V = Third ventricle.

total amount of *Rfrp* mRNA in the DMN increased with postnatal development, beginning to rise around PND 6 and peaking at PND 12 before dropping again at subsequent older ages (fig. 6b; $p < 0.05$). This robust increase in total *Rfrp* mRNA during the 2nd week of postnatal life appears to reflect a large increase in the number of HE *Rfrp* cells, which are virtually absent at birth, but substantially increase in abundance around PND 6 and peak at maximal levels at PND 12 before dropping slightly at subsequent older ages (fig. 6c; $p < 0.05$). Unlike HE *Rfrp* cells, the number of LE cells decreases slowly and steadily throughout these postnatal ages, with the lowest number of LE cells in adulthood (fig. 6d; $p < 0.05$). Virtually identical results were found in similarly aged male mice (data not shown).

Serum leptin levels were measured in each animal to see if they correlated with *Rfrp* levels. For both sexes, mean serum leptin levels were essentially undetectable at birth (PND 1) and on PND 3, and were first readily detectable at PND 6. Leptin levels increased robustly dur-

ing the second week of life, peaking at PND 10 and 12 (the leptin surge), and subsequently began to fall on PND 14 and 16, being low again in adult animals (fig. 6e; $p < 0.05$). Leptin levels were significantly higher than adult female (diestrous) levels at all postnatal ages examined, except for PND 1 and 3 ($p < 0.05$). Interestingly, these postnatal changes in leptin levels correlated well with the observed increases in the number of HE *Rfrp* cells and total *Rfrp* mRNA, with each starting to increase around PND 6 and reaching maximal levels around PND 10 and 12 (fig. 6).

Experiment 6: Juvenile Rfrp Expression Is Not Dependent on Leptin during Development

The dramatic developmental increase in HE *Rfrp* neurons and total *Rfrp* mRNA observed around PND 10 and 12 correlates with the zenith of the juvenile leptin surge (experiment 5), suggesting that the two events may be related. To test this possibility, we determined whether *Rfrp* neurons coexpress *LepRb* during specific postnatal ages

Fig. 6. Summary of changes in *Rfrp* expression over the first 16 days of postnatal life in female mice from single-label ISH. **a** Quantification of total number of *Rfrp* neurons during the first 2 weeks of life. Different letters indicate significantly different groups ($p < 0.05$). **b** Quantification of total *Rfrp* mRNA during the first 2 weeks of life. Different letters indicate significantly different groups ($p < 0.05$). **c** Quantification of the number of HE *Rfrp* neurons during

the first 2 weeks of life. Different letters indicate significantly different groups ($p < 0.05$). **d** Quantification of the number of LE *Rfrp* neurons during the first 2 weeks of life. Different letters indicate significantly different groups ($p < 0.05$). **e** Average serum leptin levels of mice sacrificed across development and in a cohort of diestrous female adult mice. Different letters indicate significantly different groups ($p < 0.05$).

and whether the normal developmental pattern and levels of *Rfrp* expression are altered in juvenile mice lacking leptin. In postnatal brains examined for *Rfrp*/*LepRb* coexpression, a subset of *Rfrp* cells clearly coexpressed notable *LepRb* mRNA (fig. 7a); the degree of coexpression did not vary with age, being 15–20% from birth through adulthood (fig. 7b).

To determine if leptin signaling is required, directly or indirectly, for the normal pattern of *Rfrp* neuron development in juveniles, female Ob and WT brains were analyzed for *Rfrp* expression on PND 1, 6, 10, 12, and 16 (fig. 8, 9). Despite the strong correlation between developmental *Rfrp* and leptin changes observed in normal mice, there were no significant differences in the number

Fig. 7. *LepRb* expression in some *Rfrp* neurons during postnatal life. **a** Left: representative photomicrographs of double-label ISH for *LepRb* (silver grains) in *Rfrp* neurons (red fluorescence) in the DMN. A single-labeled *Rfrp* neuron is identified with a blue arrow and a single-labeled *LepRb* neuron is marked with a yellow arrow. Double-labeled *LepRb/Rfrp* is marked with a green arrowhead. Right: strong *LepRb* mRNA staining in the ARC nucleus (left side of nucleus shown) in the same double-label *LepRb/Rfrp* assay. 3V = Third ventricle. **b** Quantification of the percent coexpression of *LepRb* and *Rfrp* during postnatal development. No significant differences were seen between ages. The leptin surge peaked around PND 10 and 12, as indicated by the horizontal red bar.

of *Rfrp* neurons, total amount of *Rfrp* mRNA, or the number of HE and LE *Rfrp* cells between Ob and WT mice at any developmental age (fig. 9a–d). The same developmental pattern in *Rfrp* expression seen in experiment 5 was observed in both genotypes, with identical magnitude increases detected in total *Rfrp* mRNA and the number of HE cells. Moreover, all *Rfrp* expression increases occurred at the same specific postnatal ages, regardless of genotype. Confirming the genotypes and leptin milieu, serum leptin was significantly elevated between PND 6 and 12 (representing the leptin surge), and lower on PND 16 than 12 in WT animals ($p < 0.05$), but remained undetectable in Ob animals at all ages (fig. 9e).

Discussion

The physiological roles RFRP-3 plays and how the hypothalamic RFRP-3 system develops and is regulated are still poorly understood. In addition to providing the first

demonstration of the ability of peripherally administered RFRP-3 to potentially inhibit LH secretion in vivo in mice, our current findings also show that a subset of RFRP-3 neurons express the long form of the leptin receptor and that *Rfrp* mRNA expression is diminished in Ob (leptin-deficient) animals of both sexes. We also demonstrate that *Rfrp* expression changes in a dramatic manner in postnatal development, mirroring age-specific juvenile changes in serum leptin. However, examination of the developmental maturation of *Rfrp* neurons in Ob pups revealed that neither the timing nor magnitude of these developmental changes in *Rfrp* expression are dependent on leptin.

Despite the reported effect of the rat variant of RFRP-3 to inhibit LH secretion in many species, including rodents (rats and hamsters), the effect of the murine variant of RFRP-3 on LH has not been reported in any species, including mice themselves. As the RFRP-3 field moves towards transgenic and knockout animals [26, 50], mouse models will become increasingly important and under-

Fig. 8. *Rfrp* expression over postnatal life in female Ob and WT littermates. Representative photomicrographs of single-label ISH for *Rfrp* mRNA on PND 1, 10, and 12 are shown. 3V = Third ventricle.

standing murine RFRP-3 pharmacology and physiology is essential. Our present finding of markedly suppressed LH secretion 20 min after peripheral injection of two doses of murine RFRP-3 in adult female mice matches the results found in rats, hamsters, and sheep using the rat RFRP-3 peptide [2, 51–54]. Until just recently, RFRP-3 of any variant had surprisingly never been administered to mice. León et al. [26] reported that a truncated rat RFRP-3 peptide given via intracerebroventricular injection modestly decreased serum LH (by ~25%) in WT GDX female mice. That reduction in LH was considerably less than the suppressive effects of peripheral murine RFRP-3 observed in the present study, in which we found ~70% decrease in serum LH after peripheral murine RFRP-3 injections at two different doses. Besides potential species differences in murine versus rat RFRP-3 peptide efficacy, the difference in magnitude of effect might be due to dosing or the route of administration. Interestingly, the 100- and 500-ng doses were equally effective in suppressing LH secretion, suggesting that the suppressive effect of

RFRP-3 may already be maximal near the 100-ng dose. As our RFRP-3 was administered peripherally, we cannot discern at what level(s) of the HPG axis the lowered LH secretion is attributable to. However, most data in rodents suggests that RFRP-3's effects on LH are induced in the brain, via diminished GnRH signaling, rather than at the pituitary [8]. Yet, one study recently described in vitro effects of RFRP-3 on the pituitary in murine gonadotrope L β T2 cells [55].

Several studies have demonstrated an orexigenic effect of RFRP-3 [7, 24, 49], suggesting that RFRP-3 may be involved in receiving and/or transmitting energy balance signals. Both neuropeptide Y and thyrotropin-releasing hormone (TRH) are neuropeptide cell populations that reside in the DMN, and are even expressed in a similar pattern as *Rfrp* [34, 45]. However, *Rfrp* did not colocalize with either of these neuropeptides, demonstrating that it is a unique neuropeptide population with. We also examined two metabolic hormone receptors, the long-form leptin receptor, *LepRb*, and a melanocortin receptor,

Fig. 9. Summary of changes in *Rfrp* expression on PND 1, 6, 10, 12, and 16 in female Ob and WT animals. **a** Quantification of total number of *Rfrp* neurons during the leptin surge. There were no significant changes between genotypes. **b** Quantification of total *Rfrp* mRNA during the leptin surge. *Rfrp* expression is significantly higher on PND 10, 12, and 16 than on PND 1 or 6 ($p < 0.05$). **c** Quantification of the number of HE *Rfrp* neurons during the leptin surge. The number of *Rfrp* neurons is significantly higher on PND 10, 12, and 16 than on PND 1 or 6 ($p < 0.05$). **d** Quantifica-

tion of the number of LE *Rfrp* neurons during the leptin surge. Different letters indicate significantly different groups ($p < 0.05$). **e** Serum leptin concentration in Ob and WT female mice as measured by ELISA. Leptin is highest on PND 10 and 12, dropping on 16, demonstrating the postnatal leptin surge. Leptin concentration in WT mice were significantly elevated between PND 6 and 16, peaking around PND 12 ($p < 0.05$). Leptin was not detectable on the day of birth in WT mice or in Ob animals at any age.

Mc4r, which are also expressed in the DMN. A small subset of *Rfrp* neurons expressed each of these receptors, with the leptin receptor being the more prominently expressed, suggesting that some *Rfrp* neurons may be a direct target of leptin signaling. However, most *Rfrp* neurons did not express LepR, suggesting that any major effects of leptin

signaling – or its absence – on *Rfrp* neurons may in fact be indirect.

Ob mice are infertile due to a lack of gonadotropin secretion [27], and since RFRP-3 is known to inhibit LH secretion, and some *Rfrp* neurons express the leptin receptor, we hypothesized that *Rfrp* expression would be higher in

Ob mice. However, using GDX adult Ob males and females, we found the opposite outcome: *Rfrp* expression is lower in adult Ob mice than their WT littermates. This was true for both sexes. During the writing of this manuscript, Rizwan et al. [56] published data suggesting that RFRP-3 neurons are not regulated by leptin. That study's immunohistochemical data showed no difference between the number of RFRP-3-immunoreactive neurons in GDX, E₂-replaced Ob mice and WT littermates, and minimal phosphorylation of signal transducer and activator of transcription 3 in RFRP-3 neurons after leptin injections. Those results suggest that RFRP-3 neurons are not directly regulated by leptin and that RFRP-3-immunoreactive levels are unaffected by leptin deficiency, differing from our present results in which *Rfrp* levels were significantly diminished in Ob mice. It is possible that the exogenous E₂ replacement in the Rizwan study may have masked differences between WT and Ob animals, as E₂ moderately represses *Rfrp* [9, 31], and may have done so similarly in both genotypes. Our present data quantifying gene expression (rather than protein) levels demonstrates that total *Rfrp* mRNA levels are significantly repressed in GDX Ob animals (no E₂ given, so *Rfrp* expression is not being inhibited by sex steroids). Additionally, in our Ob mice, we identified a primary deficit in *Rfrp* expression specifically in the HE *Rfrp* cell population, which is a cellular distinction that has not yet been analyzed using immunohistochemical data. The functional significance of the lower *Rfrp* levels in Ob mice is not yet known; RFRP-3 has proposed roles in several other physiological/behavioral systems beyond reproduction, such as feeding, stress, and anxiety, and the observed decreases in *Rfrp* levels in Ob mice may reflect RFRP-3's involvement in these processes rather than the reproductive impairments of these animals.

Although we show a notable decrease in overall *Rfrp* levels in adult Ob mice of both sexes, it remains unclear if this is due to changes in direct or indirect leptin signaling to RFRP-3 cells. Rizwan et al. [56] showed minimal phosphorylation of signal transducer and activator of transcription 3 in RFRP-3 neurons after leptin treatment, ranging from ~3% in mice to 7–13% in rats, suggesting that any effects of leptin would likely be indirect. In the present study, using double-label techniques, we found direct anatomical evidence for a slightly higher degree of *LepRb* expression in *Rfrp* neurons, 15–20% in female and male mice, suggesting that some effects of leptin could in fact be directly on a subset of *Rfrp* neurons. Yet, based on the overall low-to-moderate percentage of total *Rfrp* neurons expressing *LepRb*, we concur with Rizwan et al., that the majority of RFRP-3 neurons in rodents are likely not

directly responsive to leptin signaling. If so, the diminished *Rfrp* levels observed in Ob mice may reflect any of the following: (1) direct leptin signaling in only a small subset (~15–20%) of *Rfrp* cells, causing only those cells to change gene expression, (2) indirect leptin signaling on upstream leptin-responsive circuits that themselves directly regulate RFRP-3 neurons, and (3) influence of non-leptin metabolic signals that are secondarily altered in the obese state of Ob mice. This last possibility seems likely since *Rfrp* expression was normal in young juvenile Ob mice, in which leptin signaling is absent but obesity has yet to emerge. Of note, despite the observed low-to-moderate levels of *LepR-Rfrp* coexpression, very strong *LepR* expression was observed in the ARC and other areas of the same brain sections, demonstrating the robustness of *LepR* probe staining in our assays.

Central injections of RFRP-3 reportedly stimulate food intake [23, 24], and RFRP-3 neurons are activated by chronic mild (20%) food restriction in hamsters [25]. Moreover, a recent report demonstrated that LH secretion is not suppressed after 12 h of FD in *Gpr147* knockout mice (lacking the RFRP-3 receptor), suggesting that RFRP-3 is involved in the metabolic regulation of LH [26]. We therefore examined whether *Rfrp* expression levels or *Rfrp* neuronal activation is altered in adult female mice subjected to 12-hour FD. This short-term FD had no effect on *Rfrp* expression or *Rfrp* neuronal activation. These results are surprising, as body weight and LH levels were significantly lower after FD, reflecting effective metabolic challenge and inhibition of the reproductive system. Thus, the lower LH levels observed in our metabolically challenged females, which we hypothesized to be a direct or indirect effect of enhanced RFRP-3, appear to be decreased by some other mechanism. The previously observed increase in *Rfrp* neuronal activation in hamsters that were food restricted to 80% of normal diet for much longer periods of time (8 days or more [25]) may suggest that metabolic alteration of *Rfrp* neurons may be a part of either chronic or mild nutritional stress, rather than acute, severe changes in energy availability, as in our 12-hour FD paradigm.

Over the course of postnatal development, there are drastic age-specific changes in neural *Rfrp* expression, especially in the late infantile and juvenile phases. Previously, we demonstrated that both the number of HE *Rfrp* cells and the total amount of *Rfrp* mRNA per cell are considerably lower at birth than on PND 10, at which time the number of LE *Rfrp* cells is lower than at birth [9]. Here, we determined a more detailed time-line during the first 2 weeks of life. In normal C57BL6 mice, the develop-

mental increase in HE *Rfrp* cells and total *Rfrp* mRNA appears around PND 6–8, with a peak in these measures around PND 12. This was in striking synchronous alignment with the occurrence of the postnatal leptin surge. However, using Ob mice, we found that neither the timing nor magnitude of this developmental pattern of the *Rfrp* population is leptin dependent. The developmental changes observed in *Rfrp* expression may be due to other metabolic hormones, such as ghrelin or insulin [57–59], or reproductive hormones from maturing gonads [60–63], all of which have been shown to modify gene expression in the developing hypothalamus. Identifying the cause of the dramatic *Rfrp* changes during development, and just as importantly, their functional significance, will be an important avenue of future inquiry.

In conclusion, the data presented here provide further insight into the regulation and development of RFRP-3 neurons in mice. We show for the first time that peripherally administered murine RFRP-3 peptide can strongly inhibit LH secretion in adult female mice, consistent with reports for other RFRP-3 variants in other species. We also show that although *Rfrp* neurons do not express neuropeptide Y or TRH, there does appear to be a small subpopulation of *Rfrp* neurons that coexpress *LepRb*, allowing for the possibility of direct leptin signaling in these cells. This finding is extended by the novel observation that adult Ob animals have less *Rfrp* expression than their WT littermates, though whether this decrease in *Rfrp* expression is due to direct or indirect effect of leptin signaling, or other

metabolic cues associated with obesity, remains to be determined. Additionally, we demonstrate that *Rfrp* expression in female mice is resistant to short-term FD, despite previous reports demonstrating effects of long-term mild food restriction on RFRP-3 neurons in other species. Thus, the observed decrease in LH after short-term FD in mice is unlikely to be due to changes in RFRP-3 neurons. Lastly, we characterize the detailed developmental pattern of *Rfrp* expression in the rodent brain during infantile and juvenile life, and though this pattern correlates with leptin levels, we find it is independent of leptin signaling. The mechanism and physiological significance of these dramatic *Rfrp* changes during juvenile development remains to be determined.

Acknowledgements

This research was supported by NIH grant R01 HD065856. Additional support was provided by the Eunice Kennedy Shriver NICHD/NIH (SCCPIR) grant U54-HD012303 (U.C. San Diego, Calif., USA) and U54 HD-28934 (University of Virginia Ligand Assay and Analysis Core). The authors thank Kristen Tolson and Joshua Kim for technical assistance and comments on this manuscript.

Disclosure Statement

The authors have nothing to disclose.

References

- 1 Tsutsui K, Saigoh E, Ukena K, Teranishi H, Fujisawa Y, Kikuchi M, Ishii S, Sharp PJ: A novel avian hypothalamic peptide inhibiting gonadotropin release. *Biochem Biophys Res Commun* 2000;275:661–667.
- 2 Kriegsfeld LJ, Mei DF, Bentley GE, Ubuka T, Mason AO, Inoue K, Ukena K, Tsutsui K, Silver R: Identification and characterization of a gonadotropin-inhibitory system in the brains of mammals. *Proc Natl Acad Sci U S A* 2006;103:2410–2415.
- 3 Anderson GM, Relf HL, Rizwan MZ, Evans JJ: Central and peripheral effects of RFamide-related peptide-3 on luteinizing hormone and prolactin secretion in rats. *Endocrinology* 2009;150:1834–1840.
- 4 Ducret E, Anderson GM, Herbison AE: RFamide-related peptide-3, a mammalian gonadotropin-inhibitory hormone ortholog, regulates gonadotropin-releasing hormone neuron firing in the mouse. *Endocrinology* 2009;150:2799–2804.
- 5 Wu M, Dumalska I, Morozova E, van den Pol AN, Alreja M: Gonadotropin inhibitory hormone inhibits basal forebrain vGluT2-gonadotropin-releasing hormone neurons via a direct postsynaptic mechanism. *J Physiol* 2009;587:1401–1411.
- 6 Khan AR, Kauffman AS: The role of kisspeptin and RFRP-3 neurons in the circadian-timed preovulatory luteinizing hormone surge. *J Neuroendocrinology* 2012;24:131–143.
- 7 Johnson MA, Tsutsui K, Fraley GS: Rat RFamide-related peptide-3 stimulates GH secretion, inhibits LH secretion, and has variable effects on sex behavior in the adult male rat. *Horm Behav* 2007;51:171–180.
- 8 Rizwan MZ, Poling MC, Corr M, Cornes PA, Augustine RA, Quennell JH, Kauffman AS, Anderson GM: RFamide-related peptide-3 receptor gene expression in GnRH and kisspeptin neurons and GnRH-dependent mechanism of action. *Endocrinology* 2012;153:3770–3779.
- 9 Poling MC, Kim J, Dhamija S, Kauffman AS: Development, sex steroid regulation, and phenotypic characterization of RFamide-related peptide (Rfrp) gene expression and RFamide receptors in the mouse hypothalamus. *Endocrinology* 2012;153:1827–1840.
- 10 Hinuma S, Shintani Y, Fukusumi S, Iijima N, Matsumoto Y, Hosoya M, Fujii R, Watanabe T, Kikuchi K, Terao Y, Yano T, Yamamoto T, Kawamata Y, Habata Y, Asada M, Kitada C, Kurokawa T, Onda H, Nishimura O, Tanaka M, Ibata Y, Fujino M: New neuropeptides containing carboxy-terminal RFamide and their receptor in mammals. *Nat Cell Biol* 2000;2:703–708.
- 11 Liu Q, Guan XM, Martin WJ, McDonald TP, Clements MK, Jiang Q, Zeng Z, Jacobson M, Williams DL Jr, Yu H, Bomford D, Figueroa D, Mallee J, Wang R, Evans J, Gould R, Austin CP: Identification and characterization of novel mammalian neuropeptide FF-like peptides that attenuate morphine-induced antinociception. *J Biol Chem* 2001;276:36961–36969.

- 12 Yano T, Iijima N, Kakihara K, Hinuma S, Tanaka M, Iyata Y: Localization and neuronal response of RFamide related peptides in the rat central nervous system. *Brain Res* 2003; 982:156–167.
- 13 Bellinger LL, Bernardis LL: The dorsomedial hypothalamic nucleus and its role in ingestive behavior and body weight regulation: lessons learned from lesioning studies. *Physiol Behav* 2002;76:431–442.
- 14 Mieda M, Williams SC, Richardson JA, Tanaka K, Yanagisawa M: The dorsomedial hypothalamic nucleus as a putative food-entrainable circadian pacemaker. *Proc Natl Acad Sci U S A* 2006;103:12150–12155.
- 15 Bi S: Role of dorsomedial hypothalamic neuropeptide Y in energy homeostasis. *Peptides* 2007;28:352–356.
- 16 Ingalls AM, Dickie MM, Snell GD: Obese, a new mutation in the house mouse. *J Hered* 1950;41:317–318.
- 17 Zhang Y, Proenca R, Maffei M, Barone M, Leopold L, Friedman JM: Positional cloning of the mouse obese gene and its human homologue. *Nature* 1994;372:425–432.
- 18 Quenell JH, Mulligan AC, Tups A, Liu X, Phipps SJ, Kemp CJ, Herbison AE, Grattan DR, Anderson GM: Leptin indirectly regulates gonadotropin-releasing hormone neuronal function. *Endocrinology* 2009;150:2805–2812.
- 19 Louis GW, Greenwald-Yarnell M, Phillips R, Coolen LM, Lehman MN, Myers MG Jr: Molecular mapping of the neural pathways linking leptin to the neuroendocrine reproductive axis. *Endocrinology* 2011;152:2302–2310.
- 20 Elmquist JK, Ahima RS, Maratos-Flier E, Flier JS, Saper CB: Leptin activates neurons in ventrobasal hypothalamus and brainstem. *Endocrinology* 1997;138:839–842.
- 21 Oliveira VX Jr, Fazio MA, Miranda MT, da Silva JM, Bittencourt JC, Elias CF, Miranda A: Leptin fragments induce Fos immunoreactivity in rat hypothalamus. *Regul Pept* 2005;127:123–132.
- 22 Myers MG Jr, Münzberg H, Leininger GM, Leshan RL: The geometry of leptin action in the brain: more complicated than a simple ARC. *Cell Metab* 2009;9:117–123.
- 23 Johnson MA, Fraley GS: Rat RFRP-3 alters hypothalamic GHRH expression and growth hormone secretion but does not affect Kiss-1 gene expression or the onset of puberty in male rats. *Neuroendocrinology* 2008;88:305–315.
- 24 Clarke IJ, Smith JT, Henry BA, Oldfield BJ, Stefanidis A, Millar RP, Sari IP, Chng K, Fabre-Nys C, Caraty A, Ang BT, Chan L, Fraley GS: Gonadotropin-inhibitory hormone is a hypothalamic peptide that provides a molecular switch between reproduction and feeding. *Neuroendocrinology* 2012;95:305–316.
- 25 Klingerman CM, Williams WP 3rd, Simberlund J, Brahme N, Prasad A, Schneider JE, Kriegsfeld LJ: Food restriction-induced changes in gonadotropin-inhibiting hormone cells are associated with changes in sexual motivation and food hoarding, but not sexual performance and food intake. *Front Endocrinol (Lausanne)* 2012;2:101.
- 26 León S, García-Galiano D, Ruiz-Pino F, Barroso A, Manfredi-Lozano M, Romero-Ruiz A, Roa J, Vazquez MJ, Gaytan F, Blumenrohr M, van Duin M, Pinilla L, Tena-Sempere M: Physiological roles of gonadotropin-inhibitory hormone signaling in the control of mammalian reproductive axis: studies in the NPFF1 receptor null mouse. *Endocrinology* 2014;155:2953–2965.
- 27 Swerdloff RS, Batt RA, Bray GA: Reproductive hormonal function in the genetically obese (ob/ob) mouse. *Endocrinology* 1976; 98:1359–1364.
- 28 Ahima RS, Prabakaran D, Flier JS: Postnatal leptin surge and regulation of circadian rhythm of leptin by feeding. Implications for energy homeostasis and neuroendocrine function. *J Clin Invest* 1998;101:1020–1027.
- 29 Bouret SG, Draper SJ, Simerly RB: Trophic action of leptin on hypothalamic neurons that regulate feeding. *Science* 2004;304:108–110.
- 30 Bouret SG, Draper SJ, Simerly RB: Formation of projection pathways from the arcuate nucleus of the hypothalamus to hypothalamic regions implicated in the neural control of feeding behavior in mice. *J Neurosci* 2004;24: 2797–2805.
- 31 Molnar CS, Kallo I, Liposits Z, Hrabovszky E: Estradiol down-regulates RF-amide-related peptide (RFRP) expression in the mouse hypothalamus. *Endocrinology* 2011;152:1684–1690.
- 32 Mercer JG, Hoggard N, Williams LM, Lawrence CB, Hannah LT, Trayhurn P: Localization of leptin receptor mRNA and the long form splice variant (Ob-Rb) in mouse hypothalamus and adjacent brain regions by in situ hybridization. *FEBS Lett* 1996;387:113–116.
- 33 Finn PD, Steiner RA, Clifton DK: Temporal patterns of gonadotropin-releasing hormone (GnRH), c-fos, and galanin gene expression in GnRH neurons relative to the luteinizing hormone surge in the rat. *J Neurosci* 1998;18: 713–719.
- 34 Sugrue ML, Vella KR, Morales C, Lopez ME, Hollenberg AN: The thyrotropin-releasing hormone gene is regulated by thyroid hormone at the level of transcription in vivo. *Endocrinology* 2010;151:793–801.
- 35 Dror T, Franks J, Kauffman AS: Analysis of multiple positive feedback paradigms demonstrates a complete absence of LH surges and GnRH activation in mice lacking kisspeptin signaling. *Biol Reprod* 2013;88:146.
- 36 Poling MC, Quenell JH, Anderson GM, Kauffman AS: Kisspeptin neurones do not directly signal to RFRP-3 neurones but RFRP-3 may directly modulate a subset of hypothalamic kisspeptin cells in mice. *J Neuroendocrinol* 2013;25:876–886.
- 37 Chowen JA, Clifton DK: Semiquantitative analysis of cellular somatostatin mRNA levels by in situ hybridization histochemistry. *Method Neurosci* 1991;5:137–158.
- 38 Kauffman AS, Navarro VM, Kim J, Clifton D, Steiner RA: Sex differences in the regulation of Kiss1/NKB neurons in juvenile mice: implications for the timing of puberty. *Am J Physiol Endocrinol Metab* 2009;297:E1212–E1221.
- 39 Semaan SJ, Murray EK, Poling MC, Dhamija S, Forger NG, Kauffman AS: BAX-dependent and BAX-independent regulation of Kiss1 neuron development in mice. *Endocrinology* 2010;151:5807–5817.
- 40 Kim J, Semaan SJ, Clifton DK, Steiner RA, Dhamija S, Kauffman AS: Regulation of Kiss1 expression by sex steroids in the amygdala of the rat and mouse. *Endocrinology* 2011;152: 2020–2030.
- 41 Kauffman A, Sun Y, Kim J, Khan AR, Shu J, Neal-Perry G: Vasoactive intestinal peptide modulation of the steroid-induced LH surge involves kisspeptin signaling in young but not in middle-aged female rats. *Endocrinology* 2014;155:2222–2232.
- 42 Di Giorgio NP, Semaan SJ, Kim J, Lopez PV, Bettler B, Libertun C, Lux-Lantos VA, Kauffman AS: Impaired GABAB receptor signaling dramatically up-regulates Kiss1 expression selectively in nonhypothalamic brain regions of adult but not prepubertal mice. *Endocrinology* 2014;155:1033–1044.
- 43 Sahu A, Kalra PS, Kalra SP: Food deprivation and ingestion induce reciprocal changes in neuropeptide Y concentrations in the paraventricular nucleus. *Peptides* 1988;9: 83–86.
- 44 Chan YY, Steiner RA, Clifton DK: Regulation of hypothalamic neuropeptide-Y neurons by growth hormone in the rat. *Endocrinology* 1996;137:1319–1325.
- 45 Li C, Chen P, Smith MS: The acute suckling stimulus induces expression of neuropeptide Y (NPY) in cells in the dorsomedial hypothalamus and increases NPY expression in the arcuate nucleus. *Endocrinology* 1998; 139:1645–1652.
- 46 Lechan RM, Jackson IM: Immunohistochemical localization of thyrotropin-releasing hormone in the rat hypothalamus and pituitary. *Endocrinology* 1982;111:55–65.
- 47 Elmquist JK, Bjorbaek C, Ahima RS, Flier JS, Saper CB: Distributions of leptin receptor mRNA isoforms in the rat brain. *J Comp Neurol* 1998;395:535–547.
- 48 Kishi T, Aschkenasi CJ, Lee CE, Mountjoy KG, Saper CB, Elmquist JK: Expression of melanocortin 4 receptor mRNA in the central nervous system of the rat. *J Comp Neurol* 2003;457:213–235.
- 49 Murakami M, Matsuzaki T, Iwasa T, Yasui T, Irahara M, Osugi T, Tsutsui K: Hypophysiotropic role of RFamide-related peptide-3 in the inhibition of LH secretion in female rats. *J Endocrinol* 2008;199:105–112.
- 50 Soga T, Kitahashi T, Clarke IJ, Parhar IS: Gonadotropin-inhibitory hormone promoter-driven enhanced green fluorescent protein expression decreases during aging in female rats. *Endocrinology* 2014;155:1944–1955.

- 51 Clarke IJ, Sari IP, Qi Y, Smith JT, Parkington HC, Ubuka T, Iqbal J, Li Q, Tilbrook A, Morgan K, Pawson AJ, Tsutsui K, Millar RP, Bentley GE: Potent action of RFamide-related peptide-3 on pituitary gonadotropes indicative of a hypophysiotropic role in the negative regulation of gonadotropin secretion. *Endocrinology* 2008;149:5811–5821.
- 52 Rizwan MZ, Porteous R, Herbison AE, Anderson GM: Cells expressing RFamide-related peptide-1/3, the mammalian gonadotropin-inhibitory hormone orthologs, are not hypophysiotropic neuroendocrine neurons in the rat. *Endocrinology* 2009;150:1413–1420.
- 53 Pineda R, Garcia-Galiano D, Sanchez-Garrido MA, Romero M, Ruiz-Pino F, Aguilar E, Dijcks FA, Blomenrohr M, Pinilla L, van Noort PI, Tena-Sempere M: Characterization of the inhibitory roles of RFRP3, the mammalian ortholog of GnIH, in the control of gonadotropin secretion in the rat: in vivo and in vitro studies. *Am J Physiol Endocrinol Metab* 2010;299:E39–E46.
- 54 Ancel C, Bentsen AH, Sebert ME, Tena-Sempere M, Mikkelsen JD, Simonneaux V: Stimulatory effect of RFRP-3 on the gonadotrophic axis in the male Syrian hamster: the exception proves the rule. *Endocrinology* 2012;153:1352–1363.
- 55 Son YL, Ubuka T, Millar RP, Kanasaki H, Tsutsui K: Gonadotropin-inhibitory hormone inhibits GnRH-induced gonadotropin subunit gene transcriptions by inhibiting AC/cAMP/PKA-dependent ERK pathway in L β T2 cells. *Endocrinology* 2012;153:2332–2343.
- 56 Rizwan MZ, Harbid AA, Inglis MA, Quennell JH, Anderson GM: Evidence that hypothalamic RFamide related peptide-3 neurones are not leptin-responsive in mice and rats. *J Neuroendocrinol* 2014;26:247–257.
- 57 Bouret SG: Organizational actions of metabolic hormones. *Front Neuroendocrinol* 2013;34:18–26.
- 58 Hayashida T, Nakahara K, Mondal MS, Date Y, Nakazato M, Kojima M, Kangawa K, Murakami N: Ghrelin in neonatal rats: distribution in stomach and its possible role. *J Endocrinol* 2002;173:239–245.
- 59 Steculorum SM, Bouret SG: Maternal diabetes compromises the organization of hypothalamic feeding circuits and impairs leptin sensitivity in offspring. *Endocrinology* 2011;152:4171–4179.
- 60 Mannan MA, O'Shaughnessy PJ: Ovarian steroid metabolism during post-natal development in the normal mouse and in the adult hypogonadal (hpg) mouse. *J Reprod Fertil* 1988;82:727–734.
- 61 Simerly RB: Wired for reproduction: organization and development of sexually dimorphic circuits in the mammalian forebrain. *Annu Rev Neurosci* 2002;25:507–536.
- 62 Scott HM, Mason JI, Sharpe RM: Steroidogenesis in the fetal testis and its susceptibility to disruption by exogenous compounds. *Endocr Rev* 2009;30:883–925.
- 63 Poling MC, Kauffman AS: Organizational and activational effects of sex steroids on kisspeptin neuron development. *Front Neuroendocrinol* 2013;34:3–17.