A Fresh Look at the Relationship between TSH and Free Thyroxine in Cross-Sectional Data

Linde A.C. De Grande Katleen Van Uytfanghe Linda M. Thienpont

Laboratory for Analytical Chemistry, Faculty of Pharmaceutical Sciences, Ghent University, Ghent, Belgium

Dear Editor,

The classical log-linear relationship between thyroid-stimulating hormone (TSH) and free thyroxine (FT₄) generally reflects on the reasonable correlation between both hormones, while particularly emphasizing that small changes in FT₄ are accompanied by larger changes in TSH [1]. Recently, refining this relationship in cross-sectional data has received new interest [2–4]. Different from the classical monotonous relationship over the whole thyroid function range, the referred authors propose three different relationships in the hypo-, eu-, and hyperthyroid range. While they use different mathematical models to describe the relationships, they have in common that they work with functions that interconnect at the transition from the hypo-/euthyroid and eu-/hyperthyroid states. Although the real clinical relevance of knowing the exact mathematical relationship may be debated, it cannot be denied that the relation between TSH and FT₄ is discussed in view of more precisely defining the subclinical state of thyroid dysfunction, and/or even brought up by some scientists as evaluation criterion for the validity of a FT₄ assay [2, 5].

Here we take a fresh look at the TSH/FT₄ relationship, based on previously described data [1]. They were from 8,152 unselected patients (median age: 61 years, range: 18–100) from the Department of Endocrinology or Nuclear Medicine of the Klinikum Lüdenschied in Germany, who presented with various thyroid disorders. Data from pregnant women and patients

with pituitary or hypothalamic disorders were excluded, as well as from patients with conditions that potentially interfere with thyroid testing. Note that corresponding to the limits of the Abbott enzyme immunoassay, which was used for measuring both

TSH and FT₄, the TSH data are truncated at 0.001 and 100 mIU/l, respectively. Regression and correlation results were calculated by Microsoft Excel 2010.

Different from current practice, we plotted log TSH on the x-axis and FT₄ on

Fig. 1. log TSH is plotted on the x-axis and FT₄ on the y-axis. This reveals three clusters depending on the TSH-concentration: □: 'low', –: 'mid', and Δ: 'high'. The blue line is the linear regression line for the complete concentration range; the red lines are the linear regression lines for the different clusters. A clear break with little correlation in the TSH/FT₄ relationship for the 'high'-trophic hormone data group can be seen.

the y-axis (fig. 1). We did not define upfront the different concentration categories because we preferred to inspect the correlation of the complete dataset without any prejudice. Visual inspection of the plot reveals three clusters of TSH values from 0.001 to <0.045 mIU/l ('low'), 0.045 to <23 mIU/l ('mid'), and 23 to 100 mIU/l ('high'). Instead of a continuum, a clear break in the TSH/FT₄ relationship for the high-TSH data group can be seen. This is also reflected in the partial regression equations ($y = -2.50x + 14.9$, 'low'; $y = -2.60x + 14.1$, 'mid'; $y = -1.11x + 6.5$, 'high'). The regression equation over the whole range ($y = -3.04x + 13.9$, 'all') is similar to the low- and mid-range data. The correlation data, in our opinion, are misleading and are mainly influenced by the TSH range (note: correlation over the whole range is $r = -0.577$, while it is $r = -0.044$ in the high range). Although the precision and accu-

racy of the FT₄ measurement results in the low concentration range might be jeopardized by the limit of quantitation of the used assay, we consider that the effect of increased random error would result in a higher scatter around the line representing the TSH/FT₄ relationship, though without affecting the regression coefficients. Regarding the inaccuracy of measurement, we know from a previous study that most FT₄ assays tend to have a positive calibration bias in the low concentration range versus a significantly negative one in the mid- to high-concentration range [6]. This allows us to infer that after correction of this bias, the break in the TSH/FT₄ relationship would become even more obvious.

Our 'fresh' presentation of TSH/FT₄ data suggests that the log-linear relationship between both hormones holds also in cross-sectional data up to TSH concentra-

tions of ~23 mIU/l (FT₄ of ~10 pmol/l; note: the actual values will depend on the assays used). However, from this concentration on, there is a clear 'break' in the relationship with little correlation between TSH and FT₄. Consequently, curve fittings for the TSH/FT₄ relationship in both ranges should not be interconnected.

Acknowledgments

We thank Professors Rudolf Hoermann and Rolf Larisch (Klinikum Lüdenschied) for the provision of the raw data. We thank Dr. Dietmar Stöckl (STT Consulting) for helpful discussions and advice in statistics.

Disclosure Statement

The authors have nothing to disclose.

References

- 1 Demers LM, Spencer CA: Laboratory Medicine Practice Guidelines: Laboratory Support for the Diagnosis and Monitoring of Thyroid Disease. Washington, National Academy of Clinical Biochemistry (NACB), 2002.
- 2 Hoermann R, Eckl W, Hoermann C, Larisch R: Complex relationship between free thyroxine and TSH in the regulation of thyroid function. *Eur J Endocrinol* 2010;162:1123–1129.
- 3 Clark PM, Holder RL, Haque SM, Hobbs FD, Roberts LM, Franklyn JA: The relationship between serum TSH and free T₄ in older people. *J Clin Pathol* 2012;65:463–465.
- 4 Hadlow NC, Rothacker KM, Wardrop R, Brown SJ, Lim EM, Walsh JP: The relationship between TSH and free T₄ in a large population is complex and nonlinear and differs by age and sex. *J Clin Endocrinol Metab* 2013;98:2936–2943.
- 5 Jonklaas J, Kahric-Janovic N, Soldin OP, Soldin SJ: Correlations of free thyroid hormones measured by tandem mass spectrometry and immunoassay with thyroid-stimulating hormone across 4 patient populations. *Clin Chem* 2009;55:1380–1388.
- 6 Thienpont LM, Van Uytendanghe K, Van Houcke S, Das B, Faix JD, MacKenzie F, Quinn FA, Rottmann M, Van den Bruel A; IFCC Committee for Standardization of Thyroid Function Tests (C-STFT): A progress report of the IFCC Committee for standardization of thyroid function tests. *Eur Thyroid J* 2014;3:109–116.