PPAR γ Agonist-Induced Fluid Retention Depends on α ENaC Expression in Connecting Tubules

Yiling Fu^{a, c} Maria Gerasimova^c Falk Batz^{a, c} Alexander Kuczkowski^{a, c}
Yasaman Alam^{a, c} Paul W. Sanders^{d, e} Caroline Ronzaud^f Edith Hummler^f
Volker Vallon^{a–c}

Departments of ^aMedicine and ^bPharmacology, University of California, and ^cDepartment of Veterans Affairs, San Diego Healthcare System, San Diego, Calif., ^dDepartment of Medicine/Nephrology, University of Alabama at Birmingham, and ^eDepartment of Veterans Affairs Medical Center, Birmingham, Ala., USA; ^fDepartment of Pharmacology, University of Lausanne, Lausanne, Switzerland

Key Words

Peroxisome proliferator-activated receptor γ · Edema · Kidney · Body weight · Epithelial sodium channel

Abstract

Background/Aims: Thiazolidinediones (TZDs, like rosiglitazone (RGZ)) are peroxisome proliferator-activated receptor γ (PPAR γ) agonists used to treat type 2 diabetes. Clinical limitations include TZD-induced fluid retention and body weight (BW) increase, which are inhibited by amiloride, an epithelial-sodium channel (ENaC) blocker. RGZ-induced fluid retention is maintained in mice with α ENaC knockdown in the collecting duct (CD). Since ENaC in the connecting tubule (CNT) rather than in CD appears to be critical for normal NaCl retention, we aimed to further explore the role of ENaC in CNT in RGZ-induced fluid retention. **Methods:** Mice with conditional inactivation of α ENaC in both CNT and CD were used (α ENaC lox/lox AQP2-Cre; ' α ENaC-CNT/CD-KO') and compared with littermate controls (α ENaC lox/lox mice; 'WT'). BW was monitored and total body water (TBW) and extracellular fluid volume (ECF) were determined by bioelectri-

cal impedance spectroscopy (BIS) before and after RGZ (320 mg/kg diet for 10 days). **Results:** On regular NaCl diet, α ENaC-CNT/CD-KO had normal BW, TBW, ECF, hematocrit, and plasma Na⁺, K⁺, and creatinine, associated with an increase in plasma aldosterone compared with WT. Challenging α ENaC-CNT/CD-KO with a low NaCl diet unmasked impaired NaCl and K homeostasis, consistent with effective knockdown of α ENaC. In WT, RGZ increased BW (+6.1%), TBW (+8.4%) and ECF (+10%), consistent with fluid retention. These changes were significantly attenuated in α ENaC-CNT/CD-KO (+3.4, 1.3, and 4.3%). **Conclusion:** Together with the previous studies, the current results are consistent with a role of α ENaC in CNT in RGZ-induced fluid retention, which dovetails with the physiological relevance of ENaC in this segment.

© 2014 S. Karger AG, Basel

Introduction

PPARs are ligand-activated transcription factors that regulate a large number of genes by transcriptional activation and repression [1]. Thiazolidinediones (TZDs),

such as pioglitazone or rosiglitazone (RGZ), are potent ligands and activators of PPAR γ and are insulin sensitizers used to treat type 2 diabetes [2]. Fluid retention is an important clinical limitation for TZDs, which occurs in up to 5% of treated diabetic patients, and, as a consequence, these agents are contraindicated in patients with New York Heart Association class III and IV congestive heart failure [3, 4]. TZD-induced fluid retention and edema formation has been proposed to involve increases in vascular permeability [4–7] and vasodilation [8, 9] as well as primary effects on the kidney. The latter may include an increase in sodium and water reabsorption due to the upregulation of various renal transport regulating systems including the Na⁺-H⁺ exchanger NHE3, the Na-K-2Cl cotransporter NKCC2, the serum/glucocorticoid-regulated kinase 1, the α subunit of Na-K-ATPase, a non-selective cation channel, and/or the water channels aquaporin-2 (AQP-2) and AQP-3 [10–13].

Conditional knock-down of PPAR γ in the connecting tubules (CNT) and collecting ducts (CD) of mice (using Cre expression driven by the AQP-2 promoter) abrogated the ability of the TZDs – pioglitazone and RGZ – to increase plasma volume and body weight [14, 15], indicating that the site of action may primarily reside in these segments. The epithelial sodium channel ENaC is expressed in the apical membrane of the CNT and CD and serves as a primary effector of regulated sodium reabsorption. Studies in mice [15] and in patients with type 2 diabetes [16] found that TZD-induced weight gain and fluid retention were attenuated by the ENaC inhibitor, amiloride, indicating a potential role of ENaC in this process.

ENaC is a heteromultimer and the α ENaC subunit appears critical for exporting the intact ENaC complex to the apical membrane [17]. Mice with conditional knockdown of α ENaC (*Scnn1a*) in both CNT and CD (using AQP2-promoter driven Cre) or in CD only (using Hox7-promoter driven Cre) have previously been characterized and used to determine the quantitative role of ENaC in these segments for renal Na and K homeostasis. The studies found that Na and K homeostasis were impaired only when α ENaC was deleted in both CNT and CD, whereas the sole knockdown in CD was ineffective [17, 18]. This is consistent with the concept that it is ENaC in the CNT rather than in the CD that appears critical for normal NaCl retention and K excretion [19, 20]. Potentially along these lines, RGZ-induced fluid retention still fully developed in mice in which α ENaC has been knocked down only in the CD (using Cre expression driven by the Hox7 promoter) [12].

Therefore, in the present study we used mice with AQP-2 promoter driven knockdown of α ENaC to test for a potential role of α ENaC in the CNT in TZD-induced fluid retention.

Methods

Animals

All animal experimentation was conducted in accordance with the Guide for Care and Use of Laboratory Animals (National Institutes of Health, Bethesda, Md., USA) and was approved by the Institutional Animal Care and Use Committee of the Veterans Affairs San Diego Healthcare System. Female *Scnn1a*^{lox/lox} mice were bred with male *Scnn1a*^{lox/lox}/APQ2:cre+/- mice to generate experimental *Scnn1a*^{lox/lox} mice (WT, used as control) and *Scnn1a*^{lox/lox}/APQ2:cre+/- mice (α ENaC CNT/CD KO) as previously described [18]. Age-matched adult male mice (22–30 weeks of age) were selected and housed in standard rodent cages on a 12:12 h light-dark cycle with free access to food (1% K⁺, 0.4% Na⁺, 4.4% fat; Harlan Teklad TD.7001) and water.

Body Weight and Fluid Retention in Response to Rosiglitazone (RGZ)

Basal body weight (BW), daily food intake and water consumption were determined while mice were kept in standard rodent cages. Mice were then anesthetized with ketamine (33.3 mg/ml, 2.5 ml/kg BW i.p.) and xylazine (3.33 mg/dl, 2.5 ml/kg BW i.p.) to determine total body water (TBW), extracellular fluid (ECF) and intracellular fluid (ICF) by bioimpedance spectroscopy (BIS) using the ImpediVet BIS1 system (ImpediMed, San Diego, Calif., USA) as previously described [21]. Using a set of carefully positioned subcutaneous needle electrodes, BIS determines body composition based on its electrical characteristics in response to the application of low amplitude alternating electrical currents [22]. The procedure and measurement takes about 5 min. After completion of BIS and while still under anesthesia, blood was collected by retro-orbital bleeding to determine hematocrit (Hct). Mice were subsequently allowed to recover for 5–7 days. Mice were then fed with repellet diet containing RGZ (320 mg/kg diet [12, 14]) for 10 days. BW, food and water intake were measured every other day. On the last day, the mice were anesthetized with ketamine and xylazine to determine body fluid volumes by BIS and subsequently hematocrit.

Functional Confirmation of ENaC Knockdown – Response to Low NaCl Diet

Mice were maintained first on control diet (0.275% NaCl, 1% K⁺; Harlan Teklad TD.140039) and were then switched to low-NaCl diet (0.01% NaCl, 1% K⁺; Harlan Teklad TD.08601) for another 5 days. BW was monitored daily. Blood was withdrawn under isoflurane anesthesia before switching diets and on the 5th day of low salt diet to determine plasma Na⁺ and K⁺ (by flame photometer; Cole-Parmer, Vernon Hills, Ill., USA), creatinine (by isotope dilution LC-MS/MS in core laboratory of UAB-UCSD O'Brien Center for Acute Kidney Injury Research), and plasma aldosterone (by radioimmuno assay; DSL-8600; Diagnostic Systems Laboratories, Webster, Tex., USA).

Table 1. Basal measurements on regular salt diet

Parameters	WT (n = 12)	α ENaC CNT/ CD KO (n = 20)
BW, g	33.4 $\pm$ 0.6	33.6 $\pm$ 0.4
Daily food intake, g	4.2 $\pm$ 0.2	4.4 $\pm$ 0.1
Daily water intake, ml	6.4 $\pm$ 0.6	7.5 $\pm$ 0.3
Hematocrit, %	42.8 $\pm$ 0.3	43.6 $\pm$ 0.3
Plasma creatinine, mg/dl	0.086 $\pm$ 0.002	0.089 $\pm$ 0.001
TBW, ml	18.8 $\pm$ 0.6	20.3 $\pm$ 0.6
ECF, ml	6.5 $\pm$ 0.2	6.9 $\pm$ 0.2
ICF, ml	12.3 $\pm$ 0.4	13.4 $\pm$ 0.4

BW = Body weight; TBW = total body water; ECF = extracellular fluid; ICF = intracellular fluid.

Statistical Analysis

Data are reported as means $\pm$ SEM. Data from WT and KO were compared by ANOVA followed by the two-tailed t test. When comparing parameters within genotype before and after treatment, the paired t test was used. $p < 0.05$ was considered statistically significant.

Results

Basal Phenotype of α ENaC CNT/CD KO Mice and Response to Low Salt Diet

When fed a normal salt diet, BW and daily food intake and water consumption as well as hematocrit and plasma creatinine were not significantly different between WT mice and α ENaC CNT/CD KO mice (table 1). BIS showed that basal TBW, ECF and ICF did not differ in WT and KO (table 1), as were plasma concentrations of Na^+ and K^+ (fig. 1b, c). This was associated with increased plasma aldosterone concentrations in α ENaC CNT/CD KO compared with WT (fig. 1d). These data indicate that α ENaC CNT/CD KO were able to maintain relatively normal kidney function and Na^+ and K^+ homeostasis and body fluid volumes under normal salt intake, at least in part by up-regulating the aldosterone levels. When mice were placed on a low salt diet, α ENaC CNT/CD KO lost significantly more BW (fig. 1a) and showed significantly higher plasma creatinine levels than WT (0.125 ± 0.014 vs. 0.063 ± 0.002 mg/dl; $p < 0.001$). Whereas WT maintained plasma Na^+ or K^+ levels in response to a low salt diet, plasma Na^+ significantly decreased and plasma K^+ increased in α ENaC CNT/CD KO (fig. 1b, c). Aldosterone was slightly increased by low salt diet in WT but showed a strong further increase in α ENaC CNT/CD KO. These data in-

dicate that low salt diet unmasked the impaired Na^+ and K^+ homeostasis in α ENaC CNT/CD KO consistent with previous studies performed in this mouse model [18] and support the concept that Na^+ retention and K^+ secretion depended on α ENaC expression in CNTs rather than CDs [17, 18].

α ENaC CNT/CD KO Attenuated RGZ-Induced BW Gain and Fluid Retention

In WT mice, RGZ increased BW by 6.1% at day 10 (fig. 2a). This was associated with a significant reduction in hematocrit (fig. 2b) and increases in TBW by 8.4% and in ECF by 10%, while the increase in ICF by 7.6% did not quite reach statistical significance ($p = 0.052$) (fig. 3). α ENaC CNT/CD KO did not affect the RGZ reduction in hematocrit (fig. 2b) but significantly attenuated the RGZ-induced increase in BW (3.4%) (fig. 2a), TBW (1.3%) and ECF (4.3%) (each $p < 0.05$ vs. WT), and tended to reduce the increase in ICF when compared with WT (-0.1% , $p = 0.07$ vs. WT) (fig. 3). Daily food and water intake were not significantly affected by RGZ in either group (data not shown).

Discussion

Previous studies showed that the RGZ-induced fluid retention and increase in body weight were unaffected in mice with inactivation of α ENaC activity only in the CD [12]. Here we report that the RGZ-induced increase in body weight, TBW and ECF observed in WT mice were attenuated in a mouse model with inactivation of α ENaC activity in CNT and CD. Together, these studies provided evidence that RGZ-induced fluid retention and increase in body weight depends in part on ENaC activity in the CNT of the kidney. The current studies employed α ENaC knockdown driven by the AQP-2 promoter, that is, the same promoter that has previously been used to show the critical role of PPAR γ in these segments for TZD-induced fluid retention [14, 15]. A role for ENaC in CNT in TZD-induced fluid retention dovetails with the physiological relevance of ENaC in this segment for Na^+ reabsorption [17–20]. The findings are also in agreement with studies that found that the ENaC blocker, amiloride, and the aldosterone-antagonist, spironolactone, attenuated TZD-induced fluid retention in diabetic patients [16, 23].

While α ENaC knockdown in CNT and CD attenuated the TZD-induced increase in BW, TBW and ECF, it did not affect the fall in hematocrit. Thus, the TZD-induced fluid retention was only partially prevented by the α ENaC

Fig. 1. When challenged with a low salt diet, αENaC CNT/CD KO mice lost more body weight than WT mice (**a**). Plasma Na⁺ and K⁺ levels were similar between genotypes under normal salt diet. A low salt diet reduced plasma Na⁺ and increased plasma K⁺ levels only in αENaC CNT/CD KO (**b**, **c**). Plasma aldosterone was mod-

erately elevated in αENaC CNT/CD KO versus WT under normal salt diet; this difference was strongly increased in response to a low salt diet (**d**). n = 6 in each group. * p < 0.05 versus WT; # p < 0.05 versus normal salt in same genotype.

knockdown in CNT and CD. This may be due to the incomplete knockdown of αENaC in the CNT in this model (αENaC positive principal cells in late CNT reduced by about 70% – for details see [18]), potential effects of TZDs on ENaC in the late distal convoluted tubule, which together with ENaC in CNT are the primary sites of ENaC-dependent NaCl retention [20], and/or ENaC-independent mechanisms of TZD-induced fluid retention (see above).

It remains to be determined how PPARγ activation enhances ENaC activity. Lower blood pressure could reduce

renal perfusion pressure, thereby inducing ENaC and increasing renal reabsorption of Na⁺ and fluid [13, 24, 25]. Moreover, TZDs increased amiloride-sensitive Na⁺ absorption as well as γENaC expression when applied to cultured WT mouse inner medullary CD (IMCD) cells in vitro [15], indicating a potential direct effect of TZDs on ENaC. Long-term TZD treatment in patients with type 2 diabetes resulted in a decrease in plasma aldosterone concentrations and an increase in plasma atrial natriuretic peptide levels; these changes are consistent with a primary increase in sodium reabsorption and central venous

Fig. 2. α ENaC CNT/CD KO attenuated the RGZ-induced increase in body weight versus WT (a), whereas the reduction in hematocrit was not affected (b). * $p < 0.05$ versus WT; # $p < 0.05$ versus basal within genotype. $n = 12$ in WT; $n = 18$ – 20 in KO.

Fig. 3. RGZ significantly increased TBW and ECF and tended to increase ICF in WT; this response was blunted in α ENaC CNT/CD KO. * $p < 0.05$ versus WT; # $p < 0.05$ versus basal within genotype. $n = 11$ in WT; $n = 20$ in KO.

volume expansion [23]. However, significant increases in the renal expression of the ENaC subunits α , β or γ could not be detected in response to TZDs [10, 12, 13] and one study even described a suppression of the γ -subunit of ENaC at the mRNA and protein level in the mouse kidney cortex and in a cortical collecting duct cell line [26]. Furthermore, in vitro electrophysiological studies in the A6, M-1, and mpkCCD(cl4) cell lines, which express viable and functional PPAR γ , found that both the basal and the insulin-stimulated Na⁺ flux via ENaC were insensitive to the PPAR γ agonists, pioglitazone and GW7845 [27]. Considering all these results together with the current data, one may speculate that the TZD-induced activation of ENaC occurs independently of a change in ENaC protein expression and may require the complexity of the in vivo setting that is not mimicked in cell lines. Further studies are needed to explore these hypotheses.

In summary and in the context of previous studies [12], the presented in vivo studies are consistent with the

concept that ENaC-mediated Na⁺ reabsorption in the CNTs contributes to TZD-induced fluid retention and weight gain.

Acknowledgements

This work was supported by the American Heart Association GRNT3440038, the National Institutes of Health R01DK56248, R01HL94728, the UAB/UCSD O'Brien Center of Acute Kidney Injury (NIH-P30DK079337), and the Department of Veterans Affairs (to VV), and the Swiss National Science Foundation (31003A_144198/1) and the National Center of Competence in Research (NCCR, Kidney.CH) (to EH).

Disclosure Statement

Dr. Vallon has received within the past 12 months research grant support for basic science studies from Boehringer Ingelheim Pharma GmbH & Co.KG and AstraZeneca.

References

- Isseman I, Green S: Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature* 1990; 347:645–650.
- Sharma AM, Staels B: Review: Peroxisome proliferator-activated receptor gamma and adipose tissue – understanding obesity-related changes in regulation of lipid and glucose metabolism. *J Clin Endocrinol Metab* 2007; 92:386–395.
- Nesto RW, Bell D, Bonow RO, Fonseca V, Grundy SM, Horton ES, Le Winter M, Porte D, Semenkovich CF, Smith S, Young LH, Kahn R: Thiazolidinedione use, fluid retention, and congestive heart failure: a consensus statement from the American Heart Association and American Diabetes Association. October 7, 2003. *Circulation* 2003;108:2941–2948.
- Vasudevan AR, Balasubramanyam A: Thiazolidinediones: a review of their mechanisms of insulin sensitization, therapeutic potential, clinical efficacy, and tolerability. *Diabetes Technol Ther* 2004;6:850–863.
- Niemeyer NV, Janney LM: Thiazolidinedione-induced edema. *Pharmacotherapy* 2002; 22:924–929.
- Sotiropoulos KB, Clermont A, Yasuda Y, Rask-Madsen C, Mastumoto M, Takahashi J, Della Vecchia K, Kondo T, Aiello LP, King GL: Adipose-specific effect of rosiglitazone on vascular permeability and protein kinase C activation: novel mechanism for PPARgamma agonist's effects on edema and weight gain. *FASEB J* 2006;20:1203–1205.
- Emoto M, Anno T, Sato Y, Tanabe K, Okuya S, Tanizawa Y, Matsutani A, Oka Y: Troglitazone treatment increases plasma vascular endothelial growth factor in diabetic patients and its mRNA in 3T3-L1 adipocytes. *Diabetes* 2001;50:1166–1170.
- Gardiner SM, Nunez DJ, Baer PG, Brown KK, Bennett T: Regional hemodynamic effects of the N-(2-benzoylphenyl)-L-tyrosine peroxisome proliferator-activated receptor-gamma ligand, GI 262570 [(S)-2-(2-benzoylphenyl amino)-3-[4-[2-(5-methyl-2-phenyl-2-oxazol-4-yl)ethoxy]phenyl]propionic acid], in conscious rats. *J Pharmacol Exp Ther* 2004; 310:1226–1233.
- Parulkar AA, Pendergrass ML, Granda-Ayala R, Lee TR, Fonseca VA: Nonhypoglycemic effects of thiazolidinediones. *Ann Intern Med* 2001;134:61–71.
- Chen L, Yang B, McNulty JA, Clifton LG, Binz JG, Grimes AM, Strum JC, Harrington WW, Chen Z, Balon TW, Stimpson SA, Brown KK: GI262570, a peroxisome proliferator-activated receptor {gamma} agonist, changes electrolytes and water reabsorption from the distal nephron in rats. *J Pharmacol Exp Ther* 2005;312:718–725.
- Artunc F, Sandulache D, Nasir O, Boini KM, Friedrich B, Beier N, Dicks E, Potzsch S, Klingel K, Amann K, Blazer-Yost BL, Scholz W, Risler T, Kuhl D, Lang F: Lack of the serum and glucocorticoid-inducible kinase SGK1 attenuates the volume retention after treatment with the PPARgamma agonist pioglitazone. *Pflugers Arch* 2008;456:425–436.
- Vallon V, Hummler E, Rieg T, Pochynyuk O, Bugaj V, Schroth J, Dechenes G, Rossier B, Cunard R, Stockand J: Thiazolidinedione-induced fluid retention is independent of collecting duct alphaENaC activity. *J Am Soc Nephrol* 2009;20:721–729.
- Song J, Knepper MA, Hu X, Verbalis JG, Ecelbarger CA: Rosiglitazone activates renal sodium- and water-reabsorptive pathways and lowers blood pressure in normal rats. *J Pharmacol Exp Ther* 2004;308:426–433.
- Zhang H, Zhang A, Kohan DE, Nelson RD, Gonzalez FJ, Yang T: Collecting duct-specific deletion of peroxisome proliferator-activated receptor gamma blocks thiazolidinedione-induced fluid retention. *Proc Natl Acad Sci U S A* 2005;102:9406–9411.
- Guan Y, Hao C, Cha DR, Rao R, Lu W, Kohan DE, Magnuson MA, Redha R, Zhang Y, Breyer MD: Thiazolidinediones expand body fluid volume through PPARgamma stimulation of ENaC-mediated renal salt absorption. *Nat Med* 2005;11:861–866.
- Viswanathan V, Mohan V, Subramani P, Parthasarathy N, Subramaniam G, Manoharan D, Sundaramoorthy C, Gnudi L, Karalliedde J, Viberti G: Effect of spironolactone and amiloride on thiazolidinedione-induced fluid retention in South Indian patients with type 2 diabetes. *Clin J Am Soc Nephrol* 2013;8:225–232.
- Rubera I, Loffing J, Palmer LG, Frindt G, Fowler-Jaeger N, Sauter D, Carroll T, McMahon A, Hummler E, Rossier BC: Collecting duct-specific gene inactivation of alphaENaC in the mouse kidney does not impair sodium and potassium balance. *J Clin Invest* 2003; 112:554–565.

- 18 Christensen BM, Perrier R, Wang Q, Zuber AM, Maillard M, Mordasini D, Malsure S, Ronzaud C, Stehle JC, Rossier BC, Hummler E: Sodium and potassium balance depends on alphaENaC expression in connecting tubule. *J Am Soc Nephrol* 2010;21:1942–1951.
- 19 Frindt G, Palmer LG: Na channels in the rat connecting tubule. *Am J Physiol Renal Physiol* 2004;286:F669–F674.
- 20 Meneton P, Loffing J, Warnock DG: Sodium and potassium handling by the aldosterone-sensitive distal nephron: the pivotal role of the distal and connecting tubule. *Am J Physiol Renal Physiol* 2004;287:F593–F601.
- 21 Masuda T, Fu Y, Eguchi A, Czogalla J, Rose MA, Kuczkowski A, Gerasimova M, Feldstein AE, Scadeng M, Vallon V: Dipeptidyl peptidase IV inhibitor lowers PPARgamma agonist-induced body weight gain by affecting food intake, fat mass, and beige/brown fat but not fluid retention. *Am J Physiol Endocrinol Metab* 2014;306:E388–E398.
- 22 Chapman ME, Hu L, Plato CF, Kohan DE: Bioimpedance spectroscopy for the estimation of body fluid volumes in mice. *Am J Physiol Renal Physiol* 2010;299:F280–F283.
- 23 Karalliedde J, Buckingham R, Starkie M, Lorand D, Stewart M, Viberti G: Effect of various diuretic treatments on rosiglitazone-induced fluid retention. *J Am Soc Nephrol* 2006;17:3482–3490.
- 24 Buchanan TA, Meehan WP, Jeng YY, Yang D, Chan TM, Nadler JL, Scott S, Rude RK, Hsueh WA: Blood pressure lowering by pioglitazone. Evidence for a direct vascular effect. *J Clin Invest* 1995;96:354–360.
- 25 Yang B, Clifton LG, McNulty JA, Chen L, Brown KK, Baer PG: Effects of a PPARgamma agonist, GI262570, on renal filtration fraction and nitric oxide level in conscious rats. *J Cardiovasc Pharmacol* 2003;42:436–441.
- 26 Borsting E, Cheng VP, Glass CK, Vallon V, Cunard R: Peroxisome proliferator-activated receptor-gamma agonists repress epithelial sodium channel expression in the kidney. *Am J Physiol Renal Physiol* 2012;302:F540–F551.
- 27 Nofziger C, Chen L, Shane MA, Smith CD, Brown KK, Blazer-Yost BL: PPARgamma agonists do not directly enhance basal or insulin-stimulated Na(+) transport via the epithelial Na(+) channel. *Pflugers Arch* 2005;451:445–453.