Renal C3 Complement Component: Feed Forward to Diabetic Kidney Disease

Katherine J. Kelly^a Yunlong Liu^b Jizhong Zhang^a Jesus H. Dominguez^{a,c}

Departments of ^aMedicine, ^bMedical and Molecular Genetics, Indiana University School of Medicine, and ^cRoudebush Veterans' Affairs Medical Center, Indianapolis, Ind., USA

Key Words

Complement · Diabetic nephropathies · Ischemia · Kidney failure, chronic

Abstract

Background: Diabetic nephropathy is the main cause of end-stage renal disease and has reached epidemic proportions. **Methods:** Comprehensive genomic profiling (RNAseq) was employed in the ZS (F₁ hybrids of Zucker and spontaneously hypertensive heart failure) model of diabetic nephropathy. Controls were lean littermates. **Results:** Diabetic nephropathy in obese, diabetic ZS was accelerated by a single episode of renal ischemia (DI). This rapid renal decline was accompanied by the activation of the renal complement system in DI, and to a lesser extent in sham-operated diabetic rats (DS). In DI there were significant increases in renal mRNA encoding C3, C4, C5, C6, C8, and C9 over sham-operated lean normal controls (LS). Moreover, mRNAs encoding the receptors for the anaphylatoxins C3a and C5a were also significantly increased in DI compared to LS. The classic complement pathway was activated in diabetic kidneys with significant increases of C1qa, C1qb, and C1qc mRNAs in DI over LS. In addition, critical regulators of complement activation were significantly attenuated in DI and DS. These included mRNAs encoding CD55, decay accelerating factor, and CD59, which inhibit the membrane attack complex. C3, C4, and C9 proteins were demonstrated in renal tubules and glomeruli. The complement RNAseq data were incorporated into a

gene network showing interactions among C3-generating renal tubular cells and other immune competent migratory cells. **Conclusions:** We conclude that local activation of the complement system mediates renal injury in diabetic nephropathy.

© 2015 S. Karger AG, Basel

Introduction

Diabetic nephropathy is the main reason for the vast number of patients enrolling in end-stage renal disease (ESRD) programs [1]. Unfortunately, current therapies of diabetic nephropathy fail to stop progression to ESRD [2]. To gain insight into its progressive pathophysiology, we have studied diabetic nephropathy in the ZS rat, a model of obesity and diabetes that progresses to ESRD [3], with features similar to human diabetic nephropathy [4, 5]. ZS rats suffer an accelerated decline following renal ischemia with associated inflammation. These renal responses include leukocyte infiltration and broad activation of pro-inflammatory genes [6]. We hypothesized that this complexity can be dissected by systems biology, and used deep sequencing with advanced bioinformatics tools to study disease mechanisms of diabetic nephropathy [7]. We found that obese, diabetic rats with renal ischemia exhibited general and prominent activation of the renal complement system along with interacting pro-inflammatory gene networks. The complement system is a

mainstay of systemic innate immunity comprising several interacting components [8]. The master element is complement component 3 (C3), which is in a perpetual state of contained activity [9], restrained by specific proteins [8, 9]. Failure of the regulatory proteins leads to uncontrolled activation and injury [8].

The presence of renal C3 in human diabetic nephropathy was reported from the beginning of immunofluorescence, but was thought secondary to nonspecific trapping of plasma C3 [10]. Experiments on diabetic animals also found renal C3; and again, it was thought to be blood-derived [11]. In our studies of renal transcriptomes in diabetes [7] we found major activation of the renal complement system in rats with diabetic nephropathy.

Subjects and Methods

Animals

The three groups of rats included here, and their core renal transcript networks, including inflammation, have been reported elsewhere [7]. Lean and obese, diabetic male ZS rats (Charles River, Wilmington, Mass., USA) were acquired at 8 weeks of age and fed Purina diet #5008. Their body weights were measured and sera plus urine samples analyzed at biweekly intervals. One group of obese/diabetic rats was subjected to bilateral renal ischemia at 10 weeks of age as described (DI, $n = 11$) [7]. The lean rats (LS, $n = 6$) and a second obese/diabetic group (DS, $n = 7$) were subjected to sham surgery. These rats were terminated at 28 weeks of age, their kidneys removed, immediately frozen in liquid nitrogen, and RNA extracted (below). In addition, renal tubular cells from 12 week-old normal Sprague Dawley male rats ($n = 4$) were isolated as previously described [12].

Histology and Immunohistochemistry

Kidney sections were stained for histology and changes quantified in blinded sections as described [7, 12]. Immunostaining was performed as described [12] using goat anti-complement component 3 (CAT #c312-A, Alpha Diagnostics, San Antonio, Tex., USA), rabbit anti-complement component 4 (CAT #hp8023, Hycult Biotech, Plymouth Meeting, Pa., USA) and mouse anti-complement component 9 (CAT #ab17931, Abcam, Cambridge, Mass., USA) and then Texas Red-conjugated donkey anti-rabbit (CAT# 111-075-045, Jackson ImmunoResearch, West Grove, Pa., USA) secondary antibody.

Hypoxia Chamber

Kidney tubular cells, derived from normal Sprague Dawley rats [12] were grown in 38% O₂/5% CO₂ until confluent and then subjected to anoxia (1% O₂/5% CO₂ in a hypoxia chamber (Hypoxia workstation, Sci-Tive-Dual, Ruskinn Technology, Bridgend, UK)) for periods of 8–48 h ('ischemia'). The cells were then switched back to 38% O₂/5% CO₂ for 16–24 h ('reperfusion'). Control cells were always maintained in 38% O₂/5% CO₂. Total RNA were isolated from these cells and mRNAs encoding C3, C4, C5, C8, and C9 were measured by RT²PCR.

RNAseq and RT²PCR

Total kidney RNA isolation was performed as described [7]. RNA, 3 µg, was fragmented with RNAase III, cDNA libraries constructed with SOLiD adaptors by reverse transcription (RT), and then sequenced by strand specific RNAseq of short 50 bp reads using the SOLiD 4 platform (Center for Medical Genomics at Indiana University School of Medicine) as described [7]. Sequence alignment to the UCSC rat genome database was performed using BFAST [13]. Gene expression was calculated in the form of Reads per Kilobase Exon Model per million mapped reads (RPKM) [14]. To identify differentially expressed genes, we conducted the Student t-test on the logarithmically transformed RPKM values, comparing DI versus LS and DS versus LS. Network analysis was performed using Metacore Software (GeneGo, Carlsbad, Calif., USA).

RT²PCR was performed [7] using primers from Qiagen, Valencia, Calif., USA. The samples were run in quadruplicate, normalized to β-actin mRNA and are reported as amount of mRNA compared to control samples [15].

Animal Use Statement

The experiments were conducted in conformity with the 'Guiding Principles for Research Involving Animals and Human Beings'. The investigations were approved by the Institutional Animal Care and Use Committee of Indiana University School of Medicine.

Statistics

Results are expressed as means ± 1 standard error. Differences in renal and metabolic parameters were determined by 1 way ANOVA with subsequent t-tests (when ANOVA indicated statistical significance, GraphPad Prism, LaJolla, Calif., USA). Bonferroni correction was used for multiple comparisons. The null hypothesis was rejected at $p < 0.05$.

Results and Discussion

The abnormal metabolic and renal phenotypes in the ZS rats included in this report were described in more detail in an earlier article [7]. There were 3 groups of ZS rats: lean sham-operated (LS), obese/diabetic sham-operated (DS), and obese/diabetic subjected to a short period of bilateral renal ischemia (DI). At termination, DS and DI rats were heavier, 539 ± 28 and 526 ± 26 than the LS controls, 466 ± 67 (weight in g $p < 0.05$ for both). The blood glucose levels were also higher in DS and DI, 29.4 ± 16 and 30.5 ± 1.7 , respectively, when compared to LS controls, 7.8 ± 0.6 (glucose in mM, significantly higher for both, $p < 0.05$). Creatinine clearance rates were also depressed in DS and DI rats when compared to LS controls (fig. 1). Proteinuria, normal in LS, was increased 27 and 30 fold in DS and DI, respectively [7]. Renal histology in ZS rats included the following stains: Leder's stain to identify leukocytes, PAS to outline cell structure and extracellular matrix, and Mason's trichrome to label fibro-

(For legend see next page.)

sis, figure 1. Leder's stain in DS and DI rats revealed clusters of densely packed cellular infiltrates that contained neutrophils. PAS demonstrated increases in tubular damage plus interstitial and glomerular extracellular matrix, particularly in the DI rats (vs. LS). Trichrome stain showed early stages of peritubular fibrosis in DS rats, and more advanced in DI rats (vs. LS), figure 1.

Renal histology revealed the stark complexity of diabetic nephropathy; migrating inflammatory cells could be found in the vicinity of stressed and damaged renal epithelial and endothelial cells. These intricate relations were approached by systems biology, and we analyzed the interactions among expressed renal transcriptomes in the three conditions: LS, DS, and DI. Among the many up-regulated renal transcripts in diabetic nephropathy [7], a subset of stimulated mRNAs stood out because it contained most elements of the complement system. The up-regulation of these transcripts is shown on the top in figure 2a. We verified these RNAseq results with RT²PCR as an independent measurement. The correlations of mRNA results obtained between the two methods, RNAseq and RTPCR, were excellent and are shown in figure 2b.

The renal protein expression of the complement components C3, C4, and C9 was then examined by immunofluorescence. There was very faint expression of these proteins in LS rats. However in DS, and particularly in DI, C3, C4, and C9 were broadly expressed in renal tubules and glomeruli. C3, nearly absent in lean group glomeruli (LS), was robustly expressed in diabetes/sham (DS) and more so in diabetes/ischemia (DI) kidneys, figure 3.

The kidney has been viewed as a passive target of circulating complement components of hepatic origin which result in immune complex-mediated glomerular injury. It is now recognized that vast local expansion of complement can occur in injured kidneys [16] and, in animal models, can affect progression of renal failure and survival [17]. Our data showing increased expression of complement components in the renal transcriptome are

consistent with local complement activation in diabetic nephropathy. To test the hypothesis that C3 is derived from the tubular epithelium and that ischemia is a critical mediator of the C3 production, primary cultures of normal Sprague Dawley rat renal tubular cells were exposed to acute anoxic conditions, followed by re-oxygenation, that is, in vitro 'ischemia/reperfusion'. Following anoxia, mRNA encoding C3 was markedly elevated, 2.34 ± 0.05 -fold over normoxic cells ($p < 0.05$; $n = 4$), while mRNAs encoding C1, C3, C5, C8, and C9 were slightly depressed (not shown). C3 protein was also increased by anoxia (fig. 3). In addition to ischemia, inflammatory cytokines and loss or decrease in regulatory proteins have been shown to be important in local complement expression [18, 19].

Transcripts of the complement system were assembled in a gene network (fig. 4). These transcripts were differentially expressed in both conditions, DS and DI, when compared to controls (LS). The center piece of the network is complement component C3 (C3); activated 5.2 fold in chronic diabetic nephropathy with ischemia over lean rat controls, and 2.3-fold after 24 h of acute anoxia/reoxygenation in isolated renal tubular cells. These results confirm intrinsic renal expression of C3, as previously suggested [20, 21]. C3b, derived from upregulated renal C3, generates the classical pathway C5 convertase (C4bC2aC3b) that cleaves C5. Renal C5 and C5aR1 mRNAs were not among the 13,453 detected and measured renal transcripts reported by RNAseq, and we resorted to RT²-PCR to measure these transcripts in the three groups of rats: C5 mRNA in DS and DI was 58 and 35% higher than in LS ($p < 0.03$). C5aR1 mRNA in DS and DI were 2.2-fold ($p < 0.07$) and 3.4 fold higher than LS ($p < 0.04$). The C1 complex was well represented in the RNAseq analysis, with upregulation of C1q, C1s, and C1r, and C2. In the network, C1q is acting on C2 and C4, forming the C3 convertase (C4b C2a), leading to the assembly of the classical pathway C5 convertase, C4bC2aC3b [9]. C1q also stimulates the re-

Fig. 1. ZS rat diabetic nephropathy. Representative renal sections (from top to bottom: Leder's stain, Periodic Acid Schiff (PAS) stain, and Masson's trichrome stain) in the three groups of rats (from left to right, lean controls (lean), diabetic sham operated (DM/Sham), and diabetic postischemic rats (DM/Isch)) are presented. Leder's stain (**a**) labeled clusters of leukocytes (arrows, pink in online version), nearly absent in the Lean (LS) group, clearly present in Diabetes/Sham group (DS) and very prominent in Diabetes/Ischemia rats group (DI). The inset shows a neutrophil with typical nuclear morphology. PAS stain (**b**) demonstrated normal glomerular and tubular structures in LS. The Diabetes/

Sham group exhibited mild increases of interstitial and glomerular extracellular matrix, which were far more pronounced in the Diabetes/Ischemia group. Asterisk indicates examples of atrophic tubules. Masson's trichrome (**c**) stained connective tissue (arrowheads, blue in online version) was normal in LS. Increases in connective tissue were evident in the glomeruli and interstitium of the Diabetes/Sham group, and more severe in the Diabetes/Ischemia group. Mean creatinine clearance and quantification of histological parameters are shown in the graphs (**d-g**). * $p < 0.05$ vs. LS; § $p < 0.05$ vs. DS. Scale bar = 50 μ m (color figure in online version).

Fig. 2. Complement system renal transcripts detected by RNAseq and RT²PCR. Transcript levels by RNAseq (**a**) are expressed as Reads per Kilobase Exon Model per Million mapped reads (RPKM, see text) and represent mRNAs encoding complement components (C1qa, C1s, C2–4, C6–9), as well as complement factor D (cfd), complement receptor type 2 (CR2) and c4 binding protein (C4b). These renal transcripts are significantly elevated in Diabetes/Sham and Diabetes/Ischemia groups. CD55, decay accelerating factor for complement, and CD59, encode complement regulatory proteins that limit the complement cascade. The two regulatory renal genes are suppressed in both Diabetes/Sham and Diabetes/Ischemia groups. Significant correlations between transcript levels measured by RNAseq (horizontal axis) and quantitative PCR (QPCR) vertical axis are shown in (**b**). These comparisons were made to verify the accuracy of the RNAseq determinations. To facilitate visualization on one axis, RNAseq data are presented as follows: C1s/2, C2*2, C4/3, C6*10, C8*50, C9*10, cfd*100, CD55*3, CD59/10, CR2*50, C4b*10.

lease of interleukin-12 (IL-12) [22], also activated by other components (below), and the pro-inflammatory homing receptor CCR7, which enhances the effects of its up-regulated ligand, CCL19 via PI3K [23].

CD55, or decay accelerating factor (DAF), is restricted to cell membranes [24] where it antagonizes both C3 and C5 convertases. This transcript encodes a critical controller of activation, and it is suppressed in kidneys from DI rats 35% ($p < 0.004$). The attenuation of renal DAF may cede the modulatory role to upregulated complement factor I (CFI) [25] (increased 10.5 fold, $p = 1.25 \times 10^{-5}$) and complement factor H (CFH) [26] (increased 1.8 fold, $p < 0.002$). However, the upregulation of CFI and CFH failed to prevent complement activation. It is likely that complement was activated by up-regulation of complement

factor D (CFD) and unchanged factor B (CFB) that cause the formation of the C3 amplification convertase C3bBb [27]. These events lead to the formation of the membrane attack complex (MAC). The cytotoxic activity of MAC in diabetic nephropathy was further enhanced by the attenuation of CD59, which promotes the clearance of MAC [28]. Renal C3 activation generates the anaphylatoxins C3a and C5a, and the portal for C3a, the receptor C3ar1, is also upregulated in diabetic nephropathy. Activated CD80 (or B7-1), the receptor for cytotoxic T lymphocyte antigen 4 (CTLA4), is a downstream target for C3a [29]. C3a and C5a both act on unchanged renal extracellular signal-regulated kinases 1/2 (ERK), in accord with a role for ERK signaling in complement activation [30–32]. IL-12 is another downstream target for activated C3a [33, 34]

Fig. 3. Renal complement components in ZS rat diabetic nephropathy. Complement components 3, 4, and 9 are shown (**a**) in (left to right) Lean control rats (control), Diabetic/Sham group (DM Sham) and Diabetes/Ischemia group (DM Isch). Complement (3, 4, 9) protein expression was barely detectable in the Lean control group, but, in contrast, was robustly expressed in the two diabetic groups. Similarly, glomerular C3 levels (**b**) were nearly undetectable in the Lean group (left), and rose prominently in the two diabetic groups (right). To examine complement generation in tubule cells, primary renal cells (**c**) were cultured in 38% O₂ and 5% CO₂ and transferred to 1% O₂ and 5% CO₂ for the designated times. The cells were then returned to 38% O₂ and 5% CO₂ for 24 h. C3 mRNA was increased at 24 h, whereas levels of immunoreactive C3 increased at 48 h of anoxia. Scale bar = 50 μ m; * $p < 0.05$ vs. time 0 h (color figure in online version.)

Fig. 4. Complement system gene pathways in diabetic nephropathy. The numerical table on the upper left corner shows the fold changes in the different transcripts of the complement system. These values were all statistically significant ($p < 0.05$). The network on the right upper corner contains upregulated (normal font, red in online version), inhibited (italics, blue in online version) or unchanged transcripts (serif font, black). Solid arrows (green in online version) show positive interactions. The main elements of this network were organized to show activation of the local classic and alternative pathways. The end result was the production of the anaphylatoxins C3a and C5a and the formation of the membrane attack complex (MAC). The bottom network delineates critical interactions among the anaphylatoxins C3a and C5a with pro-inflammatory and pro-fibrotic transcripts. These morbid interactions might be responsible, at least in part, for the development and eventual progression of diabetic nephropathy (color figure in online version).

and C5a [35]. The network also shows C3a acting on ICAM-1, a relationship previously reported [36]. C3a also modulates T cell maturation [37]. The importance of the renal IL-12 gene in diabetic nephropathy is further emphasized by additional IL-12 activation from upregulated C1q [38] and C5a [39]. C3a also acts on TGF β [40] a fibrogenic interaction exemplified by the upregulation of Collagen 4.

C5a works through its receptor C5a1, found in most renal cells [41]. Renal C5a acts on upregulated p-selectin (Selp) [42], which also amplifies complement activation [43] and may account for grievous clinical states [44].

Selp is also activated by C5b when integrated as the MAC [45]. Renal ICAM-1 was up-regulated in DI/DS rats, and it is shown as a downstream target for activation not only by C3a1 [36] but also by C5a1 [46]. In addition, C3a and C5a activate PI3K via independent pathways [47]. The gene encoding the co-stimulatory receptor CD40 is also upregulated in diabetic nephropathy, and it is shown interacting with both upregulated C3a and C5a, suggesting that adaptive immunity, or a form of autoimmunity, might occur in diabetic nephropathy [31]. Some of these pro-inflammatory actions are mediated by upregulated NF κ B, which is a downstream target of both C3a and C5a

[31], and may include upregulated CD40 [48]. NF κ B activates upregulated CXCR4, suggesting a role for the CXCR4/CXCL12 signaling axis in diabetic nephropathy [49]. The network also shows NF- κ B acting on upregulated CCR2, a receptor involved in monocyte activation [50], on upregulated CXCL1, the prototypical neutrophil chemokine [51], on upregulated CXCL10, a leukocyte chemoattractant [52], and on upregulated CCL19, a chemokine for dendritic cells and T lymphocytes [53]. NF- κ B, activated by IL1R, the receptor for IL-1 [54], acts on upregulated collagen I gene, a redundant effect mediated by C3a, C5a, and IL-1 [55]. C5a also acts on the fibronectin gene [56], and presumably its cognate protein, which can then activate upregulated BMP1 and promote the formation of collagens I and III [57]. C5a also acts on upregulated TLR4 gene [58], potentially acting CASP1, which is required to cleave Pro-IL-1b and generate renal IL-1b, the master cytokine [59].

In conclusion, our data point to renal C3 activation in diabetic nephropathy of rats. C3 reactivity in tubules leads to broader renal activation of the complement system, which is sustained by suppression of complement regulators, and contributes to renal inflammation, impaired function, and fibrosis.

Acknowledgment

This work was supported by an NIH grant R01 DK 082739 and a research grant from DCI Paul Teschan Research and Development Fund to KJK and VA Merit Review funds and DoD support to JHD.

Disclosure Statement

None of the authors has a conflict of interest to declare.

References

- 1 US Renal Data System: USRDS 2013 Annual Data Report: Atlas of Chronic Kidney Disease and End-Stage Renal Disease in the United States. Bethesda, MD, National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, 2013.
- 2 Kelly KJ, Dominguez JH: Rapid progression of diabetic nephropathy is linked to inflammation and episodes of acute renal failure. *Am J Nephrol* 2010;32:469–475.
- 3 Dominguez J, Wu P, Packer CS, Temm C, Kelly KJ: Lipotoxic and inflammatory phenotypes in rats with uncontrolled metabolic syndrome and nephropathy. *Am J Physiol Renal Physiol* 2007;293:F670–F679.
- 4 Temm C, Dominguez JH: Microcirculation: nexus of comorbidities in diabetes. *Am J Physiol Renal Physiol* 2007;293:F486–F493.
- 5 Breyer MD, Bottinger E, Brosius FC, Coffman TM, Fogo A, Harris RC, Heilig CW, Sharma K: Diabetic nephropathy: of mice and men. *Adv Chronic Kidney Dis* 2005;12:128–145.
- 6 Kelly KJ, Burford JL, Dominguez JH: Post-ischemic inflammatory syndrome: a critical mechanism of progression in diabetic nephropathy. *Am J Physiol Renal Physiol* 2009;297:F923–F931.
- 7 Kelly KJ, Liu Y, Zhang J, Goswami C, Lin H, Dominguez JH: Comprehensive genomic profiling in diabetic nephropathy reveals the predominance of proinflammatory pathways. *Physiol Genomics* 2013;45:710–719.
- 8 Ricklin D, Hajishengallis G, Yang K, Lambris JD: Complement: a key system for immune surveillance and homeostasis. *Nat Immunol* 2010;11:785–797.
- 9 Muller-Eberhard HJ: Molecular organization and function of the complement system. *Annu Rev Biochem* 1988;57:321–347.
- 10 Ainsworth SK, Hirsch HZ, Brackett NC Jr, Brissie RM, Williams AV Jr, Hennigar GR: Diabetic glomerulonephropathy: histopathologic, immunofluorescent, and ultrastructural studies of 16 cases. *Hum Pathol* 1982;13:470–478.
- 11 Fujita T, Ohi H, Komatsu K, Endo M, Ohsawa I, Kanmatsuse K: Complement activation accelerates glomerular injury in diabetic rats. *Nephron* 1999;81:208–214.
- 12 Kelly KJ, Zhang J, Han L, Wang M, Zhang S, Dominguez JH: Intravenous renal cell transplantation with SAA1-positive cells prevents progression of chronic renal failure in rats with ischemic-diabetic nephropathy. *Am J Physiol Renal Physiol* 2013;305:F1804–F1812.
- 13 Homer N, Merriman B, Nelson SF: BFAST: an alignment tool for large scale genome sequencing. *PLoS One* 2009;4:e7767.
- 14 Mortazavi A, Williams BA, McCue K, Schaefer L, Wold B: Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nat Methods* 2008;5:621–628.
- 15 Cohen CD, Lindenmeyer MT, Eichinger F, Hahn A, Seifert M, Moll AG, Schmid H, Kiss E, Grone E, Grone HJ, Kretzler M, Werner T, Nelson PJ: Improved elucidation of biological processes linked to diabetic nephropathy by single probe-based microarray data analysis. *PLoS One* 2008;3:e2937.
- 16 Tang S, Zhou W, Sheerin NS, Vaughan RW, Sacks SH: Contribution of renal secreted complement C3 to the circulating pool in humans. *J Immunol* 1999;162:4336–4341.
- 17 Sheerin NS, Risley P, Abe K, Tang Z, Wong W, Lin T, Sacks SH: Synthesis of complement protein C3 in the kidney is an important mediator of local tissue injury. *FASEB J* 2008;22:1065–1072.
- 18 Sacks S, Zhou W: New boundaries for complement in renal disease. *J Am Soc Nephrol* 2008;19:1865–1869.
- 19 Yamada K, Miwa T, Liu J, Nangaku M, Song WC: Critical protection from renal ischemia reperfusion injury by CD55 and CD59. *J Immunol* 2004;172:3869–3875.
- 20 Sacks SH, Zhou W, Pani A, Campbell RD, Martin J: Complement C3 gene expression and regulation in human glomerular epithelial cells. *Immunology* 1993;79:348–354.
- 21 Tang S, Sheerin NS, Zhou W, Brown W, Sacks SH: Apical proteins stimulate complement synthesis by cultured human proximal tubular epithelial cells. *J Am Soc Nephrol* 1999;10:69–76.
- 22 Baruah P, Dumitriu IE, Peri G, Russo V, Mantovani A, Manfredi AA, Rovere-Querini P: The tissue pentraxin PTX3 limits C1q-mediated complement activation and phagocytosis of apoptotic cells by dendritic cells. *J Leukoc Biol* 2006;80:87–95.
- 23 Liu S, Wu J, Zhang T, Qian B, Wu P, Li L, Yu Y, Cao X: Complement C1q chemoattracts human dendritic cells and enhances migration of mature dendritic cells to CCL19 via activation of AKT and MAPK pathways. *Mol Immunol* 2008;46:242–249.
- 24 Kirkitadze MD, Barlow PN: Structure and flexibility of the multiple domain proteins that regulate complement activation. *Immunol Rev* 2001;180:146–161.

- 25 Nilsson SC, Sim RB, Lea SM, Fremeaux-Bacchi V, Blom AM: Complement factor I in health and disease. *Mol Immunol* 2011;48:1611–1620.
- 26 Makou E, Herbert AP, Barlow PN: Functional anatomy of complement factor H. *Biochemistry* 2013;52:3949–3962.
- 27 Esterbauer H, Krempler F, Oberkofler H, Patsch W: The complement system: a pathway linking host defence and adipocyte biology. *Eur J Clin Invest* 1999;29:653–656.
- 28 Tegli CA, Cudrici C, Patel S, Trippe R 3rd, Rus V, Niculescu F, Rus H: Membrane attack by complement: the assembly and biology of terminal complement complexes. *Immunol Res* 2011;51:45–60.
- 29 Chen L, Flies DB: Molecular mechanisms of T cell co-stimulation and co-inhibition. *Nat Rev Immunol* 2013;13:227–242.
- 30 Cao Q, McIsaac SM, Stadnyk AW: Human colonic epithelial cells detect and respond to C5a via apically expressed C5aR through the ERK pathway. *Am J Physiol Cell Physiol* 2012;302:C1731–C1740.
- 31 Li K, Fazekasova H, Wang N, Peng Q, Sacks SH, Lombardi G, Zhou W: Functional modulation of human monocytes derived DCs by anaphylatoxins C3a and C5a. *Immunobiology* 2012;217:65–73.
- 32 Vibhuti A, Gupta K, Subramanian H, Guo Q, Ali H: Distinct and shared roles of beta-arrestin-1 and beta-arrestin-2 on the regulation of C3a receptor signaling in human mast cells. *PLoS One* 2011;6:e19585.
- 33 Kawamoto S, Yalcindag A, Laouini D, Brodeur S, Bryce P, Lu B, Humbles AA, Oettgen H, Gerard C, Geha RS: The anaphylatoxin C3a downregulates the Th2 response to epicutaneously introduced antigen. *J Clin Invest* 2004;114:399–407.
- 34 Reis ES, Barbutto JA, Kohl J, Isaac L: Impaired dendritic cell differentiation and maturation in the absence of C3. *Mol Immunol* 2008;45:1952–1962.
- 35 Peng Q, Li K, Wang N, Li Q, Asgari E, Lu B, Woodruff TM, Sacks SH, Zhou W: Dendritic cell function in allostimulation is modulated by C5aR signaling. *J Immunol* 2009;183:6058–6068.
- 36 Ducruet AF, Hassid BG, Mack WJ, Sosunov SA, Otten ML, Fusco DJ, Hickman ZL, Kim GH, Komotar RJ, Mocco J, Connolly ES: C3a receptor modulation of granulocyte infiltration after murine focal cerebral ischemia is reperfusion dependent. *J Cereb Blood Flow Metab* 2008;28:1048–1058.
- 37 Peng Q, Li K, Patel H, Sacks SH, Zhou W: Dendritic cell synthesis of C3 is required for full T cell activation and development of a Th1 phenotype. *J Immunol* 2006;176:3330–3341.
- 38 Baruah P, Dumitriu IE, Malik TH, Cook HT, Dyson J, Scott D, Simpson E, Botto M: C1q enhances IFN-gamma production by antigen-specific T cells via the CD40 costimulatory pathway on dendritic cells. *Blood* 2009;113:3485–3493.
- 39 Lalli PN, Strainic MG, Lin F, Medof ME, Heeger PS: Decay accelerating factor can control T cell differentiation into IFN-gamma-producing effector cells via regulating local C5a-induced IL-12 production. *J Immunol* 2007;179:5793–5802.
- 40 Bora PS, Sohn JH, Cruz JM, Jha P, Nishihori H, Wang Y, Kaliappan S, Kaplan HJ, Bora NS: Role of complement and complement membrane attack complex in laser-induced choroidal neovascularization. *J Immunol* 2005;174:491–497.
- 41 Klos A, Tenner AJ, Johswich KO, Ager RR, Reis ES, Kohl J: The role of the anaphylatoxins in health and disease. *Mol Immunol* 2009;46:2753–2766.
- 42 Foreman KE, Vaporciyan AA, Bonish BK, Jones ML, Johnson KJ, Glovsky MM, Eddy SM, Ward PA: C5a-induced expression of P-selectin in endothelial cells. *J Clin Invest* 1994;94:1147–1155.
- 43 Del Conde I, Cruz MA, Zhang H, Lopez JA, Afshar-Kharghan V: Platelet activation leads to activation and propagation of the complement system. *J Exp Med* 2005;201:871–879.
- 44 Noris M, Mescia F, Remuzzi G: STEC-HUS, atypical HUS and TTP are all diseases of complement activation. *Nat Rev Nephrol* 2012;8:622–633.
- 45 Kilgore KS, Ward PA, Warren JS: Neutrophil adhesion to human endothelial cells is induced by the membrane attack complex: the roles of P-selectin and platelet activating factor. *Inflammation* 1998;22:583–598.
- 46 Floreani AA, Wyatt TA, Stoner J, Sanderson SD, Thompson EG, Allen-Gipson D, Heires AJ: Smoke and C5a induce airway epithelial intercellular adhesion molecule-1 and cell adhesion. *Am J Respir Cell Mol Biol* 2003;29:472–482.
- 47 Venkatesha RT, Berla Thangam E, Zaidi AK, Ali H: Distinct regulation of C3a-induced MCP-1/CCL2 and RANTES/CCL5 production in human mast cells by extracellular signal regulated kinase and PI3 kinase. *Mol Immunol* 2005;42:581–587.
- 48 Brasier AR: The nuclear factor-kappaB-interleukin-6 signalling pathway mediating vascular inflammation. *Cardiovasc Res* 2010;86:211–218.
- 49 Arora S, Bhardwaj A, Singh S, Srivastava SK, McClellan S, Nirodi CS, Piazza GA, Grizzle WE, Owen LB, Singh AP: An undesired effect of chemotherapy: gemcitabine promotes pancreatic cancer cell invasiveness through reactive oxygen species-dependent, nuclear factor kappaB- and hypoxia-inducible factor 1alpha-mediated up-regulation of CXCR4. *J Biol Chem* 2013;288:21197–21207.
- 50 Lisi S, Sisto M, Lofrumento DD, D'Amore M: Sjogren's syndrome autoantibodies provoke changes in gene expression profiles of inflammatory cytokines triggering a pathway involving TACE/NF-kappaB. *Lab Invest* 2012;92:615–624.
- 51 Cowley MJ, Weinberg A, Zammit NW, Walters SN, Hawthorne WJ, Loudovaris T, Thomas H, Kay T, Gunton JE, Alexander SI, Kaplan W, Chapman J, O'Connell PJ, Grey ST: Human islets express a marked proinflammatory molecular signature prior to transplantation. *Cell Transplant* 2012;21:2063–2078.
- 52 Burke SJ, Goff MR, Lu D, Proud D, Karlstad MD, Collier JJ: Synergistic expression of the CXCL10 gene in response to IL-1beta and IFN-gamma involves NF-kappaB, phosphorylation of STAT1 at Tyr701, and acetylation of histones H3 and H4. *J Immunol* 2013;191:323–336.
- 53 Pietila TE, Veckman V, Lehtonen A, Lin R, Hiscott J, Julkunen I: Multiple NF-kappaB and IFN regulatory factor family transcription factors regulate CCL19 gene expression in human monocyte-derived dendritic cells. *J Immunol* 2007;178:253–261.
- 54 Burke SJ, Lu D, Sparer TE, Masi T, Goff MR, Karlstad MD, Collier JJ: NF-kappaB and STAT1 control CXCL1 and CXCL2 gene transcription. *Am J Physiol Endocrinol Metab* 2014;306:E131–E149.
- 55 Parthasarathy A, Gopi V, Umadevi S, Simna A, Sheik MJ, Divya H, Vellaichamy E: Suppression of atrial natriuretic peptide/natriuretic peptide receptor-A-mediated signaling upregulates angiotensin-II-induced collagen synthesis in adult cardiac fibroblasts. *Mol Cell Biochem* 2013;378:217–228.
- 56 Boor P, Konieczny A, Villa L, Schult AL, Bucher E, Rong S, Kunter U, van Roeyen CR, Polakowski T, Hawlisch H, Hillebrandt S, Lammert F, Eitner F, Floege J, Ostendorf T: Complement C5 mediates experimental tubulointerstitial fibrosis. *J Am Soc Nephrol* 2007;18:1508–1515.
- 57 Huang G, Zhang Y, Kim B, Ge G, Annis DS, Mosher DF, Greenspan DS: Fibronectin binds and enhances the activity of bone morphogenetic protein 1. *J Biol Chem* 2009;284:25879–25888.
- 58 Stevens MG, Van Poucke M, Peelman LJ, Rainard P, De Spiegeleer B, Rogiers C, Van de Walle GR, Duchateau L, Burvenich C: Anaphylatoxin C5a-induced toll-like receptor 4 signaling in bovine neutrophils. *J Dairy Sci* 2011;94:152–164.
- 59 Brown GT, Narayanan P, Li W, Silverstein RL, McIntyre TM: Lipopolysaccharide stimulates platelets through an IL-1beta autocrine loop. *J Immunol* 2013;191:5196–5203.