Original Article

Aspects of Exercise before or after Bariatric Surgery: A Systematic Review

Sjaak Pouwels^{a, b} Marjon Wit^c Joep A.W. Tejjink^{a, b} Simon W. Nienhuijs^a

^aDepartment of Surgery, Catharina Hospital, Eindhoven, the Netherlands; ^bDepartment of Epidemiology, CAPHRI Research School, Maastricht University, Maastricht, the Netherlands;

^cDepartment of Physiotherapy, Catharina Hospital, Eindhoven, the Netherlands

Key Words

Bariatric surgery · Physical activity · Exercise · Weight loss

Abstract

Background: Bariatric surgery has a considerable effect on weight loss. A positive relation of exercise and weight loss has been described before. However, the mode of exercise and its timing pre- or postoperatively or a combination remains unclear. **Methods:** A multi-database search was conducted. Identified articles were reviewed on description of exercise, timing around a bariatric intervention, and outcome. Methodological quality of the included studies was rated using the Physiotherapy Evidence Database scale. A Cohen's kappa score assessed the level of agreement. Outcome measurements were improvement of anthropometric and physical fitness variables, operation related complications, weight regain, and quality of life. **Results:** A total of 8 prospective studies were included. Four focused on training before and 4 on training after a bariatric procedure. Details of exercises varied from 45 min treadmill up to full descriptive programs. Supervision was frequently included. Significant improvement was encountered for biometric results physical fitness variables. **Conclusion:** In the majority of reports on exercising in a (future) bariatric population, positive effects on anthropometrics, cardiovascular risk factors and physical fitness were described. However, the results were not unanimous, with a wide range of exercise programs and perioperative timing, therefore hampering adequate practical guidance.

© 2015 S. Karger GmbH, Freiburg

Introduction

Obesity is a major contributor to a wide variety of diseases, including diabetes, hypertension, dyslipidemia, and obstructive sleep apnea syndrome [1, 2]. Over the last decades, dietary and exercise interventions had increasing public health and governmental attention.

Simon W. Nienhuijs, MD, PhD
Department of Surgery
Catharina Hospital
Michelangelolaan 2, P.O. Box 1350, 5602 ZA Eindhoven, the Netherlands
simon.nienhuijs@catharinaziekenhuis.nl

Nevertheless, the proportion of the population with morbid obesity gained every year [3]. The only treatment with a longstanding effect is bariatric surgery [4, 5]. However, these interventions are no universal remedy and a considerable proportion of patients has insufficient weight loss or regain in weight over time after the operation [6].

The key elements for durable weight loss are preoperative screening, an appropriate algorithm of bariatric techniques, and, most important, a strict regime of follow-up. Follow-up should include medical and psychological guidance, including continuing dietary advice and maintenance of a physical exercise regimen.

There is extensive knowledge that exercise helps to gain a better physical fitness and quality of life (QoL). A Cochrane review showed a reduction of 1.5 kg that was contributed to exercise [7]. Another systematic review focusing on physical training and obesity found a 4% excess weight loss [8]. In their review, Egberts et al. [9] found a reduction of 3.6 kg related to physical exercise. In each of these three reviews, only prospective cohort studies were included as no randomized controlled trials could be identified. Weight measurement was merely self-reported, and in the majority of studies the mode of physical activity was not defined. However, the positive relationship between exercise and weight loss seems evident. The problem is; what kind/type/mode of exercise should an obese patient be advised and what about its timing – pre- or postoperatively or a combination of both? These questions were the objective of the present systematic review.

Material and Methods

A systematic search on the qualitative aspects and timing of physical exercise in bariatric surgery patients was conducted. The population of interest consisted of obese patients with a BMI ≥ 30 kg/m². The intervention of interest was exercise training compared to regular care (no specific physical exercise training program). Also included were the different modes of exercise training; those were compared to each other. Outcome measurements were: improvement of anthropometric and physical fitness variables, effect on weight, and quality of life.

Pubmed, Embase, and CINAHL were searched from the earliest date of each database up to July 2014. The search for publications was performed using the following search string: (('Bariatrics'[Mesh] OR bariatrics OR bariatric surgery OR 'Obesity, Morbid'[Mesh] OR morbid obesity)) AND ('Physical Therapy Modalities'[Mesh] OR physical therapy OR physiotherapy OR 'Exercise'[Mesh] OR exercise).

The authors MW and SP, blinded for authors and journals, separately screened and selected the studies on the basis of title and abstract. After primary selection, both authors reviewed the full text of the selected studies to determine suitability for inclusion, based on the established selection criteria.

- Randomized controlled trial or prospective trial.
- Patients awaiting bariatric surgical intervention or with bariatric surgery in the past.
- A description of an exercise program, defined as a regimen of physical activities, included.
- Modes of exercise training were defined as follows: strength training and/or endurance training or a combination of both. Also multimodal programs with exercise components were included.
- Registration of anthropometric and physical fitness variables, complications, effects on weight, and QoL.
 - Physical fitness variables are defined as a percentage of VO₂ max and/or heart rate reserve / heart rate kinetics.

In addition cross-references were screened for further eligible studies. Disagreements between the two authors were resolved by discussion and by consultation of the senior authors (JT, SN), until consensus was reached. It was presupposed to be confronted with a significant clinical heterogeneity for which a systematic review could be performed, but not a meta-analysis

The methodological quality of the included studies was rated using the Physiotherapy Evidence Database (PEDro) scale [10], which has acceptable reliability [11]. The PEDro scale consists of 11 criteria, with a maximum score of 10 (range 0–10) since the first item (the specification of the eligibility criteria) is not included in the total score. Two authors (MW and SP) separately assigned PEDro scores to each included study. For rating the methodological quality, the following classification was used: a PEDro score of 4 indicated a poor methodological quality, a score of 4–5 fair quality, a score of 6–8 good quality, and a score of

Fig. 1. PRISMA flowchart.

9–10 excellent quality [12]. A Cohen's kappa score assessed the level of agreement between the authors. The score was determined as follows: <0.20 poor agreement, 0.21–0.40 fair agreement, 0.41–0.60 moderate agreement, 0.61–0.80 good agreement, and 0.81–1.00 very good agreement [13].

Results

The primary search produced 1,023 results, including 81 duplicates. 16 studies were identified as possibly relevant and underwent a full text critical appraisal, resulting in 8 exclusions. One study investigated exercise programs for Prader-Willi patients [14], 3 studies investigated pulmonary physiotherapy in the bariatric population [15–17], 1 publication appeared to be a case report [18], 1 was a research protocol [19], 1 article was only available in Swedish [20], and 1 article was only available in Spanish [21].

Table 1. Methodological quality of included studies using the PEDro score [10]

	Criteria ^a											total
	1	2	3	4	5	6	7	8	9	10	11	
Castello-Simoes et al., 2013 [27]	x	x	x	x	–	–	–	x	x	x	x	7
Funderburk and Callis, 2010 [23]	x	x	x	–	–	–	–	x	x	x	x	6
Hickey et al., 1999 [24]	x	–	–	x	–	–	–	x	x	x	x	5
Stegen et al., 2011 [28]	x	–	–	x	–	–	–	x	x	x	x	5
Baillot et al., 2013 [22]	x	–	–	x	–	–	–	x	x	x	x	5
Shah et al., 2011 [26]	x	x	x	x	–	–	–	x	x	x	x	7
Marcon et al., 2011 [25]	x	–	–	x	–	–	–	–	x	x	x	4
Berggren et al., 2008 [29]	x	–	–	x	–	–	–	–	x	x	x	4

^aPEDro criteria: (1) Specification of eligibility criteria (this criterion was not counted for the final score), (2) random allocation, (3) concealed allocation, (4) prognostic similarity at baseline, (5) participant blinding, (6) therapist blinding, (7) outcome assessor blinding, (8) more than 85% follow-up of at least one key outcome, (9) intention-to-treat analysis, (10) between or within-group statistical comparison, (11) point estimates of variability provided (10).

Figure 1 summarizes the search results. The methodological quality of the included studies ranged from moderate to good quality, as indicated by the PEDro scale (table 1). The level of agreement between reviewers (MW and SP) was reflected by a Cohen's kappa 0.78, which represents a good agreement. The key findings of the included studies are shown in table 2 for those with exercise before and in table 3 for those with exercise after bariatric surgery.

Study Designs and Patient Population

In total 4 studies with exercise before bariatric [22–25] and 4 studies with exercise after bariatric surgery [26–29] were included. There were 3 randomized controlled trials [23, 26, 27] and 5 prospective trials [22, 24, 25, 28, 29], including a patient population with a mean age ranging from 28 to 54 years and a mean BMI ranging from 40.4 to 48.7 kg/m². In the 4 studies with exercise before surgery, follow-up was done preoperatively and 7 days after surgery [24], preoperatively and 12 weeks after surgery [22, 23], or preoperatively and 24 weeks after surgery [25]. Of the 4 studies with exercise after surgery, 1 had postoperative follow-up only at 12 weeks [26] and 3 had preoperative and postoperative follow-up (1 week preoperatively and 4 weeks postoperatively [27], preoperatively (unknown how long) and 4 months postoperatively [28], or 1 week preoperatively and 10 days postoperatively [29]).

Exercise before Bariatric Surgery

Anthropometric Variables

Four studies reported on the effects of exercise training on the anthropometric variables. Baillot et al. [22] reported a significant reduction in weight (144.3 vs. 140.2 kg; $p = 0.07$), BMI (51.4 vs. 47.2 kg/m²; $p = 0.004$), neck circumference (42.2 vs. 41.0 cm; $p = 0.016$), and fat mass (72.1 vs. 69.1 kg; $p = 0.026$) after supervised exercise training in 12 patients awaiting bariatric surgery.

Table 2. Pre-bariatric surgery exercise regimes

Study	Exercise regime	Outcome
Baillet et al., 2013 [22]	<p>The physical exercise sessions consisted of: 10 min of warm up, 30 min of endurance activity (treadmill, walking circuit), 20–30 min of strength exercises, 10 min of cooling-down. Supervision by a physical activity specialist and/or completed independently at home. The intensity of the endurance training was determined using the HRR method and using data from the previously performed symptom limited cardiac treadmill test. The progression of endurance intensity and duration according to physical fitness and physical activity. Electronic pulse monitors (Polar F4™) were used to control endurance exercise intensity during supervised sessions and at home.</p> <p><i>Strength exercises:</i> Performed with small equipment easily available at home: dumbbells, elastic bands, and sticks. Three mini-circuits (upper body, lower body, and trunk) repeated by subjects according to their prescription.</p>	<p><i>6MWT distance, m:</i> before: 464 (412–501) after: 492 (450–531) (p = 0.05) <i>Arm Curl Test (n):</i> before: 22 (16.7–26) after: 26 (21–31) (p = 0.01)</p>
Funderburk and Callis, 2010 [23]	The intervention group did two sessions of 60 min of aquatic exercises including endurance and strength exercises for 12 weeks	<p><i>Weight loss:</i> I: +5.0 kg C: –2.3kg <i>SBP and DBP:</i> I: +34 mm Hg and –4 mm Hg C: –57 mm Hg and 0 mm Hg <i>6MWT:</i> I: +10.4 m C: +40.2 m <i>QoL and depression score:</i> No significant post intervention difference between groups I: decrease in bodily pain and depression score (p < 0.05) C: Increase in mental health (p < 0.05)</p>
Hickey et al., 1999 [24]	7 days of supervised endurance training at 60% of VO ₂ peak. Each session was 60 min	<p>Decrease of fasting plasma insulin (–41.7 pmol/l; p < 0.05) No significant change in body weight, % fat mass, glucose and blood lipid concentration after intervention No significant change in VO₂ peak after exercise intervention</p>

Table 2 continued on next page

Funderburk and Callis [23] showed a reduction in body weight of 5.0 kg after 12 weeks of supervised aquatic exercises in the intervention group and of 2.3 kg in the control group. It was not reported whether this is a significant decrease. There was no significant change in body weight and free fat mass after a 7-day exercise regimen in a study by Hickey et al. [24]. In contrast, Marcon et al. [25] found a significant decrease in body weight (–5.3 kg; p < 0.001) and BMI (–1.9 kg/m²; p < 0.001) after a 24-week supervised low-intensity endurance training.

Table 2. Continued

Study	Exercise regime	Outcome
Marcon et al., 2011 [25]	24 weeks of supervised low intensity endurance training. One session per week, consisting of 209 min of exercise and 10 min stretching	<p><i>Body weight and BMI</i> decreased body weight (–5.3 kg; $p < 0.001$) decreased BMI (–1.9 kg/m²; $p < 0.001$)</p> <p><i>SBP and DBP:</i> decrease in SBP (–23.8 mm Hg; $p = 0.007$) and DBP (–14.4 mm Hg; $p = 0.001$)</p> <p><i>Lipids and glucose:</i> improvement of TC (–23.0 mg/dl), HDL-C (+1.7 mg/dl), LDL-C (–2.3 mg/dl), TG (–26.7 mg/dl) and glucose (–17.2 mg/dl): for all $p < 0.007$</p> <p><i>6MWT:</i> improved walking distance(+69.8 m; $p < 0.0001$)</p>

HRR = Heart rate reserve; I = intervention group; C = control group; 6MWT = 6-min walking test; n = number, SBP = systolic blood pressure; DBP = diastolic blood pressure; VO₂ peak = maximum oxygen consumption; TC = total cholesterol, TG = triglycerides.

Table 3. Post-bariatric surgery exercise regimes

Study	Exercise regime	Outcome
Shah et al., 2011 [26]	<p>Energy expenditure $\geq 2,000$ kcal/week in moderate-intensity aerobic exercise at 60–70% of maximal oxygen consumption (VO₂ max). The subjects were instructed to achieve these goals gradually and were asked to expend 500 kcal during the first week and increase by 500 kcal every week until they achieved their goal of $\geq 2,000$ kcal/week. Each subject was asked to exercise on the treadmill at a certain speed and grade and on the cycle ergometer or rowing machine at a particular wattage that would correspond to 60–70% of her/his measured VO₂ max. Subjects who preferred to walk outside or on the walking track were asked to measure the distance that they walked and asked to complete this distance in a time period to achieve a pace associated with an intensity of 60–70% of the measured VO₂ max. The subjects were asked to exercise at least 5 days a week and to increase the intensity of exercise to remain the same level of exertion after each training. The exercise was partially supervised and the subjects were asked to come to the fitness center at least once or twice a week. Energy expenditure per week was calculated approximately from the work data shown on the exercise equipment and from the duration and distance of walking relative to body weight. Exercise away from the fitness center was monitored by asking the subjects to keep an exercise diary and/or using heart rate monitors.</p>	<p><i>Energy expenditure (moderate physical activity) (kcal/kg/day):</i> IG: baseline 1.1 (0–5.8), 6 weeks 3.5 (0.4–8.2); $p < 0.02$, 12 weeks 4.1 (0–15.0); $p < 0.0001$</p> <p><i>Time spent (moderate physical activity) (h/day):</i> IG: baseline 0.3 (0–1.5), 6 weeks 0.9 (0.1–2.1); $p < 0.02$ 12 weeks 1.0 (0–3.8), $p < 0.0001$</p>

Table 3 continued on next page

Table 3. Continued

Study	Exercise regime	Outcome
Castello-Simoes et al., 2013 [27]	<p>Start of aerobic training 48 h after maximal exercise test.</p> <p>Session duration 1 h on alternate days, 3 times a week, for 12 weeks, totaling 36 sessions.</p> <p>A session consisted of the following:</p> <p>initial 5-min stretching of upper and lower limbs and diaphragmatic breathing.</p> <p>5-min warm up on a treadmill at 3 km/h.</p> <p>40 min of exercise on a treadmill with speed and inclination varying according to the behavior of HR, separated in 4 steps;</p> <p>Step 1: intensity of exercise in which the HR remained at 50% of HR peak, reached in maximal exercise testing.</p> <p>Step 2: 60% of HR peak.</p> <p>Step 3: 70% of HR peak.</p> <p>Step 4: Maintaining 70% of the HR peak.</p> <p>1 min recovery at 3 km/h.</p> <p>10 min of the same initial stretching and diaphragmatic breathing.</p> <p>HR and blood pressure were obtained at the beginning of the session, at the end of each step, recovery and at the end of the session.</p>	<p><i>Lung function (FVC)</i></p> <p>TG:</p> <p>BGBS 94.0 ± 3.1</p> <p>4GBS 101.0 ± 2.5</p> <p>CG:</p> <p>BGBS 99.0 ± 2.4</p> <p>4GBS 100.0 ± 3.7</p> <p>EG: 100 ± 4.0</p> <p><i>Mean HR (beats/min)</i></p> <p>TG:</p> <p>BGBS 132.3 ± 6.0</p> <p>4GBS 124 ± 5.9</p> <p>CG:</p> <p>BGBS 138.2 ± 3.7</p> <p>4GBS 125 ± 3.9</p> <p>EG: 135.5 ± 3.5</p> <p><i>6MWT walking distance, m</i></p> <p>TG:</p> <p>BGBS 470.0 ± 23.9</p> <p>4GBS 515.0 ± 14.0</p> <p>CG:</p> <p>BGBS 453.0 ± 29.0</p> <p>4GBS 505.0 ± 10.9</p> <p>EG: 641.0 ± 10.4</p> <p><i>Dyspnea (0–10)</i></p> <p>TG:</p> <p>BGBS 5.8 ± 0.6</p> <p>4GBS 2.7 ± 0.8</p> <p>CG:</p> <p>BGBS 5.9 ± 0.5</p> <p>4GBS 4.5 ± 0.9</p> <p>EG: 2.8 ± 0.9</p>

Table 3 continued on next page

Compliance

Baillot et al. [22] had a high compliance to the preoperative exercise training program. They proposed 35 supervised exercise sessions in 12 weeks. Seven patients completed the full program, and 5 had bariatric surgery before the end of the program. With median 57.3% (32.5–77.6%) and 64.5% (47.5–79.9%), the attendance to the supervised exercise sessions and the total exercise sessions, respectively, was satisfactory.

Blood Pressure

After 12 weeks of aquatic exercises, Funderburk and Callis [23] found an increase in systolic blood pressure and a decrease in diastolic blood pressure in the intervention group (+34 mm Hg and –4 mm Hg, respectively). In the control group, there was a decrease of the systolic blood pressure (–57 mm Hg) while the diastolic blood pressure remained unchanged [23]. After 24 weeks of supervised endurance training, Marcon et al. [25] reported a significant decrease in systolic and diastolic blood pressure (–23.8 mm Hg ($p = 0.007$) and –14.4 mm Hg ($p = 0.001$) respectively).

Table 3. Continued

Study	Exercise regime	Outcome
Stegen et al., 2011 [28]	<p><i>Strength exercises</i> were performed using stack-weight equipment (elbow flexion and extension and knee flexion and extension). Intensity of strength training was calculated from one repetition maximum (1RM) values. The starting level was 60% of 1RM, which increased up to 75% of 1RM. Patients were habituated to the exercises and equipment during the first three training sessions. In the next 15 sessions, two sets of 15 repetitions at 60% of 1RM were done. In the final 18 sessions, three sets of 10 repetitions at 75% were done. Resting periods between repetition sets were approximately 60 s. <i>Endurance training</i> consisted of cycling (10 min), walking (10 min), and stepping (10 min). Exercise intensity was initially set at 60% of heart rate reserve (HRR, defined as the difference between resting heart rate and peak heart rate) and was gradually increased up to 75% of HRR at the end of the training period. Stationary bicycles, treadmills, and steppers were used in a rehabilitation training room.</p>	<p><i>Dynamic muscle strength in kg (4 months postoperatively):</i></p> <p>Quadriceps: IG: 58.0 ± 25.6 CG: 45.9 ± 25.1 (p = 0.002)</p> <p>Biceps: IG: 25.9 ± 13.0 CG: 20.8 ± 8.8 (p = 0.001)</p> <p>Triceps: IG: 30.7 ± 23.0 CG: 22.0 ± 6.6 (p = 0.038)</p>
Berggren et al., 2008 [29]	<p>10 consecutive days of endurance-oriented exercise training. An incremental, maximal stress test was initially performed to screen for underlying cardiovascular disease and determine the workload (%peak oxygen consumption and heart rate peak) for the exercise training. VO₂ peak was determined on an electronically braked cycle ergometer in the upright position, with oxygen consumption measured via open-circuit spirometry. The training program was based upon previous exercise prescriptions that improved whole body fat oxidation or increased mitochondrial content. Participants exercised 60 min/day at 70% VO₂ peak for 10 consecutive days on a cycle ergometer. All training was performed under direct supervision. Subjects were initially permitted to take a rest break (5 min) during exercise; by day 5 all subjects exercised continuously.</p>	<p>Muscle fatty acid oxidation: Lean subjects +1.7-fold increase Obese subjects +1.8-fold increase Previously extreme obese subjects after weight loss +2.6-fold increase</p>

GBS = Gastric bypass; TG = trained group; CG = control group; EG = eutrophic group; BGBS = before gastric bypass surgery; 4GBS = 4 months after gastric bypass surgery; W = weight; 6MWT = 6-min walking test; FVC = forced vital capacity; HR = heart rate; 1RM = one repetition maximum; HRR = heart rate reserve; IG = intervention group.

Cardiovascular Risk Factors and Inflammation

Hickey et al. [24] reported a significant decrease of fasting plasma insulin (−41.7 pmol/l) after 7 days of supervised exercise training, 60 min per day. There were no significant changes in glucose and blood lipid concentration after the exercise intervention [24].

In the study by Marcon et al. [25], a significant improvement was seen in blood lipid and glucose concentrations after 24 weeks of supervised endurance training (total cholesterol −23.0 mg/dl; HDL-cholesterol +1.7 mg/dl, LDL-cholesterol −2.3 mg/dl, triglycerides −26.7 mg/dl, glucose −17.2 mg/dl (for all p < 0.007)).

Table 4. Common findings

	Shah et al., [26]	Castello-Simoes et al., [27]	Stegen et al. [28]	Baillet et al., [22]	Funderburk and Callis, [23]	Berggren et al., [29]	Hickey et al., [24]	Marcon et al., [25]
Duration, weeks	12	12	16	12	12	2	1	24
Intensity	60–70% VO ₂ max	60–70% HR	60–75% HR/RM	55–85% of HRR	x	70% VO ₂ max	65% VO ₂ peak	x
Supervision	partial	whole	whole	partial	whole	x	whole	whole
Timing bariatric surgery	post	post	post	pre	pre	post	pre	pre

HR = Heart rate; HRR = heart rate reserve; RM = repeated measurements; 2KmWT = 2-km walking test; x = unknown.

Physical Fitness and Functional Capacity

Baillet et al. [22] found that all 12 patients completed the baseline symptom limited treadmill test with a median duration of 11.5 min (11.1–12.4 min). No major cardiac complication was noted during or after the test. The 6-min walk test distance, the time of the half-squat test, and the number of flexion during the arm curl test were all significantly ($p < 0.05$) increased after the preoperative exercise program [22].

Funderburk and Callis [23] showed an increase in the 6-min walking test distance of 10.4 m after 12 weeks of aquatic exercises, but strangely the control group showed an increase of 40.2 m. After 24 weeks, Marcon et al. [25] also saw an improvement of the 6-min walking test distance (+69.8 m; $p < 0.0001$).

Aerobic Capacity

Hickey and coworkers [24] reported that after 7 days of supervised exercise there was no significant difference in maximal oxygen uptake (which was measured by VO₂ peak) indicating that there is no increase in physical fitness.

QoL and Satisfaction

Baillet et al. [22] showed that each of their 12 patients who were in the exercise program improved at least one level for endurance training and two levels for strength training. This increase appeared to be greater in the patients who trained longer. All patients were satisfied by the coaching and the advices given by the exercise professionals [22].

Also the total health-related QoL significantly improved after the exercise intervention ($p = 0.012$). On the domains of emotions, social interaction and sexual life they also noted a significant improvement ($p = 0.002$, $p = 0.025$ and $p = 0.003$, respectively) [22]. Unfortunately, no significant change was seen in symptoms ($p = 0.25$), activity/mobility ($p = 0.07$), and personal hygiene/clothing ($p = 0.078$) scores. [22]

Measured by the Short Form 36 (SF-36) questionnaire and the Beck Depression inventory, Funderburk and Callis [23] found no significant postintervention difference between groups. In the aquatic exercise group, there was a significant decrease in bodily pain and a significant decrease of the depression score after 12 weeks ($p < 0.05$). In the control group, there was a significant increase in mental health ($p < 0.05$) [23].

Exercise after Bariatric Surgery

Anthropometric Variables

Four studies reported on the effects of exercise training on the anthropometric variables.

Stegen et al. [28] investigated the effect of an exercise program after gastric bypass surgery and found that both groups (gastric bypass + exercise training (GB+E) compared with

gastric bypass only (GB)) had the same decrease 4 months postoperatively with regard to weight (GB -26.6 ± 14.6 kg; GB+E -22.7 ± 5.7 kg), BMI (GB -8.3 ± 4.1 kg/m²; GB+E -8.1 ± 2.5 kg/m²), and waist circumference (GB -20.3 ± 11.6 cm; GB+E -17.2 ± 8.1 cm) [28].

Castello-Simoes et al. [27] investigated a post-bariatric training program and used three groups: a trained group (TG), a eutrophic group (EG) and a control group (CG). They found that there was a significant reduction in weight (TG: before 115.0 ± 6.9 kg; after 92.0 ± 5.1 kg, CG: before 113.0 ± 4.7 kg, after 89.0 ± 3.4 kg) and BMI (TG: before 45.5 ± 1.7 kg/m²; after 36.5 ± 1.3 kg/m², CG: before 43.6 ± 1 kg/m², after 34.2 ± 1.1 kg/m²) in TG and CG 4 months after bariatric surgery compared to the preoperative values [27].

Berggren and colleagues [29] investigated the skeletal muscle lipid oxidation before and after 10 days of endurance orientated exercise training. They found a significant weight loss after 10 days of exercise ($p < 0.05$) [29].

Shah et al. [26] investigated a high-volume exercise program in 20 patients and found that in both the exercise group and the control group there was a significant reduction in body weight and hip circumference after 12 weeks (compared to baseline values).

Cardiovascular Risk Factors and Inflammation

In contrast to the above mentioned, Shah et al. [26] found no difference in cardiovascular risk factors after a high-volume exercise training program compared to the control group.

Physical Fitness and Functional Capacity

Castello-Simoes et al. [27] reported a significant increase in the 6-min walking test distance after exercise training and bariatric surgery (470 ± 23.9 vs. 515.0 ± 14.0 m). Shah et al. [26] found a significantly increased energy expenditure and time spent on exercise in high-volume exercise group per day.

Aerobic Capacity

In a maximal bicycle ergometer test before bariatric surgery, Stegen et al. [28] found that both groups (GB (gastric bypass only) and GB+E (gastric bypass and postoperative exercise program) reached their ventilator anaerobic threshold (VAT) at the power (GB = 93 ± 24 W; GB+E = 90 ± 24 W) with an equal time of occurrence (GB = 270 ± 107 s; GB+E = 266 ± 133 s) [28]. Gastric bypass surgery did not improve time of occurrence of VAT and power at VAT. When patients followed an exercise program (GB+E group) postoperatively, the time of occurrence of VAT was delayed (349 ± 19 s) [28]. Also Stegen and colleagues [28] registered the peak exercise parameters. In the preoperative phase, morbidly obese patients reached a peak oxygen uptake of 17.4 ± 4.9 ml/kg/min (GB) and 17.6 ± 3.2 ml/kg/min (GB+E). Unfortunately, absolute peak oxygen uptake and power did not improve 4 months after an intensive exercise program, nor by bariatric surgery.

Castello-Simoes et al. [27] reported a significant increase of the predicted forced vital capacity (94.0 ± 3.1 vs. 101.0 ± 2.5) in the trained group 4 months after bariatric surgery [27]. In the same group there was a significant reduction of the dyspnea score (5.8 ± 0.6 vs. 2.7 ± 0.8) 4 months after bariatric surgery.

Muscle Strength

Stegen et al. [28] found that the untrained patients (who only had a GB) had a decrease in dynamic muscle strength 4 months postoperatively. They have lost 16% of their quadriceps strength, 36% of their biceps strength, and 39% of triceps strength. Patients who had a GB followed by a postoperative exercise program prevented this decrease. Biceps (21.8 ± 8.0 vs. 25.9 ± 13.0 kg) and triceps (24.3 ± 10.1 vs. 30.7 ± 23.0 kg) strength were preserved, and both hamstrings and quadriceps strength increased (by 72% and 27%, respectively) due

to the exercise program [28]. Static muscle strength, which is measured as handgrip strength, decreased in both groups 4 months after GB surgery (GB 18%, GB + exercise 7%). Muscle fatigue was neither influenced by GB surgery nor by an additional exercise training program [28].

Fatty Acid Oxidation

Berggren et al. [29] studied the effect of weight loss through exercise training on the muscle fatty acid oxidation. Muscle fatty acid oxidation in extremely obese women who had lost weight (approximately 50 kg) was compared with that of extremely obese and lean individuals [29]. There was no difference in muscle fatty acid oxidation between extremely obese persons and the weight loss group. Fatty acid oxidation was depressed compared with the lean individuals (–45%; $p < 0.05$). In contrast, 10 days of exercise training increased fatty acid oxidation in the skeletal muscle of lean, obese and previously extreme obese subjects after weight loss by the factor of 1.7, 1.8 and 2.6, respectively [29].

QoL and Satisfaction

Shah et al. [26] reported a significant improvement in the high-volume exercise group with respect to the health-related QoL in several domains including physical function ($p = 0.049$), self-esteem ($p = 0.0002$), sexual life ($p = 0.02$), public distress ($p = 0.003$), and the total score ($p = 0.0004$) over 12 weeks. The control group showed a significant improvement in self-esteem ($p = 0.004$), sexual life ($p = 0.04$), work (or daily activities) ($p = 0.04$), and the total score ($p = 0.012$). [26] There was no group-by-week interaction for any of the scales, except for self-esteem, which was approximately $p = 0.05$. The data from the SF-36 questionnaire showed that there was no group-by-week interaction for any of the scales, but the high-volume exercise group reported a significant improvement in emotional wellbeing ($p = 0.001$), energy levels ($p = 0.0002$), and mental QOL total score ($p = 0.006$) over 12 weeks, whereas QoL in the control group did not change [26].

Discussion

Considering the great value assigned to exercise, a search for evidence provided a relatively little number of prospective studies addressing this subject in relation to bariatric surgery. In the majority of reports a positive effect on anthropometrics, cardiovascular risk factors, and physical fitness was described. However, the results were not unanimous, with a wide range of exercise programs and perioperative timing, therefore hampering adequate practical guidance. Simplifying the common findings, an exercise program lasts for a median of 12 weeks, has an intensity of median 65% peak heart rate/ VO_2 max, and is at least partially supervised (table 4).

Two systematic reviews [30, 31] assessed the impact of physical exercise and fitness in obese individuals, but without an in-depth analysis of the perioperative timing and the type of exercise needed to improve health in the bariatric population. The results of exercise in terms of weight loss and changes of other anthropometric variables have been described before [1–3]. However the type of exercise used to achieve this weight loss was not specified/given in detail. These details could be relevant as exercise regimes can include endurance training, strength training, or combinations of both components. Unfortunately, it is difficult to assess the appropriate type and amount of physical activity for the bariatric surgical patient. It is unclear if preoperative exercise has the same effect as postoperative exercise. Also there is no uniformity with regard to the outcome variables, and little is known about the effect of exercise programs on the weight regain after bariatric surgical interventions.

Nevertheless, it is known that exercise results in significant changes of anthropometric parameters [22–29] and greater weight loss than bariatric surgery alone [7–9].

Exercise training demonstrates favorable effects on body composition, blood pressure control, insulin sensitivity, inflammation biomarkers, and subclinical carotid atherosclerosis (CIMT). These effects have been associated with evidence of removal of oxidized phospholipids from the vessel wall, stabilization of atherosclerosis and positive changes in LDL-cholesterol [32]. These changes have a strong correlation with increase in vascular function and regression in the burden of atherosclerosis [33]. Ahmadi et al. [34] showed that the CIMT decrease was significantly greater in response to a 7-month exercise program than statin or niacin therapies in the ARBITER 6 trial [35]. In other words, exercise training reduces cardiovascular risk in obese individuals.

Bariatric surgery is the only longstanding intervention when conservative therapy fails to succeed. However, bariatric surgery also has consequences. One of them is a decrease of dynamic muscle strength after the operation [28]. A possible explanation is the decrease of muscle mass after the operation. In the study of Stegen et al. [28], the untrained patients lost a total of 7.6 kg of muscle mass through bariatric surgery; that is 29.7% of the total body weight lost (–26.6 kg).

Reviews by Stiegler and Cunliffe [36] and Chaston et al. [37] confirmed the positive correlation between weight loss and fat-free mass loss (FFML). Very low caloric diets result in a greater FFML compared to moderate caloric diets. Bariatric surgery results in greater FFML than very low caloric diets. Webster et al. [38] found that the FFML should not exceed the 22% of the total weight loss because of the function in resting metabolic rate, thermoregulation, oxidative capacity of the body, and weight management. Exercise training can attenuate muscle atrophy and can maintain FFML during weight loss [36, 37], but the value of a perioperative exercise program for bariatric surgery has not been investigated.

It can be debated whether or not decrease of muscle strength is a problem for morbidly obese patients after weight loss. It is well known that obese individuals have higher absolute muscle strength compared to lean subjects, but there is lower relative muscle strength in terms of total body weight [39–42]. This gives patients an impaired functional capacity, which is due to the fact that more strength is needed to handle a heavier body. Therefore, it might be important to prevent a decrease in muscle strength after bariatric surgery. However, when investigating muscle changes before and after bariatric surgery, several factors have to be taken into account. One of them is the type of bariatric surgery. Stegen and coworkers [28] included only patients who have undergone GB surgery. To our knowledge it is unknown whether there are differences in muscle loss between different types of bariatric surgery (e.g. GB surgery compared to sleeve gastrectomy). Moreover, the patient compliance must also be studied, i.e., whether or not patients followed the dietary instructions ((re)substitution of proteins, vitamins and minerals) [36–38]. None of these data were measured in the included studies.

It is known that obese patients have a reduced functional capacity and walking distance compared to lean subjects [43]. Tompkins and colleagues [44] showed that bariatric surgery leads to an improvement of approximately 75% in functional capacity and walking capacity 6 months after the operation. Weight loss (especially loss of body fat) makes walking less exhausting which could be demonstrated by a decreased rate of exertion and heart frequency at the end of the 6-min walking test [27, 28]. Important predictors are not only the percentage of body fat but also the maximum of oxygen uptake (VO_2 max) and the quadriceps muscle strength [43, 45, 46].

Exercise capacity can be measured by the anaerobic threshold, which reflects the true aerobic capacity and by the maximal oxygen uptake (VO_2 peak). To interpret these parameters we have to keep in mind that there are differences between obese and lean subjects.

Healthy obese subjects undergoing weight loss will experience a decrease in the maximal oxygen uptake and peak work output, because of a decrease of muscle mass [47] and an elevated cardiac stress (which is training for the heart). This cardiac stress will disappear over time [48].

In contrast, many obese patients suffer from obesity-related disorders, such as impaired cardiac function [49, 50], inefficient ventilatory work [51], and a decreased fatty acid oxidation [29, 52]. These disorders are associated with a reduced exercise capacity. During weight loss, both the cardiac and pulmonary systems will restore and may exceed the decrease in muscle mass. This delicate balance might result in the stabilization or even an improvement in maximal oxygen uptake.

Our study has several limitations that have to be taken into account. First, weight loss measured in the included studies was merely self-reported. Ideally, exercise programs and the effects of them must be monitored frequently (and objectively) by visiting a bariatric clinic. Second, because of the small sample sizes of the included studies and the heterogeneity in the reported exercise programs and outcomes, it is difficult to interpret and generalize the findings for the whole bariatric population.

It is well known that exercise programs (with or without bariatric surgery) lead to a decrease in anthropometric parameters, but there are even more beneficial effects of exercise. Future research should be focused on a comparison of different exercise regimes (e.g., endurance and strength) in obese subjects. There is also an increasing need for randomized controlled trials that focus on the effect of peribariatric exercise programs on weight regain after bariatric surgery, because this is still an ongoing problem. Studies by Bond et al. [53] and Li et al. [54] investigate whether or not a preoperative exercise program has effects on the postoperative outcome. The definitive results of those studies are not published yet. Also the long-term effects of exercise programs on QoL, depressive and anxiety symptoms, and eating behavior are not well understood.

Conclusion

In the majority of reports on exercising in a (future) bariatric population, a positive effect on anthropometrics, cardiovascular risk factors, and physical fitness was described. However, the results were not unanimous, with a wide range of exercise programs and perioperative timing, therefore hampering adequate practical guidance. Summarizing our findings, a beneficial exercise program lasts for a median of 12 weeks, had an intensity of median 65% peak heart rate / VO_2 max, and is at least partially supervised.

Financial Support

None

Disclosure Statement

The authors declare that there is no conflict of interests regarding the publication of this article.

References

- 1 World Health Organization: Obesity: preventing and managing the global epidemic. Report of a WHO consultation. World Health Organization Tech Report Ser 2000;894:i–xii:1–253.
- 2 Tjepkema M: Adult obesity. Health Reports 2006;17:9–25.
- 3 Sturm R: Increases in morbid obesity in the USA: 2000–2005. Pub Health 2007;121:492–496.
- 4 Picot J, Jones J, Colquitt JL, Gospodarevskaya E, Loveman E, Baxter L, Clegg AJ: The clinical effectiveness and cost-effectiveness of bariatric (weight loss) surgery for obesity: a systematic review and economic evaluation. Health Technol Assess 2009;3:1–190, 215–357, iii–iv.
- 5 Colquitt JL, Picot J, Loveman E, Clegg AJ: Surgery for obesity. Cochrane Database Syst Rev 2009;2:CD003641.
- 6 Buchwald H: The evolution of metabolic/bariatric surgery. Obes Surg 2014;24:1126–1135.
- 7 Shaw K, Gennat H, O'Rourke P, Del Mar C: Exercise for overweight or obesity. Cochrane Database Syst Rev 2006;4:CD003817.
- 8 Livhits M, Mercado C, Yermilov I, Parikh JA, Dutson E, Mehran A, Ko CY, Gibbons MM: Exercise following bariatric surgery: SYSTEMATIC review. Obes Surg 2010;20:657–665.
- 9 Egberts K, Brown WA, Brennan L, O'Brien PE: Does exercise improve weight loss after bariatric surgery? A systematic review. Obes Surg 2012;22:335–341.
- 10 Mosely AM, Herbert RD, Sherrington C: Evidence for physiotherapy practice: a survey of the Physiotherapy Evidence Database (PEDro). Aust J Physiother 2002;48:43–49.
- 11 Maher CG, Sherrington C, Herbert RD, Moseley AM, Elkins M: Reliability of the PEDro scale for rating quality of randomized controlled trials. Phys Ther 2003;83:713–721.
- 12 Foley NC, Teasell RW, Bhogal SK, Speechley MR: Stroke rehabilitation evidence-based review: methodology. Top Stroke Rehabil 2003;10:1–7.
- 13 Altman D: Practical Statistics for Medical Research. London, Chapman and Hall, 1991.
- 14 Vismara L, Cimolin V, Grugni G, Galli M, Parisio C, Sibilia O, Capodaglio P: Effectiveness of a 6-month home-based training program in Prader-Willi patients. Res Dev Disabil 2010;31:1373–1379.
- 15 Barbalho-Moulim MC, Miguel GP, Forti EM, Campos Fdo A, Costa D: Effects of preoperative inspiratory muscle training in obese women undergoing open bariatric surgery: respiratory muscle strength, lung volumes, and diaphragmatic excursion. Clinics (Sao Paulo) 2011;66:1721–1727.
- 16 Bernhardt V, Lorenzo S, Babb TG, Zavorsky GS: Corrected end-tidal P(CO(2)) accurately estimates Pa(CO(2)) at rest and during exercise in morbidly obese adults. Chest 2013;143:471–477.
- 17 Cattano D, Altamirano A, Melnikov V, Pivalizza E, Feldman A, Hagberg C: Postoperative pulmonary inspiratory reserve volume and incentive spirometry in morbidly obese patients undergoing bariatric surgery. J Invest Med 2010;58:678–679.
- 18 Marzocchi R, Cappellari D, Dalle Grave R, Marchesini G: Massive weight loss without surgery in a super obese patient. Obes Surg 2011;21:540–545.
- 19 Zapico AG, Benito PJ, Gonzalez-Gross M, Peinado AB, Morencos E, Romero B, Rojo-Tirado MA, Cupeiro R, Szendrei B, Butragueno J, Bermejo M, Alvarez-Sanchez M, Garcia-Fuentes M, Gomez-Candela C, Bermejo LM, Fernandez-Fernandez C, Calderon FJ: Nutrition and physical activity programs for obesity treatment (PRONAF study): methodological approach of the project. BMC Pub Health 2012;12:1100.
- 20 Adolfsson G: Physical training and weight reduction in connection with operations (in Swedish). Lakartidningen 1970;67:4941–4944.
- 21 Sanchez Ortega L, Sanchez Juan C, Garcia AA: Evaluation of a structured program of physical exercise in morbidly obese patients awaiting bariatric surgery (in Spanish). Nutr Hosp 2014;29:64–72.
- 22 Baillet A, Mampuya W, Comeau E, Meziat-Burdin A, Langlois M: Feasibility and impacts of supervised exercise training in subjects with obesity awaiting bariatric surgery: a pilot study. Obes Surg 2013;23:882–891.
- 23 Funderburk JA, Callis S: Aquatic intervention effect on quality of life prior to obesity surgery: a pilot study. Annu Ther Recreat 2010;18:66–78.
- 24 Hickey MS, Gavigan KE, McGammon MR: Effects of 7 days of exercise training on insulin action in morbidly obese men. Clin Exerc Physiol 1999;1:24–28.
- 25 Marcon ER, Gus I, Neumann CR: Impact of a minimum program of supervised exercises in the cardiometabolic risk in patients with morbid obesity. Arq Bras Endocrinol Metabol 2011;55:331–338.
- 26 Shah M, Snell PG, Rao S, Adams-Huet B, Quittner C, Livingston EH, Garg A: High-volume exercise program in obese bariatric surgery patients: a randomized, controlled trial. Obesity (Silver Spring) 2011;19:1826–1834.
- 27 Castello-Simoes V, Polaquini Simoes R, Beltrame T, Bassi D, Maria Catai A, Arena R, Azambuja NC Jr, do Nascimento Ortega J, Borghi-Silva A: Effects of aerobic exercise training on variability and heart rate kinetic during submaximal exercise after gastric bypass surgery – a randomized controlled trial. Disabil Rehabil 2013;35:334–342.
- 28 Stegen S, Derave W, Calders P, Van Laethem C, Pattyn P: Physical fitness in morbidly obese patients: effect of gastric bypass surgery and exercise training. Obes Surg 2011;21:61–70.
- 29 Berggren JR, Boyle KE, Chapman WH, Houmard JA: Skeletal muscle lipid oxidation and obesity: influence of weight loss and exercise. Am J Physiol Endocrinol Metab 2008;294:E726–732.
- 30 Baillet A, Audet M, Baillargeon JP, Dionne IJ, Valiquette L, Rosa-Fortin MM, Abou Chakra CN, Comeau E, Langlois MF: Impact of physical activity and fitness in class II and III obese individuals: a systematic review. Obes Rev 2014;15:721–739.

- 31 Fonseca-Junior SJ, Sa CG, Rodrigues PA, Oliveira AJ, Fernandes-Filho J: Physical exercise and morbid obesity: a systematic review. *Arq Bras Cir Dig* 2013;26(suppl 1):67–73.
- 32 Bergmark C, Dewan A, Orsoni A: A novel function of lipoprotein [a] as a preferential carrier of oxidized phospholipids in human plasma. *J Lipid Res* 2008;49:2230–2239.
- 33 Ahmadi N, Tsimikas S, Hajsadeghi F: Relation of oxidative biomarkers, vascular dysfunction, and progression of coronary artery calcium. *Am J Cardiol* 2010;105:459–466.
- 34 Ahmadi N, Eshaghian S, Huizenga R, Sosnin K, Ebrahimi R, Siegel R: Effects of intense exercise and moderate caloric restriction on cardiovascular risk factors and inflammation. *Am J Med* 2011;124:978–982.
- 35 Taylor AJ, Villines TC, Stanek EJ: Extended-release niacin or ezetimibe and carotid intima-media thickness. *N Engl J Med* 2009;361:2113–2122.
- 36 Stieglerand P, Cunliffe A: The role of diet and exercise for the maintenance of fat-free mass and resting metabolic rate during weight loss. *Sports Med* 2006;36:239–262.
- 37 Chaston TB, Dixon JB, O'Brien PE: Changes in fat-free mass during significant weight loss: a systematic review. *Int J Obes (Lond)* 2007;31:743–750.
- 38 Webster JD, Hesp R, Garrow JS: The composition of excess weight in obese women estimated by body density, total body water and total body potassium. *Hum Nutr Clin Nutr* 1984;38:299–306.
- 39 Maffiuletti N.A, Jubeau M, Munzinger U: Differences in quadriceps muscle strength and fatigue between lean and obese subjects. *Eur J Appl Physiol* 2007;101:51–59.
- 40 Blimkie CJ, Sale DG, Bar-Or O: Voluntary strength, evoked twitch contractile properties and motor unit activation of knee extensors in obese and non-obese adolescent males. *Eur J Appl Physiol Occup Physiol* 1990;61:313–318.
- 41 Pescatello LS, Kelsey BK, Price TB: The muscle strength and size response to upper arm, unilateral resistance training among adults who are overweight and obese. *J Strength Cond Res* 2007;21:307–313.
- 42 Hulens M, Vansant G, Lysens R: Study of differences in peripheral muscle strength of lean versus obese women: an allometric approach. *Int J Obes Relat Metab Disord* 2001;25:676–681.
- 43 Hulens M, Vansant G, Claessens AL: Predictors of 6-minute walk test results in lean, obese and morbidly obese women. *Scand J Med Sci Sports* 2003;13:98–105.
- 44 Tompkins J, Bosch PR, Chenoweth R: Changes in functional walking distance and health-related quality of life after gastric bypass surgery. *Phys Ther* 2008;88:928–935.
- 45 Laukkanen R, Oja P, Pasanen M: Validity of a two kilometre walking test for estimating maximal aerobic power in overweight adults. *Int J Obes Relat Metab Disord* 1992;16:263–268.
- 46 Mattsson E, Larsson UE, Rossner S: Is walking for exercise too exhausting for obese women? *Int J Obes Relat Metab Disord* 1997;21:380–386.
- 47 Goran M, Fields DA, Hunter GR: Total body fat does not influence maximal aerobic capacity. *Int J Obes Relat Metab Disord* 2000;24:841–848.
- 48 Seres L, Lopez-Ayerbe J, Coll R: Increased exercise capacity after surgically induced weight loss in morbid obesity. *Obesity (Silver Spring)* 2006;14:273–279.
- 49 Carroll S, Marshall P, Borkoles E: Efficacy of lifestyle intervention on peak exercise cardiac power output and reserve in premenopausal obese females: a randomised pilot study. *Int J Cardiol* 2007;119:147–155.
- 50 Kanoupakis E, Michaloudis D, Fraidakis O: Left ventricular function and cardiopulmonary performance following surgical treatment of morbid obesity. *Obes Surg* 2001;11:552–558.
- 51 Gilbert R, Sipple JH, Auchincloss JH: Respiratory control and work of breathing in obese subjects. *J Appl Physiol* 1961;16:21–26.
- 52 Corpeleijn E, Saris WH, Blaak EE: Metabolic flexibility in the development of insulin resistance and type 2 diabetes: effects of lifestyle. *Obes Rev* 2009;10:178–193.
- 53 Bond D: Bari-active: a preoperative intervention to increase physical activity. *Obes Surg* 2011;21:1042.
- 54 Li C, Zavorsky GS, Kim DJ, Christou NV, Feldman LS, Carli F: Effects of a bariatric preoperative exercise program: a pilot randomized study. *Surg Endosc* 2013;27(suppl 1):S431.