A variance analysis has shown a significant decrease in estradiol levels, while testosterone, LH, SHBG, and FTI levels did not experience any significant change.

According to these findings it can be concluded that the chronic ovarian suppression with buserelin did not influence negatively the androgenic profile.

Laparoscopic Explorative Cyto-Puncture and Aspiration of Intra-Ovarian Endometriotic Cysts and Deposits 186 Cases in 22 Years

Michel Mintz
Paris, France

84% of the patients are between 20 and 40 years old, 70% are nulliparous or sterile. Among 214 punctured cysts, 62% contained between 10 and 400 ml (27-91 mm diameter). Judged by 79 surgical controls, the reliability of cytology is 84%: it eliminated corpus luteum cysts, revealed among 214 ovarian endometriosis almost 50% of intra-ovarian hidden cases, and contributed to the discovery of 3 endometriotic malign cysts of benign aspect (1 false-negative).

Among 186 patients, 24 had complementary surgical cyst resections. 139 patients bearing 155 cysts were followed-up after puncture: 80% for more than 2 years, 50% between 6 and 20 years. Most of the prescriptions (33 different gynecologists) were not correctly followed.

Among 109 second-looks and/or ultrasonic controls: 47 cases of persistence or recurrence, 62 showed no cysts, although 15 had revealed cysts from 50 to 100 ml.

Among 65 patients who wanted children, 31 bore 40 known living children, although 16 had had large punctured cysts of 50-400 ml.

Conclusions

This technique seems appropriate to young, nulliparous or sterile women, and should be used before long prescriptions and before surgical resections which do not always prevent recurrence and may cause adhesions.

For elder patients, surgical resection of endometriotic cysts should be performed, since it concerns a risk group of ovarian cancer.

Hormonal Changes during Danol Treatment of Patients

Attila Török, József Bôdis, Imre Csaba

Department of Obstetrics and Gynecology, University Medical School of Pécs,
Pécs, Hungary

Fifteen patients were treated with 400 mg/day and another 15 patients with 300 mg/day of danol. Blood samples were taken before the treatment and in monthly intervals thereafter. Serum luteinizing hormone (LH), follicle-stimulating hormone (FSH), prolactin, hydroxycortisone, estradiol and progesterone levels were determined with radioimmunoassays.