observation would agree with the suggested conversion of the precursor material to Lea substance by the addition of a-fucosyl residues.

The outlined scheme thus fits well with many of the biochemical facts which are known about the water soluble blood group mucopolysaccharides. One possible difficulty, however, is that with the sequence of gene action suggested it would be possible for a mating of an ABH non-secretor, Lea non-secretor, i.e. an individual of the genotype I'I'S'S', with an ABH non-secretor, secretor of Lea, i.e. an individual of the genotype L'Ls's' to give rise to an ABH secretor (genotype L'VS's'). This combination would presumably be very rare and whether this difficulty is sufficiently serious to make the scheme untenable we do not feel competent to judge and we would welcome the comments of the geneticists on this point. An alternative pathway, shown in table III in which it is postulated that the gene S' can act on either the precursor substance or Lea substance, would overcome this objection. Again we would very much appreciate the comments of the geneticists and serologists, and would be very much helped if, before using these schemes as a basis for further biochemical investigations, they would give us an opinion as to whether the ideas put forward are genetically sound.

208

On the Biological Importance of Group Specific Substances

A. Cresseri
Milan, Italy

Abstract

After a short review of the physical and chemical properties of group specific substances, and of their distribution throughout the body, the author briefly discusses some possible relationships with other biological properties (resistance to experimental infections, viral inhibitors, intrinsic factor, Bifidus factor, etc.). Then follow the results of a series of researches particularly performed with pure A and H group specific substances obtained from hog gastric mucosa, which have revealed the action stimulating body growth (in rats on diets containing different amounts of protein), the sparing effect on nitrogen metabolism (in cortisonized rats), the capacity to correct dysproteinaemia and dyslipidaemia alterations occurring after gastrectomy, the capacity to hasten hematologic recovery of rats made anemic with phenylhydrazine hydrochloride, the power of protection against alimentary hepatic steatosis, etc.

The importance of these observations from a physiological point of view is discussed also in the light of other observations regarding the action of the same group specific substances on in vitro cultures of HeLa strain tumoral cells.