4. Freund, E.: Ein Beitrag zur Kenntnis der Blutgerinnung, Wiener med. Jahrbuch 1886, p.46-48.
5. Bordet, J., and O. Gengou: Recherches sur la coagulation du sang. Ann. Inst. Pasteur 17, 822, 1903.
6. Neubauer, O., and H. Lampert: Ein neuer Bluttransfusionsapparat. Münch, med. Wschr. 77, 582, 1930.
7. Hirschboeck, J.: Delayed Blood Coagulation and Absence of Clot Retraction in Collodion Lined Vessels. Proc. Soc. exp. Biol. Med. 45, 122, 1940.
8. Quick, A. J.: The Physiology and Pathology of Hemostasis. Lea & Febiger, Philadelphia 1951.
9. Quick, A. J., and E. Epstein: Thromboplastic Activity in Human Blood. J. appl. Physiol. 4, 840, 1952.
10. Quick, A. J., J. G. Georgatsos and C. V. Hussey: The Clotting Activity of Human Erythrocytes: Theoretical and Clinical Implications, Amer. J. med. Sci. 228, 207, 1954.
11. Quick, A. J., and C. V. Hussey: Prothrombin and the One-Stage Prothrombin Time. Brit, med. J. 1955 11, 934.
12. Ware, A. G., and G. F. Lanchantin: Purification of Fibrinogen, Prothrombin and Thrombin, Physiol. Rev. 34, 714, 1954.
13. Quick, A. J., and C. V. Hussey: Initiation of the Clotting of Blood. Proc. Soc. Exp. Biol. Med. 80, 40, 1952.
14. Quick, A. J., C. V. Hussey and E. Epstein: Activation of Thromboplastinogen by Thrombin. Amer. J. Physiol. 174, 123, 1953.

Plasma Thromboplastin Antecedent Deficiency

Robert L. Rosenthal

New York, New York, USA

Summary

Plasma thromboplastin antecedent is a beta-globulin which is essential to normal coagulation. Its deficiency results in a hemorrhagic disease which is relatively less severe than either AHG or PTC deficiency. Transmitted as an autosomal dominant gene with variable expression and penetrance, the disease occurs in both males and females. The incidence of PTA deficiency seems to be high in comparison with the other hereditary clotting protein abnormalities. At present, our series numbers 39 cases (29 families). Since PTA is a catalytic, stable factor, which is present in serum, it is well corrected by both fresh or stored blood or plasma (15% replacement). The correction gradually disappears over the period of one week.

23 Blood Transfusion