and on the relationship between level of arousal and retention is more persuasively in favor of a rapid consolidation theory [4].

References

- 1 Allen, S.; Oswald, I.; Lewis, S., and Tagney, J.: The effect of distorted visual input on sleep. *Psychophysiology* 9: 111 (1972).
- 2 Cartwright, R. D.: Problem solving in REM, NREM, and waking. *Psychophysiology* 9: 108 (1972).
- 3 Empson, J. and Clarke, P.: Rapid eye movements and remembering. *Nature, Lond.* 227: 287-288 (1970).
- 4 Goodenough, D.; Sap an, J.; Cohen, H.; Portnoff, G., and Shapiro, A.: Some experiments concerning the effects of sleep on memory. *Psychophysiology* 8: 749-761 (1971).
- 5 Klein, G. S.: *Perception, motives, and personality* (Knopf, New York 1970).
- 6 Lewin, I. and Gombosh, D.: Increase in REM time as a function of the need for divergent thinking. *Sleep: physiology, biochemistry, psychology, pharmacology, clinical implications. 1st Europ Congr. Sleep Res., Basel 1972*, pp. 399-403 (Karger, Basel 1973).
- 7 Muzio, J. N.: *Alterations in the normal human sleep EEG patterns and retention of meaningful verbal material; diss., Columbia University* (1971).
- 8 Zimmerman, J.; Stoyva, J., and Metcalf, D.: Distorted visual feedback and augmented REM sleep. *Psychophysiology* 7: 298 (1971).

Author's address: Dr. M. Bertini, Catholic University, Via della Pineta Sacchetti 644, I-00168 Rome (Italy)

Sleep: Physiology, Biochemistry, Psychology, Pharmacology, Clinical Implications. 1st Europ. Congr. Sleep Res., Basel 1972, pp. 457-462 (Karger, Basel 1973)

Information Processing during Sleep

An Experimental Approach¹

H. Schulz

Max-Planck-Institut für Psychiatrie, Munich

The following paper describes an experimental method to investigate the role of sleep mentation in the storing process of recently collected daytime information.

Fig. 1. Number of reactions (button-presses) per night for two subjects.

Usually, information about ongoing mental processes in a sleeping person is taken from verbal reports after awakening. There are three main objections to this method:

1. The sleep of the subject must be interrupted.
2. If one is interested in the interrelation between daytime events and sleep mentation, there is no clear criterion when to awaken the subject.
3. The awakening technique cannot be used continuously throughout the night.

Therefore, another technique for collecting data was used. There is experimental evidence that a sleeping person is able to react with specific motor actions, viz. pushing a button. Experiments of this sort were first carried out by Davis et al. [1939] and later by Granda and Hammack [1961], Fischgold and Schwartz [1961], Weinberg [1964] and Williams et al. [1966],

We adapted this experimental technique for our purposes in the following manner. After some baseline nights, a series of 7 pairs of emotionarousing slides was shown to the subject before he went to bed. During the following and all other nights he was told to push a button fixed to his

Fig. 2. Mean percentage of reactions during sleep in 6 consecutive nights after a test session (n = 4 subjects).

hand every time he was mentally occupied with test pictures during the night.

The subjects were unmarried male students, who were paid for their participation in the experiment. They slept about 8 h/night; polygraphic recordings were done. In a first experiment two of the subjects were studied for 60 nights each. In a second experiment, the three remaining subjects stayed in the laboratory for 10 nights each under the same conditions. After the test session both subjects from the first (long-term) experiment slept for the rest of the first month of the experiment without any awakenings. Only the number of reactions by pushing the button was recorded. As can be seen from figure 1, in the first night after the test session the

number of motor reactions, i.e. button-pushing, was high but decreased in the following nights (nights 3 and 4). However, the decrease was not a continuous one as the number of reactions of both subjects would again periodically increase and decrease during the whole observation period. This multiphasic time course of the number of reactions was found as well for the other 3 subjects who slept for 10 nights each in the laboratory. One of these 3 men reacted so little that he had to be excluded from statistical analysis of the data.

Figure 2 shows the time course of reactions for the first 6 nights after

Free Communications: G. Memory and Information Processing during Sleep 460

Fig. 3. Mean number of reactions during different sleep stages (subject H.K., $n = 19$ nights). REM = rapid eye movements, NREM = non-REM, W = waking, MT = movement time.

the test session for all four remaining subjects. A trend analysis revealed two significant trend components ($p < 0.05$), a linear and a quadratic one. This suggests that the pattern is characterized not only by a continuously decreasing, but also by a more differentiated function.

Next, we studied the interrelation between sleep stages and frequency of reactions.

In the course of the experiment the subject (H.K.) revealed continuously decreasing numbers of reactions during rapid eye movement (REM) sleep and an increase during non-REM (NREM) sleep (fig. 3).

The second sleeper (W.B.) produced only very few reactions during REM sleep; but he showed an increase of reactions during slow-wave sleep (NREM) in the course of the experiment (fig. 4).

Both subjects of the first (long-term) experiment came back to the sleep laboratory after an interval of one month during which they slept at home. During their second month in the sleep laboratory they were again exposed to the test stimuli after some readaptation nights. Each subject was awakened once per night. Half of the total number of awakenings followed immediately after a reaction. Control awakenings were done not earlier than 20 min after a reaction.

The awakening reports were inconclusive. No correspondence between the test stimuli and the awakening reports was found, with the exception

Schulz Information Processing during Sleep 461

Fig. 4. Mean number of reactions during different sleep stages (subject W.B., $n = 18$ nights). Abbreviations as for figure 3.

tion of some poor reports in short awakening periods where the sleepers remembered the test situation or one of the pictures.

Our experimental results may be summarized as follows:

1. The motor reactions of the sleepers change systematically in time.

However, the number of reactions does not decrease asymptotically. It shows a nonlinear trend.

2. In the beginning of the experiment, reactions are most frequently found in connection with a fast, desynchronized EEG. Later-on the number of reactions during desynchronized phases decreases and the reactions correlated with slow-wave, synchronized EEG increase in number.

3. No clear influence of the test stimuli on the verbal reports after forced awakening was found.

Two completely opposite conclusions may be drawn from our results.

We may either stress the negative results from the awakening reports and conclude that the test stimuli had no effect on sleep mentation, or we may stress the typical reaction curve and the stage shifting and conclude that the test stimuli played a role in the cognitive processes during sleep.

A tentative model may be proposed to explain the conflicting results.

It is clear that during the day there are two modes of effective action, one with and one without attentive consciousness. For example, we may drive a car for some miles without making any serious error but also without

Free Communications: G. Memory and Information Processing during Sleep 462

having any conscious recognition of the route, the landscape and of our driving behavior, and we can be sure that we were fully awake. In turn, we may make the same trip another day and do it with full attentive consciousness. Evidently one can act effectively whilst awake sometimes with and sometimes without attentive consciousness. Thus, one could hypothesize that both these modes of cognition are also available during the night. It may be that the subjects use the reaction button according to the instruction but without being conscious of it.

References

Davis, H.; Davis, P. A.; Loomis, A. L.; Harvey, E. N., and Hobart, G.: Electrical reactions of the human brain to auditory stimulation during sleep. *J. Neurophysiol.* 2: 500-514 (1939).

Fischgold, H. and Schwartz, B. A.: A clinical, electroencephalographic and polygraphic study of sleep in the human adult; in Wolstenholme and O'Connor Ciba Foundation Symposium on the Nature of Sleep, pp. 209-231 (Little, Brown, Boston

1961).

Granda, A. M. and Hammack, J. T.: Operant behavior during sleep. *Science* 133: 1485-1486 (1961).

Weinberg, H.: Evidence suggesting the acquisition of a simple discrimination during sleep. *Amer. J. Psychol.* 19: 538 (1964).

Williams, H. L.; Morlock, H. C., jr., and Morlock, J. V.: Instrumental behavior during sleep. *Psychophysiology* 2: 208-216 (1966).

Author's address: Dr. H. Schulz, Max-Planck-Institute for Psychiatry, D 8000 Munich (FRG)