Recurrent Nonconvulsive Status Epilepticus in a Patient with Coffin-Lowry Syndrome

Markus Gschwind^a Giovanni Foletti^{a, b} Alessandra Baumer^c Armand Bottani^d
Jan Novy^a

^aDepartment of Clinical Neuroscience, Centre Hospitalier Universitaire Vaudois, University of Lausanne, Lausanne,

^bInstitution de Lavigny, Lavigny, ^cInstitute of Medical Genetics, University of Zurich, Schlieren-Zurich, and

^dService of Genetic Medicine, Geneva University Hospitals, Geneva, Switzerland

Key Words

Coffin-Lowry syndrome · Nonconvulsive status epilepticus · *RPS6KA3* mutation c.575G>C (p.R192T) · Stimulus-induced drop episodes

Abstract

Coffin-Lowry syndrome (CLS) is a rare neurodevelopmental condition caused by heterogeneous mutations in the *RPS6KA3* gene on the X chromosome, leading to severe intellectual disability and dysmorphism in men, while women are carriers and only weakly affected. CLS is well known for stimulus-induced drop episodes; however, epilepsy is not commonly reported in this condition. We report on a CLS patient presenting with recurrent episodes of nonconvulsive status epilepticus (NCSE) with generalized epileptic activity, for which investigations did not find any other cause than the patient's genetic condition. This case underlines that the possibility of nonconvulsive epileptic seizures and status epilepticus should, therefore, be considered in those patients. The treatable diagnosis of NCSE may easily be overlooked, as symptoms can be unspecific.

© 2015 S. Karger AG, Basel

Coffin-Lowry syndrome (CLS) is an X-linked dominant syndrome characterized in males by severe intellectual disability, short stature, facial dysmorphism, peculiar hands (short, soft and fleshy), and progressive kyphoscoliosis [Lowry et al., 1971]. It is caused by highly heterogeneous loss-of-function mutations in the *RPS6KA3* gene located on Xp22.12 [Pereira et al., 2010]. Heterozygous females are less severely affected. About 20% of individuals with CLS present stimulus-induced drop episodes (SIDEs) which typically start from middle childhood to the teens [Stephenson et al., 2005], and 5% of individuals are also reported to be affected by epileptic seizures [Touraine et al., 2002; Hahn and Hanauer, 2012]. To the best of our knowledge, this is the first description of status epilepticus in CLS. We report on an adult patient presenting with recurrent episodes of nonconvulsive status epilepticus (NCSE).

Clinical Report

The male patient was first admitted to hospital at the age of 45 years because of a sudden episode of fluctuating decreased alertness; a second, very similar episode recurred one year later. He is an orphan living in an institution, and both his family and

Fig. 1. **a** The patient at the age of 45 years. Note the facial dysmorphisms with coarse facial traits, slightly downslanting palpebral fissures, a long, folded tongue, and a thick everted lower lip. **b** Postaxial polydactyly of the patient's left foot. **c** X-rays of the patient's hands showing the so-called tufting of all distal phalanges (circles).

past medical history are very poorly documented. Allegedly, he suffered from marked learning disabilities since early childhood. The first available medical records at the age of 37 years report severe mental handicap, but no other medical history. Upon examination, the patient presented with a happy and quiet demeanor, was of short stature (158 cm), had a severe kyphoscoliosis, and had no verbal communication. He had dysmorphic features with coarse facial traits, slightly downslanting palpebral fissures, a short philtrum, a long tongue, hypodontia, and a thick everted lower lip. (fig. 1a). Postaxial polydactyly was noted on his left foot (fig. 1b), and his fingers were neither particularly stubby nor tapering, as often described in textbooks. For several years, the patient had been experiencing attacks of very sudden lower limb weakness, especially when he got surprised, making him fall down. He needed walking assistance because of an ataxic gait. He never experienced epileptic seizures previously. Both episodes presented with repeated losses of contact of ~30 s, occurring 20 times over 2 h, with tonic head movement to the left, subtle hand myoclonus and purposeless arm movements, profuse sweating and salivation, and associated with urinary incontinence. At admission, he continued to experience episodes of head movements to the left with decreased awareness, as he underwent a video-EEG (fig. 2a). The recording showed generalized continuous ictal activity (discharges of polyspikes) which resolved after administration of 1 mg of intravenous clonazepam (fig. 2b). The patient was then loaded with 1,000 mg of levetiracetam (20 mg/kg intra-

venous). The work-up did not disclose any other known cause of status epilepticus. Brain MRI showed no major abnormality, but some minor unspecific punctiform T2-hyperintensities in the frontal white matter and moderately enlarged lateral ventricles (most notably on the posterior horns). The patient was discharged on levetiracetam 750 mg per os daily. A year later, while the treatment of levetiracetam was continued at the same dosage, very similar episodes suddenly occurred again with varying intensity. Upon admission to the hospital, he was administered an intravenous loading dose of 1,000 mg levetiracetam, which stopped the seizures. The patient recovered rapidly, and the following day, his EEG was devoid of epileptiform activity. Ictal levetiracetam serum level upon admission was detectable at 32 $\mu\text{mol/l}$, i.e. below the reference range, but in keeping with the relatively low dose he had been taking. Consequently, the dosage was increased to 1,250 mg daily. When the patient experienced another similar episode of NCSE while on monotherapy, and no triggering factor was identified for this second episode, the treatment was intensified with the addition of another wide-spectrum antiepileptic drug (valproate 500 mg daily). After the second episode, the diagnosis of CLS was postulated upon review of the patient's photographs. X-rays of the hands were performed and showed the typical tufted drumstick appearance of distal phalanges (fig. 1c). Genetic analysis of the *RPS6KA3* gene confirmed the diagnosis.

nonepileptic collapses with atonia, or exaggerated startle responses [Havaligi et al., 2007]. Others have described them as cataplexy-like events, characterized by sudden quadriceps hypotonia with subsequent collapse, or hyperekplexia-like episodes, with a startle response to mostly auditory or tactile stimuli [Hahn and Hanauer, 2012]. SIDEs are widely considered as nonepileptic episodes [Nelson and Hahn, 2003; Hahn and Hanauer, 2012].

Epileptic seizures in CLS are rarely reported. In large series, only 5% of patients with CLS were described to have epileptic seizures [Hunter, 2002; Stephenson et al., 2005], while other authors question if this truly represents an increased prevalence, as the occurrence of epilepsy is commonly reported in more than 5% of persons with mild learning disabilities and in up to 24% of persons with severe learning disabilities [Steffenburg et al., 1995; Lhatoo and Sander, 2001; Hahn and Hanauer, 2012], compared to a prevalence of around 1% in the general population [Picot et al., 2008]. It is, however, commonly thought that in individuals with learning disabilities, the increased prevalence of epilepsy is related to the fact that both epilepsy and learning disabilities are symptoms of a common underlying condition [Lhatoo and Sander, 2001]. In the specific case of CLS, mutations of *RPS6KA3* have wide implications [Sassone-Corsi et al., 1999] but without any direct link to known epileptogenic pathways [Ottman et al., 2010; Allen et al., 2013]. An association with epilepsy was however reported in a family with an Xp22.12 microduplication encompassing *RPS6KA3* [Matsumoto et al., 2013]. *RPS6KA3*, expressed in the brain, although less markedly in the cortex (www.genecards.org) [Rappaport et al., 2014], is the major generator of epileptic seizures. One of the pathways, modulated by the gene, is NMDA receptor activation, a func-

tion conceivably involved in the occurrence of seizures [Moldrich et al., 2003].

Our patient presented with recurrent unprovoked NCSE with an EEG showing a peculiar generalized pattern (long runs of polyspikes), typically seen in symptomatic generalized epilepsies [Sutter and Kaplan, 2012], the etiology of which is increasingly demonstrated to be genetic [Allen et al., 2013]. Although the short duration of the repeated clinical episodes may have suggested repeated absence seizures, we underline that recurrent motor manifestations suggestive of focal seizures as well as generalized continuous epileptic activity usually associated with tonic seizures make the best description of the patient as NCSE [Sutter and Kaplan, 2012], further highlighting the peculiarity of those episodes. Investigations did not disclose any alternative cause than the patient's underlying genetic condition. The relatively unspecific semiology of these episodes and absence of convulsions may easily have been overlooked and misdiagnosed as behavioral changes, or as a nonspecific deterioration in the general state of health occurring in parallel to medical complications in a patient with severe learning disabilities. This case highlights that prolonged nonconvulsive seizures and status epilepticus can occur in CLS. Therefore, this possibility should be considered in those patients in cases of altered behavior or a decreased general state of health, even though seizures per se are not a frequently reported feature of CLS.

Acknowledgements

We are grateful to the patient and his caregivers for their cooperation and would like to especially thank Prof. Anita Rauch from the Institute of Medical Genetics in Zurich for making the molecular analysis of *RPS6KA3* possible.

References

- Allen AS, Berkovic SF, Cossette P, Delanty N, Dlugos D, et al: De novo mutations in epileptic encephalopathies. *Nature* 501:217–221 (2013).
- Hahn JS, Hanauer A: Stimulus-induced drop episodes in Coffin-Lowry syndrome. *Eur J Med Genet* 55:335–337 (2012).
- Havaligi N, Matadeen-Ali C, Khurana DS, Marks H, Kothare SV: Treatment of drop attacks in Coffin-Lowry syndrome with the use of sodium oxybate. *Pediatr Neurol* 37:373–374 (2007).
- Hunter AG: Coffin-Lowry syndrome: a 20-year follow-up and review of long-term outcomes. *Am J Med Genet* 111:345–355 (2002).
- Lhatoo SD, Sander JW: The epidemiology of epilepsy and learning disability. *Epilepsia* 42(suppl 1):6–9, discussion 19–20 (2001).
- Lowry B, Miller JR, Fraser FC: A new dominant gene mental retardation syndrome. Association with small stature, tapering fingers, characteristic facies, and possible hydrocephalus. *Am J Dis Child* 121:496–500 (1971).

- Matsumoto A, Kuwajima M, Miyake K, Kojima K, Nakashima N, et al: An Xp22.12 microduplication including RPS6KA3 identified in a family with variably affected intellectual and behavioral disabilities. *J Hum Genet* 58:755–757 (2013).
- Moldrich RX, Chapman AG, De Sarro G, Meldrum BS: Glutamate metabotropic receptors as targets for drug therapy in epilepsy. *Eur J Pharmacol* 476:3–16 (2003).
- Nelson GB, Hahn JS: Stimulus-induced drop episodes in Coffin-Lowry syndrome. *Pediatrics* 111:e197–e202 (2003).
- O’Riordan S, Patton M, Schon F: Treatment of drop episodes in Coffin-Lowry syndrome. *J Neurol* 253:109–110 (2006).
- Ottman R, Hirose S, Jain S, Lerche H, Lopes-Cendes I, et al: Genetic testing in the epilepsies—report of the ILAE Genetics Commission. *Epilepsia* 51:655–670 (2010).
- Pereira PM, Schneider A, Pannetier S, Heron D, Hanauer A: Coffin-Lowry syndrome. *Eur J Hum Genet* 18:627–633 (2010).
- Picot MC, Baldy-Moulinier M, Daurès JP, Dujols P, Crespel A: The prevalence of epilepsy and pharmaco-resistant epilepsy in adults: a population-based study in a Western European country. *Epilepsia* 49:1230–1238 (2008).
- Rappaport N, Twik M, Nativ N, Stelzer G, Bahir I, et al: MalaCards: A Comprehensive Automatically-Mined Database of Human Diseases. *Curr Protoc Bioinformatics* 47:1.24.1–1.24.19 (2014).
- Sassone-Corsi P, Mizzen CA, Cheung P, Crosio C, Monaco L, et al: Requirement of Rsk-2 for epidermal growth factor-activated phosphorylation of histone H3. *Science* 285:886–891 (1999).
- Steffenburg U, Hagberg G, Kyllerman M: Active epilepsy in mentally retarded children. II. Etiology and reduced pre- and perinatal optimality. *Acta Paediatr* 84:1153–1159 (1995).
- Stephenson JB, Hoffman MC, Russell AJ, Falconer J, Beach RC, et al: The movement disorders of Coffin-Lowry syndrome. *Brain Dev* 27:108–113 (2005).
- Sutter R, Kaplan PW: Electroencephalographic criteria for nonconvulsive status epilepticus: synopsis and comprehensive survey. *Epilepsia* 53(suppl 3):1–51 (2012).
- Touraine RL, Zeniou M, Hanauer A: A syndromic form of X-linked mental retardation: the Coffin-Lowry syndrome. *Eur J Pediatr* 161:179–187 (2002).
- Wang Y, Martinez JE, Wilson GL, He XY, Tuck-Muller CM, et al: A novel *RSK2* (*RPS6KA3*) gene mutation associated with abnormal brain MRI findings in a family with Coffin-Lowry syndrome. *Am J Med Genet A* 140:1274–1279 (2006).