Original Paper

Plasma Catecholamine Profile of Subarachnoid Hemorrhage Patients with Neurogenic Cardiomyopathy

Michael Moussouttas^a Elizabeth Mearns^b Arthur Walters^d
Matthew DeCaro^c

^aNeurocritical Care Division, Institute for Neurosciences, Capital Regional Medical Center, Trenton, N.J., ^bPharmacy Department and ^cDivision of Cardiology, Department of Medicine, Thomas Jefferson Medical Center, Philadelphia, Pa., and ^dDepartment of Neurology, Vanderbilt Medical Center, Nashville, Tenn., USA

Key Words

Neurogenic cardiomyopathy · Norepinephrine · Catecholamines · Sympathetic function · Subarachnoid hemorrhage

Abstract

Purpose: To investigate the connection between sympathetic function and neurogenic cardiomyopathy (NC), and to determine whether NC is mediated primarily by circulating adrenal epinephrine (EPI) or neuronally transmitted norepinephrine (NE), following subarachnoid hemorrhage (SAH). **Methods:** This is a prospective observational investigation of consecutive severe-grade SAH patients. All participants had transthoracic echocardiography and serological assays for catecholamine levels – dopamine (DA), NE and EPI – within 48 h of hemorrhage onset. Clinical and serological independent predictors of NC were determined using multivariate logistic regression analyses, and the accuracy of predictors was assessed by receiver operating characteristic (ROC) curves. Multivariate linear regression analyses were used to evaluate correlations among the catecholamines. **Results:** The investigation included a total of 94 subjects: the mean age was 55 years, 81% were female and 57% were Caucasian. NC was identified in approximately 10% (9/94) of cases. Univariate analyses revealed associations between NC and worse clinical severity ($p = 0.019$), plasma DA ($p = 0.018$) and NE levels ($p = 0.024$). Plasma NE correlated with DA levels ($p = 0.206$, $p = 0.046$) and EPI levels ($p = 0.392$, $p < 0.001$), but was predicted only by plasma EPI in bivariate [parameter estimate (PE) = 1.95, $p < 0.001$] and multivariate (PE = 1.89, $p < 0.001$) linear regression models. Multivariate logistic regression analyses consistently demonstrated the predictive value of clinical grade for NC ($p < 0.05$ for all analyses) except in models incorporating plasma NE, where NC was independently predicted by NE level (OR 1.25, 95% CI 1.01–1.55) over clinical grade (OR 4.19, 95% CI 0.874–20.1). ROC curves similarly revealed the greater accuracy of plasma NE [area under the

Michael Moussouttas, MD
750 Brunswick Ave
Trenton, NJ 08638 (USA)
E-Mail moussouttas@hotmail.com

curve (AUC) 0.727, 95% CI 0.56–0.90, $p = 0.02$] over clinical grade (AUC 0.704, 95% CI 0.55–0.86, $p = 0.05$) for identifying the presence or absence of NC. **Conclusions:** Following SAH, the development of NC is primarily related to elevated plasma NE levels. Findings implicate a predominantly neurogenic process mediated by neuronal NE (and not adrenal EPI), but cannot exclude synergy between the catecholamines.

© 2015 S. Karger AG, Basel

Introduction

Subarachnoid hemorrhage (SAH) occurs in approximately 9.1/100,000 persons per year, and is associated with a 30% mortality rate [1]. Risk factors for mortality from SAH include age [2, 3], clinical grade [2, 3] and neurogenic cardiomyopathy (NC) [4]. SAH is associated with a marked acute increase in sympathetic activity [5], which is related to clinical severity [6], and which has been linked to the development of NC [7] and to disability/mortality [4].

NC is believed to result from fulminant catecholamine activation, causing excessive influx of calcium into myocytes, hypercontraction of myofibrils, myocytolysis from energy depletion, and myocardial dysfunction [8]. Uncertainty exists as to whether the process is primarily mediated by adrenal epinephrine (EPI) release into the systemic circulation, or by direct neuronal norepinephrine (NE) innervation of myocardial fibers [9, 10].

The purpose of this project is to investigate the connection between sympathetic function and the occurrence of NC, and specifically to determine which catecholamine hormone/transmitter is primarily responsible for the development of NC in patients with severe-grade SAH.

Methods

Inclusion/Exclusion Criteria

This is a prospective observational study of consecutive, primary/spontaneous, nonrecurrent, Hunt-Hess (H/H) grades 3–5 SAH patients. Patients with severe preexisting medical or cardiopulmonary disease, those requiring cardiovascular or hemodynamic resuscitation with vasopressors or inotropes, and those experiencing SAH due to secondary causes (arteriovenous malformation, trauma, intracerebral hemorrhage, dissection, infection or neoplasm) were excluded. Cases with a severe clinical grade were selected due to the greater probability of identifying patients with NC and/or catecholamine elevations [11, 12].

Investigative Overview

Demographic, clinical, radiological and angiographic data were abstracted from the medical record or obtained from family interviews. During the first 48 h following the ictus, all patients had serological samples collected for analysis of catecholamine levels, and all patients underwent transthoracic echocardiography (TTE). Eligible and enrolled patients were admitted to the Neurological Intensive Care Unit at the Hospital for Neurosciences between May 27, 2010 and June 13, 2012. The study was approved by the Institutional Research Review Board at the Thomas Jefferson Medical Center and was performed in accordance with the standards laid down by the 1964 Declaration of Helsinki and later amendments. Consent for participation was obtained from all patients or family representatives.

Sample Collection

Prior to endovascular or surgical intervention, serological samples (3 ml) were collected in lavender EDTA-containing tubes, and immediately centrifuged at 2,000 rpm for 5 min at

4°C. Supernatant was aspirated by a pipette and injected into an unused red-top tube for storage at –80°C, and the precipitant was discarded. After defrosting and mixing, the supernatant was assayed by high-performance liquid chromatography for catecholamine levels – dopamine (DA), NE and EPI – using standard techniques [13]. Assays were performed by an experienced scientist unfamiliar with the patients or details of the clinical scenario, and unaware of the TTE findings.

Cardiac Evaluation

TTE examinations were performed using the Phillips iE33 ultrasound system by experienced echocardiographers, and reviewed by a cardiologist certified in the interpretation of echocardiograms (M.D.). The reviewer was unfamiliar with the patients or details of the clinical scenario, and was unaware of any cerebrospinal fluid (CSF) assay results. NC was defined as regional or global mural cardiac dysfunction whose distribution was considered characteristic for NC (and not characteristic of cardiac ischemic disease), and which was explainable only by the inciting SAH.

Variable Categorization

Due to the low number of H/H grade 5 patients, H/H grades 4 and 5 were combined, and the clinical variable was dichotomized between grade 3 and grades 4/5 for all calculations. Admission CT scans were scored using the system proposed by Frontera et al. [14], which allows for improved linear quantification of the hemorrhage extent. Given an uneven distribution in CT scores, scores 1 and 2 and scores 3 and 4 were combined, and the radiological variable dichotomized between scores 1/2 and 3/4 for all calculations. The outcome measure evaluated was the presence or absence of NC.

Statistical Analysis

For univariate analyses, categorical proportions were compared using Fisher's exact or χ^2 tests, continuous values were correlated using Pearson's coefficient, and comparisons of continuous values were performed using the rank-sum or (two-sided) t test. For the outcome of NC, variables indicating a possible association in univariate analysis ($p < 0.1$) were reevaluated by bivariate logistic regression testing. Predictor variables from the bivariate analyses ($p < 0.05$) were cumulatively entered into multivariate logistic regression models using the backward stepwise elimination technique. For any catecholamine variable emerging as an independent predictor of NC, associations evident from correlation analyses ($p < 0.1$) were reassessed by bivariate linear regression testing. Catecholamine(s) demonstrating a predictive value in bivariate analyses ($p < 0.05$) were entered into multivariate linear regression models. Based on the results of multivariate logistic regression analyses, receiver operating characteristic (ROC) curves were created for every individual independent predictor of NC, and for combined models incorporating all independent predictors cumulatively. For all analyses, a p value < 0.05 was considered statistically meaningful. The SPSS program v.12 was used for all computations.

Results

Patient Selection

To obtain the study group, a total of 312 patients were screened. Of these patients, 134 were excluded due to low clinical severity (H/H grades 1/2), leaving a candidate study population of 178 patients. From these, 80 were excluded due to discontinuation of support (22/80), late presentation (21/80), delayed enrollment (10/80), lack of patient or family

Moussouttas et al.: Plasma Catecholamine Profile of Subarachnoid Hemorrhage Patients with Neurogenic Cardiomyopathy

Table 1. Demographic, clinical and radiological characteristics of the two study groups

	NC (n = 9)	No NC (n = 85)	p value
Demographics			
Age, years	49 ± 17	55 ± 14	0.348
Female	79	81	0.547
Caucasian	67	57	0.730
Medical history			
Hypertension	38	62	0.260
Tobacco use	44	49	1.00
EtOH use	22	18	0.664
Drug use	0	7	0.537
CAD	0	5	0.693
SAH features			
H/H grade 4/5	67	26	0.019
CT score 3/4	56	41	0.490
Aneurysm location			
A1-ACom/MCA	50/17	40/7	0.690/0.920
Posterior/VA	0/0	15/9	0.590/1.00
Treatment modality			
Endovascular	100	93	0.537

Data reveal the association between clinical severity (H/H grade) and the presence of NC. For age, values represent means ± SD; the remaining values represent percentages. Drug use refers to sympathomimetic agents of abuse. EtOH use = Ethanol use; CAD = coronary artery disease; A1-ACom = proximal anterior cerebral/communicating artery complex; MCA = middle cerebral artery; posterior = posterior circulation; VA = vertebral artery.

Table 2. Catecholamine levels and timing of serological acquisition in the two study groups

	NC (n = 9)	No NC (n = 85)	p value
Sample acquisition			
Collection time, h	23 ± 10	21 ± 9	0.580
Catecholamine level, pg/μl			
DA	7.93 ± 22.5	4.03 ± 36.4	0.018
NE	7.91 ± 11.3	1.82 ± 2.36	0.024
EPI	0.645 ± 0.986	0.428 ± 0.878	0.289

Data reveal the association between plasma NE levels and NC. For all variables, values represent means ± SD. p values <0.1 are bolded.

consent (9/80), cerebral circulatory arrest (5/80), early rehemorrhage (4/80), cardiopulmonary arrest (3/80), SAH secondary to trauma (2/80) or infectious endocarditis (2/80), a previous history of SAH (1/80), and serious medical illness of metastatic carcinoma (1/80).

Subject Characteristics

Of the 98 originally enrolled subjects, 3 did not undergo TTE due to anticipated mortality, and in 1 patient TTE imaging quality was inadequate for interpretation. Therefore, the final study group consisted of 94 patients: the mean age was 55 years, 81% were female and 57% were Caucasian. NC was identified in approximately 10% (9/94) of cases. Demographic, clinical, radiological and laboratory details for the two groups are listed in tables 1 and 2.

Table 3. Correlation analyses and bivariate/multivariate linear regression analyses for plasma NE levels

	Correlation analyses		Bivariate analyses		Multivariate analyses	
	coefficient (ρ)	p value	PE	p value	PE	p value
H/H grade	NA		1.83	0.06	NA	
EPI level	0.392	<0.001	1.95	<0.001	1.89	<0.001
DA level	0.206	0.046	n.s.		NA	

Data from correlation analyses and linear regressions reveal the association between/predictive value of plasma EPI and/or plasma NE levels. p values <0.05 are bolded. NA = Not applicable; n.s. = not significant.

Table 4. Bivariate and multivariate logistic regression analyses for the main outcome measure NC

	Bivariate analyses		Multivariate analyses	
	OR (95% CI)	p value	OR (95% CI)	p value
H/H grade	5.73 (1.32–24.9)	0.020	4.19 (0.874–20.1)	0.073 ^a
NE level	1.28 (1.04–1.56)	0.017	1.25 (1.01–1.55)	0.039
DA level	0.998 (0.983–1.01)	0.730	NA	

Data from bivariate analyses reveal the predictive value of clinical severity (H/H grade) and plasma NE levels for the occurrence of NC. Data also reveal NE as an independent predictor of NC, over and above H/H grade in multivariate analyses. p values <0.05 are bolded. NA = Not applicable.

^a Alternate models including H/H grade, CT score, gender and age consistently validate the independent predictive value of H/H grade for NC (p < 0.05 for all analyses).

Univariate and Logistic Regression Analyses

From the univariate analyses (tables 1, 2), NC was associated with a worse H/H grade (p = 0.019), plasma NE levels (p = 0.024) and plasma DA levels (p = 0.018). Correlation analyses (table 3) demonstrated a positive association between plasma NE and EPI levels (ρ = 0.392, p < 0.001), and between plasma NE and DA levels (p = 0.206, p = 0.046). Bivariate logistic regression (table 4) identified H/H grade (OR 5.73, 95% CI 1.32–24.9) and plasma NE levels (OR 1.28, 95% CI 1.04–1.56), but not plasma DA levels (p = 0.730), as predictors of NC. In the multivariate logistic regression analysis (table 4), NC was independently predicted by plasma NE level (OR 1.25, 95% CI 1.01–1.55), more so and to the exclusion of H/H grade (OR 4.19, 95% CI 0.874–20.1). Alternate models (including H/H grade, gender and age) consistently confirmed the independent predictive value of H/H grade for NC (p < 0.05 for all analyses).

Linear Regression and ROC Curve Analyses

In the bivariate linear regression analyses (table 3), plasma EPI was a predictor of NE level (p < 0.001), and multivariate linear regression models (variably including H/H grade, gender and age) identified plasma EPI as an independent predictor of the NE level (p < 0.001 for all analyses). ROC curves for the variables H/H grade and NE level, individually and in combination, were constructed for the primary outcome of NC (fig. 1). Moderate accuracy was observed for H/H grade [area under the curve (AUC) 0.704, 95% CI 0.55–0.86, p = 0.05] and for plasma NE level (AUC 0.727, 95% CI 0.56–0.90, p = 0.02), as well as for the combined model (AUC 0.77, 95% CI 0.62–0.92, p = 0.01).

Fig. 1. ROC curves for the accuracy of clinical H/H grade and plasma NE levels in predicting NC. The graphs illustrate the accuracy of clinical H/H grade (H/H) and plasma NE levels (NE) (**a**) and that of a combined model in predicting the development of NC (**b**). Moderate accuracy was observed for H/H grade [AUC 0.704, 95% CI 0.55–0.86, $p = 0.05$] and plasma NE level [AUC 0.727, 95% CI 0.56–0.90, $p = 0.02$], as well as for the combined model [AUC 0.77, 95% CI 0.62–0.92, $p = 0.01$] in predicting NC.

Discussion

Catecholamine synthesis begins with the conversion of L-3,4-dihydroxyphenylalanine to DA by L-3,4-dihydroxyphenylalanine decarboxylase. DA is then converted by DA β -hydroxylase to NE in sympathetic nerve terminals, which in adrenal gland cells is transformed into EPI by phenylethanolamine N-methyltransferase. Serological quantification of catecholamine levels provides a valuable measure of separate aspects of sympathoadrenal function, in that plasma NE represents neuronal innervation and activation, whereas plasma EPI reflects adrenal endocrine secretion and activity [15].

The major finding of our investigation is that the development of NC is related to plasma catecholamine activity, and that the main catecholamine associated with NC is NE. Findings therefore implicate a predominantly neuronal process in the production of NC via direct innervation of the myocardium by sympathetic nerves, over an endocrine process involving adrenally secreted EPI that circulates systemically before activating myocyte β -receptors. The results of our investigation are consistent with prior investigations, including two retrospective studies demonstrating increased NE (but not EPI) levels among SAH patients with myocardial dyskinesias [16, 17], studies revealing a greater frequency of α_2 -receptor polymorphisms predisposing to an increased presynaptic release of NE among SAH patients developing NC [18], and studies identifying NE (and not EPI) as the main catecholamine elevated following acute SAH [5, 19, 20]. However, the observed associations between plasma NE and EPI levels do not exclude the possibility of synergy between the two catecholamines.

The findings contradict hypotheses implicating EPI as the predominant catecholamine involved in producing NC [10], prior investigations demonstrating greater elevations of EPI over NE among non-SAH patients presenting with NC [9], and some animal models of SAH-induced myocardial injury [21]. Such theories cite greater myocardial apical concentrations of β_2 -receptors with particular affinity towards EPI, and decreased sympathetic NE inner-

vation at the apex of the myocardium, as explanations for why ‘typical’ NC characterized by apical paralysis may be mediated primarily by EPI [10]. However, such theories fail to consider the proven direct connection between hypothalamic excitation and myocardial injury [22, 23], and cannot explain experiments revealing greater cardioprotection from sympathetic lesioning than from adrenal resection in animal models of NC [24–26]. However, our investigation cannot exclude the possibility of a predominantly adrenal EPI-mediated process for NC in nonintracranial conditions.

Similar to previous investigations, no connection was demonstrated between NC and age [11, 12], gender [12, 27] or aneurysm location [12, 20]. Conversely, our investigation confirms prior reports that NC is related to clinical severity [11, 12, 28, 29] and, importantly, demonstrates that this association may be mediated by neuronal NE transmission. This conclusion is based on the logistic regression findings which reveal the prominence of NE level over H/H grade as an independent predictor of NC. Findings provide parsimony between various investigations identifying clinical grade as a/the principle determinant of central catecholaminergic activation [6], investigations implicating (nor)adrenergic function in the production of NC [7, 16, 17], and investigations identifying clinical severity as a major determinant of NC among patients with SAH [11, 12, 28, 29].

The results are analogous to our previous report demonstrating a tendency towards greater CSF NE levels among patients with NC, yet they are also discrepant to the prior report in which clinical severity emerged as a greater predictor of NC than CSF NE [28]. Differences between the current and original investigations may be explained by inadequate statistical power to demonstrate consistently comparable results, and by the influence of unmeasured variables such as prehemorrhage antiadrenergic medications [30], hormonal factors and genetic polymorphisms [16, 18]. In addition, plasma NE may provide a more accurate measure of sympathetic discharge reaching the myocardium, and CSF contamination by hemorrhagic products may produce dilutional and chemical alterations with less precise measures. Alternatively, clinical severity and NE release may represent independent parallel sequelae of a common central apoplectic event, such as acute severe intracranial hypertension [21], so that inclusion of clinical grade into the multivariate model may confound analyses regarding the impact of NE upon NC. In the absence of statistical correction/adjustment for clinical grade, our investigations become entirely consistent, with CSF and plasma NE levels both independently predicting NC.

Probable anatomic and physiological structures and processes involved in central NC are illustrated in figure 2. On a molecular level, NE activation of the β_1 -adrenergic receptor (β_1 AR) initiates guanosine protein induction of adenylate cyclase, converting adenosine triphosphate to cyclic adenosine monophosphate, which in turn induces protein kinase A (PKA) activation of cellular membrane calcium channels, cytoplasmic calcium influx and actin/myosin filament contraction. However, guanosine protein also activates the phosphoinositide second-messenger system, which produces phosphoinositide 3-kinases composed of P110 γ and P84/87 subunits, and ultimately the formation of phosphatidylinositol (3,4,5)-triphosphate [31]. Normally, PKA functions as the main cellular mediator of (nor) adrenergic function by inhibiting P110 γ counteraxis activity, while at the same time modulating phosphodiesterase 3B activity, which in turn regulates cyclic adenosine monophosphate and PKA [31]. Conversely, in chronic heart failure (a condition characterized by chronically elevated catecholaminergic activity), PKA inhibition of P110 γ is impaired, allowing the P110 γ activation of β -adrenergic receptor kinase (β ARK) [G-protein-coupled receptor kinase 2 (GRK2)] enhancement of β -arrestin, which ultimately causes desensitization and internalization of β_1 AR, rendering the myocardium refractory to sympathetic stimulation [31, 32]. The importance of β ARK (GRK2) and β -arrestin in regulating the sympathetic control of cardiac contractility is evident in various clinical circumstances including myocardial

Fig. 2. Proposed anatomic and physiologic structures and processes involved in central NC.

infarction [33, 34] and chronic heart failure [32], as well as in structural reparative/reactive processes such as remodeling [33] and ventricular hypertrophy (for GRK5) [35]. In addition to cytosolic functions, the influence of GRK2 may be mediated by the modification of mitochondrial gene expression, and thus metabolic function and energy production [36].

Excessive NE release, causing myofibrillary hypercontraction with adenosine triphosphate depletion and myocytolysis, represents one potential means by which myocardial dyskinesia may occur in NC [8]. Secondary damaging consequences of marked calcium release, including free radical formation, cellular enzyme degradation, and cytoskeletal destruction, have also been proposed [37]. However, NC is also associated with acute functional sympathetic denervation [7], possibly analogous to the β 1AR catecholamine resistance present in heart failure, which may theoretically also contribute to the inotropic deficiency observed. Similar to the situation in heart failure, it has been hypothesized that predominant activation of the phosphoinositide 3-kinase system by fulminant (nor)adrenergic release may inactivate β 1ARs and lead to temporary catecholamine resistance, and to the myocardial dyskinesia of NC [10]. Given the known functions of the principal myocytic sympathetic mediators [PKA, P110 γ , β ARK (GRK2) and β -arrestin] in normal conditions and in disease states, it is interesting to speculate on the role these compounds may play in the production of NC, and whether the manipulation of these compounds may eventually lead to successful treatments for NC.

Study limitations include the small number of subjects enrolled, the exclusion of many patients who originally presented and were screened, and possibly the delay in the acqui-

sition of serological specimens. However, statistically relevant and physiologically plausible findings were revealed despite the limited number of subjects, and elevated catecholamine levels are known to persist for several days following acute cardiac and intracranial events [5, 9, 38, 39]. Regarding the exclusion of candidate patients, every attempt was made by the researchers to enroll as many participants as possible and as quickly as possible into the investigation. The exclusion of critical patients who died early, who may have exhibited greater elevations in catecholamine levels and therefore greater propensity for developing NC, may have led to an underestimation of the impact of the (nor)adrenergic response upon the development of NC. An additional limitation is the inability to account for variable usage and differing doses of pre/posthemorrhage $\alpha\beta$ -receptor modulating and sedative-analgesic medications [30], or the impact of such medications upon the sympathoadrenal response of any given individual, and thus the influence of these agents upon our results. Finally, the modest accuracy of sensitivity/specificity models in estimating the occurrence of NC indicates the influence of factors not evaluated by our work, such as catecholamine receptor polymorphisms and hormonal interactions [16, 18].

Attributes include analogous findings to those of previous investigators [7, 16, 17, 24–26] and the consistency of our findings with prevailing contemporary theories regarding a sympathetically mediated neurogenic process in the development of NC [37]. Additionally, results are internally consistent given the positive associations between plasma catecholamine levels, and externally validated by reinforcing the connection between clinical SAH severity and the presence of NC [11, 12, 28]. The prospective nature of this investigation and the masking of investigators regarding echocardiography and laboratory results minimized the likelihood of bias and adds to the validity of the results. Finally, we believe our research represents the only prospective investigation relating NC to plasma NE levels and presents the novel finding that clinical severity likely mediates NC via plasma NE elevations.

Conclusion

In SAH, and potentially any intracranial process, the development of NC is related to increased catecholamine activity, and in particular to elevations in plasma NE levels. Findings implicate a primarily neuronal NE-mediated (and not adrenal EPI-mediated) process, which is thus neurogenic in origin. Results cannot exclude the potential for synergy between NE and EPI in the development of NC following SAH, nor a predominantly EPI-mediated process in nonintracranial conditions.

Acknowledgments

The authors express their gratitude to the nurses of the Neurological Intensive Care Unit from the Jefferson Hospital of Neurosciences at Thomas Jefferson University Medical Center for their assistance in this project.

References

- 1 Gonzalez-Perez A, Gaist D, Wallander MA, McFeat G, Garcia-Rodriguez LA: Mortality after hemorrhagic stroke: data from general practice (the health improvement network). *Neurology* 2013;81:559–565.
- 2 Rosengart A, Schultheiss K, Tolentino J, Macdonald L: Prognostic factors for outcome in patients with aneurysmal subarachnoid hemorrhage. *Stroke* 2007;38:2315–2321.

- 3 Shirao S, Yoneda H, Kunitsugu I, Ishihara H, Koizumi H, Suehrio E, et al: Preoperative prediction of outcome in 283 poor-grade patients with subarachnoid hemorrhage – a project of the Chugoku-Shikoku Division of the Japan Neurosurgical Society. *Cerebrovasc Dis* 2010;30:105–113.
- 4 Moussouttas M, Huynh T, Khoury J, Lai E, Pello S, Dombrowski K, et al: Cerebrospinal fluid catecholamine levels as predictors of outcome in subarachnoid hemorrhage. *Cerebrovasc Dis* 2012;33:173–181.
- 5 Naredi S, Lambert G, Eden E, Zall S, Runnerstam M, Rydenhag B, et al: Increased sympathetic nervous activity in patients with nontraumatic subarachnoid hemorrhage. *Stroke* 2000;31:901–906.
- 6 Moussouttas M, Lai E, Khoury J, Huynh T, Dombrowski K, Pacak K: Determinants of central sympathetic activation in spontaneous primary subarachnoid hemorrhage. *Neurocrit Care* 2012;16:381–388.
- 7 Banki N, Kopelnik A, Dae M, Miss J, Tung P, Lawton M, et al: Acute neurocardiogenic injury after subarachnoid hemorrhage. *Stroke* 2005;112:3314–3319.
- 8 Bybee KA, Prasad A: Stress related cardiomyopathy syndromes. *Circulation* 2008;118:397–409.
- 9 Wittstein I, Thieman D, Lima J, Baughman K, Schulman S, Gerstenblith G, et al: Neurohumoral features of myocardial stunning due to sudden emotional stress. *N Engl J Med* 2005;352:539–548.
- 10 Lyon A, Rees P, Prasad S, Poole-Wilson P, Harding S: Stress (Takotsubo) cardiomyopathy – a novel pathophysiological hypothesis to explain catecholamine induced acute myocardial stunning. *Nat Clin Pract Cardiovasc Med* 2008;5:22–29.
- 11 Tung P, Kopelnik A, Banki N, Ong K, Ko N, Lawton M, et al: Predictors of neurocardiogenic injury after subarachnoid hemorrhage. *Stroke* 2004;35:548–553.
- 12 Kothavale A, Banki N, Kopelnik A, Yarlagadda S, Lawton M, Ko N, et al: Predictors of left ventricular regional wall motion abnormalities after subarachnoid hemorrhage. *Neurocrit Care* 2006;4:199–205.
- 13 Eisenhofer G, Goldstein D, Stull R, Keiser H, Sunderland T, Murphy D, et al: Simultaneous liquid chromatographic determination of 3,4-dihydroxyphenylglycol, catecholamines, and 3,4-dihydroxyphenylalanine in plasma, and their responses to inhibition of monoamine oxidase. *Clin Chem* 1986;32:2030–2033.
- 14 Frontera J, Claassen J, Schmidt J, Warternberg K, Temes R, Connolly S, et al: Prediction of symptomatic vasospasm after subarachnoid hemorrhage. *Neurosurgery* 2006;58:21–27.
- 15 Goldstein D, Eisenhofer G, Kopin I: Sources and significance of plasma levels of catechols and their metabolites in humans. *J Pharmacol Exp Ther* 2003;305:800–811.
- 16 Sugimoto K, Inamasu J, Hirose Y, Kato Y, Ito K, Iwase M, et al: The role of norepinephrine and estradiol in the pathogenesis of cardiac wall motion abnormality associated with subarachnoid hemorrhage. *Stroke* 2012;43:1897–1903.
- 17 Sugimoto K, Inamasu J, Kato Y, Yamada Y, Ganaha T, Oheda M, et al: Association between elevated plasma norepinephrine levels and cardiac wall motion abnormality in poor-grade subarachnoid hemorrhage patients. *Neurosurg Rev* 2013;36:259–266.
- 18 Zaroff J, Pawlikowska L, Miss J, Yarlagadda S, Ha C, Achrol A, et al: Adrenoceptor polymorphisms and the risk of cardiac injury and dysfunction after subarachnoid hemorrhage. *Stroke* 2006;37:1680–1685.
- 19 Benedict CR, Loach AB: Clinical significance of plasma adrenaline and noradrenaline concentrations in patients with subarachnoid haemorrhage. *J Neurol Neurosurg Psychiatry* 1978;41:113–117.
- 20 Lambert G, Naredi S, Eden E, Rydenhag B, Friberg P: Monoamine metabolism and sympathetic nervous system activation following subarachnoid hemorrhage – influence of gender and hydrocephalus. *Brain Res Bull* 2002;58:77–82.
- 21 Masuda T, Sato K, Yamamoto S, Matsuyama N, Shimohama T, Matsunaga A, et al: Sympathetic nervous activity and myocardial damage immediately after subarachnoid hemorrhage in a unique animal model. *Stroke* 2002;33:1671–1676.
- 22 Melville K, Blum B, Shister H, Silver M: Cardiac ischemic changes and arrhythmias induced by hypothalamic stimulation. *Am J Cardiol* 1963;12:781–791.
- 23 Doshi R, Neil-Dwyer G: Hypothalamic and myocardial lesions after subarachnoid hemorrhage. *J Neurol Neurosurg Psychiatry* 1977;40:821–826.
- 24 Selye H: *The Chemical Prevention of Cardiac Necrosis*. New York, Ronald Press, 1958.
- 25 Raab W, Stark E, Macmillan WH, Gigue WR: Sympathogenic origin and antiadrenergic prevention of stress-induced myocardial lesions. *Am J Cardiol* 1961;8:203–211.
- 26 Greenhoot JH, Reichenbach DD: Cardiac injury and subarachnoid hemorrhage. A clinical, pathological, and physiological correlation. *J Neurosurg* 1969;30:521–531.
- 27 Kopelnik A, Fisher L, Miss J, Banki N, Tung P, Lawton M, et al: Prevalence and implications of diastolic dysfunction after subarachnoid hemorrhage. *Neurocrit Care* 2005;3:132–138.
- 28 Moussouttas M, Lai E, Dombrowski K, Huynh T, Khoury J, Carmona G, et al: CSF catecholamine profile in subarachnoid hemorrhage patients with neurogenic cardiomyopathy. *Neurocrit Care* 2011;14:401–406.
- 29 Inamasu J, Hayashi T, Oheda M, Yamashiro K, Tateyama S, Kogame H, et al: Cardiac wall motion abnormality after bleeding from vertebral artery aneurysms. *Clin Auton Res* 2014;24:259–264.
- 30 Liang CW, Chen R, Macri E, Naval N: Preadmission beta-blockers are associated with decreased incidence of neurogenic stunned myocardium in aneurysmal subarachnoid hemorrhage. *J Stroke Cerebrovasc Dis* 2013;22:601–607.
- 31 Perino A, Ghigo A, Ferrero E, Morello F, Santulli G, Baillie GS, et al: Integrating cardiac PIP3 and cAMP signaling through a PKA anchoring function of p110γ. *Mol Cell* 2011;42:84–95.

- 32 Santulli G: Adrenal signaling in heart failure: something more than a distant ship's smoke on the horizon. *Hypertension* 2014;63:215–216.
- 33 Santulli G, Campanile A, Spinelli L, Assante di Panzillo E, Ciccarelli M, Trimarco B, et al: G protein-coupled receptor kinase 2 in patients with acute myocardial infarction. *Am J Cardiol* 2011;107:1125–1130.
- 34 Santulli G, Trimarco B, Iaccarino G: G-protein-coupled receptor kinase 2 and hypertension: molecular insights and pathophysiological mechanisms. *High Blood Press Cardiovasc Prev* 2013;20:5–12.
- 35 Sorriento D, Santulli G, Fusco A, Anastasio A, Trimarco B, Iaccarino G: Intracardiac injection of AdGRK5-NT reduces left ventricular hypertrophy by inhibiting NF-kappaB-dependent hypertrophic gene expression. *Hypertension* 2010;56:696–704.
- 36 Fusco A, Santulli G, Sorriento D, Cipolletta E, Garbi C, Dorn GW 2nd, et al: Mitochondrial localization unveils a novel role for GRK2 in organelle biogenesis. *Cell Signal* 2012;24:468–475.
- 37 Samuels MA: The brain-heart connection. *Circulation* 2007;116:77–84.
- 38 Espiner EA, Leikis R, Ferch RD, MacFarlane MR, Bonkowski JA, Frampton CM, et al: The neuro-cardio-endocrine response to acute subarachnoid hemorrhage. *Clin Endocrinol* 2002;56:629–635.
- 39 Naredi S, Lambert G, Friberg P, Zall S, Eden E, Rydenhag B, et al: Sympathetic activation and inflammatory response in patients with subarachnoid hemorrhage. *Intensive Care Med* 2006;32:1955–1961.