The Effects of Thyroid Hormones on Gene Expression of Acyl-Coenzyme A Thioesterases in Adipose Tissue and Liver of Mice

Kerstin Krause^a Juliane Weiner^a Sebastian Hönes^d Nora Klöting^{a, c} Eddy Rijntjes^e
John T. Heiker^a Claudia Gebhardt^a Josef Köhrle^e Dagmar Führer^d Karen Steinhoff^b
Swen Hesse^{b, c} Lars C. Moeller^d Anke Tönjes^a

^aDivision of Endocrinology and Nephrology, Department of Medicine, and ^bDepartment of Nuclear Medicine, University of Leipzig, and ^cIFB Adiposity Diseases, Leipzig University Medical Centre, Leipzig, ^dDepartment of Endocrinology and Metabolism, University of Duisburg-Essen, Essen, and ^eInstitute for Experimental Endocrinology, Charité-Universitätsmedizin Berlin, Berlin, Germany

Key Words

Thyroid hormones · Acyl-CoA thioesterases · Fatty acids · Brown adipose tissue

Abstract

Background: Thyroid hormones (TH) exert pleiotropic effects on glucose and lipid homeostasis. However, it is as yet unclear how TH regulate lipid storage and utilization in order to adapt to metabolic needs. Acyl-CoA thioesterases (ACOTs) have been proposed to play a regulatory role in the metabolism of fatty acids. **Objectives:** We investigated the interaction between thyroid dysfunction and *Acot* expression in adipose tissues and livers of thyrotoxic and hypothyroid mice. **Methods:** Ten-week-old female C57BL/6NTac mice ($n = 10/\text{group}$) were made hyperthyroid by the application of L-thyroxine (2 $\mu\text{g}/\text{ml}$ in drinking water) for 4 weeks. Hypothyroidism was induced in 10-week-old mice by feeding an iodine-free chow supplemented with 0.15% PTU for 4 weeks. We measured mRNA expression levels of *Acot8*, *11* and *13* in the liver and epididymal and inguinal white and brown adipose tissues (BAT). Furthermore, we investigated hepatic *Acot* gene expression in TR α - and TR β -deficient mice. **Results:** We showed that the expression of *Acot8*, *11* and *13* is

predominantly stimulated by a thyrotoxic state in the epididymal white adipose tissue. In contrast, hypothyroidism predominantly induces the expression of *Acot8* in BAT in comparison with BAT of thyrotoxic and euthyroid mice ($p < 0.01$). However, no significant changes in *Acot* expression were observed in inguinal white adipose tissue. In liver, *Acot* gene expression is collectively elicited by a thyrotoxic state. **Conclusions:** These data suggest that ACOTs are targets of TH and are likely to influence 3,5,3'-triiodo-L-thyronine-orchestrated mechanisms of lipid uptake, storage and utilization to adapt the regulation of metabolic demands.

© 2015 European Thyroid Association
Published by S. Karger AG, Basel

Introduction

Thyroid hormones (TH) regulate a variety of physiological processes, including growth and development, and are potent regulators of metabolism throughout life [1, 2]. TH status correlates with body weight and energy

K.K. and J.W. contributed equally to this work.

expenditure [3–5]. Elevated circulating concentrations of TH lead to a hypermetabolic state which is characterized by increased resting energy expenditure, weight loss, accelerated lipolysis and gluconeogenesis, and reduced serum cholesterol concentrations. In contrast, the manifestation of a hypothyroid state, i.e. decreased serum TH concentration, is associated with hypometabolism characterized by cold intolerance, weight gain and increased serum cholesterol concentrations [1].

It is well established that TH stimulate both a lipogenesis/lipolysis ‘futile cycle’ and that elevated TH concentrations lead to fat loss [6]. This is in particular regulated by the liver-specific crosstalk of the active TH 3,5,3'-triiodo-L-thyronine (T₃) by binding to its cognate receptors TR α and TR β and by TR crosstalk with other nuclear receptors, for instance PPAR α , PGC1 α and LXR [7]. Lipid storage and synthesis are regulated by T₃ via an increased expression of lipogenic genes such as fatty acid (FA) synthase (*Fas*), Thrsp (*Spot14*) and acetyl-coA-carboxylase (*Acc1*) [8]. Moreover, T₃ stimulates the shuttling of free FA (FFA) into mitochondria by increasing the expression and activity of CPT1 α , which is the rate-limiting step for β -oxidation of FA in the mitochondria [9].

A prerequisite for nearly all FA metabolic pathways is the activation of ligation of FA to acyl-CoA by acyl-coA synthetases [10]. The high-energy acyl-CoA thioesters subsequently serve as substrates for numerous processes, in particular lipid synthesis and β -oxidation [11]. This process is counterbalanced by the hydrolysis of fatty acyl-CoA thioesters to FFA and coenzyme A (CoASH) by Acyl-CoA thioesterases (ACOTs) [12–14]. Although their reactions are well known, the cellular and metabolic effects of ACOTs are incompletely understood.

ACOTs are localized in different intracellular compartments, e.g. cytoplasm (*Acot1*), peroxisomes (*Acot8*) and in mitochondria (*Acot2*, *11* and *13*). *Acot11* is predominantly expressed in brown adipose tissue (BAT), a tissue characterized by its high number of mitochondria, in which the uncoupling of energy metabolism and ATP production by UCP-1 (uncoupling protein-1) provides the basis for nonshivering thermogenesis. In a recent study on cultured brown adipocytes it was demonstrated that there is a switch of isoform expression from *Acot1* to *Acot2* during brown adipocyte differentiation, thereby shifting the availability of an intracellular pool of Acyl-CoA (restricted by *Acot1*) towards increased mitochondrial acyl-CoA uptake for β -oxidation and lipid synthesis [15]. Peroxisomal *Acot8* is abundantly expressed and has a broad CoA substrate specificity. Interestingly, it has been demonstrated by Hunt et al. [16] that *Acot8* activity

is inhibited by CoASH. Thus, it is anticipated that *Acot8* plays a role in the regulation of intraperoxisomal CoA/CoASH concentrations in order to balance the availability of FA for β -oxidation [16].

In view of the unique role of ACOTs in directing lipid storage and utilization we investigated the interaction between thyroid dysfunction and *Acot* gene expression in adipose tissue and liver of mice. We show that gene expression of *Acot8*, *11* and *13* is stimulated by a thyrotoxic state predominantly in epididymal white adipose tissue, whereas a hypothyroid state enhances the expression of *Acot8*, in particular in BAT. Likewise, hepatic *Acot* expression is elicited by thyrotoxicosis. These data suggest that ACOTs are targets of TH and may provide a fundamental means by which T₃ integrates the mechanisms of lipid uptake, storage and utilization to gauge the regulation of metabolic demands.

Materials and Methods

Animal Care

Breeding and husbandry of all C57BL/6NTac mice (female, n = 10 per group) was done in the Medical Experimental Center of the University of Leipzig (Leipzig, Germany). All mice were maintained in a room under pathogen-free conditions with a controlled temperature (21 $\pm$ 1 °C) on a 12/12 h light/dark cycle. Water and chow were provided ad libitum. The local ethics committee (Regierungspräsidium Leipzig) of the state of Saxony (Germany) approved the protocol of the animal experiments (approval No. TVV04/12). TR α - and TR β -deficient mice (male, 15 weeks old) were obtained from the European Mutant Mouse Archive [17] and the local ethics committee approved the study (Landesamt für Umwelt, Natur und Verbraucherschutz Nordrhein-Westfalen Az. 84-02.04.2014.A092).

Animal Treatment

For the induction of thyrotoxicosis, mice received L-thyroxine at a dose of 2 μ g/ml diluted in drinking water for 4 weeks. Hypothyroidism was induced in 10-week-old mice by feeding iodine-free chow supplemented with 0.15% PTU (catalog TD 97061; Harlan Teklan, Madison, Wis., USA) for 4 weeks. Euthyroid mice fed a normal chow diet (Altromin GmbH, Lage, Germany) served as controls.

T₄ Serum Measurements

Total T₄ concentrations in serum were determined by radioimmunoassay using commercially available kits (RIA-4524; DRG Instruments GmbH, Marburg, Germany). The samples and calibrators were incubated in duplicate with radiolabeled tracer in antibody-coated tubes according to the manufacturer's instructions. After incubation the liquid was aspirated and the antibody-bound radiolabeled tracer was counted in a gamma counter (1277 Gammamaster; LKB Wallac, Turku, Finland). The limit of quantification was 10 nM with an intra-assay coefficient of variation of 5.5% at 36 nM, 5.9% at 81 nM and 2.2% at 169 nM.

Table 1. Primer sequences used for qPCR

Gene	Forward primer (5'→3')	Reverse primer (3'→5')
<i>Acot8</i>	GGTCTGGGAGATGCTCATGG	CCCAGTAATGCCTTCCTCTGT
<i>Acot11</i>	CAGAATGTGGGCAACCACTTG	GCATGCCTCTCCGCTGAT
<i>Acot13</i>	GAGTTTTGGAAAAGGTGACGCT	TGGAGCGTGCCCAGTTTATT
<i>Acox1</i>	TCGCAGACCCTGAAGAAATC	CCTGATTACAGCAAGGTAGGG
<i>Acc1</i>	TACAGGATGGTTTGGCCTTT	CAAATTCTGCTGGAGAAGCC
<i>Cpt1a</i>	TGGATGGCTATGGTCAAGGT	TCTCCCTCCTTCATCAGTGG
<i>Cpt2</i>	TGGCTGAGTGCTCCAAATACC	GCCAGATACCGTAGAGCAAACA
<i>Scd1</i>	GAGGCCTGTACGGGATCATA	CAGCCGAGCCTTGTAAGTTC
<i>Fasn</i>	AGATCCTGGAACGAGAACACGAT	GAGACGTGTCACTCCTGGACTTG

RNA Extraction and RT-PCR

RNA was isolated from snap-frozen livers and adipose tissue using Trizol reagent (Invitrogen, Carlsbad, Calif., USA). In total, 1 µg of RNA per sample was reverse transcribed in a final mixture of 5× first-strand buffer (250 mM Tris-HCl, pH 8.3, 375 mM KCl, 15 mM MgCl₂), 0.5 mM of dNTPs, 5 mM of DTT (Promega, Madison, Wis., USA), 15 U of Prime RNase Inhibitor (ThermoFisher, Waltham, Mass., USA), 0.5 µg of random hexamer primers and 200 U of Moloney murine leukemia virus reverse transcriptase (Promega). Reverse transcription was performed at 37°C for 60 min and 94°C for 5 min.

Quantitative Real-Time PCR

Exon/intron spanning primers were designed to quantitatively calculate the expression of the *Acot8*, *11* and *13* isoforms in mouse liver and adipose tissue. The primer sequences are presented in table 1. Gene expression was analyzed in duplicate by reverse-transcriptase real-time PCR by using the commercial LightCycler-DNA Master SYBR Green I Kit (Roche, Mannheim, Germany) as previously described [18]. Gene expression was analyzed according to the $\Delta\Delta C_t$ method normalized to housekeeping genes *36B4* (adipose tissue) or *18s* (liver), and expression levels were calculated as $2^{-\Delta\Delta C_t}$ [19]. Data are expressed as the mean $\pm$ SEM. All statistical analyses and final presentations were performed using GraphPad Prism 6.0 software (GraphPad, San Diego, Calif., USA). One-way ANOVA followed by the Bonferroni's post hoc test was used for multiple comparisons between groups. *p* values ≤ 0.05 were considered to be significant.

Histology and Oil Red O Staining

For oil red O staining, livers were shock frozen in liquid nitrogen, embedded in OCT cryostat sectioning medium (Fisher Scientific, Schwerte, Germany) and cut at 8 µm. Sections were subjected to oil red O staining as described and counterstained with hematoxylin [20].

Results

Characterization of Thyrotoxic and Hypothyroid Mice

Thyroid dysfunction was established as described in Materials and Methods. Hypothyroid female mice had

significantly reduced T_4 concentrations (14.00 ± 1.2 nM) compared to thyrotoxic (41.33 ± 3.67 nM, $p < 0.01$) and euthyroid mice (37.5 ± 0.5 nM, $p < 0.001$; fig. 1a).

In liver, the expression of genes known to be TH responsive, *Dio1* and *Pdk4*, were significantly downregulated in hypothyroid mice and upregulated in thyrotoxic mice compared to euthyroid controls (fig. 1b). Thyroxine-binding globulin, *Tbg*, was studied as an example of a hepatic gene that is negatively regulated by TH. There was a 12-fold increase in *Tbg* mRNA expression in hypothyroid mice (fig. 1b). Altogether, in addition to the T_4 serum concentration, the gene expression analysis of TH-responsive genes further confirms the thyrotoxic and hypothyroid phenotype of the T_4 - and PTU-treated mouse cohorts.

Distinct Tissue-Specific *Acot* Gene Expression in Mice with Thyroid Dysfunction

To address the impact of thyroid dysfunction on *Acot* gene expression in major tissues relevant for lipid metabolism, we first investigated *Acot* gene expression in epididymal (eWAT) and inguinal (iWAT) white adipose tissue and interscapular BAT in thyrotoxic, hypo- and euthyroid mice. Interestingly, we found both (i) fat depot-specific differences in the expression of *Acot* and (ii) a switch of *Acot* expression associated with thyroid status (fig. 2). The *Acot11* gene reached the significantly highest values in white fat depots, eWAT and inguinal iWAT of thyrotoxic mice (fig. 2a). There was a 6-fold increase in eWAT in these mice when compared to eWATs of hypothyroid and euthyroid mice ($p < 0.001$; fig. 2a). The same regulation was found for *Acot13*, with predominant expression in eWAT of thyrotoxic mice with a 5-fold higher expression compared to eWATs of hypothyroid and euthyroid mice ($p < 0.001$, respectively; fig. 2b). Similarly, *Acot8* was overexpressed in the eWAT of thyrotoxic mice. However, in a hypothyroid state *Acot8* expression was highest in the

Fig. 1. Characterization of thyrotoxic and hypothyroid mice. Serum T₄-concentrations in thyrotoxic, hypothyroid and euthyroid female mice (a) and mRNA expression of TH responsive genes in liver (b) confirm the thyrotoxic and hypothyroid phenotype of

the T₄- and PTU-treated mouse cohorts. Statistical analysis was performed by one-way ANOVA followed by Bonferroni's post hoc test. Data are presented as means ± SEM. * p < 0.05, ** p < 0.01, *** p < 0.001.

Fig. 2. Effect of altered TH status on *Acot* gene expression in adipose tissue. The mRNA expression of *Acot11* (a), *Acot13* (b) and *Acot8* (c) genes was determined in WAT (eWAT, iWAT) and BAT of thyrotoxic, hypothyroid and euthyroid female mice. Statistical analysis was performed by one-way ANOVA followed by Bonferroni's post hoc test. Data are presented as means ± SEM. * p < 0.05, ** p < 0.01, *** p < 0.001.

Fig. 3. Effect of altered TH status on liver lipid metabolism. **a** Quantitative RT-PCR of genes involved in FA oxidation and lipogenesis. Euthyroid values were defined as 1 and changes in thyrotoxic and hypothyroid animals were expressed as relative amounts compared with euthyroid controls (n = 5 per group). **b** Representative oil red O stainings and hematoxylin stainings in

euthyroid, thyrotoxic and hypothyroid mice (n = 5 per group). **c** Gene expression of *Acot8*, *11* and *13* analyzed in livers of thyrotoxic, hypothyroid and euthyroid mice. Statistical analysis was performed by one-way ANOVA followed by Bonferroni's post hoc test. Data are presented as means $\pm$ SEM. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

BAT, which was 3-fold higher compared to *Acot8* expression in the BAT of thyrotoxic and euthyroid mice ($p < 0.01$; fig. 2c). Collectively, the data suggest that gene expression of *Acot8*, *11* and *13* are stimulated by a thyrotoxic state predominantly in eWAT, whereas low TH concentrations elevate expression of the *Acot8* isoform in BAT.

Increased Hepatic *Acot* Gene Expression in Thyrotoxic Mice

In order to elucidate a potential role of TH-induced *Acot* expression on hepatic lipid metabolism, we first examined the mRNA expression of genes associated with lipid metabolism in livers of thyrotoxic or hypo- and euthyroid female mice (fig. 3). In the livers of hypothyroid mice, expression of genes involved in FA oxidation as well as lipogenic genes is decreased ($p < 0.001$ for *Cpt1α* and

$p < 0.05$ for *Scd1*; fig. 3a). Although these genes encode for proteins which promote or reduce hepatic triglyceride content, no triglyceride accumulation in the liver was observed in the thyrotoxic, hypo- or euthyroid mice (fig. 3b). Similar to the observed increase in transcript concentrations of *Acot* in eWAT in a thyrotoxic state, in livers of thyrotoxic mice a trend towards collective upregulation of all analyzed *Acot* transcripts was evident when compared to the livers of hypothyroid and euthyroid mice (fig. 3c). Collectively, these data suggest the induction of hepatic *Acot* gene expression by TH.

Hepatic *Acot* Expression in *TRα* and *TRβ* Knockout Mice

In liver, T_3 exerts its effects on lipid and carbohydrate metabolism by binding to its cognate nuclear receptors

TR α and TR β . Prolonged treatment with T₃ promotes FA catabolism [6]. Having shown that expression of ACOTs is coordinately upregulated by TH in livers in vivo, we next aimed to assess whether this effect is mediated by TR. We used male TR α and TR β knockout (KO) mice and compared the hepatic *Acot* gene expression to that in wild-type mice. As shown in figure 4, deficiency of TR α had no effect on hepatic *Acot* expression. However, in livers of TR β -deficient mice there was a significant upregulation of *Acot8* compared to the wild-type mice (1.5-fold, $p < 0.05$). This effect was even stronger for *Acot11* with a 5-fold increase in TR β -deficient mice versus wild-type controls ($p < 0.05$; fig. 4). The highly elevated serum and liver TH concentrations in TR β -deficient mice [8, 21] possibly increase the hepatic expression of *Acot8* and *11* via compensation by TR α . It therefore appears as if both TRs can mediate *Acot* expression in liver, at least for *Acot8* and *11*.

Discussion

TH influence a wide variety of physiological processes, including growth and differentiation as well as thermogenesis. To date little is known about the molecular mechanisms by which TH orchestrates energy storage and expenditure within the cell. Lipogenesis as well as lipolysis are regulated by T₃, and it is well known that T₃ stimulates the shuttling of FFA into mitochondria [22]. In recent years, a number of ACOTs have been identified with distinct roles in lipid metabolism. It has been demonstrated that *Acot11* is upregulated in BAT when mice are exposed to cold and it has been proposed that *Acot11* supports the transition of this tissue towards increased metabolic activity, most likely through alteration of intracellular fatty acyl-CoA concentration [23]. In our present work we found a significant upregulation of *Acot11* in eWAT from thyrotoxic mice but not in BAT. Depot-specific differences in adipose tissue lipolysis are well known, for instance in the context of FFA release from the adipocyte. As an example, visceral (omental) cultured differentiated preadipocytes display a greater FFA flux than subcutaneous adipocytes, suggesting the contribution of a partially cell-autonomous mechanism [24]. Whether or not visceral *Acot11* expression in thyrotoxic mice represents such a regulatory point in visceral FA metabolism by balancing lipid storage and FA oxidation needs to be addressed in further functional studies.

A second finding of the present work was that the lack of TH triggers *Acot8* expression in BAT. *Acot8* is a per-

Fig. 4. Effect of TR KO on hepatic *Acot* expression. Gene expression of *Acot8*, *11* and *13* analyzed in livers of TR α or TR β KO mice and wild-types under euthyroid conditions. Wild-type values were defined as 1 and changes in TR α and TR β KO mice were expressed as relative amounts compared with the euthyroid wild-type controls ($n = 5$ per group). Statistical analysis was performed by one-way ANOVA followed by Bonferroni's post hoc test. Data are presented as means $\pm$ SEM. * $p < 0.05$.

oxisomal ACOT whose activity is strongly inhibited by CoASH, suggesting a role of *Acot8* in the regulation of peroxisomal CoA/CoASH concentrations. Peroxisomes optimize the flux of FA through β -oxidation and also participate in the synthesis of bile acids [14, 25]. It is worth noting that bile acids stimulate deiodinase 2 activity in BAT, thereby promoting energy expenditure and preventing resistance to insulin and obesity [26]. However, the contribution of *Acot8* in the circuit of bile acids/cAMP/deiodinase 2 in BAT is thus far elusive. Additionally, it has been demonstrated that in BAT *Dio2* may regulate the local availability of T₃. This results not only in an increase of the intracellular but also the extracellular T₃ pool [27–30]. In fact, 30–40% of T₃ is produced by PTU-insensitive pathways in adult athyroid T₄-supplemented [29], cold-exposed [28], preweaning [31] or hypothyroid rats [29, 30]. Since *Dio2* is insensitive to PTU [29], it is likely that the compensatory local BAT T₃ concentration contributes to enhanced *Acot8* gene expression in hypothyroid BAT.

TH regulation of lipid metabolism in liver is primarily dependent on direct actions of T₃ on TR as well as indirect crosstalks with nutrient-activated nuclear receptors, such as PPAR α and LXR. LXR and TR β control the expression of key enzymes involved in FA oxidation as well as lipogenesis, such as *Cpt1 α* , *Fasn* and *Acc1* [1]. In the livers of our hypothyroid mice these enzymes were

decreased, but were upregulated in thyrotoxic mice (fig. 2a). There were no differences in hepatic lipid stores between thyrotoxic and hypothyroid animals, as indicated by oil red O staining (fig. 2b). However, expression of *Acot8*, *11* and *13* were significantly higher in thyrotoxic compared to livers of hypothyroid mice (fig. 2c). In view of the upregulation of genes involved in β -oxidation, the overexpression of ACOT might represent a regulatory trait in hepatic FA metabolism by limiting the access of acyl-CoA thioesters to FA oxidation. Genetic inactivation of TR α does not alter hepatic *Acot* expression compared to wild-type mice. However, elevated transcript concentrations of both *Acot8* and *11*, but not *Acot13*, in TR β KO mice compared to wild-type livers suggest a TR β -independent stimulation of gene expression. This might be related to nonclassical effects of TH known to be markedly elevated in the serum and livers of TH-resistant TR β mice [8, 21]. Alternatively, both hepatic *Acot8* and *11*, but not *Acot13*, gene expression might be regulated indirectly by yet unidentified hepatic and/or systemic factors, hormones or metabolites altered in TR β -deficient TH-resistant mice. For instance, in mice with TR mutations crosstalks between TH and PPAR α [32] as well as LXR [33] have been demonstrated.

In summary, our study provides the first evidence that ACOTs are targets of TH and that a disturbed TH status is associated with alterations in *Acot* gene expression in the liver and both white and brown fat depots in mice. In the liver the changes in *Acot* expression are not primarily dependent on the direct actions of TH on TR. However, our observations may add to the knowledge of TH-regulated hepatic lipid metabolism. Finally, the finding of increased expression of *Acot8* in the BAT of hypothyroid mice might provide further insight into the oxidative metabolism in BAT under TH-deprived conditions.

Acknowledgement

This work was supported by grants from the German Research Council (DFG Priority Programme 1629 and SFB 1052 'obesity mechanisms' C01, C07, B4), from the German Diabetes Association and from the DHFD (Diabetes Hilfs- und Forschungsfonds Deutschland).

Disclosure Statement

The authors have nothing to disclose.

References

- 1 Brent GA: Mechanisms of thyroid hormone action. *J Clin Invest* 2012;122:3035–3043.
- 2 McAninch EA, Bianco AC: Thyroid hormone signaling in energy homeostasis and energy metabolism. *Ann NY Acad Sci* 2014;1311:77–87.
- 3 Iwen KA, Schröder E, Brabant G: Thyroid hormones and the metabolic syndrome. *Eur Thyroid J* 2013;2:83–92.
- 4 Jourdan C, Linseisen J, Meisinger C, Petersen A-K, Gieger C, Rawal R, et al: Associations between thyroid hormones and serum metabolite profiles in an euthyroid population. *Metabolomics* 2014;10:152–164.
- 5 Vergani L: Lipid lowering effects of iodothyronines: in vivo and in vitro studies on rat liver. *World J Hepatol* 2014;6:169–177.
- 6 Oppenheimer JH, Schwartz HL, Lane JT, Thompson MP: Functional relationship of thyroid hormone-induced lipogenesis, lipolysis, and thermogenesis in the rat. *J Clin Invest* 1991;87:125–132.
- 7 Coppola M, Glinni D, Moreno M, Cioffi F, Silvestri E, Goglia F: Thyroid hormone analogues and derivatives: actions in fatty liver. *World J Hepatol* 2014;6:114–129.
- 8 Flores-Morales A, Gullberg H, Fernandez L, Ståhlberg N, Lee NH, Vennström B, et al: Patterns of liver gene expression governed by TR β . *Mol Endocrinol* 2002;16:1257–1268.
- 9 Jackson-Hayes L, Song S, Lavrentyev EN, Jansen MS, Hillgartner FB, Tian L, et al: A thyroid hormone response unit formed between the promoter and first intron of the carnitine palmitoyltransferase-1 α gene mediates the liver-specific induction by thyroid hormone. *J Biol Chem* 2003;278:7964–7972.
- 10 Watkins PA, Ellis JM: Peroxisomal acyl-CoA synthetases. *Biochim Biophys Acta* 2012;1822:1411–1420.
- 11 Watkins PA: Fatty acid activation. *Prog Lipid Res* 1997;36:55–83.
- 12 Cohen DE: New players on the metabolic stage. *Adipocyte* 2013;2:3–6.
- 13 Li LO, Klett EL, Coleman RA: Acyl-CoA synthesis, lipid metabolism and lipotoxicity. *Biochim Biophys Acta* 2010;1801:246–251.
- 14 Hunt MC, Alexson SEH: The role Acyl-CoA thioesterases play in mediating intracellular lipid metabolism. *Prog Lipid Res* 2002;41:99–130.
- 15 Momose A, Fujita M, Ohtomo T, Umemoto N, Tanonaka K, Toyoda H, et al: Regulated expression of acyl-CoA thioesterases in the differentiation of cultured rat brown adipocytes. *Biochem Biophys Res Commun* 2011;404:74–78.
- 16 Hunt MC, Solaas K, Kase BF, Alexson SEH: Characterization of an Acyl-CoA thioesterase that functions as a major regulator of peroxisomal lipid metabolism. *J Biol Chem* 2002;277:1128–1138.
- 17 Gauthier K, Chassande O, Plateroti M, Roux JP, Legrand C, Pain B, et al: Different functions for the thyroid hormone receptors TR α and TR β in the control of thyroid hormone production and post-natal development. *EMBO J* 1999;18:623–631.
- 18 Krause K, Eszlinger M, Gimm O, Karger S, Engelhardt C, Dralle H, et al: TFF3-based candidate gene discrimination of benign and malignant thyroid tumors in a region with borderline iodine deficiency. *J Clin Endocrinol Metab* 2008;93:1390–1393.
- 19 Livak KJ, Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta C_T}$ method. *Methods* 2001;25:402–408.

- 20 Ruschke K, Illes M, Kern M, Klötting I, Fasshauer M, Schön MR, et al: Repin1 maybe involved in the regulation of cell size and glucose transport in adipocytes. *Biochem Biophys Res Commun* 2010;400:246–251.
- 21 Flamant F, Samarut J: Thyroid hormone receptors: lessons from knockout and knock-in mutant mice. *Trends Endocrinol Metab* 2003;14:85–90.
- 22 Elks ML, Manganiello VC: Effects of thyroid hormone on regulation of lipolysis and adenosine 3',5'-monophosphate metabolism in 3T3-L1 adipocytes. *Endocrinology* 1985;117:947–953.
- 23 Adams SH, Chui C, Schilbach SL, Yu XX, Goddard AD, Grimaldi JC, et al: BFIT, a unique acyl-CoA thioesterase induced in thermogenic brown adipose tissue: cloning, organization of the human gene and assessment of a potential link to obesity. *Biochem J* 2001;360:135–142.
- 24 Caserta F, Tchkonja T, Civelek VN, Prentki M, Brown NF, McGarry JD, et al: Fat depot origin affects fatty acid handling in cultured rat and human preadipocytes. *Am J Physiol Endocrinol Metab* 2001;280:E238–E247.
- 25 Wanders RJA, Waterham HR: Peroxisomal disorders: the single peroxisomal enzyme deficiencies. *Biochim Biophys Acta* 2006;1763:1707–1720.
- 26 Watanabe M, Houten SM, Matak C, Christoffolete MA, Kim BW, Sato H, et al: Bile acids induce energy expenditure by promoting intracellular thyroid hormone activation. *Nature* 2006;439:484–489.
- 27 Silva JE, Gordon MB, Crantz FR, Leonard JL, Larsen PR: Qualitative and quantitative differences in the pathways of extrathyroidal triiodothyronine generation between euthyroid and hypothyroid rats. *J Clin Invest* 1984;73:898–907.
- 28 Silva JE, Larsen PR: Potential of brown adipose tissue type II thyroxine 5'-deiodinase as a local and systemic source of triiodothyronine in rats. *J Clin Invest* 1985;76:2296–2305.
- 29 Silva JE, Leonard JL, Crantz FR, Larsen PR: Evidence for two tissue-specific pathways for in vivo thyroxine 5'-deiodination in the rat. *J Clin Invest* 1982;69:1176–1184.
- 30 Christoffolete MA, Linardi CCG, Jesus L de, Ebina KN, Carvalho SD, Ribeiro MO, et al: Mice with targeted disruption of the Dio2 gene have cold-induced overexpression of the Uncoupling Protein 1 gene but fail to increase brown adipose tissue lipogenesis and adaptive thermogenesis. *Diabetes* 2004;53:577–584.
- 31 Silva JE, Matthews P: Thyroid hormone metabolism and the source of plasma triiodothyronine in 2-week-old rats: effects of thyroid status. *Endocrinology* 1984;114:2394–2405.
- 32 Liu Y-Y, Heymann RS, Moatamed F, Schultz JJ, Sobel D, Brent GA: A mutant thyroid hormone receptor alpha antagonizes peroxisome proliferator-activated receptor alpha signaling in vivo and impairs fatty acid oxidation. *Endocrinology* 2007;148:1206–1217.
- 33 Hashimoto K, Cohen RN, Yamada M, Markan KR, Monden T, Satoh T, et al: Cross-talk between thyroid hormone receptor and liver X receptor regulatory pathways is revealed in a thyroid hormone resistance mouse model. *J Biol Chem* 2006;281:295–302.