'Is Going through Clinical Test a Headache?' An HRV Study and Descriptive Report of Subjective Experience of Undergoing EEG Testing

Rajesh Kathrotia Yogesh Singh Arun Goel Prashant Patil

Department of Physiology, All India Institute of Medical Sciences (AIIMS), Rishikesh, India

Key Words

EEG electrode placement · 10–20 system · Subjective report · Mental stress and fatigue · Massage therapy · Heart rate variability

Abstract

To explore the heart rate variability (HRV) changes and subjective perception of undergoing electroencephalography (EEG). We conducted a study on 35 healthy male volunteers. The intervention consisted of placing 23 disc-type EEG electrodes of 5–7 mm diameter with long flexible lead according to international 10–20 system for the duration of 30 min, in a sitting position, on the scalp. The outcome measures were time and frequency domain parameters of HRV analysis and descriptive report of subjective experiences on a 3-point Likert scale. The perception of undergoing EEG ranged from pleasant to uneasy. For 13 (37%) participants it was soothing and relaxing, for 11 (31.5%) it was neutral and for the rest 11 (31.5%) it was uneasy and restrictive in nature. However, HRV analysis of the pre and post EEG, showed no statistically significant difference. In our study, the mixed subjective experience of undergoing EEG may be due to individual variation in the perception of the intervention. No difference in HRV parameters may be because of 2 possibilities. The first possibility is varied experiences of procedure with temporal progression. Same participants may have experienced 2 oppo-

site extremes of experiences over and over again, which may have cancelled out sympathetic and parasympathetic responses. The second possibility may be that no stress is generated during clinical test.

© 2016 S. Karger AG, Basel

Introduction

In the past few decades, there has been a rapid escalation in use of diagnostic and clinical testing facilities for various diseases and health checkups [1]. Clinical tests may vary from a simple venesection to prolonged procedures like electroencephalography (EEG). This prolonged duration during investigations may produce additional stress over and above the stress of disease.

EEG is routinely being used for epilepsy, seizures [2] and cognitive neuroscience research. The EEG requires placement of many disc electrodes with long flexible lead on the scalp. The prerequisites for good signal acquisition include moderate rubbing, debridement of scalp with scrubbing agent and cleaning with alcohol swab at each site. The procedure is routinely done in a sitting posture and preparation for testing protocol usually takes 30–60 min. Alternatively, the cap system of EEG signal acquisition can be used but it has its own limitations. The common notion is that EEG preparation and procedure is ex-

haustive and tiring but the possibility that it may be perceived as soothing massage needs to be explored.

Recently, massage therapy has emerged as one of the alternative treatments [3] and is often considered as a technique of relaxation [4]. Studies involving processes like gentle stretching and moderate cutaneous pressure have shown to decrease tension and create a state of tranquility [5].

The stress and relaxation responses have been associated with discrete changes in autonomic nervous system (ANS), which can be measured by heart rate variability (HRV) [6, 7]. Measuring HRV is a simple, sensitive and non-invasive technique based on beat-to-beat alteration in heart rate [8]. HRV is supposed to be an indicator of the level of stimulation of ANS and dynamic interaction and balance between the sympathetic nervous system and the parasympathetic nervous system [6, 9–11].

HRV has been used to measure changes in sympathovagal activity during massage therapy [12], emotional states (such as stress and anxiety) and pain studies [6, 13–16]. The findings from meta-analysis of several studies have established a link between emotions, such as anxiety and hostility, and a reduced HRV [8]. Therefore, exploring the emotional variables that may affect HRV may be beneficial to the heart and health research [17].

We assessed subjective perception and HRV changes while undergoing EEG. To the best of our knowledge, no such prior study is available in literature.

Methods

Study design: prospective cohort study.

Study Population

We conducted a study on 35 healthy male volunteers who were randomly selected and recruited from an ongoing research study in a cognitive neuroscience laboratory. Only male participants were included because women tend to have different perceptions of relaxation responses, mental stress and stress reactivity during the different phases of menstrual cycle [18]. Participants with history of cardiac diseases, high blood pressure, diabetes or prior treatment for any medical condition or those indulging in substance abuse, smoking and alcohol were excluded.

We briefed the participants about the study through a subject information sheet after obtaining written informed consent. Participants were requested to report in the cognitive neuroscience lab at 10 a.m., to not eat for 4 h prior to the experiment, refrain from undue body movements, talking and falling asleep and specifically instructed not to voluntarily change their breathing pattern during electrocardiogram (ECG).

Ethical clearance for the study was taken from the institutional ethics committee for the human subjects.

Study Protocol

EEG Electrodes Placement Protocol

The participants were instructed to clean their hair with shampoo without the application of conditioner or oil 1 day prior to the testing. A total of 23 silver-silver chloride (Ag-AgCl) disc-type electrodes of 5–7 mm diameter with long flexible leads (21 scalp electrode and 2 ear electrodes) were placed on the scalp according to the 10–20 international electrode placement system with the help of ‘Elefix’ paste for EEG (supplied by Nihon Kohden). An electrode impedance of less than 5 k Ω at each position was achieved by cleaning scalp with alcohol swab followed by rubbing with skin preparation gel ‘SKINPURE’ (supplied by Nihon Kohden). The whole procedure was performed in a sitting posture and lasted 30 min. EEG electrodes of RMS EEG-19 Superspec system were used.

ECG Signal Acquisition and HRV Analysis

ECG signals were acquired using the lead-2 at sample rate of 200/s and gain X 2,000 from power lab data recording system (ADInstruments, Australia). HRV analysis was based on a 5-minute period of ECG signal using LabChart software (ADInstruments, Australia). The heart rates were calculated using the R-waves from artefact-free graph for both stages of recordings. Time domain and frequency domain analysis of HRV of both sessions was done offline.

The parameters used were set as follows: total power frequency (TPF) (0.00–0.40 Hz), low frequency (LF) power (0.04–0.15 Hz) and high frequency (HF) power (0.15–0.40 Hz). The LF/HF ratio was calculated based on the outcome of power spectral analysis.

The following HRV parameters were analyzed.

Time domain parameters:

- Average heart rate (bpm)
- SD heart rate (bpm)
- SDS

- RMSSD
- pRR50%
- Frequency domain parameters:
- LF (nu)
- HF (nu)
- LF/HF ratio

Statistical Analysis

The statistical analysis was done using the SPSS software version 20. The distribution of the data was found to be non-Gaussian type; therefore, non-parametric test (Wilcoxon signed-rank test) was applied. HRV comparison of pre- and post-EEG electrode placement was done, and p values <0.05 were considered significant.

Results

The mean age of the participants in the study was 30.15 years. The detailed anthropometric parameters of the participants are given in the table 1.

Self-Assessment Description

Out of the 35 participants, 13 (37%) participants perceived the experience of EEG electrode placement as relaxing. For 11 (31.5%) participants, the experience was neutral. The remaining 11 (31.5%) participants complained of uneasiness.

The participants who perceived the procedure as a gentle massage, soothing in nature and conducive for induction of sleep reported the experience as relaxing in the 3-point Likert scale. The reason for uneasiness was discomfort, tiredness and a feeling of constrain that was due to prolonged sitting in the same posture. The mixed feeling was reported as neutral.

HRV Results

The comparison of before and after 5 min of each fast Fourier transform applied epoch of ECG showed no significant differences in any of the time and frequency domain parameters of the HRV.

The results of subjective perception and HRV analysis are summarized in figure 1 and table 2.

Discussion

The present study was carried to find out the HRV changes and subjective perceptions of undergoing EEG. The subjective experiences of the procedure ranged from relaxing to uneasy. These differences may be because of varied temperament, mood and personality types of the volunteers. Stress response is dependent on the environ-

Table 1. Anthropometric parameters of the study participants (n = 35)

Parameters	Mean $\pm$ SD
Age, years	30.15 $\pm$ 6.30
Height, cm	166 $\pm$ 25.32
Weight, kg	62.64 $\pm$ 9.01
BMI, kg/m ²	22.69 $\pm$ 3.19

BMI = Body mass index.

Table 2. Parameters during pre- and post-electrode placement

	HRV frequency		
	pre	post	p value
LF (nu)	58.90 $\pm$ 19.80	62.82 $\pm$ 19.28	0.235
HF (nu)	40.51 $\pm$ 19.16	36.25 $\pm$ 18.21	0.172
LF/HF	2.49 $\pm$ 2.98	2.68 $\pm$ 2.19	0.140
	HRV time domain		
	pre	post	p value
Average heart rate, bpm	81.08 $\pm$ 10.84	79.82 $\pm$ 9.76	0.32
SD heart rate, bpm	5.01 $\pm$ 1.83	4.46 $\pm$ 1.64	0.067
SDSD	34.21 $\pm$ 23.96	31.81 $\pm$ 25.59	0.067
RMSSD	34.17 $\pm$ 23.93	31.77 $\pm$ 25.55	0.07
pRR50%	13.38 $\pm$ 17.21	11.11 $\pm$ 16.89	0.08

All values are mean $\pm$ SD.

Fig. 1. Descriptive self-report of EEG electrode placement experience (n = 35).

mental and personal characteristics of the individual [19]. Another possibility is the mobilization of different cognitive faculties as the key players in stress appraisal response. The stress response depends not only on the specific type of stressor but it commences after the cognitive appraisal of stimulus as a threat [20].

Scientific studies to assess the effects of massage on the ANS are limited, yet evidences suggest varied findings. An increase in both parasympathetic [21] and sympathetic [22] activities have been reported with massages. Subjective relaxation is a common finding with different types of massage although a physiological relaxation response depends on the type, duration or site of massage. These varied results may be dependent on increased or decreased arousal as indicated by heart rate and EEG activity depending on the type (light, moderate or vibratory) of massage [23].

In the present study, we found no significant differences in time and frequency parameters of HRV. Although HRV parameters are useful and it is a sensitive tool to assess ANS activity during various emotional states, anxiety, therapeutic touch, massage [12] and pain [6, 13–16], the lack of clear cut patterns in HRV may be because of varied reasons. The first possibility is the differences in individual perception of the intervention. The second possibility is varied experiences with temporal progression, with the same participant having experienced 2 opposite extremes of experiences over and over again, which may have cancelled out sympathetic and parasympathetic responses. The procedure of EEG electrodes placement may have evoked relaxation response due to parasympathetic activation resulting from gentle soothing massage and touch or stress response resulting in sympathetic activation due to fatigue, exhaustion and restriction or mixed typed response. Third is that the intervention was gentle and hence failed to activate any physiological autonomic response. Though this is quite unlikely, yet it cannot be ruled out. Another alternative explanation may be that the effect was too small to be measured or the instrumentation system not sensitive enough to detect the effects.

The scenario may also point at the effect being masked by the interplay of unknown variables or the effect of responders in the study population being masked by opposite effects from non-responders. The variations in response of individual may have first evoked sympathetic and later on parasympathetic responses or vice versa and a more complex cyclical response with the temporal progression of the procedure over a period.

A variety of variables ranging from physical to mental have been reported in literature to significantly alter HRV and they can be quantified by average LF/HF ratio. It may

range from controlled breathing at 15 breaths per minute to performing mental tasks [24, 25]. Therefore, during the study, all confounding factors like physical, mental and others that may affect HRV should be taken into consideration through randomization [25]. Still stringently randomized controlled studies may get confounded by participant's mental activity with the temporal course of the study and mental activity is deemed as the factor having an utmost effect on the HRV. Controlling mental activity, however, is nearly impossible, with the possible exception of study during sleep [25], which was not feasible in our experimental settings.

In summary, undergoing EEG has varied experience and no significant change in HRV may be due to the mix heterogeneous autonomic response, different individual personality types (sympathetic type, parasympathetic type and non-responder) and complex mental states originating with temporal progression of task. However, the possibility may be that no stress is generated during clinical test.

The results of our study may be specific to our participants and their perceptions of intervention in our laboratory settings.

Authorship Contributions

R.K.: data collection, drafting of manuscript, statistical analysis; Y.S.: data collection, drafting, interpretation of result; A.G.: critical revision of draft and valuable input; P.P.: design of study.

Source of Funding

Intramural research grant, AIIMS, Rishikesh.

Disclosure Statement

None.

References

- 1 Krogstøll LT, Jørgensen KJ, Grønhoj Larsen C, Gøtzsche PC: General health checks in adults for reducing morbidity and mortality from disease: Cochrane systematic review and meta-analysis. *BMJ* 2012;345:e7191.
- 2 Noachtar S, Rémi J: The role of EEG in epilepsy: a critical review. *Epilepsy Behav* 2009; 15:22–33.
- 3 Eisenberg DM, Davis RB, Ettner SL, Appel S, Wilkey S, Van Rompay M, et al: Trends in alternative medicine use in the United States, 1990–1997: results of a follow-up national survey. *JAMA* 1998;280:1569–1575.

- 4 Moyer CA, Rounds J, Hannum JW: A meta-analysis of massage therapy research. *Psychol Bull* 2004;130:3–18.
- 5 Tapanya S: Traditional Thai Massage. Bangkok, Duang Kamol, 1993.
- 6 Terathongkum S, Pickler RH: Relationships among heart rate variability, hypertension, and relaxation techniques. *J Vasc Nurs* 2004;22:78–82; quiz 83–84.
- 7 Benson H, Beary JF, Carol MP: The relaxation response. *Psychiatry* 1974;37:37–46.
- 8 MacArthur JD, MacArthur CT: Heart Rate Variability. 2000. <http://www.macses.ucsf.edu/Research/Allostatic/notebook/heart.rate.html> (retrieved May 30, 2015).
- 9 Berntson GG, Bigger JT Jr, Eckberg DL, Grossman P, Kaufmann PG, Malik M, et al: Heart rate variability: origins, methods, and interpretive caveats. *Psychophysiology* 1997;34:623–648.
- 10 Heart rate variability: standards of measurement, physiological interpretation and clinical use. Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. *Circulation* 1996;93:1043–1065.
- 11 Heart rate variability. Standards of measurement, physiological interpretation, and clinical use. Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. *Eur Heart J* 1996;17:354–381.
- 12 Buttagat V, Eungpinichpong W, Chatchawan U, Kharmwan S: The immediate effects of traditional Thai massage on heart rate variability and stress-related parameters in patients with back pain associated with myofascial trigger points. *J Bodyw Mov Ther* 2011;15:15–23.
- 13 Carney RM, Saunders RD, Freedland KE, Stein P, Rich MW, Jaffe AS: Association of depression with reduced heart rate variability in coronary artery disease. *Am J Cardiol* 1995;76:562–564.
- 14 Stauss HM: Heart rate variability. *Am J Physiol Regul Integr Comp Physiol* 2003;285:R927–R931.
- 15 Tousignant-Laflamme Y, Marchand S: Sex differences in cardiac and autonomic response to clinical and experimental pain in LBP patients. *Eur J Pain* 2006;10:603–614.
- 16 Hummel P, van Dijk M: Pain assessment: current status and challenges. *Semin Fetal Neonatal Med* 2006;11:237–245.
- 17 Tulppo M, Huikuri HV: Origin and significance of heart rate variability. *J Am Coll Cardiol* 2004;43:2278–2280.
- 18 Madhyastha S, Latha KS, Kamath A: Stress, coping and gender differences in third year medical students. *J Health Manag* 2014;16:315–326.
- 19 Hisli Şahin N, Güler M, Basim HN: [The relationship between cognitive intelligence, emotional intelligence, coping and stress symptoms in the context of type A personality pattern]. *Turk Psychiatry Derg* 2009;20:243–254.
- 20 Lazarus RS: From psychological stress to the emotions: a history of changing outlooks. *Annu Rev Psychol* 1993;44:1–21.
- 21 Hatayama T, Kitamura S, Tamura C, Nagano M, Ohnuki K: The facial massage reduced anxiety and negative mood status, and increased sympathetic nervous activity. *Biomed Res* 2008;29:317–320.
- 22 Diego MA, Field T, Sanders C, Hernandez-Reif M: Massage therapy of moderate and light pressure and vibrator effects on EEG and heart rate. *Int J Neurosci* 2004;114:31–44.
- 23 Bernardi L, Wdowczyk-Szulc J, Valenti C, Castoldi S, Passino C, Spadacini G, et al: Effects of controlled breathing, mental activity and mental stress with or without verbalization on heart rate variability. *J Am Coll Cardiol* 2000;35:1462–1469.
- 24 Clifford GD: *Signal Processing Methods for Heart Rate Variability*. Oxford, University of Oxford, 2002.
- 25 Clifford GD, Tarassenko L: Quantifying errors in spectral estimates of HRV due to beat replacement and resampling. *IEEE Trans Biomed Eng* 2005;52:630–638.