Effects of Fragment and Vegetation Structure on the Population Abundance of *Ateles hybridus*, *Alouatta seniculus* and *Cebus albifrons* in Magdalena Valley, Colombia

Christopher Marsh^a Andres Link^{c, d} Gillian King-Bailey^e
Giuseppe Donati^b

^aFaculty of Science and Technology, Bournemouth University, Poole, and ^bDepartment of Social Sciences, Oxford Brookes University, Oxford, UK; ^cDepartamento de Ciencias Biológicas and ^dFacultad de Administración, Universidad de Los Andes, Bogotá, Colombia; ^eDepartment of Anthropology, Tulane University, New Orleans, La., USA

Key Words

Endangered species · Fragmentation · Population abundance · Vegetation structure

Abstract

Many primate species currently subsist in fragmented and anthropogenically disturbed habitats. Different threats arise depending on the species' life history strategy, dietary requirements and habitat preference. Additionally, anthropogenic disturbance is far from uniform and may affect individual forest fragments in a single landscape in differing ways. We studied the effects of fragmentation on three species of diurnal primate, *Cebus albifrons*, *Alouatta seniculus* and *Ateles hybridus*, in Magdalena Valley, Colombia. We tested the assumption that generalist species are more resilient than specialist species to habitat degradation by examining the fragments' vegetation and spatial structure and how these affected primate presence and abundance patterns. We found *C. albifrons*, a generalist, to be the most abundant species in 9 of 10 forest fragments, regardless of the level of habitat disturbance. *A. hybridus*, a large-bodied primate with a specialist diet, was either absent or low in abundance in fragments that had experienced recent disturbances and was found only in higher-quality fragments, regardless of the fragment size. *A. seniculus*, a species considered to have a highly flexible diet and the ability to survive in degraded habitat, was found in intermediate abundances between those of *Cebus* spp. and *Ateles* spp., and was more frequently found in high-quality fragments.

© 2016 S. Karger AG, Basel

Introduction

Anthropogenic effects, including habitat loss and fragmentation, hunting and disease transfer, are leading causes of the pervasive loss of global biodiversity [Chapman and Peres, 2001; Michalski and Peres, 2005; Goldberg et al., 2008]. Primate species in particular are affected by habitat fragmentation [Arroyo-Rodríguez et al., 2013; Benchimol and Peres, 2014]. Despite this, continuous forest is rapidly being divided into small and isolated fragments in order to create farmlands and extract wood, minerals and oil from the land [Cowlshaw and Dunbar, 2000; Marsh and Chapman, 2013]. This creates a drastically different landscape in the habitat, and any organisms that cannot quickly adapt their behaviour to these emerging changes are ultimately faced with extinction [Chapman and Peres, 2001]. Despite numerous studies on the effects of habitat fragmentation and anthropogenic disturbance on primate populations [Marsh and Chapman, 2013], new findings frequently arise with each additional study [Benchimol and Peres, 2014]. The growing understanding of primates' flexibility within and across species necessitates further investigation of the effects of habitat fragmentation on individual primate species [Ewers and Didham, 2006].

While some primate species exhibit morphological adaptations and behavioural plasticity that allow them to continue to subsist in these habitats (howler monkeys, *Alouatta palliata* [Cristobal-Azkarate and Arroyo-Rodríguez, 2007]; collared lemur, *Eulemur collaris* [Donati et al., 2011]; diademed sifaka, *Propithecus diadema* [Irwin 2008a, b]; black and white colobus, *Colobus guereza* [Gillespie and Chapman, 2008]; moustached guenon, *Cercopithecus* [Tutin, 1999]), others appear to be more vulnerable due to greater specialization both morphologically and behaviourally (red colobus, *Procolobus rufomitratus* [Gillespie and Chapman, 2008]; bearded saki monkeys, *Chiropotes satanas chiropotes* [Boyle and Smith, 2010; Schwitzer et al., 2011]). Factors that enable some species to persist in fragmented forests include a small home range, broad-ranged diet and small group size [Purvis et al., 2000] (howler monkeys, genus *Alouatta* [Bicca-Marques, 2003]; mantled howler monkeys, *Alouatta palliata* [Estrada, 1999]; white-faced capuchins, *Cebus capucinus* [Panger et al., 2002]), whereas primate species with large home ranges, specialist diets and large group sizes can be more severely and immediately affected by forest fragmentation [Andrén, 1994] (southern gentle lemurs, *Hapalemur meridionalis* [Eppley et al., 2011]; brown spider monkeys, *Ateles hybridus* [Link et al., 2012]; spider monkeys, family Atelidae [Rimbach et al., 2012]). However, degradation can vary greatly, and fragment size and quality are important factors to consider [Arroyo-Rodríguez et al., 2007].

Since fragmentation is at the landscape level, studies focussing on single fragments, often compared with continuous forest, are unable to take into consideration the continuous, rather than discrete, variation that occurs between fragments within a single landscape [Arroyo-Rodríguez et al., 2013]. Fragment quality can vary greatly based on the amount of time that has passed since isolation, the shape and size of the fragment, and the degree to which it has been exposed to other anthropogenic disturbances [Fahrig, 1999; Benchimol and Peres, 2014]. The 'quality' of the fragment, as measured through vegetation structure, is predicted to impact its ability to sustain larger populations or even multiple primate species [Mbora and Meikle, 2004; Wiczowski, 2004]. Thus, by measuring population response to fragment geometry, vegetation structure and anthropogenic effects, we can predict species' responses in other habitats, aiding the development of targeted conservation efforts in the future.

In this study we assessed the effects of fragmentation on the abundance of three diurnal sympatric primates: brown spider monkeys (*A. hybridus*), red howler monkeys (*Alouatta seniculus*) and white-fronted capuchins (*Cebus albifrons*) in a fragmented landscape in Magdalena Valley, Colombia. *A. hybridus*, a relatively understudied primate, is considered to be one of the 25 most endangered primates, necessitating an immediate and better understanding of the species' response to habitat disturbance [Mittermeier et al., 2012]. We analysed the influence of fragment quality, fragment spatial structure and anthropogenic disturbance (selective logging) on the abundance of these three platyrrhines. Given that high-quality habitat has been empirically linked to high fruit production [Chapman et al., 1992], we hypothesized that omnivorous, generalist *C. albifrons* and folivorous *A. seniculus* would be more resilient to habitat degradation. In contrast, ripe-fruit specialist *A. hybridus* would require larger and less degraded fragments and be less abundant in low-quality and small fragments.

Methods

Study System

Our study took place near the Magdalena River, between the eastern and central cordilleras of Andean Colombia in and around the private cattle ranch 'Hacienda San Juan del Carare' (hereafter 'San Juan'; 06°43' N, 74°09' W). The area is seasonally flooded tropical rain forest, with a mean annual rainfall of 2,070 mm [Link et al., 2010], and the average temperature is around 28°C. The primates have been studied there since 2007. Most forest fragments are inaccessible during the wet season (October to December) due to flooding, making it most efficient to conduct censuses during the dry months, in which this study took place. The area is made up of a matrix of forest fragments within naturally occurring and cleared pasture for cattle. Selective logging took place in and around the fragments, though large trees (e.g. *Ficus insipida*) are still present [Link et al., 2010].

We investigated 10 forest fragments (F1, F2, F3, FA, FT, FN, FC, FTF, FP, FDU) ranging in size from 0.7 to 171 ha (mean = 51.9 ha; table 1). Each fragment exists in isolation from both continuous forest and other fragments. The edges are defined by naturally occurring savanna, pasture for cattle ranching, and the San Juan River itself. The study area supports three diurnal primate species (*A. hybridus*, *A. seniculus* and *C. albifrons*) and one nocturnal species (*Aotus griseimembra*) [Link et al., 2010].

The white-fronted capuchin, *C. albifrons*, is a medium-sized primate (mean = 3.2 kg, $n = 12$ [Russo and Young, 2011]) with highly adaptable behaviour. Untufted capuchin monkeys (*Cebus* spp.) engage in both arboreal and terrestrial foraging [Terborgh, 1983] and exploit an ample variety of food resources including ripe fruits, roots and flowers, insects, arthropods, eggs and mammalian prey, and engage in crop raiding in some areas [Buckley, 1983; Panger et al., 2002; Sabbatini et al., 2006]. These primates have medium-sized home ranges (median = 130 ha, $n = 4$ [Defler, 1979; Terborgh, 1983; Matthews, 2009]) and are found in groups ranging in size from 8 to 35 individuals (mean = 13.8, $n = 5$ [Defler, 1979; Terborgh, 1983]). *Cebus* spp. have relatively long interbirth intervals (interbirth interval unknown in *C. albifrons*; 2.25 years in wild *C. capucinus* [Fedigan et al., 2008]) and slow rates of maturation when controlling for body size [Fedigan and Rose, 1995]. This slow development is theorized to be part of a strategy to increase individual learning and ability to adjust to environmental variation [Janson, 1998]. *C. albifrons* is currently listed as 'Least Concern' by the IUCN [2014].

Red howler monkeys, *A. seniculus*, are large-bodied (mean = 6.75 kg [Peres, 1997]), slow-moving primates that use relatively small home ranges (median = 21 ha, $n = 12$ [Barbisan-Fortes et al., 2014]) and are characterized as energy minimizers [Strier, 1992]. *Alouatta* spp. have high levels of dietary flexibility, opportunistically eating fruit when it is available, with leaves serving as their primary food source when preferred foods are not available [Marshall et al., 2009]. They

Table 1. Vegetation structure results

Fragment ID	Ecological vegetation variables				Spatial structural variables			
	edge proportion	tree density, trees/ha	mean DBH, cm	canopy density, %	fragment size, ha	fragment isolation	C-to-C coefficient distance, m	BZ land type coefficient proportion
FP	432.84	2,007	9.9±4.2	70.9	0.67	673	1,464	1.11
F1	170.45	1,718	16.8±17.5	76.1	13.4	578	925	1.22
F2	138.92	1,749	12.6±8.3	72.8	16.6	588	925	1.21
FT	200.12	1,740	11.6±11.2	73.8	16.8	584	829	1.22
FDU*	74.62	2,941	10.7±8.6	68.4	39.8	984	815	1.41
F3	102.72	1,685	12.2±7.4	74.3	48.6	566	1,238	1.27
FA	114.97	1,805	13.4±8.5	71.8	58	606	1,238	1.43
FC*	76.57	2,230	9.0±4.7	74.0	70.9	747	1,897	1.33
FN	117.50	1,393	16.5±11.0	72.6	83.3	470	3,460	1.25
FTF*	39.60	1,043	9.1±3.5	75.3	171	351	1,553	1.40

Asterisks denote the presence of selective logging within fragments. C-to-C = Target fragment centroid to nearest fragment centroid; BZ = buffer zone; ± = standard deviation.

have shorter interbirth intervals (17 months [Crockett and Sekulic, 1982; Crockett and Rudran, 1987]) compared to *Cebus* spp. and other ateline primates, and exhibit faster maturation rates [Ross, 1989]. *A. seniculus* is currently classified as 'Least Concern' by the IUCN [2014].

Brown spider monkeys, *A. hybridus*, are large-bodied (mean = 8.9 kg [Di Fiore and Campbell, 2007]) energy maximizers [Strier, 1992]. They are fast-moving, wide-ranging, highly arboreal frugivores. Their high degree of fission-fusion sociality allows large groups to disperse widely in the pursuit of high-energy ripe fruit, whilst minimizing intragroup feeding competition [Symington, 1988, 2000; Asensio et al., 2008; Link et al., 2012]. Spider monkeys have long interbirth intervals (~36 months [Shimooka et al., 2008, Vick, 2008]). As a result of deforestation and fragmentation, populations have declined in a number of *Ateles* spp., and many populations are now limited to forest fragments and degraded habitats [Ramos-Fernández and Ayala-Orozco, 2002; Ramos-Fernández and Wallace, 2008; Rangel-Negrín et al., 2009; Link et al., 2010; Abondano and Link, 2012; Rimbach et al., 2012]. *A. hybridus* is currently classified as 'Critically Endangered' by the IUCN [Urbani et al., 2008].

Data Collection

Ecological Vegetation Variables

Ecological variables of each fragment were measured using the methods suggested by Ganzhorn et al. [2011]. The fragment size (hectares), total length of the perimeter of the fragment (metres), tree density (trees per hectare), the mean diameter at breast height (DBH; centimetres), and the canopy density (percent; table 1) were measured for each fragment. Fragment size and length of fragment perimeter were determined using Landsat 5 imagery and processed with GIS (ArcGIS 10.1). The edge proportion was calculated by dividing the fragment perimeter length (metres) by the size of the fragment (hectares). Tree density, DBH and canopy density were measured using the quarter-point plotless sampling method [Ganzhorn et al., 2011]. Sampling points were randomly selected within each fragment using a dice and grid system. At each starting point we took a digital photograph of the tree canopy from breast height for canopy density analysis. CanopyDigi software [Goodenough and Goodenough, 2012] was used to estimate the percent of canopy coverage. After the initial point had been selected using dice, subsequent points were randomly selected in a direction dictated by the second hand of an analogue watch 5 m from the previous point. The distance to the nearest 4 trees (≥ 5 cm DBH), with 1 tree within each quarter point of a grid system, was measured along with their DBH. We used a DBH threshold lower than

the standard 10 cm since the low occurrence of large trees would have prevented us from carrying out quantitative habitat surveys in heavily disturbed fragments. Fifty sampling points were completed in each fragment [Ganzhorn et al., 2011].

Fragment Spatial Structure

Using spatial data collected from satellite imagery which were ground-truthed and processed in ArcGIS, three separate landscape variables were generated. Isolation of forest fragments was examined in terms of their proximity to surrounding forests. A single multidirectional proximity figure was calculated for each fragment within ArcGIS. The figure was derived as the mean distance from the target fragment edge to the next nearest fragment edge as calculated across 4 quadrants delimited by the cardinal compass directions [Hill and Curran, 2004]. The data ranged from 140 to 515 m, and the error was defined by the GIS raster resolution as 5 m. The surrounding matrix of each fragment was measured to quantify the ease of primate dispersal using two separate methods. Firstly, a 250-metre 'buffer zone' was measured from the target fragment edge and the total area of land type (forest, pasture, water) was calculated [Bender et al., 2003; Tischendorf et al., 2003]. To estimate the relative ease of primate travel through these differing land types [Arroyo-Rodríguez and Mandujano, 2009], the area of these differing land types was multiplied by coefficient values (forest = coefficient 1, pasture = coefficient 2, water = coefficient 3). The proportion of these values to the total area within 250 m of the fragment edge, or buffer zone, was a product of this calculation. Secondly, the distance from the target fragment centroid (obtained using ArcGIS 'calculate geometry' function) to the nearest fragment centroid was measured in metres, and the same coefficient values were applied to the total distance. We included this second measurement because, whilst similar to the land type buffer zone coefficient proportion value, it is also an approximation of the likelihood that primate dispersals are occurring. For instance, FTF is a large, degraded fragment that satellite imagery shows has become isolated from the surrounding forest at some point between the years of 1970 and 2011. Because of its relatively large size, the likelihood that a primate would disperse from this area to a smaller fragment is low. This is reflected in the 'centroid-to-centroid coefficient distance', but not the 'land type buffer zone coefficient proportion' as the distance between two fragment centroids will be large, but the area surrounding the fragment is largely pasture with some forest cover.

We determined whether fragments had recently been reduced in size using LANDSAT 5 and LANDSAT 7 mosaic images from 1970 and 2011. Fragments were denoted as degraded if they had been reduced in size and/or connectivity since 1970 (fig. 1). We assessed recent logging activity in individual fragments based on visual evidence of cut wood found in the fragment (table 1).

Line Transects

Three researchers conducted surveys between May and August of 2014. In order to assess the density of *A. hybridus*, *A. seniculus* and *C. albifrons*, we used the line transect method, following recommendations suggested by Buckland et al., [2001, 2010] and Peres [1999]. Five of the studied fragments (F1, F2, F3, FA, FT) had previously been used for research, and the established trails were used for this study. We randomly superimposed sets of equally spaced (>100 m) parallel transect lines on a map of each of these 5 fragments, where fragment size allowed [Buckland et al. 2010]. We then uploaded these maps and used GPS devices (Garmin GPSMAP 64s GPS) to follow the lines in order to cut and mark the trails.

Overall, we sampled 127 km of line transects, with individual line transect length ranging from 96 to 1,440 m (mean transect length $\pm$ SD = 753.9 ± 277.1 m, $n = 19$). Transects were walked between 6:00 and 18:00 h at speeds of approximately 1 km/h stopping every 100 m and waiting 5 min in silence in order to detect any signs of movement or sound. Upon detecting a primate group, we waited 5–20 min if audio cues (movement sounds, calls) suggested that unseen primates were in close proximity in order to increase the chances of detecting all members of the group. Primate group centre location (distance and angle from line transect) was taken from first sighting of a primate, then amended to include accurate group centre estimates [Marshall et al., 2008]. Due to their variable size, transect effort was unequal between fragments, and ranged from 1,056 m (FP) to 20,282 m (FA), with a mean of 12,958 m. A single transect in each fragment was

Fig. 1. Description: composite vegetation classification map of study area comparing forest structure from 1970 and 2011.

walked approximately once every 3 days. Upon encountering primates we recorded the time of day, the number of individuals, the perpendicular distance of the group's centre to the line transect, the location of the encounter, and the sex-age category of each individual, when possible. Cluster size was defined as the total number of individuals in a group. For *C. albifrons* and *A. seniculus*, cluster size corresponds to the size of the cohesive social group, while for *A. hybridus* it corresponds to the size of the subgroup (given their fission-fusion social dynamics). Group spread was recorded for each cluster, but not for sightings of individual primates. Before data recording took place, interobserver bias was tested and found to be within a 95% threshold [Rovero et al., 2006].

Due to a low number of observations per fragment (mean = 28.6 ± 14.7 , range = 0–50, $n = 250$ observations) population density for each of the three primate species was calculated only for the entire area using Distance software [Thomas et al., 2006]. We selected the best fit model using the lowest Akaike's information criterion [Buckland et al., 2001]. Cluster size was calculated using the cluster-biased regression method. Data were truncated at 5% as suggested by Buckland et al. [2001] to exclude potential outliers. Encounter rate was then calculated for each fragment by dividing the entire number of encounters for a single species by the number of kilometres of transect walked (table 2). Primate encounter rates were grouped into 4 time periods of 3 h each (6–9 a.m., 9 a.m. to 12 p.m., 12–3 p.m., 3–6 p.m.) for later analysis.

Table 2. Encounter rates per kilometre of primate species by fragment

Fragment ID	<i>Ateles</i>	<i>Alouatta</i>	<i>Cebus</i>
FP	0	0	0
F1	0.98	1.16	0.89
F2	0	1.18	0.79
FT	0.31	0.93	1.96
FDU	0	0.29	0.68
F3	0.14	0.82	0.77
FA	0.15	0.35	0.84
FC	0	0.28	1.67
FN	0.31	0.69	0.46
FTF	0	0.09	1.63

Statistical Analyses

We used multiple linear regressions to predict population abundance using ecological vegetation variables (canopy density, mean DBH, tree density, edge proportion) and spatial/structural vegetation variables (fragment size, fragment isolation, centroid-to-centroid coefficient distance, buffer zone land type coefficient proportion) separately and selected the best predictors. Since we used the fragment as the unit of analysis, we ran two separate models due to sample size limitations. Before this model was run, vegetation variables and spatial/structural vegetation variables were checked for multicollinearity using Pearson's correlation. No variables were found to be significantly correlated ($p > 0.05$). Residual errors of this analysis were tested for normality using the Kolmogorov-Smirnov test in order to check the assumptions of general linear models, and the residual errors were found to be normally distributed. Vegetation structure differences between fragments were tested using a Kruskal-Wallis test, and qualitative disturbance cues (binary; presence of logging) were tested against quantitative vegetation data using a Mann-Whitney U test. Temporal differences in encounter rates of each primate species were tested for significant differences with a non-parametric Friedman's ANOVA test. All statistical analyses were performed with SPSS 20.

Results

Vegetation Structure

Mean DBH was highly variable between fragments (K-W: $\chi^2 = 31.55$, d.f. = 9, $p < 0.001$). Canopy density also varied significantly between fragments (K-W: $\chi^2 = 20.35$, d.f. = 9, $p = 0.016$). We found evidence of selective logging (cut and prepared wood ready for extraction) in 3 of the study fragments (FC, FTF, FDU) and a single fragment had indications that hunting had taken place (FDU, hunting platforms), though we cannot quantify the use of the platforms or the targeted prey. We found that fragments which had recently been reduced in size (FC, FTF, FDU; fig. 1) and showed visual evidence of recent logging exhibited lower mean DBH values (Mann-Whitney U test, $Z = -3.437$, $p = 0.001$, $n = 1,880$).

Density Estimates

In total, we recorded 250 sightings of primate groups. We encountered *C. albifrons* the most ($n = 130$ encounters), followed by *A. seniculus* ($n = 87$ encounters) and *A. hybridus* ($n = 33$). No differences between the number of sightings between differ-

ent time periods were found for any species (*Alouatta*, $\chi^2 = 4.744$, d.f. = 3, $p = 0.192$; *Ateles*, $\chi^2 = 0.625$, d.f. = 3, $p = 0.891$; *Cebus*, $\chi^2 = 3.989$, d.f. = 3, $p = 0.263$).

C. albifrons mean group size was 3.2 ± 2.4 individuals and ranged from 1 to 11 individuals. Average cluster/group spread was 16.1 m, and highest group spread was recorded as 40 m. They were found in all but the smallest fragment (FP; table 2). Density of *C. albifrons* was 62.2 individuals/km² (95% CI = 42.5–90.9). No ecological or spatial/structure vegetation variables were found to be significantly good predictors of encounter rates of *C. albifrons*.

A. seniculus had the largest group size, with an average of 3.9 ± 2.0 individuals, and it ranged from 1 to 12 individuals. Average cluster/group spread was 11.9 m, and highest group spread was recorded as 36 m. They were found in all but the smallest fragment (FP; table 2). Density of *A. seniculus* was 26.2 individuals/km² (95% CI = 17.4–39.3). We found a positive trend between number of encounters of *A. seniculus* and mean tree DBH ($\beta = 0.096$, adj. $R^2 = 0.252$, $p = 0.096$).

A. hybridus were found in subgroups that had an average size of 2.2 ± 1.2 individuals, and it ranged from 1 to 6 individuals. Average cluster/group spread was 15.38 m, and highest group spread was recorded as 28 m. They were found in 5 of the 10 fragments (F1, FT, F3, FA, FN; table 2). Density of *A. hybridus* was 8.2 individuals/km² (95% CI = 4.3–15.6). There was a positive correlation between number of encounters of *A. hybridus* and both mean tree DBH and canopy density (β DBH = 0.079, $p = 0.011$; β canopy density = 0.1, $p = 0.04$; model adj. $R^2 = 0.718$).

A possible confounding factor to density estimates was varying habituation levels of primates in separate fragments, some having been previously studied. However, the relatively uniform encounter rates of *C. albifrons* across fragments (table 2) regardless of degree of habituation suggest that the methodology used prevented bias.

Discussion

Our results indicate that the three diurnal primate species at San Juan differ in their ability to cope with recent fragmentation, and fragment quality may be more important than fragment size or isolation for some large-bodied primates. For *A. hybridus*, higher fragment quality, described in this study as high values in the ecological variables measured, corresponded with higher abundance, while fragment spatial structure variables had no apparent effect on the presence or absence of any primate species. However, all three primate species were found to be absent from the smallest fragment (FN), suggesting there is a minimum fragment size threshold for primate persistence. *C. albifrons* and *A. seniculus* showed no apparent dependence on either vegetation variables or spatial structure variables.

The results of this study support findings of previous research on the effect of forest fragmentation on primates [Garber et al., 2006], namely that ecomorphological adaptations and social structures govern species' ability to survive in disturbed habitat. In Madagascar, the diademed sifaka (*P. diadema*) occurring in forest fragments utilized fallback food sources and reduced daily path length, thus minimizing energy expenditure, when favoured fruit resources were absent [Irwin, 2008a, b]. Conversely, the collared lemur (*E. collaris*) increased time spent travelling and feeding, and reduced group size in response to fragmentation [Donati et al., 2011]. In Gabon, the moustached guenon (*Cercopithecus cephus*) exhibited a high degree of fission-fusion

dynamics to decrease feeding competition when inhabiting forest fragments [Tutin, 1999]. These studies suggest that dietary and behavioural flexibility of some primate species may guarantee some resilience to anthropogenic disturbance.

Given that forest fragmentation in this area is relatively recent [Link et al., 2010] (fig. 1) and all the species that we studied have the ability to cross the intervening matrix [Link, unpubl. data], we can assume that primate species 'choose' favourable fragments which to inhabit, possibly over generational time periods. As hypothesized, the far-ranging, arboreal frugivores *A. hybridus* are adversely affected by the constraints of living in degraded and fragmented forest. The overall population density of *A. hybridus* in this study (8.2 individuals/km²) is low when compared to other studies conducted in fragmented habitats (median = 19.6 individuals/km², n = 20 [Ramos-Fernández and Wallace, 2008]) despite the 'condensing' effects of habitat fragmentation [Link et al., 2010], suggesting that individuals are widely spaced within the available quality habitat.

Ateles spp. exhibit a social system with a high degree of fission-fusion dynamics, and under normal conditions subgroup size is highly flexible. In small fragments with limited food resources, scramble and contest competition may prevent some individuals from gaining adequate nutrition, resulting in physiological stress [Rimbach et al., 2013] and limiting the number of individuals that can subsist in each fragment. In this study *A. hybridus* subgroup sizes were found to be consistently low, suggesting that only 'subgroups' were found in isolated fragments and few fusion events took place. Reduced group size may be used by this species to reduce scramble and contest competition over food resources, and may be successful for the short-term survival of the population. However, the effects of small group sizes may have negative consequences for the long-term survival of this population, potentially leading to inbreeding and limited gene flow [Knapp, 2013].

The two ecological variables found to predict *A. hybridus* encounter rate in individual forest fragments were canopy density and mean DBH values. These variables showed lower values in fragments which showed evidence of selective logging. Whether this is due to selective logging or 'edge effects' related to fragmentation events is unclear [Harper et al., 2005].

Canopy density may be a key factor in allowing highly arboreal brachiators such as *Ateles* spp. to locate food resources efficiently [Skorupa, 1988; Dew, 2005]. In contrast, *Alouatta* spp. are quadrupedal climbers with small home ranges and highly available food resources, thus loss of canopy density may not limit their ability to survive in degraded forest fragments [Bernstein et al., 1976; Youlatos, 1993]. This may enable them to subsist in both small fragments and in fragments where large trees have been removed [Estrada and Coates-Estrada, 1996]. However, this may happen at a cost to overall group survivorship, as suggested by our findings that they still occur in degraded fragments but at reduced numbers (as seen in FC, FTF, FDU) [Schwarzkopf and Rylands, 1989; Palacios and Rodriguez, 2001].

No relationship was found between *A. seniculus* encounter rates, ecological vegetation variables or fragment structure variables. This suggests that *A. seniculus* may not be severely affected by fragmentation and habitat degradation in the short term, a similar result to previous studies at this site [Rimbach et al., 2013]. *Alouatta* spp. exhibit morphological adaptations that allow them to utilize leaves when fruit availability is low [Anapol and Lee, 1994; Fleagle, 2013]. Previous studies have shown *Alouatta* spp. display a broad range of behavioural adaptations to inhabit anthropo-

genically disturbed habitats: from adopting extreme energy-minimizing foraging patterns [Milton, 1980], expanding the range of plant species used as food sources [Bicca-Marques, 2003], even using anthropogenic elements (live fences) to move to new food sources [Asensio et al., 2008]. Adaptations such as these may enable the *A. seniculus* population in this study to subsist in degraded fragments.

C. albifrons were found in relatively high numbers in all but the smallest fragment (FN). Fragment quality and fragment spatial structure had no apparent effect on *C. albifrons*, suggesting that this species is not strongly affected by anthropogenic disturbance. Their ability to travel and forage terrestrially, and exploit numerous food sources is likely a contributing factor [Miller, 2002]. During our study we observed groups of *Cebus* travelling terrestrially between forest fragments several times, indicating that their flexible locomotion allows them to exploit a larger range of habitats [De Freitas et al., 2008]. We also found that *C. albifrons* abundance was high in fragments where *A. hybridus* and *A. seniculus* abundance was low, indicating that lower interspecific competition in these fragments may potentially enable this species to fully exploit the resources available in these fragments. This is a phenomenon that has been recorded previously in areas experiencing high levels of hunting activity [Rosin and Swamy, 2013].

Our study builds on the current understanding of how primates are affected by habitat degradation and fragmentation, which is of particular importance for the critically endangered brown spider monkey (*A. hybridus*). Given the long lifespan of *Ateles* spp., and the relatively short time scale in which anthropogenic disturbance can markedly affect primate populations, understanding the needs of the species may prove useful for the implementation of conservation strategies. While it appears in our study that *A. hybridus* has initially fared the worst as a result of the fragmentation of the habitat, *C. albifrons versicolor* and *A. seniculus* will likely experience long-term negative effects that are not yet fully evident.

Acknowledgements

We thank Gabriella Robayo for help with field data collection, the De Greiff and Lalinde families who allowed us to work on their land and for their support with the long-term research project at Hacienda San Juan de Carare in the last 7 years. Humberto Rueda, Hugo Soto and Rodolfo Alza also collaborated with us in the field and allowed us to work on their properties. We especially thank Fundación Proyecto Primates and in particular Anthony Di Fiore and Gabriela de Luna who helped us with logistics in the field. We would also like to thank Anna Nekaris, Vincent Nijman, Catherine Hill, Susan Cheyne, Rachel Sawyer, Stephanie Poindexter, Kathleen Reinhardt, Magda Svensson and Helena Bersacola for their support and Sarah Hammond for help with fund-raising.

References

- Abondano LA, Link A (2012). The social behavior of brown spider monkeys (*Ateles hybridus*) in a fragmented forest in Colombia. *International Journal of Primatology* 33: 769–783.
- Anapol F, Lee S (1994). Morphological adaptation to diet in platyrrhine primates. *American Journal of Physical Anthropology* 94: 239–261.
- Andrén H (1994). Effects of habitat fragmentation on birds and mammals in landscapes with different proportions of suitable habitat: a review. *Oikos* 71: 355–366.

- Arroyo-Rodríguez V, Mandujano S (2009). Conceptualization and measurement of habitat fragmentation from the primates' perspective. *International Journal of Primatology* 30: 497–514.
- Arroyo-Rodríguez V, Cuesta-del Moral E, Mandujano S, Chapman CA, Reyna-Hurtado R, Fahrig L (2013). Assessing habitat fragmentation effects on primates: the importance of evaluating questions at the correct scale. In *Primates in Fragments: Ecology and Conservation* (Marsh LK, Chapman CA, eds.), pp 13–28. New York, Springer.
- Arroyo-Rodríguez V, Mandujano S, Benítez-Malvido J, Cuende-Fanton C (2007). The influence of large tree density on howler monkey (*Alouatta palliata mexicana*) presence in very small rain forest fragments. *Biotropica* 39: 760–766.
- Asensio N, Korstjens AH, Schaffner CM, Aureli F (2008). Intragroup aggression, fission-fusion dynamics and feeding competition in spider monkeys. *Behaviour* 145: 983–1001.
- Barbisan-Fortes V, Bicca-Marques JC, Urbani B, Fernández VA, da Silva Pereira T (2014). Ranging behavior and spatial cognition of howler monkeys. In *Howler Monkeys: Behavior, Ecology and Conservation* (Kowalewski M, Garber PA, Cortés-Ortiz L, Urbani B, Youlatos D, eds.), pp 219–255. New York, Springer.
- Benchimol M, Peres CA (2014). Predicting primate local extinctions with 'real-world' forest fragments: a pan-Neotropical analysis. *American Journal of Primatology* 76: 289–302.
- Bender DJ, Tischendorf L, Fahrig L (2003). Using patch isolation metrics to predict animal movement in binary landscapes. *Landscape Ecology* 18: 17–39.
- Bernstein IS, Balcaen P, Dresdale L, Gouzoules H, Kavanagh M, Patterson T, Neyman-Warner P (1976). Differential effects of forest degradation on primate populations. *Primates* 17: 401–411.
- Bicca-Marques JC (2003). How do howler monkeys cope with habitat fragmentation? In *Primates in Fragments: Ecology and Conservation* (Marsh LK, ed.), pp 283–303. New York, Springer.
- Boyle SA, Smith AT (2010). Behavioral modifications in northern bearded saki monkeys (*Chiropotes satanas chiropotes*) in forest fragments of central Amazonia. *Primates* 51: 43–51.
- Buckland ST, Anderson DR, Burnham KP, Laake JL, Borchers DL, Thomas L (2001). *Introduction to Distance Sampling: Estimating Abundance of Biological Populations*. Oxford, Oxford University Press.
- Buckland ST, Plumptre AJ, Thomas L, Rexstad EA (2010). Design and analysis of line transect surveys for primates. *International Journal of Primatology* 31: 833–847.
- Buckley JS (1983). *The Feeding Behavior, Social Behavior, and Ecology of the White-Faced Monkey, Cebus capucinus, at Trujillo, Northern Honduras, Central America*. PhD thesis, University of Texas, Austin (University Microfilms International).
- Chapman CA, Peres CA (2001). Primate conservation in the new millennium: the role of scientists. *Evolutionary Anthropology Issues* 10: 16–33.
- Chapman CA, Wrangham R, Hunt K, Gardner L (1992). Estimators of fruit abundance of tropical trees. *Biotropica* 24: 527–531.
- Cowlishaw G, Dunbar R (2000). *Primate Conservation Biology*. Chicago, University of Chicago Press.
- Cristóbal-Azkarate J, Arroyo-Rodríguez V (2007). Diet and activity pattern of howler monkeys (*Alouatta palliata*) in Los Tuxtlas, Mexico: effects of habitat fragmentation and implications for conservation. *American Journal of Primatology* 69: 1013–1029.
- Crockett CM, Rudran R (1987). Red howler monkey birth data. II. Interannual, habitat, and sex comparisons. *American Journal of Primatology* 13: 369–384.
- Crockett CM, Sekulic R (1982). Gestation length in red howler monkeys. *American Journal of Primatology* 3: 291–294.
- Defler TR (1979). On the ecology and behavior of *Cebus albifrons* in eastern Columbia. I. Ecology. *Primates* 20: 475–490.
- De Freitas CH, Setz EZF, Araújo ARB, Gobbi N (2008). Agricultural crops in the diet of bearded capuchin monkeys, *Cebus libidinosus* Spix (Primates: Cebidae), in forest fragments in southeast Brazil. *Revista Brasileira de Zoologia* 25: 32–39.
- Dew JL (2005). Foraging, food choice, and food processing by sympatric ripe-fruit specialists: *Lagothrix lagotricha poeppigii* and *Ateles belzebuth belzebuth*. *International Journal of Primatology* 26: 1107–1135.
- Di Fiore A, Campbell CJ (2007). The atelines: variation in ecology, behavior, and social organization. In *Primates in Perspective* (Campbell CJ, Fuentes A, Mackinnon KC, Panger M, Beader SK, eds.), pp 155–185. New York, Oxford University Press.
- Donati G, Kesch K, Ndremifidy K, Schmidt SL, Ramanamanjato JB, Borgognini-Tarli SM, Ganzhorn JU (2011). Better few than hungry: flexible feeding ecology of collared lemurs *Eulemur collaris* in littoral forest fragments. *PLoS One* 6: e19807.
- Eppley TM, Verjans E, Donati G (2011). Coping with low-quality diets: a first account of the feeding ecology of the southern gentle lemur, *Hapalemur meridionalis*, in the Mandena littoral forest, southeast Madagascar. *Primates* 52: 7–13.
- Estrada A, Coates-Estrada R (1996). Tropical rain forest fragmentation and wild populations of primates at Los Tuxtlas, Mexico. *International Journal of Primatology* 17: 759–783.

- Estrada R (1999). Feeding and general activity patterns of a howler monkey (*Alouatta palliata*) troop living in a forest fragment at Los Tuxtlas, Mexico. *American Journal of Primatology* 48: 167–183.
- Ewers RM, Didham RK (2006). Confounding factors in the detection of species responses to habitat fragmentation. *Biology Reviews* 81: 117–142.
- Fahrig L (1999). Forest loss and fragmentation: which has the greater effect on persistence of forest-dwelling animals? In *Forest Fragmentation: Wildlife and Management Implications* (Rochelle L, Lehman L, eds.), pp 87–95. Leiden, Koninklijke Brill NV.
- Fedigan LM, Rose LM (1995). Interbirth interval variation in three sympatric species of neotropical monkey. *American Journal of Primatology* 37: 9–24.
- Fedigan LM, Carnegie SD, Jack KM (2008). Predictors of reproductive success in female white-faced capuchins (*Cebus capucinus*). *American Journal of Physical Anthropology* 137: 82–90.
- Fleagle JG (2013). *Primate Adaptation and Evolution*, 3rd ed. San Diego, Academic Press.
- Ganzhorn JU, Rakotonirany SJ, Rakotonirany YR (2011). Habitat description and phenology. In *Field and Laboratory Methods in Primatology: A Practical Guide* (Setchell JM, Curtis DJ, eds.), pp 40–56. Cambridge, Cambridge University Press.
- Garber PA, Estrada A, Pavelka MS (2006). New perspectives in the study of Mesoamerican primates: concluding comments and conservation priorities. In *New Perspectives in the Study of Mesoamerican Primates: Distribution, Ecology, Behavior, and Conservation* (Estrada A, Garber PA, Pavelka MSM, Luecke L, eds.), pp 563–584. New York, Springer.
- Gillespie TR, Chapman CA (2008). Forest fragmentation, the decline of an endangered primate, and changes in host-parasite interactions relative to an unfragmented forest. *American Journal of Primatology* 70: 222–230.
- Goldberg TL, Gillespie TR, Rwegu IB, Esoff EL, Chapman CA (2008). Forest fragmentation as cause of bacterial transmission among nonhuman primates, humans and livestock, Uganda. *Emerging Infectious Diseases* 14: 1375–1384.
- Goodenough AE, Goodenough AS (2012). Development of a rapid and precise method of digital image analysis to quantify canopy density and structural complexity. *ISRN Ecology* 2012: 1–11.
- Harper KA, Macdonald SE, Burton PJ, Chen J, Euskirchen NIES, Broszofski KD, Saunders SC, Eug E, Roberts DAR, Jaiteh MS, Esseen P (2005). Edge influence on forest structure and composition in fragmented landscapes. *Conservation Biology* 19: 768–782.
- Hill J, Curran PJ (2004). Area, shape and isolation of tropical forest fragments: effects on tree species diversity and implications for conservation. *Journal of Biogeography* 30: 1391–1403.
- Irwin MT (2008a). Feeding ecology of diademed sifakas (*Propithecus diadema*) in forest fragments and continuous forest. *International Journal of Primatology* 29: 95–115.
- Irwin MT (2008b). Diademed sifaka (*Propithecus diadema*) ranging and habitat use in continuous and fragmented forest: higher density but lower viability in fragments? *Biotropica* 40: 231–240.
- IUCN (2014). The IUCN Red List of Threatened Species. Version 2014.3. <http://www.iucnredlist.org> (accessed February 18, 2015).
- Janson CH (1998). Experimental evidence for spatial memory in foraging wild capuchin monkeys (*Cebus apella*). *Animal Behaviour* 55: 1229–1243.
- Knapp LA (2013). Molecular genetic tools for evaluating the consequences of habitat fragmentation. In *Primates in Fragments: Complexity and Resilience* (Marsh LK, ed.), pp 389–398. New York, Springer.
- Link A, de Luna AG, Alfonso F, Giraldo-Beltran P, Ramirez F (2010). Initial effects of fragmentation on the density of three neotropical primate species in two lowland forests of Colombia. *Endangered Species Research* 13: 41–50.
- Link A, Galvis N, Marquez M, Guerrero J, Solano C, Stevenson PR (2012). Diet of the critically endangered brown spider monkey (*Ateles hybridus*) in an inter-Andean lowland rainforest in Colombia. *American Journal of Primatology* 74: 1097–1105.
- Marsh LK, Chapman CA (2013). *Primates in Fragments: Complexity and Resilience*. New York, Springer.
- Marshall AJ, Boyko CM, Feilen KL, Boyko RH, Leighton M (2009). Defining fallback foods and assessing their importance in primate ecology and evolution. *American Journal of Physical Anthropology* 140: 603–614.
- Marshall AR, Lovett JC, White PCL (2008). Selection of line-transect methods for estimating the density of group-living animals: lessons from the primates. *American Journal of Primatology* 70: 452–462.
- Matthews LJ (2009). Activity patterns, home range size, and intergroup encounters in *Cebus albifrons* support existing models of capuchin socioecology. *International Journal of Primatology* 30: 709–728.
- Mbora DNM, Meikle DB (2004). Forest fragmentation and the distribution, abundance and conservation of the Tana River red colobus (*Procolobus rufomitratus*). *Biological Conservation* 118: 67–77.
- Michalski F, Peres CA (2005). Anthropogenic determinants of primate and carnivore local extinctions in a fragmented forest landscape of southern Amazonia. *Biological Conservation* 124: 383–396.
- Miller LE (2002). The role of group size in predator sensitive foraging decisions in wedge-capped capuchin monkeys (*Cebus olivaceus*). In *Eat or Be Eaten: Predator Sensitive Foraging among Primates* (Miller L, ed.), pp 95–106. Cambridge, Cambridge University Press.

- Milton K (1980). *The Foraging Strategy of Howler Monkeys: A Study in Primate Economics*. New York, Columbia University Press.
- Mittermeier RA, Rylands AB, Taylor LA, Chiozza F, Williamson EA, Wallis J, Schwitzer C (eds.) (2012). *Primates in Peril: The World's 25 Most Endangered Primates 2012–2014*. Bristol, IUCN/SSC Primate Specialist Group (PSG), International Primatological Society (IPS), Conservation International (CI), Bristol Conservation and Science Foundation.
- Palacios E, Rodríguez A (2001). Ranging pattern and use of space in a group of red howler monkeys (*Alouatta seniculus*) in a southeastern Colombian rainforest. *American Journal of Primatology* 55: 233–251.
- Panger MA, Perry S, Rose L, Gros-Lewis J, Vogel E, Mackinnon K, Baker M (2002). Cross-site differences in foraging behavior of white-faced capuchins (*Cebus capucinus*). *American Journal of Physical Anthropology* 119: 52–66.
- Peres CA (1997). Effects of habitat quality and hunting pressure on arboreal folivore densities in Neotropical forests: A case study of howler monkeys (*Alouatta* spp.). *Folia Primatologica* 68: 199–222.
- Peres CA (1999). General guidelines for standardizing line-transect surveys of tropical forest primates. *Neotropical Primates* 7: 11–16.
- Purvis A, Gittleman JL, Cowlishaw G, Mace GM (2000). Predicting extinction risk in declining species. *Proceedings of the Royal Society of London Biological Sciences* 267: 1947–1952.
- Ramos-Fernández G, Ayala-Orozco B (2002). Population size and habitat use of spider monkeys at Punta Laguna, Mexico. In *Primates in Fragments: Ecology and Conservation* (Marsh LK, ed.), pp 191–210. New York, Plenum-Kluwer.
- Ramos-Fernández G, Wallace RB (2008). Spider monkeys conservation in the twenty-first century: recognizing risks and opportunities. In *Spider Monkeys: The Behaviour, Ecology and Evolution of the Genus Ateles* (Campbell CJ, ed.), pp 351–376. Cambridge, Cambridge University Press.
- Rangel-Negrin A, Alfaro JL, Valdez RA, Romano MC, Serio-Silva JC (2009). Stress in Yucatan spider monkeys: effects of environmental conditions on fecal cortisol levels in wild and captive populations. *Animal Conservation* 12: 496–502.
- Rimbach R, Link A, Heistermann M, Gómez-Posada C, Galvis N, Heymann EW (2013). Effects of logging, hunting, and forest fragment size on physiological stress levels of two sympatric ateline primates in Colombia. *Conservation Physiology* 1: 1–11.
- Rimbach R, Pardo-Martínez A, Montes-Rojas A, Di Fiore A, Link A (2012). Interspecific infanticide and infant-directed aggression by spider monkeys (*Ateles hybridus*) in a fragmented forest in Colombia. *American Journal of Primatology* 8: 1–8.
- Rosin C, Swamy V (2013). Variable density responses of primate communities to hunting pressure in a western Amazonian river basin. *Neotropical Primates* 20: 25–31.
- Ross CA (1989). *Life-History Strategies of Primates*. Doctoral dissertation, University of London.
- Rovero F, Struhsaker TT, Marshall AR, Rinne TA, Pedersen UB, Butynski TM, Mtui AS (2006). Abundance of diurnal primates in Mwanihana Forest, Udzungwa Mountains, Tanzania: a multi-observer comparison of line-transect data. *International Journal of Primatology* 27: 675–697.
- Russo GA, Young JW (2011). Tail growth tracks the ontogeny of prehensile tail use in capuchin monkeys (*Cebus albifrons* and *C. apella*). *American Journal of Physical Anthropology* 146: 465–473.
- Sabbatini G, Stammati M, Tavares MCH, Giuliani MV, Visalberghi E (2006). Interactions between humans and capuchin monkeys (*Cebus libidinosus*) in the Parque Nacional de Brasília, Brazil. *Applied Animal Behaviour Science* 97: 272–283.
- Schwarzkopf L, Rylands AB (1989). Primate species richness in relation to habitat structure in Amazonian rainforest fragments. *Biological Conservation* 48: 1–12.
- Schwitzer C, Glatt L, Nekaris KAI, Ganzhorn JU (2011). Responses of animals to habitat alteration: an overview focussing on primates. *Endangered Species Research* 14: 31–38.
- Shimooka Y, Campbell CJ, Di Fiore A, Felton AM, Izawa K, Link A, Nishimura A, Ramos-Fernández G, Wallace R (2008). Demography and group composition of *Ateles*. In *Spider Monkeys: Behavior Ecology and Evolution of the Genus Ateles* (Campbell C, ed.), pp 329–348. Cambridge, Cambridge University Press.
- Skorupa JP (1988). *The Effect of Selective Timber Harvesting on Rain-Forest Primates in Kibale Forest, Uganda*. PhD dissertation, University of California at Davis.
- Strier KB (1992). Atelinae adaptations: behavioral strategies and ecological constraints. *American Journal of Physical Anthropology* 88: 515–524.
- Symington MM (1988). Demography, ranging patterns, and activity budgets of black spider monkeys (*Ateles paniscus chamek*) in the Manu National Park, Peru. *American Journal of Primatology* 15: 45–67.
- Symington MM (2000). Sex ratio and maternal rank in wild spider monkeys: when daughters disperse. *Behavioral Ecology and Sociobiology* 20: 421–425.
- Terborgh J (1983). Ecology of mixed troops. In *Five New World Primates: A Study in Comparative Ecology* (Terborgh J, ed.), pp 155–176. Princeton, Princeton University Press.

- Thomas L, Lake JL, Strindberg S, Marques FFC, Borchers DL, Buckland ST, Anderson DR, Burnham KP, Hedley SL, Pollard JH (2006). *Distance 4.0. Research Unit for Wildlife Population Assessment*. St Andrews, University of St Andrews.
- Tischendorf L, Bender DJ, Fahrig L (2003). Evaluation of patch isolation metrics in mosaic landscapes for specialist vs generalist dispersers. *Landscape Ecology* 18: 41–50.
- Tutin CE (1999). Fragmented living: behavioural ecology of primates in a forest fragment in the Lopé Reserve, Gabon. *Primates* 40: 249–265.
- Urbani B, Morales AL, Link A, Stevenson P (2008). *Ateles hybridus*. In *2008 IUCN Red List of Threatened Species*. www.iucnredlist.org.
- Vick L (2008). Immaturity in spider monkeys: a risky business. In *Spider Monkeys: Behavior, Ecology and Evolution of the Genus Ateles* (Campbell CJ, ed.), pp 288–328. Cambridge, Cambridge University Press.
- Wieczkowski J (2004). Ecological correlates of abundance in the Tana mangabey (*Cercocebus galeritus*). *American Journal of Primatology* 63: 125–138.
- Youlatos D (1993). Passages within a discontinuous canopy: bridging in the red howler monkey (*Alouatta seniculus*). *Folia Primatologica* 61: 144–147.