As a group, these carbohydrates are termed fermentable oligosaccharides, disaccharides, monosaccharides, and polyols (FODMAP) carbohydrates

Ann Nutr Metab 2016;68(suppl 1):8–17

Dietary Carbohydrates and Childhood Functional Abdominal Pain

by Bruno P. Chumpitazi and Robert J. Shulman

Key insights

Carbohydrates have been implicated as causative factors in the pathophysiology of childhood abdominal pain-related functional gastrointestinal disorders (AP-FGIDs). Certain carbohydrates (including lactose and fructose) may be malabsorbed. Carbohydrate malabsorption results in both bacterial fermentation (gas production) and increased gut osmotic activity which may generate pain and bloating. As a group, these carbohydrates are termed fermentable oligosaccharides, disaccharides, monosaccharides, and polyols (FODMAP) carbohydrates.

Current knowledge

Childhood AP-FGIDs affect up to 20% of children worldwide. These disorders are now classified as irritable bowel syndrome (IBS), functional dyspepsia, functional abdominal pain, and abdominal migraine. Many potential factors have been linked to AP-FGIDs, including psychosocial stress, inflammation, and alterations in gut homeostasis and function. The role of diet is particularly relevant, as the symptoms of AP-FGIDs are associated with intake of certain foods.

Practical implications

Each individual malabsorbed carbohydrate (i.e. lactose or fructose) exerts distinct physiological effects. Other factors which may affect symptom severity include the presence of low-grade gut inflammation and visceral hypersensitivity. Dietary restriction of FODMAP carbohydrates is a potential strategy for mitigating gastrointestinal symptoms in childhood IBS. Data from

The role of FODMAP carbohydrates in the etiology of AP-FGIDs.

randomized controlled trials also indicate that soluble fiber supplementation can reduce abdominal pain in children with AP-FGIDs.

Recommended reading

Gibson PR, Varney J, Malakar S, Muir JG: Food components and irritable bowel syndrome. *Gastroenterology* 2015;148:1158–1174.

Dietary Carbohydrates and Childhood Functional Abdominal Pain

Bruno P. Chumpitazi^{a, b} Robert J. Shulman^{a–c}

^aDepartment of Pediatrics, Baylor College of Medicine, ^bSection of Pediatric Gastroenterology, Hepatology, and Nutrition, Texas Children's Hospital, and ^cChildren's Nutrition Research Center, Houston, Tex., USA

Key Messages

- Carbohydrate malabsorption may cause gastrointestinal symptoms (e.g., bloating) via the physiologic effects of both increased osmotic activity and increased gas production from bacterial fermentation.
- There is a general lack of strong evidence supporting a restriction of individual carbohydrates (e.g., lactose) for childhood functional gastrointestinal disorders; however, emerging evidence suggests that a restriction of fermentable oligosaccharides, disaccharides, monosaccharides and polyols (FODMAP) may be effective.
- Soluble fiber (a complex carbohydrate) supplementation via randomized controlled intervention trials in childhood functional gastrointestinal disorders has demonstrated efficacy.

Key Words

Lactose · Fructose · FODMAP · Irritable bowel syndrome · Fiber · Recurrent abdominal pain

Abstract

Childhood functional gastrointestinal disorders (FGIDs) affect a large number of children throughout the world. Carbohydrates (which provide the majority of calories consumed in the Western diet) have been implicated both as culprits for the etiology of symptoms and as potential therapeutic agents (e.g., fiber) in childhood FGIDs. In this review, we detail how carbohydrate malabsorption may cause gastrointestinal symptoms (e.g., bloating) via the physiologic effects of both increased osmotic activity and increased gas production from bacterial fermentation. Several factors may play a role, including: (1) the amount of carbohydrate ingested; (2) whether ingestion is accompanied by a meal or other food; (3) the rate of gastric emptying (how quickly the meal enters the small intestine); (4) small intestinal transit time (the time it takes for a meal to enter the large intestine after first entering the small intestine); (5) whether the meal contains bacteria with enzymes capable of breaking down the carbohydrate; (6) colonic bacterial adaptation to one's diet, and (7) host factors such as the presence or absence of visceral hypersensitivity. By detailing controlled and uncontrolled trials, we describe how there is a general lack of strong evidence supporting restriction of individual carbohydrates (e.g., lactose, fructose) for childhood FGIDs. We review emerging evidence suggesting that a more comprehensive restriction of fermentable oligosaccharides, disaccharides,

monosaccharides and polyols (FODMAP) may be effective. Finally, we review how soluble fiber (a complex carbohydrate) supplementation via randomized controlled intervention trials in childhood functional gastrointestinal disorders has demonstrated efficacy.

© 2016 S. Karger AG, Basel

Introduction

Childhood abdominal pain-related functional gastrointestinal disorders (AP-FGIDs) affect up to 20% of children worldwide and account for at least 5% of all pediatric office visits in the United States [1, 2]. These disorders do not have an identifiable organic etiology based on conventional diagnostic testing; nevertheless, children with AP-FGIDs have both decreased quality of life and increased school absences as compared to their peers [3]. The gastrointestinal (GI) symptoms (e.g., abdominal pain) associated with these disorders may persist for years and into adulthood [4]. Successful interventions that ameliorate these symptoms in childhood AP-FGIDs may have an impact into adulthood [4]. Unfortunately, de-

spite growing interest and research, current conventional clinical interventions for these disorders are often ineffective.

Though previously defined under the broad category of recurrent abdominal pain (RAP), using Rome III criteria, AP-FGIDs are now classified into: irritable bowel syndrome (IBS), functional dyspepsia, functional abdominal pain, and abdominal migraine [5]. The etiology of AP-FGIDs is believed to be multifactorial and is best approached using the biopsychosocial model (fig. 1). Potential factors include: psychosocial distress (e.g., somatization, anxiety); alterations in the composition and function of the gut microbiome; low-grade gut inflammation; increased gut permeability; visceral hypersensitivity; altered GI motor function, and diet. Diet in particular has generated interest as a culprit given that children with AP-FGIDs often associate symptoms with intake of particular foods [6].

Carbohydrates provide the majority of calories consumed in the Western diet [7]. For many decades, individual carbohydrates (e.g., lactose) have been implicated both as culprits for the etiology of symptoms and as po-

Fig. 1. Biopsychosocial model of FGIDs. Modified from Rodriguez-Fandino et al. [45] with permission from the *Journal of Neurogastroenterology and Motility*.

Table 1. Listing of carbohydrates and examples of foods which contain them within the FODMAP group

Carbohydrate	Structure	Common foods
Fructose	Monosaccharide	Certain fruits: apples, pears; honey; juices
Lactose	Disaccharide	Dairy products: cow's milk, cheese
Fructans	Fructose polymers	Wheat, onions, rye
Galactans	Galactose polymers	Beans, legumes, asparagus
Polyols	Sugar alcohols	Certain fruits and vegetables: apricots, cherries, pears

Fig. 2. Proposed pathophysiologic mechanism of malabsorbed carbohydrates in FGIDs. Modified from Barrett et al. [46] with permission from *Practical Gastroenterology*.

tential therapeutic agents (e.g., fiber) in children with AP-FGIDs. This review will examine the proposed pathophysiology of carbohydrate intolerance and the current evidence implicating individual carbohydrates and groupings of carbohydrates in causing or improving symptoms in childhood AP-FGIDs. It will also briefly examine the proposed digestive physiology of fiber and the current evidence pointing toward the use of fiber in children with AP-FGIDs.

Carbohydrate Intolerance

Proposed Common Pathophysiologic Mechanisms

In children with AP-FGIDs, several individual carbohydrates (including lactose and fructose) have been implicated as exacerbating GI symptoms [8, 9]. As a group, these individual carbohydrates in combination with fructans, galactans, and polyols are termed fermentable oligosaccharides, disaccharides, monosaccharides and polyols (FODMAP) carbohydrates (table 1). FODMAP carbohy-

drates are rapidly fermented by colonic bacteria and are osmotically active (fig. 2). These physiologic effects can cause colonic distention from influx of water into the lumen and may lead to symptoms of abdominal pain, flatus, bloating, and loose and/or more frequent stools. These nonimmunologic adverse reactions to carbohydrates are termed carbohydrate intolerance.

Though grouped together as FODMAP carbohydrates, each individual malabsorbed carbohydrate may have a different physiologic effect. Magnetic resonance imaging evaluation in healthy adults demonstrates that fructose significantly increases small bowel water content; however, fructans increase small bowel water content to a much smaller degree [10]. In comparison, fructans significantly distend the colonic lumen; however, fructose increases colonic distention to a much smaller degree [10]. The individual effects of lactose, galactans, and polyols on GI physiologic function as measured by magnetic resonance imaging remain to be elucidated. In addition, whether these same physiologic changes occur following FODMAP carbohydrate ingestion in children (or adults) with AP-FGIDs is unknown.

Additional factors which may play a role in the generation of symptoms in subjects with lactose intolerance include low-grade gut inflammation and visceral hypersensitivity [11]. This was demonstrated by Yang et al. [11] in adults with IBS following lactose challenge and breath hydrogen testing to identify lactose malabsorbers (increased hydrogen production following the lactose challenge). Those with IBS who had lactose malabsorption were categorized into one of two groups: lactose malabsorption with lactose intolerance (concomitant increase in GI symptoms with the lactose challenge) and lactose malabsorption without lactose intolerance (no increase in GI symptoms). Yang et al. [11] found that adults with IBS with lactose malabsorption and intolerance (in comparison to those with malabsorption alone) had both increased mast cells in ileocolonic biopsies and increased visceral hypersensitivity.

Table 2. Prospective randomized controlled trials of lactose in children with AP-FGIDs

Authors [Ref.], country (year)	Design	Results
Lebenthal et al. [20], USA (1981)	Children with AP-FGIDs (n = 38: 21 LHBT positive/17 LHBT negative) 6-week baseline period followed by double-blind crossover challenges × 6 weeks (chocolate cow's milk vs. chocolate soy milk)	10/21 with a positive LHBT versus 4/17 (p = 0.13) with a negative LHBT had worsening of pain with cow's milk versus regular diet 7/21 with a positive LHBT versus 4/17 (p = 0.51) with a negative LHBT had worsening of pain with soy milk versus regular diet
Dearlove et al. [22], UK (1983)	Children with AP-FGIDs (n = 21: 8 lactose intolerant and 13 lactose tolerant) 2-week baseline followed by 2-week lactose-free diet followed by double-blind crossover challenge × 2 weeks (tonic with vs. without lactose)	1/8 lactose-intolerant versus 4/13 (p = 0.34) lactose-tolerant improved with lactose-free diet 1/8 lactose-intolerant versus 2/13 (p = 1.0) lactose-tolerant worsened with the lactose tonic
Gremse et al. [21] USA (2003)	Children with AP-FGIDs (n = 30) all with lactose malabsorption by LHBT 14-day double-blind crossover challenge (lactose-containing vs. lactose-hydrolyzed milk, 12 g/240 ml); subjects had been instructed to otherwise maintain a lactose-free diet throughout	Abdominal pain scores were significantly lower on the lactose hydrolyzed milk (p = 0.02); bloating, diarrhea, flatulence scores were lower but not statistically significantly different on the lactose-hydrolyzed milk

LHBT = Lactose hydrogen breath test.

Whether these factors also play a role in carbohydrate intolerance in childhood AP-FGIDs is currently unknown.

Lactose

Hippocrates first described lactose intolerance around 400 years BC, but the clinical symptoms have only become recognized in the past 50 years [12]. Lactose is a disaccharide (two conjoined sugars) that is unique in that it is only present in mammalian milk with estimates of 5.5–8.0 g/100 ml in human breast milk and 4.5–5.0 g/100 ml in cow's milk [13]. In order to be utilized, lactose needs to be broken down (hydrolyzed) by the enzyme lactase. Lactase is found on the tips of the villi of the small intestine and breaks down lactose into two monosaccharides – galactose and glucose. These monosaccharides are then absorbed by the small intestine and metabolized.

At the time of birth in humans, lactase activity is at its peak. However, lactase activity begins to decrease in early childhood in approximately 70% of humans; by adulthood, lactase activity is very low or undetectable [14, 15]. Approximately 30% of the population has lactase persistence whereby lactase activity remains beyond weaning and into adulthood [14, 15]. Lactase persistence occurs primarily in people of northern European descent.

The amount of lactose needed to induce symptoms in someone who is lactose intolerant varies depending on numerous factors. These include: (1) the amount of lactose ingested; (2) whether ingestion is accompanied by a meal or other food; (3) the rate of gastric emptying (how quickly the meal enters the small intestine); (4) small intestinal transit time (the time it takes for a meal to enter the large intestine after first entering the small intestine); (5) whether the meal contains bacteria with enzymes (beta-galactosidase) capable of breaking down lactase, and (6) colonic bacterial adaptation to one's previous diet [16, 17]. For example, in one study, single lactose loads of ≥ 15 g produced symptoms in the majority of lactase-deficient adults; however, lactose loads of up to 12 g (particularly if spread throughout the day) resulted in minimal or no symptoms [18]. Similarly, feeding subjects dairy-based yogurts with viable microbial cultures that contain bacteria with beta-galactosidase activity resulted in fewer signs of lactose malabsorption than feeding them pasteurized yogurts with little beta-galactosidase activity [17]. The amount of lactose needed to induce symptoms in a child with lactose malabsorption (either healthy or with an AP-FGID) is currently unknown.

Two other studies support a dose-response gradient of lactose exposure and symptoms. One study challenged 13 healthy adults with lactose maldigestion with varying

Table 3. Observational or uncontrolled lactose-related studies in children with AP-FGIDs

Authors [Ref.], country (year)	Methods	Findings
Liebman et al. [47], USA, (1979)	Children with AP-FGIDs (n = 38) versus controls (n = 29) completed lactose dietary challenges 17 children with AP-FGIDs (n = 11 malabsorbers and n = 6 absorbers) underwent lactose-free diet	11/38 AP-FGIDs versus 1/29 controls with positive lactose challenge (malabsorbers) ($p < 0.01$) 10/11 malabsorbers versus 0/6 absorbers had total relief of symptoms on diet ($p < 0.001$)
Barr et al. [8], USA (1979)	Children with AP-FGIDs (n = 80) completed LHBT 28 with positive LHBT placed on 6-week non-blinded dietary trial (2 weeks lactose free, 2 weeks lactose, 2 weeks lactose free)	32/80 (40%) LHBT positive 20/28 (71%) exacerbation of pain on lactose versus lactose-free diet ($p < 0.01$)
Christensen [41], Denmark (1986)	Children with AP-FGIDs (n = 50) versus controls (n = 40) completed lactose tolerance testing (2 g/kg); determined positive if abdominal pain and/or diarrhea developed	1/50 AP-FGID children found to be lactose intolerant versus 0/40 controls ($p = 1.0$)
Lebenthal et al. [20], USA (1981)	Children with AP-FGIDs (n = 69 underwent LHBT) 12-month milk elimination diet in positive LHBT (n = 15) versus negative LHBT (n = 13)	21/69 (30%) LHBT positive 6/15 positive LHBT versus 5/13 negative LHBT had elimination of pain ($p = 0.93$)
Blumenthal et al. [48], UK (1981)	Children with AP-FGIDs (n = 26) given LHBT (2 g/kg up to 50g, 20% solution)	3/26 (12%) LHBT positive 1 of 2 children with positive LHBT improved on lactose-free diet
Wald et al. [49], USA (1982)	Children with FGIDs (n = 40) completed LHBT (2 g/kg, up to 50 g) Dietary intervention (total 6 weeks): 2 weeks lactose elimination, 2 weeks lactose-containing, 2 weeks lactose elimination	12/40 (30%) LHBT positive at 2 g/kg 3/12 (25%) positive LHBT versus 5/28 (18%) negative LHBT ($p = 0.68$) with significant improvement on lactose-free diet
Bhan et al. [50], India (1982)	Children with AP-FGIDs (n = 70) versus controls (n = 50) lactose tolerance testing All AP-FGID children started on 4-week lactose-free diet	33/70 with AP-FGIDs (47%) versus 9/50 (18%) controls were lactose malabsorbers ($p < 0.01$) 11/33 (33%) lactose malabsorbers improved versus 6/37 (16%) lactose absorbers on diet ($p < 0.01$)
Webster et al. [51], USA (1995)	Children with AP-FGIDs (n = 137) given LHBT (1 g/kg, 10% solution) Those with positive LHBT recommended to have lactose elimination diets; telephone follow-up in 115/137 (84%)	LHBT positive in 33/137 (24%) In follow-up, 20/27 (74%) of lactose malabsorbers had less frequent abdominal pain versus 28/88 (32%) lactose absorbers ($p < 0.001$)
Ceriani et al. [52], Italy (1988)	Children with AP-FGIDs (n = 32) given LHBT (2 g/kg up to 50 g) 18 LHBT positive placed on lactose-free diet	24/32 (75%) LHBT positive 14/18 (78%) LHBT positive improved on the diet
Gremse et al. [53], USA (1999)	Children with AP-FGIDs (n = 146) given LHBT (1 g/kg, 10% solution)	LHBT positive in 50/146 (34%)
Boey [54], Malaysia (2001)	Children with AP-FGIDs (n = 24) given LHBT (2 g/kg) All participants started on lactose-free diet	17/24 (71%) LHBT positive None responded to lactose-free diet
Gijsbers et al. [55], The Netherlands (2012)	Children with AP-FGIDs (n = 220) given LHBT (2 g/kg) In those with positive LHBT: initial lactose elimination, provocation with lactose if symptoms resolved, followed by double-blind placebo controlled challenge undertaken for proof of causal relationship	57/210 (27%) LHBT positive 24/38 (63%) with positive LHBT improved on lactose elimination diet 7/23 (30%) of those who improved had worsening symptoms with provocation None of the 6 subsequently investigated via double-blind challenge had specific lactose-related symptoms

Table 3 (continued)

Authors [Ref.], country (year)	Methods	Findings
Ockeloen et al. [56], The Netherlands (2012)	Retrospective review of LHBT testing in children with chronic abdominal pain (n = 91); subsequent follow-up following lactose-free diet in those with positive LHBT	22/91 (24%) LHBT positive 21/22 (95%) improved on lactose free-diet at 5 months, and 14/22 (64%) continued with improvement at 15 months' follow-up
Dabritz et al. [27], Germany (2014)	Children with AP-FGIDs (n = 161) completed LHBT	35/161 (22%) LHBT positive Unclear how many specifically improved on a lactose free-diet

LHBT = Lactose hydrogen breath test.

amounts of lactose (0, 2, 6, 12, and 20 g) [16]. This study found that doses of up to 6 g (representing 120 ml of milk) were well tolerated, but symptoms began to emerge at 12 g of exposure [16]. Similar findings were noted in adults challenged with increasing quantities of lactose in a lactose-hydrolyzed milk preparation; the severity of symptoms was primarily dependent on the amount of lactose present [19].

Lactose-Related Studies in Children with AP-FGIDs

In children with AP-FGIDs, three randomized controlled trials have been completed, though none evaluated a restricted lactose diet in a controlled fashion (table 2). There have been a greater number of observational or uncontrolled trials (table 3). Lebenthal et al. [20] used both a randomized controlled challenge and an uncontrolled treatment component. The majority of the trials employed either: a lactose challenge (to identify those who were lactose intolerant based on the development of GI symptoms); or a lactose hydrogen breath test (to identify children who were malabsorbing lactose by demonstrating an excessive amount of hydrogen production within a specified time after lactose ingestion).

The randomized controlled trials provide mixed results regarding the ability of lactose to worsen symptoms in children with AP-FGIDs. One trial identified worsening abdominal pain (but not other symptoms) in children with AP-FGIDs with lactose malabsorption when using nonhydrolyzed lactose milk versus hydrolyzed lactose milk [21]. However, the other two trials did not find that lactose significantly worsened symptoms in children with AP-FGIDs and lactose malabsorption [20, 22]. Similarly, the uncontrolled studies have had mixed results with respect to both the proportion of children with AP-FGIDs

who have lactose malabsorption and the efficacy of a low-lactose diet in these children. Further prospective randomized controlled studies are needed, particularly with respect to evaluation of a lactose-restricted diet in children with AP-FGIDs who are lactose malabsorbers. Additionally, the potential role of low-grade gut inflammation in producing symptoms in lactose malabsorbers needs to be investigated in children.

Fructose

Fructose is a monosaccharide of which American children consume a mean of 54.7 g/day (representing approx. 10% of their daily caloric intake) [23]. Fructose is dependent on the glucose transporter 5 (GLUT5) and glucose transporter 2 (GLUT2) for passive absorption.

Fructose Studies in Children with AP-FGIDs

One prospective randomized controlled trial using a fructose-restricted diet has been conducted in children with AP-FGIDs. Wirth et al. [24] randomized 103 children with AP-FGIDs to either a fructose-restricted diet (n = 51) or a no dietary intervention group (n = 52) for 2 weeks. Those on the fructose-restricted diet (irrespective of their fructose hydrogen breath test result) had less pain intensity; however, they did not have a decrease in pain frequency [24]. In a prospective observational study, Wintermeyer et al. [25] placed 75 children with AP-FGIDs, all of whom had a positive fructose breath test, on a restricted fructose diet. Overall pain frequency and pain severity decreased while on the diet [25].

Three retrospective fructose-related studies have been completed. In 32 children with an AP-FGID, Gomara et al. [9] completed fructose hydrogen breath testing using various doses of fructose, including 1, 15, and 45 g. They

found that 11 (34%) of the 32 studied children had fructose malabsorption at either the 15 or 45 g doses. Of these 11, 9 (82%) had a significant improvement on a 2-week dietitian-recommended fructose-restricted diet [9]. Escobar et al. [26] completed fructose breath testing using 1 g/kg (up to 25 g) in 222 children with AP-FGIDs. Of these, 121 (55%) had fructose malabsorption, of whom 93 (77%) had improvement on a dietitian-recommended low-fructose diet. Dabritz et al. [27] included fructose hydrogen breath testing in their retrospective review and found that 55/142 (39%) children had fructose malabsorption. As several in the Dabritz cohort had multiple positive carbohydrate tests, the number of children who specifically responded to a fructose-restricted diet is unclear.

Sorbitol

No prospective studies regarding sorbitol restriction have been completed in children with AP-FGIDs. Hyams [28] published a case report describing a 15-year-old girl with chronic abdominal pain attributed to sorbitol ingestion from sugar-free gum which improved with elimination of the sorbitol source. In their retrospective study of hydrogen breath testing in children with RAP, Dabritz et al. [27] found that 109/146 (75%) children had sorbitol malabsorption; 27/31 (87%) improved on a sorbitol-restricted diet.

FODMAP Carbohydrates (table 1)

FODMAP carbohydrates include fructose and lactose, fructans, galactans, and polyols (such as sorbitol). Fructans are consumed primarily from wheat in the Western diet, with an average consumption ranging from approximately 4.4 to 6.7 g/day in children in the United States [29]. In the Western diet, galactans are consumed primarily from legumes and beans. Polyols are sugar alcohols such as sorbitol and xylitol.

Fructans, galactans, and polyols have unique bonds which are not able to be hydrolyzed by human enzymes. As such, following ingestion, the vast majority of these sugars enter the human colon essentially intact [30]. Within the colon, they may be metabolized by the gut bacteria, which contain enzymes with the ability to metabolize complex carbohydrates [31].

FODMAP Evidence in Childhood AP-FGIDs

Two prospective studies – one an open-label pilot study and the second one a randomized controlled trial – have evaluated a low-FODMAP diet in children with IBS. In the open-label study, 8 children with IBS were instructed by a dietitian to follow a low-FODMAP diet for 1 week [32]. The group as a whole had a decrease

in abdominal pain frequency, with 4 children having a $\geq 50\%$ decrease in abdominal pain frequency as compared to baseline [32]. In the randomized double-blind cross-over trial, children with IBS ($n = 33$) received a low-FODMAP or typical American childhood diet for 48 h [33]. The group as a whole had fewer abdominal pain episodes during the low-FODMAP diet. Those who had significant improvement ($>50\%$ decrease in abdominal pain frequency) on the low-FODMAP diet, compared to those who did not, had a gut microbiome composition that was enriched in bacteria with high saccharolytic potential (e.g., *Faecalibacterium prausnitzii*) [33].

Sucrose and Starch

Sucrose and starch are normally easily hydrolyzed and absorbed in the small intestine through the activity of sucrase-isomaltase (which hydrolyzes both sucrose and starch) and maltase-glucoamylase (which hydrolyzes starch) [34]. The activities of these enzymes may be evaluated using the Dahlqvist method from duodenal biopsy tissue obtained during upper endoscopy [35].

Carbohydrate enzyme deficiencies related to sucrose and starch digestion have been investigated in children with AP-FGIDs. Children with AP-FGID ($n = 44$) undergoing upper GI endoscopy for the evaluation of abdominal pain, vomiting, or gastroesophageal reflux were evaluated for lactase, sucrase, and glucoamylase activities [36]. Low enzyme activities were found in 32, 34, and 28% of children, respectively, with some children having a combination of enzyme deficiencies [36]. These findings are supported by those of El-Chammas et al. [37] who evaluated disaccharidase activity in 203 children with AP-FGIDs. Low enzyme levels for lactase, sucrose, glucoamylase, and palatinase (a measure of isomaltase) were found in 37, 21, 25, and 8%, respectively [37]. However, neither study found a correlation between enzyme activity levels and GI symptoms. In addition, the studies did not have a dietary intervention component. Future studies are needed to further investigate the significance of disaccharidase deficiencies and potential dietary or other interventions in children with AP-FGIDs.

Future studies are needed to further investigate the significance of disaccharidase deficiencies and potential dietary or other interventions in children with AP-FGIDs

Fiber Therapy in Childhood AP-FGIDs

Proposed Mechanisms

The rationale for fiber therapy for AP-FGIDs primarily relates to using either soluble fiber or insoluble fiber. Soluble fiber is able to absorb water and maintain the hydration of stool [38]. Insoluble fiber has the potential ability to mechanically stimulate/irritate the gut mucosa and induce a laxative effect through secretion of mucous and water, resulting in more rapid transit through the colon [38]. Fiber that is not easily fermented and is able to maintain its gel-like property, such as psyllium, provides increased water-holding capacity and regulation of stool form [38]. In contrast to soluble fiber, insoluble fiber (e.g., bran) has been shown to potentially exacerbate GI symptoms in adults with IBS [39].

Evidence in Childhood AP-FGIDs

Soluble fiber has demonstrated efficacy in randomized controlled trials in children with AP-FGIDs. In a double-blind randomized controlled trial with 52 children with AP-FGIDs, Feldman et al. [40] compared 10 g of corn fiber (a type of soluble fiber) supplementation for 2 weeks to placebo. With the primary endpoint of a 50% reduction in abdominal pain frequency, 13/26 (50%) of those with fiber supplementation achieved the endpoint compared to 7/26 (27%) of those receiving placebo ($p < 0.04$) [40]. In a double-blind randomized controlled trial with 32 children with RAP who were hospitalized for their abdominal pain, Christensen [41] reported no benefit using 7 weeks of ispaghula husk (psyllium – containing greater soluble than insoluble fiber properties) [38]. However, as this report is in the form of a letter to the editor, important information is missing (e.g., p values, standard deviations), making interpretation of the results difficult [41]. A prospective randomized controlled trial recently published as an abstract suggests that psyllium may be beneficial in children with IBS, though full details are still not published

as of the time of this review [42]. Retrospective studies suggest a benefit of increased fiber intake in reducing the risk of abdominal pain in children [43, 44].

Conclusions/Future Opportunities

Carbohydrate malabsorption can lead to GI symptoms through direct physiologic effects within the GI tract. However, current evidence does not strongly support restriction of single carbohydrates in children with AP-FGIDs. Rather, as in adults with IBS, FODMAP restriction is emerging as a better clinical strategy. In addition, soluble fiber supplementation appears to be an effective therapy in childhood AP-FGID.

Future research opportunities related to carbohydrates in childhood AP-FGIDs include: further elucidating the mechanisms of action of both FODMAP carbohydrates to induce symptoms and soluble fiber to ameliorate symptoms; determining the role of the gut microbiome in carbohydrate metabolism and associated symptoms; understanding the long-term effects of dietary carbohydrate-related interventions on the gut microbiome composition and function; and identifying those children with AP-FGIDs who would benefit most from dietary carbohydrate-related interventions.

Disclosure Statement

B.P.C. has received research support from the National Institutes of Health and QOL Medical, Inc., and was a consultant for Mead Johnson Nutrition. R.J.S. has received research support from the National Institutes of Health and is a consultant for Mead Johnson Nutrition.

Financial and/or intellectual support during the completion of this review was provided by NIH K23 DK101688 (B.P.C.) and NIH R01 NR013497 as well as the Daffy's Foundation (R.J.S.), the USDA/ARS under Cooperative Agreement No. 6250-51000-043 (R.J.S.), and P30 DK56338, which funds the Texas Medical Center Digestive Disease Center (B.P.C., R.J.S.).

The writing of this article was supported by Nestlé Nutrition Institute.

Soluble fiber has demonstrated efficacy in children with AP-FGIDs

References

- 1 Chitkara DK, Rawat DJ, Talley NJ: The epidemiology of childhood recurrent abdominal pain in Western countries: a systematic review. *Am J Gastroenterol* 2005;100:1868–1875.
- 2 Hyams JS, Burke G, Davis PM, Rzepski B, Andrulis PA: Abdominal pain and irritable bowel syndrome in adolescents: a community-based study. *J Pediatr* 1996;129:220–226.
- 3 Youssef NN, Murphy TG, Langseder AL, Rosh JR: Quality of life for children with functional abdominal pain: a comparison study of patients' and parents' perceptions. *Pediatrics* 2006;117:54–59.
- 4 Chitkara DK, van Tilburg MA, Blois-Martin N, Whitehead WE: Early life risk factors that contribute to irritable bowel syndrome in adults: a systematic review. *Am J Gastroenterol* 2008;103:765–774.

- 5 Rasquin A, DiLorenzo C, Forbes D, Guiraldes E, Hyams JS, Staiano A, Walker LS: Childhood functional gastrointestinal disorders: child/adolescent. *Gastroenterology* 2006;130:1527–1537.
- 6 Carlson MJ, Moore CE, Tsai CM, Shulman RJ, Chumpitazi BP: Child and parent perceived food-induced gastrointestinal symptoms and quality of life in children with functional gastrointestinal disorders. *J Acad Nutr Diet* 2014;114:403–413.
- 7 Austin GL, Ogden LG, Hill JO: Trends in carbohydrate, fat, and protein intakes and association with energy intake in normal-weight, overweight, and obese individuals: 1971–2006. *Am J Clin Nutr* 2011;93:836–843.
- 8 Barr RG, Levine MD, Watkins JB: Recurrent abdominal pain of childhood due to lactose intolerance. *N Engl J Med* 1979;300:1449–1452.
- 9 Gomara RE, Halata MS, Newman LJ, Bostwick HE, Berezin SH, Cukaj L, et al: Fructose intolerance in children presenting with abdominal pain. *J Pediatr Gastroenterol Nutr* 2008;47:303–308.
- 10 Murray K, Wilkinson-Smith V, Hoad C, Costigan C, Cox E, Lam C, et al: Differential effects of FODMAPs (fermentable oligo-, di-, mono-saccharides and polyols) on small and large intestinal contents in healthy subjects shown by MRI. *Am J Gastroenterol* 2014;109:110–119.
- 11 Yang J, Deng Y, Chu H, Cong Y, Zhao J, Pohl D, et al: Prevalence and presentation of lactose intolerance and effects on dairy product intake in healthy subjects and patients with irritable bowel syndrome. *Clin Gastroenterol Hepatol* 2013;11:262–268.
- 12 Matthews SB, Waud JP, Roberts AG, Campbell AK: Systemic lactose intolerance: a new perspective on an old problem. *Postgrad Med J* 2005;81:167–173.
- 13 Fusch G, Choi A, Rochow N, Fusch C: Quantification of lactose content in human and cow's milk using UPLC-tandem mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci* 2011;879:3759–3762.
- 14 Deng Y, Misselwitz B, Dai N, Fox M: Lactose intolerance in adults: biological mechanism and dietary management. *Nutrients* 2015;7:8020–8035.
- 15 Misselwitz B, Pohl D, Fruhauf H, Fried M, Vavricka SR, Fox M: Lactose malabsorption and intolerance: pathogenesis, diagnosis and treatment. *United European Gastroenterol J* 2013;1:151–159.
- 16 Hertzler SR, Savaiano DA: Colonic adaptation to daily lactose feeding in lactose maldigesters reduces lactose intolerance. *Am J Clin Nutr* 1996;64:232–236.
- 17 Martini MC, Smith DE, Savaiano DA: Lactose digestion from flavored and frozen yogurts, ice milk, and ice cream by lactase-deficient persons. *Am J Clin Nutr* 1987;46:636–640.
- 18 Savaiano DA, Levitt MD: Milk intolerance and microbe-containing dairy foods. *J Dairy Sci* 1987;70:397–406.
- 19 Gudmand-Hoyer E, Simony K: Individual sensitivity to lactose in lactose malabsorption. *Am J Dig Dis* 1977;22:177–181.
- 20 Lebenthal E, Rossi TM, Nord SK, Branski D: Recurrent abdominal pain and lactose absorption in children. *Pediatrics* 1981;67:828–832.
- 21 Gremse DA, Greer AS, Vacik J, DiPalma JA: Abdominal pain associated with lactose ingestion in children with lactose intolerance. *Clin Pediatr (Phila)* 2003;42:341–345.
- 22 Dearlove J, Dearlove B, Pearl K, Primavesi R: Dietary lactose and the child with abdominal pain. *Br Med J* 1983;286:1936–1936.
- 23 Vos MB, Kimmons JE, Gillespie C, Welsh J, Blanck HM: Dietary fructose consumption among US children and adults: the Third National Health and Nutrition Examination Survey. *Medscape J Med* 2008;10:160.
- 24 Wirth S, Klodt C, Wintermeyer P, Berrang J, Hensel K, Langer T, et al: Positive or negative fructose breath test results do not predict response to fructose restricted diet in children with recurrent abdominal pain: results from a prospective randomized trial. *Klin Padiatr* 2014;226:268–273.
- 25 Wintermeyer P, Baur M, Pilic D, Schmidt-Choudhury A, Zilbauer M, Wirth S: Fructose malabsorption in children with recurrent abdominal pain: positive effects of dietary treatment. *Klin Padiatr* 2012;224:17–21.
- 26 Escobar MA Jr, Lustig D, Pflugeisen BM, Amoroso PJ, Sherif D, Saeed R, et al: Fructose intolerance/malabsorption and recurrent abdominal pain in children. *J Pediatr Gastroenterol Nutr* 2014;58:498–501.
- 27 Dabritz J, Muhlbauer M, Domagk D, Voos N, Hennebohl G, Siemer ML, et al: Significance of hydrogen breath tests in children with suspected carbohydrate malabsorption. *BMC Pediatr* 2014;14:59.
- 28 Hyams JS: Chronic abdominal pain caused by sorbitol malabsorption. *J Pediatr* 1982;100:772–773.
- 29 Moshfegh AJ, Friday JE, Goldman JP, Ahuja JK: Presence of inulin and oligofructose in the diets of Americans. *J Nutr* 1999;129:1407S–1411S.
- 30 Barrett JS, Gibson PR: Fermentable oligosaccharides, disaccharides, monosaccharides and polyols (FODMAPs) and nonallergic food intolerance: FODMAPs or food chemicals? *Therap Adv Gastroenterol* 2012;5:261–268.
- 31 Flint HJ, Scott KP, Duncan SH, Louis P, Forano E: Microbial degradation of complex carbohydrates in the gut. *Gut Microbes* 2012;3:289–306.
- 32 Chumpitazi BP, Hollister EB, Oezguen N, Tsai CM, McMeans AR, Luna RA, et al: Gut microbiota influences low fermentable substrate diet efficacy in children with irritable bowel syndrome. *Gut Microbes* 2014;5:165–75.
- 33 Chumpitazi BP, Cope JL, Hollister EB, Tsai CM, McMeans AR, Luna RA, et al: Randomised clinical trial: gut microbiome biomarkers are associated with clinical response to a low FODMAP diet in children with the irritable bowel syndrome. *Aliment Pharmacol Ther* 2015;42:418–427.
- 34 Robayo-Torres CC, Quezada-Calvillo R, Nichols BL: Disaccharide digestion: clinical and molecular aspects. *Clin Gastroenterol Hepatol* 2006;4:276–287.
- 35 Dahlqvist A, Hammond JB, Crane RK, Dunphy JV, Littman A: Assay of disaccharidase activities in peroral biopsies of the small-intestinal mucosa. *Acta Gastroenterol Belg* 1964;27:543–555.
- 36 Karnsakul W, Luginbuehl U, Hahn D, Sterchi E, Avery S, Sen P, et al: Disaccharidase activities in dyspeptic children: biochemical and molecular investigations of maltase-glucosylase activity. *J Pediatr Gastroenterol Nutr* 2002;35:551–556.
- 37 El-Chammas K, Williams SE, Miranda A: Disaccharidase deficiencies in children with chronic abdominal pain. *JPEN J Parenter Enteral Nutr* 2015. Epub ahead of print.
- 38 McRorie JW Jr: Evidence-based approach to fiber supplements and clinically meaningful health benefits. Part 2. What to look for and how to recommend an effective fiber therapy. *Nutr Today* 2015;50:90–97.
- 39 Hebden JM, Blackshaw E, D'Amato M, Perkins AC, Spiller RC: Abnormalities of GI transit in bloated irritable bowel syndrome: effect of bran on transit and symptoms. *Am J Gastroenterol* 2002;97:2315–2320.
- 40 Feldman W, McGrath P, Hodgson C, Ritter H, Shipman RT: The use of dietary fiber in the management of simple, childhood, idiopathic, recurrent abdominal pain. Results in a prospective, double-blind, randomized, controlled trial. *Am J Dis Child* 1985;139:1216–1218.
- 41 Christensen MF: Recurrent abdominal pain and dietary fiber. *Am J Dis Child* 1986;140:738–739.
- 42 Shulman RJ, Cain K, Czyzewski D, Self M, Weidler E, Devaraj S, et al: Randomized, double blind trial of psyllium fiber in children with irritable bowel syndrome (IBS). *Gastroenterology* 2015;148:S120.
- 43 Huang RC, Palmer LJ, Forbes DA: Prevalence and pattern of childhood abdominal pain in an Australian general practice. *J Paediatr Child Health* 2000;36:349–353.
- 44 Paulo AZ, Amancio OM, de Moraes MB, Tabacow KM: Low-dietary fiber intake as a risk factor for recurrent abdominal pain in children. *Eur J Clin Nutr* 2006;60:823–827.
- 45 Rodriguez-Fandino O, Hernandez-Ruiz J, Schmulson M: From cytokines to toll-like receptors and beyond – current knowledge and future research needs in irritable bowel syndrome. *J Neurogastroenterol Motil* 2010;16:363–373.

- 46 Barrett JSG, Gibson PR: Clinical ramifications of malabsorption of fructose and other short-chain carbohydrates. *Pract Gastroenterol* 2007;53:51–65.
- 47 Liebman WM: Recurrent abdominal pain in children: lactose and sucrose intolerance, a prospective study. *Pediatrics* 1979;64:43–45.
- 48 Blumenthal I, Kelleher J, Littlewood JM: Recurrent abdominal pain and lactose intolerance in childhood. *Br Med J (Clin Res Ed)* 1981;282:2013–2014.
- 49 Wald A, Chandra R, Fisher SE, Gartner JC, Zitelli B: Lactose malabsorption in recurrent abdominal pain of childhood. *J Pediatr* 1982;100:65–68.
- 50 Bhan MK, Arora NK, Ghai OP, Dhamija NK, Nayyar S, Fotedar A: Lactose and milk intolerance in recurrent abdominal pain of childhood. *Indian J Pediatr* 1982;49:199–202.
- 51 Webster RB, Dipalma JA, Gremse DA: Lactose maldigestion and recurrent abdominal pain in children. *Dig Dis Sci* 1995;40:1506–1510.
- 52 Ceriani R, Zuccato E, Fontana M, Zuin G, Ferrari L, Principi N, Paccagnini S, Mussini E: Lactose malabsorption and recurrent abdominal pain in Italian children. *J Pediatr Gastroenterol Nutr* 1988;7:852–857.
- 53 Gremse DA, Nguyenduc GH, Sacks AI, Dipalma JA: Irritable bowel syndrome and lactose maldigestion in recurrent abdominal pain in childhood. *South Med J* 1999;92:778–781.
- 54 Boey CC: Lactase deficiency among Malaysian children with recurrent abdominal pain. *J Paediatr Child Health* 2001;37:157–160.
- 55 Gijsbers CF, Kneepkens CM, Buller HA: Lactose and fructose malabsorption in children with recurrent abdominal pain: results of double-blinded testing. *Acta Paediatr* 2012;101:e411–e415.
- 56 Ockeloen LE, Deckers-Kocken JM: Short- and long-term effects of a lactose-restricted diet and probiotics in children with chronic abdominal pain: a retrospective study. *Complement Ther Clin Pract* 2012;18:81–84.