The Planar Cell Polarity Transmembrane Protein Vangl2 Promotes Dendrite, Spine and Glutamatergic Synapse Formation in the Mammalian Forebrain

Nathan D. Okerlund^{a, b} Robert E. Stanley^{b, c} Benjamin N.R. Cheyette^{b–e}

^aDepartment of Neurology and Neurological Sciences, Stanford University, Palo Alto, Calif., and ^bDepartment of Psychiatry, ^cTetrad Graduate Program, ^dUCSF Weill Institute for Neurosciences, and ^eKavli Institute for Fundamental Neuroscience, University of California, San Francisco (UCSF), San Francisco, Calif., USA

Key Words

Dendrite · Forebrain · Planar cell polarity · Pyramidal neuron · Spine · Synapse · Vangl2 · Wnt

Abstract

The transmembrane protein Vangl2, a key regulator of the Wnt/planar cell polarity (PCP) pathway, is involved in dendrite arbor elaboration, dendritic spine formation and glutamatergic synapse formation in mammalian central nervous system neurons. Cultured forebrain neurons from Vangl2 knockout mice have simpler dendrite arbors, fewer total spines, less mature spines and fewer glutamatergic synapse inputs on their dendrites than control neurons. Neurons from mice heterozygous for a semidominant *Vangl2* mutation have similar but not identical phenotypes, and these phenotypes are also observed in Golgi-stained brain tissue from adult mutant mice. Given increasing evidence linking psychiatric pathophysiology to these subneuronal sites and structures, our findings underscore the relevance of core PCP proteins including Vangl2 to the underlying biology of major mental illnesses and their treatment.

© 2016 S. Karger AG, Basel

Nathan D. Okerlund and Robert E. Stanley contributed equally.

Introduction

Dendrite, dendritic spine and excitatory (glutamatergic) synapse formation and plasticity are molecularly interrelated developmental prerequisites for proper brain function and behavior. Defects in spine and synapse formation and turnover are increasingly understood to be key contributors to neuropsychiatric disorders including autism, schizophrenia and major affective disorders [1–5].

Cell communication pathways with well-established roles in other aspects of development – including both the Wnt/ β -catenin pathway [6–8] and the Wnt/planar cell polarity (PCP) pathway [9–11] – participate in these processes. The four-pass transmembrane protein Van Gogh-like 2 (Vangl2) is a key player in the PCP pathway and interacts with several proteins that influence synapse formation, including other PCP proteins such as Dishevelled (Dvl) [12] and Dapper-antagonist of catenin-1 (Dact1) [13] as well as the postsynaptic protein PSD95 [14–16]. Vangl2 also participates in signaling upstream of small GTPases [17, 18] regulating cytoskeletal dynamics crucial to dendrite and dendritic spine formation and plasticity [19–22]. *Looptail* (*Lp*) is a missense mutation in *Vangl2* [23] that causes semidominant phenotypes reflective of abnormal PCP including, in heterozygous animals, the curled or kinked tail from which the mutation gets its name, and in homozy-

gous animals, craniorachischisis, a completely open neural tube and exposed brain [23–25]. Genetically engineered *null* mutations in *Vangl2* cause similar phenotypes, but recessively and with lower penetrance in homozygous mutant animals [26, 27]. Using both allele types, we show here that *Vangl2* functions during neural differentiation in dendrite arborization, spine formation, spine maturation and glutamatergic synapse formation.

Methods

Genetics

The *Vangl2*^Δ allele [26] is here referred to as *Vangl2*KO or *Vangl2*⁻. The *Ltap*^{Lp} allele (Jackson Laboratory stock No. 000220) [23] is here referred to as *Vangl2*^{Lp} or *Lp*. All assays compared littermates of the designated experimental and control genotypes derived from *Vangl2*^{-/+} or *Vangl2*^{Lp/+} intercrosses.

Recombinant DNA

The mouse *Vangl2* cDNA clone and expression plasmid has been described previously [28].

Primary Culture and Immunostaining

Dissociated neurons were obtained from embryos, fixed, transfected with pEGFP-C1 (Clontech) and immunostained for synaptic markers as previously described [29, 30]. Hippocampal cultures were used where possible because of the ease of producing populations of predominantly pyramidal neurons and prior validation as a model system relevant to dendrite, spine and synapse formation in the forebrain [31]. The borders of the hippocampus were not consistently identifiable in mutant homozygotes (both *Vangl2*^{Δ/Δ} and *Vangl2*^{Lp/Lp}) due to craniorachischisis and neural precursor migration defects [27, 32]. Therefore, for these genotypes the entire forebrain (cortex and hippocampus) was prepared and compared to entire forebrain cultures of wild type (WT) and heterozygous littermates.

Visualization and Quantification

Cells were visualized on a Nikon CS1i upright spectral confocal at ×40 magnification or a custom-built spinning disc confocal microscope (Zeiss Axiovert 200M with Perkin-Elmer spinning disc and Melles Griot 43 series ion laser, Cascade 512B digital camera; Roper Scientific) at ×40 magnification. Golgi images were obtained on an Olympus IX51 compound inverted fluorescence microscope, also at ×40. Images were analyzed with ImageJ software (NIH). Sholl analysis, dendritic spine binning and spine and synapse quantification were performed as previously described [30].

Golgi Staining

Golgi-Cox silver staining was performed on 4-month-old littermates using the FD GolgiStain kit (FD Neurotechnologies) according to the manufacturer's instructions.

Statistics

All *p* values were calculated by unpaired parametric *t* tests (2-way comparison) or one-way ANOVA (≥3 comparisons) with Tukey's post hoc analysis using Graphpad Prism software. Each

comparison entailed ≥8 neurons and ≥11 dendrites per condition derived from multiple independent experiments; all reported differences reflect a minimal *p* ≤ 0.05 for experimental vs. control mice.

Results

Genetic Elimination of *Vangl2* Reduces Dendrite Arbor and Spine Formation

On inspection, cultured *Vangl2*^{-/-} forebrain pyramidal neurons had simpler dendrite arbors than controls (fig. 1a vs. b). Sholl analysis confirmed that mutant neurons had reduced numbers of dendrite branch crossings (fig. 1f). The maturation and density of dendritic spines were also affected (fig. 1a' vs. b'): the density of spines along dendrites was lower in *Vangl2*^{-/-} neurons compared to WT (fig. 1g). Moreover, *Vangl2*^{-/-} neurons had an increased percentage of immature (i.e. filopodial) relative to mature (i.e. thin, mushroom or stub-shaped) spines (fig. 1h).

Recombinant expression of *Vangl2* rescued the dendrite arbor phenotype in *Vangl2*^{-/-} neurons (fig. 1c, f). It also rescued spine density and spine maturity (fig. 1a', c', g, h).

Vangl2 Overexpression and Heterozygosity Do Not Alter Dendrites or Spines

To investigate effects of other genetic manipulations expected to alter (but not eliminate) *Vangl2* levels, we examined phenotypes in WT neurons recombinantly overexpressing *Vangl2*, and also in heterozygous (*Vangl2*^{-/+}) neurons. Neither of these genetic manipulations had any effect on dendrite complexity (fig. 1a vs. d, e; f), spine density (fig. 1a' vs. d', e'; g) or spine maturity (fig. 1a' vs. d', e'; h).

Vangl2^{-/-} Dendrites Have Fewer Glutamatergic Synaptic Contacts

We quantified density of glutamatergic synapses along dendrites by visualization with antibodies specific for VGlut1 (presynaptic marker) and PSD95 (postsynaptic marker). *Vangl2*^{-/-} neurons had reduced glutamatergic synapse density (fig. 1i vs. j; n). Recombinant expression of *Vangl2* in *Vangl2*^{-/-} neurons rescued glutamatergic synapse density (fig. 1j vs. k; n). Neither WT neurons recombinantly overexpressing *Vangl2* (fig. 1l) nor *Vangl2*^{-/+} neurons (fig. 1m) were significantly different from WT in this assay (fig. 1n). In contrast to this glutamatergic synapse phenotype, *Vangl2*^{-/-} neurons had no significant reduction in inhibitory (GABAergic) synapse density on their dendrites measured similarly (data not shown).

Fig. 1. Differentiation phenotypes in cultured *Vangl2*KO forebrain neurons. **a–e** EGFP-transfected cultured forebrain neurons from WT (**a**), *Vangl2*^{-/-} (**b**), *Vangl2*^{-/-} + *Vangl2* (**c**), WT + *Vangl2* (**d**) and *Vangl2*^{-/-} (**e**). **a'–e'** Corresponding dendritic segments at a higher magnification. **f–h** *Vangl2*^{-/-} neurons have simpler dendritic arbors as quantified by Sholl analysis (**f**), fewer dendritic spines

(**g**) and less mature spines (**h**) than controls. **i–m** Immunostaining for glutamatergic synapse markers along dendrite segments from WT (**i**), *Vangl2*^{-/-} (**j**), *Vangl2*^{-/-} + *Vangl2* (**k**), WT + *Vangl2* (**l**) and *Vangl2*^{-/-} (**m**) neurons. **n** Quantification. Scale bars = 30 μm (**a–e**); 5 μm (**a'–e'**, **i–m**). n.s. = *p* > 0.05, ** *p* ≤ 0.01, **** *p* ≤ 0.0001.

(For legend see next page.)

Lp Causes Mixed Effects on Dendrite Arbors, Spines and Glutamatergic Synapses

As with *Vangl2*^{-/-} neurons, upon visual inspection, *Vangl2*^{Lp/+} neurons had simpler dendrite arbors than WT (fig. 2a, b). Interestingly, although homozygous (*Vangl2*^{Lp/Lp}) mice have more severe cell polarity phenotypes than heterozygous (*Vangl2*^{Lp/+}) mice in embryonic axis elongation, inner ear epithelia, and neural precursor proliferation and migration [26, 32], neurons from *Vangl2*^{Lp/Lp} mice did not have a greater decrease in dendrite complexity than those from *Vangl2*^{Lp/+} mice (fig. 2b, c). Sholl analysis revealed that *Lp* neurons, whether heterozygous or homozygous, had similar reductions in number of dendrite branch crossings (fig. 2f). Dendritic spines were also affected by the *Lp* mutation, not in terms of density (fig. 2a', b', c; g) but in terms of maturity (fig. 2h).

Similar to *Vangl2*^{-/-} neurons, glutamatergic synapse density along dendrites of *Vangl2*^{Lp/+} and *Vangl2*^{Lp/Lp} neurons was reduced compared to WT (fig. 2i–k; m). As in *Vangl2*^{-/-} neurons, GABAergic synapse density was unaffected (data not shown).

Vangl2 Overexpression Rescues Only Some Lp Neurodevelopmental Phenotypes

As stated above, neurons carrying the *Vangl2*^{Lp} allele, whether heterozygous or homozygous, displayed similar decreases in dendrite complexity, spine maturation and glutamatergic synapse density. Interestingly, only the spine maturation phenotype was rescued by recombinant expression of *Vangl2*: recombinant overexpression of *Vangl2* did not rescue dendrite complexity (fig. 2e, f), nor did it rescue glutamatergic synapse density along den-

drites in *Lp* mutant neurons (fig. 2l, m). In contrast, overexpression of WT *Vangl2* did rescue spine maturity in these neurons (fig. 2b', d'; h).

Elimination of Dact1 Does Not Rescue Lp Neurodevelopmental Phenotypes

We previously showed that genetic loss of the Wnt signal pathway scaffold protein Dact1 can rescue embryonic phenotypes in *Vangl2*^{Lp/+} mice [13]. Dact1 is expressed in differentiating forebrain neurons, and its loss causes reductions in dendrite arbor complexity, dendritic spine maturity and glutamatergic synapse formation [30] similar to the *Vangl2* mutant phenotypes reported here. Nonetheless, neurons from *Vangl2*^{Lp/+}; *Dact1*^{-/-} mice had no rescue of dendrite complexity (fig. 2b vs. e; f) and no rescue of spine maturity (fig. 2b' vs. e'; h) relative to *Vangl2*^{Lp/+} neurons.

Golgi Staining Confirms Spine Reductions in the Vangl2 Mutant Forebrain

All the preceding assays were conducted using cultured forebrain neurons. To confirm that similar phenotypes occur in intact mammalian forebrain tissue, we analyzed the morphology of pyramidal neurons in the CA1 region of the hippocampus via Golgi-Cox staining on brains taken from adult (4- to 6-month-old) *Vangl2*^{Lp/+} and littermate control mice. (The prenatal death of *Vangl2*^{-/-} and *Vangl2*^{Lp/Lp} mice precluded such analysis.) CA1 pyramidal neurons in *Vangl2*^{Lp/+} mice had decreased spine density on apical dendrites compared to controls (fig. 2n–p). They also had an increased percentage of immature (filopodial) projections (fig. 2q).

Discussion

Loss of *Vangl2* function, whether via the semidominant *Lp* missense mutation or a targeted knockout, leads to decreased dendrite arbor complexity, spine maturity and glutamatergic synapse density without similar losses in GABAergic synapse density on forebrain pyramidal neuron dendrites. As dendritic spines are the specific subcellular site of glutamatergic synapses in pyramidal neurons, these data are consistent with previous findings that *Vangl2* localizes to the postsynaptic compartment of glutamatergic synapses, where it interacts with PSD95, trans-synaptic adhesion molecules and the PCP pathway protein Prickle2 [14–16]. Our genetic data corroborate previous reports of similar phenotypes following shRNA-mediated knockdown of *Vangl2* in cultured neu-

Fig. 2. Differentiation phenotypes in *Vangl2*^{Lp} forebrain neurons. **a–e** EGFP-transfected cultured neurons from WT (**a**), *Vangl2*^{Lp/+} (**b**), *Vangl2*^{Lp/Lp} (**c**), *Vangl2*^{Lp/+} + *Vangl2* (**d**) and *Vangl2*^{Lp/+}; *Dact1*^{-/-} (**e**). **a'–e'** Corresponding dendritic segments at a higher magnification. **f–h** Quantification. *Vangl2*^{Lp} neurons have simpler dendritic arbors (**f**), no reduction in total density of dendritic projections (spines + filopodia) (**g**), but a larger proportion of immature (filopodial) dendritic projections (**h**) than controls; only the last phenotype is rescued by recombinant expression of *Vangl2* (blue bar; color refers to the online version only). **i–l** Immunostaining for glutamatergic synapse markers along dendrite segments from WT (**i**), *Vangl2*^{Lp/+} (**j**), *Vangl2*^{Lp/Lp} (**k**) and *Vangl2*^{Lp/+} + *Vangl2* (**l**) cultured neurons. **m** Quantification. **n, o** Segments of apical dendrite from a Golgi-stained pyramidal neuron in hippocampal CA1 of WT (**n**) and *Vangl2*^{Lp/+} (**o**) littermates. Quantification of total spines (**p**) and immature (filopodial) spines (**q**). Scale bars = 30 μm (**a–e**); 5 μm (**a'–e'**, **i–l**, **n, o**). n.s. = *p* > 0.05, * *p* ≤ 0.05; ** *p* ≤ 0.01, **** *p* ≤ 0.0001.

rons [15, 33] and have allowed us to compare and contrast neurodevelopmental phenotypes induced by two molecularly distinct (engineered *null* vs. spontaneous missense) alleles at this locus.

Unlike the *KO (null)* allele, the *Lp* allele of *Vangl2* is not a simple loss of function: it causes dominant phenotypes to varying degrees in different biological contexts and exerts complex cell biological and biochemical effects on the encoded protein [12, 26, 27, 34]. Unlike homozygous *Vangl2KO* neurons, *Lp* neurons do not have reductions in spine density and their dendrite complexity and glutamatergic synapse phenotypes are not rescued by recombinant overexpression of *Vangl2*; this suggests that *Vangl2* has a molecularly distinct role in spine formation and maturation compared to dendrite arborization and glutamatergic synapse formation. The differences in rescue of these *Lp* neurodevelopmental phenotypes cannot be explained by different temporal requirements for *Vangl2* in these subneuronal compartments and processes, because recombinant expression of *Vangl2* in the same manner rescues all four phenotypes (dendrite complexity, spine number, spine maturity and glutamatergic synapse density) in *Vangl2^{-/-}* neurons.

Conclusions

PCP Pathway Proteins Play Important Divergent Roles in Neurons

Genetically altering *Vangl2* function, whether by semidominant missense (*Vangl2^{Lp}*) or engineered knockout (*Vangl2KO*), results in forebrain pyramidal neurons with simpler dendrite arbors, a larger proportion of immature spines and fewer glutamatergic synapses. Genetic disruption of other PCP genes in mammals, including *Dvl1* [35] and *Dact1* [30], causes similar phenotypes. Given the widespread neural expression of *Vangl2* and several other PCP genes during prenatal development, postnatal development and in the mature brain, it is plausible that the neural functions of these molecules are similarly widespread and continuous over the lifespan. In prior work, we have demonstrated requirements for the *Vangl2* partner *Dact1* during dendrite, spine and synapse development in both pyramidal neurons [30] and interneurons of the cerebral cortex [36, 37]. However, mutations in *Vangl2* and *Dact1* that exhibit strong mutual rescue during gastrulation [13] do not exhibit similar reciprocal functional relationships during neurodevelopment, suggesting that the molecular mechanisms underlying these phenotypes differ. Consistent with other studies [15, 38,

39] our genetic work therefore suggests that although components of the PCP pathway play important roles in the nervous system, the molecular pathways by which they function in developing neurons differ substantially from the PCP pathway established in studies of basic embryonic development.

The Goldilocks Principle in Molecular Neuropsychiatry

Evidence increasingly supports that neurodevelopmental and neuroplastic processes regulating spine and glutamatergic synapses contribute to the pathogenesis of psychiatric conditions including autism, schizophrenia and major affective disorders [2–4, 40–43]. In line with this, several PCP proteins contributing to these processes have been implicated in psychiatric pathophysiology. For example, in mice, the elimination of *Dvl1* reduces social behavior [44] and *Prickle2* sequence variants associated with autism lead to dendrite and glutamatergic synapse phenotypes [45]. Our work with *Vangl2* accords with these findings and underscores that at some genetic loci and in some biochemical pathways different molecular defects can lead to similar neural phenotypes. We refer to the general idea that similar neural and behavioral phenotypes can result from functionally different and even opposite molecular defects as the ‘Goldilocks Principle’ – i.e. either *too much* or *too little* can be deleterious in molecular neuropsychiatry. This is, in fact, a well-established theme for Wnt/PCP signaling in other developmental contexts [13, 46, 47]. A similar phenomenon has been observed for loci contributing to neuropsychiatry through entirely different mechanisms, including ion channel proteins such as *KCNA2* for which both gain- and loss-of-function mutations cause epilepsy [48] and *CACNA1C* for which both a gain-of-function mutation and reduced expression variants are associated with bipolar disorder [49]. This is now also firmly established for several copy number variants, such as 7q11.23 and 22q11.2, that contribute to psychiatric susceptibility either when deleted or duplicated [3, 50].

Acknowledgments

N.D.O. was supported in part by Autism Speaks Predoctoral Fellowship 4654. R.E.S. was supported in part by supplement R01HD55300-S1 to B.N.R.C. This research was also supported by a Simons Foundation Autism Research Initiative Grant (B.N.R.C.). We thank all members of the Cheyette laboratory for advice and support, and Yingzi Yang (NHGRI) for providing the *Vangl2^d* allele.

Statement of Ethics

All procedures involving live animals were carried out in accordance with an IACUC-approved animal use protocol at the University of California, San Francisco.

Disclosure Statement

The authors have no conflicts of interest to report.

References

- Williams AJ, Umemori H: The best-laid plans go oft awry: synaptogenic growth factor signaling in neuropsychiatric disease. *Front Synaptic Neurosci* 2014;6:4.
- Sekar A, Bialas AR, de Rivera H, Davis A, Hammond TR, Kamitaki N, Tooley K, Presumey J, Baum M, Van Doren V, Genovese G, Rose SA, Handsaker RE, Daly MJ, Carroll MC, Stevens B, McCarroll SA: Schizophrenia risk from complex variation of complement component 4. *Nature* 2016;530:177–183.
- Pinto D, Delaby E, Merico D, Barbosa M, Merikangas A, Klei L, Thiruvahindrapuram B, Xu X, Ziman R, Wang Z, Vorstman JA, Thompson A, Regan R, Pilorge M, Pellicchia G, Pagnamenta AT, Oliveira B, Marshall CR, Magalhaes TR, Lowe JK, Howe JL, Griswold AJ, Gilbert J, Duketis E, Dombroski BA, De Jonge MV, Cuccaro M, Crawford EL, Correia CT, Conroy J, et al: Convergence of genes and cellular pathways dysregulated in autism spectrum disorders. *Am J Hum Genet* 2014;94:677–694.
- Duman RS: Pathophysiology of depression and innovative treatments: remodeling glutamatergic synaptic connections. *Dialogues Clin Neurosci* 2014;16:11–27.
- Sanders SJ: First glimpses of the neurobiology of autism spectrum disorder. *Curr Opin Genet Dev* 2015;33:80–92.
- Ahmad-Annuar A, Ciani L, Simeonidis I, Herreros J, Ben Fredj N, Rosso SB, Hall A, Brickley S, Salinas PC: Signaling across the synapse: a role for Wnt and Dishevelled in presynaptic assembly and neurotransmitter release. *J Cell Biol* 2006;174:127–139.
- Cerpa W, Godoy JA, Alfaro I, Farias GG, Metcalfe MJ, Fuentealba R, Bonansco C, Inestrosa NC: Wnt-7a modulates the synaptic vesicle cycle and synaptic transmission in hippocampal neurons. *J Biol Chem* 2008;283:5918–5927.
- Farias GG, Valles AS, Colombres M, Godoy JA, Toledo EM, Lukas RJ, Barrantes FJ, Inestrosa NC: Wnt-7a induces presynaptic colocalization of $\alpha 7$ -nicotinic acetylcholine receptors and adenomatous polyposis coli in hippocampal neurons. *J Neurosci* 2007;27:5313–5325.
- Okerlund ND, Cheyette BN: Synaptic Wnt signaling – a contributor to major psychiatric disorders? *J Neurodev Disord* 2011;3:162–174.
- Farias GG, Alfaro I, Cerpa W, Grabowski CP, Godoy JA, Bonansco C, Inestrosa NC: Wnt-5a/JNK signaling promotes the clustering of PSD-95 in hippocampal neurons. *Biol Chem* 2009;284:15857–15866.
- Varela-Nallar L, Alfaro IE, Serrano FG, Parodi J, Inestrosa NC: Wingless-type family member 5a (Wnt-5a) stimulates synaptic differentiation and function of glutamatergic synapses. *Proc Natl Acad Sci USA* 2010;107:21164–21169.
- Torban E, Wang HJ, Groulx N, Gros P: Independent mutations in mouse Vangl2 that cause neural tube defects in looptail mice impair interaction with members of the Dishevelled family. *J Biol Chem* 2004;279:52703–52713.
- Suriben R, Kivimae S, Fisher DA, Moon RT, Cheyette BNR: Posterior malformations in Dact1 mutant mice arise through misregulated Vangl2 at the primitive streak. *Nat Genet* 2009;41:977–985.
- Yoshioka T, Hagiwara A, Hida Y, Ohtsuka T: Vangl2, the planar cell polarity protein, is complexed with postsynaptic density protein PSD-95 [corrected]. *FEBS Lett* 2013;587:1453–1459.
- Nagaoka T, Ohashi R, Inutsuka A, Sakai S, Fujisawa N, Yokoyama M, Huang YH, Igarashi M, Kishi M: The Wnt/planar cell polarity pathway component Vangl2 induces synapse formation through direct control of N-cadherin. *Cell Rep* 2014;6:916–927.
- Nagaoka T, Tabuchi K, Kishi M: PDZ interaction of Vangl2 links PSD-95 and Prickle2 but plays only a limited role in the synaptic localization of Vangl2. *Sci Rep* 2015;5:12916.
- Phillips HM, Murdoch JN, Chaudhry B, Copp AJ, Henderson DJ: Vangl2 acts via RhoA signaling to regulate polarized cell movement during development of the proximal outflow tract. *Circ Res* 2009;96:292–299.
- Lindqvist M, Horn Z, Bryja V, Schulte G, Papachristou P, Ajime R, Dyberg C, Arenas E, Yamaguchi TP, Lagercrantz H, Ringstedt T: Vang-like protein 2 and Rac1 interact to regulate adherens junctions. *J Cell Sci* 2009;123:472–483.
- Nakayama AY, Harms MB, Luo L: Small GTPases Rac and Rho in the maintenance of dendritic spines and branches in hippocampal pyramidal neurons. *J Neurosci* 2000;20:5329–5338.
- Negishi M, Katoh H: Rho family GTPases and dendrite plasticity. *Neuroscientist* 2005;11:187–191.
- Tashiro A, Minden A, Yuste R: Regulation of dendritic spine morphology by the Rho family of small GTPases: antagonistic roles of Rac and Rho. *Cereb Cortex* 2000;10:927–938.
- Van Aelst L, Cline HT: Rho GTPases and activity-dependent dendrite development. *Curr Opin Neurobiol* 2004;14:297–304.
- Kibar Z, Vogan KJ, Groulx N, Justice MJ, Underhill DA, Gros P: Ltap, a mammalian homolog of Drosophila Strabismus/Van Gogh, is altered in the mouse neural tube mutant Loop-tail. *Nat Genet* 2001;28:251–255.
- Strong LC, Hollander WF: Hereditary looptail in the house mouse. *J Hered* 1947;40:329–334.
- Murdoch JN, Doudney K, Paternotte C, Copp AJ, Stanier P: Severe neural tube defects in the loop-tail mouse result from mutation of Lpp1, a novel gene involved in floor plate specification. *Hum Mol Genet* 2001;10:2593–2601.
- Song H, Hu J, Chen W, Elliott G, Andre P, Gao B, Yang Y: Planar cell polarity breaks bilateral symmetry by controlling ciliary positioning. *Nature* 2010;466:378–382.
- Yin H, Copley CO, Goodrich LV, Deans MR: Comparison of phenotypes between different vangl2 mutants demonstrates dominant effects of the Looptail mutation during hair cell development. *PLoS One* 2012;7:e31988.
- Kivimae S, Yang XY, Cheyette BNR: All Dact (Dapper/Frodo) scaffold proteins dimerize and exhibit conserved interactions with Vangl, Dvl, and serine/threonine kinases. *BMC Biochem* 2011;12:33.
- Elia LP, Yamamoto M, Zang K, Reichardt LF: p120 catenin regulates dendritic spine and synapse development through Rho-family GTPases and cadherins. *Neuron* 2006;51:43–56.
- Okerlund ND, Kivimae S, Tong CK, Peng I-F, Ullian EM, Cheyette BNR: Dact1 is a postsynaptic protein required for dendrite, spine, and excitatory synapse development in the mouse forebrain. *J Neurosci* 2010;30:4362–4368.
- Banker G, Goslin K (eds): *Culturing Nerve Cells*, ed 2. Cambridge, MIT Press, 1998.
- Lake BB, Sokol SY: Strabismus regulates asymmetric cell divisions and cell fate determination in the mouse brain. *J Cell Biol* 2009;185:59–66.
- Hagiwara A, Yasumura M, Hida Y, Inoue E, Ohtsuka T: The planar cell polarity protein Vangl2 bidirectionally regulates dendritic branching in cultured hippocampal neurons. *Mol Brain* 2014;7:79.

- 34 Gravel M, Iliescu A, Horth C, Apuzzo S, Gros P: Molecular and cellular mechanisms underlying neural tube defects in the loop-tail mutant mouse. *Biochemistry* 2010;49:3445–3455.
- 35 Rosso SB, Sussman DJ, Wynshaw-Boris A, Salinas PC: Wnt signalling through Dishevelled, Rac and Jnk regulates dendritic development. *Nat Neurosci* 2005;8:34–42.
- 36 Arguello A, Yang X, Vogt D, Stanco A, Rubenstein JL, Cheyette BN: Dapper antagonist of catenin-1 cooperates with Dishevelled-1 during postsynaptic development in mouse forebrain GABAergic interneurons. *PLoS One* 2013;8:e67679.
- 37 Arguello A, Cheyette BN: Dapper Antagonist of Catenin-1 (Dact1) contributes to dendrite arborization in forebrain cortical interneurons. *Commun Integr Biol* 2014;6:e26656.
- 38 Glasco DM, Sittaramane V, Bryant W, Fritzsch B, Sawant A, Paudyal A, Stewart M, Andre P, Cadete Vilhais-Neto G, Yang Y, Song MR, Murdoch JN, Chandrasekhar A: The mouse Wnt/PCP protein Vangl2 is necessary for migration of facial branchiomotor neurons, and functions independently of Dishevelled. *Dev Biol* 2012;369:211–222.
- 39 Qu Y, Huang Y, Feng J, Alvarez-Bolado G, Grove EA, Yang Y, Tissir F, Zhou L, Goffinet AM: Genetic evidence that Celsr3 and Celsr2, together with Fzd3, regulate forebrain wiring in a Vangl-independent manner. *Proc Natl Acad Sci USA* 2014;111:E2996–E3004.
- 40 Krumm N, O’Roak BJ, Shendure J, Eichler EE: A de novo convergence of autism genetics and molecular neuroscience. *Trends Neurosci* 2014;37:95–105.
- 41 De Rubeis S, He X, Goldberg AP, Poultney CS, Samocha KE, Cicek AE, Kou Y, Liu L, Fromer M, Walker EM, Singh T, Klei L, Kosmicki J, Fu S-C: Synaptic, transcriptional and chromatin genes disrupted in autism. *Nature* 2014;515:209–215.
- 42 Musazzi L, Treccani G, Mallei A, Popoli M: The action of antidepressants on the glutamate system: regulation of glutamate release and glutamate receptors. *Biol Psychiatry* 2013;73:1180–1188.
- 43 Yi F, Danko T, Botelho SC, Patzke C, Pak C, Wernig M, Sudhof TC: Autism-associated SHANK3 haploinsufficiency causes Ih channelopathy in human neurons. *Science* 2016;352:aaf2669.
- 44 Long JM, LaPorte P, Paylor R, Wynshaw-Boris A: Expanded characterization of the social interaction abnormalities in mice lacking Dvl1. *Genes Brain Behav* 2004;3:51–62.
- 45 Sowers LP, Loo L, Wu Y, Campbell E, Ulrich JD, Wu S, Paemka L, Wassink T, Meyer K, Bing X, El-Shanti H, Usachev YM, Ueno N, Manak JR, Shepherd AJ, Ferguson PJ, Darbro BW, Richerson GB, Mohapatra DP, Wemmie JA, Bassuk AG: Disruption of the non-canonical Wnt gene PRICKLE2 leads to autism-like behaviors with evidence for hippocampal synaptic dysfunction. *Mol Psychiatry* 2013;18:1077–1089.
- 46 Park E, Kim GH, Choi SC, Han JK: Role of PKA as a negative regulator of PCP signaling pathway during *Xenopus* gastrulation movements. *Dev Biol* 2006;292:344–357.
- 47 Veeman MT, Slusarski DC, Kaykas A, Louie SH, Moon RT: Zebrafish prickle, a modulator of noncanonical Wnt/Fz signaling, regulates gastrulation movements. *Curr Biol* 2003;13:680–685.
- 48 Syrbe S, Hedrich UB, Riesch E, Djémié T, Müller S, Möller RS, Maher B, Hernandez-Hernandez L, Synofzik M, Caglayan HS, Arslan M, Serratosa JM, Nothnagel M, May P, Krause R, Löffler H, Detert K, Dorn T, Vogt H, Krämer G, Schöls L, Mullis PE, Linnankivi T, Lehesjoki AE, Sterbova K, Craiu DC, Hoffman-Zacharska D, Korff CM, Weber YG, Steinlin M, et al: De novo loss- or gain-of-function mutations in KCNA2 cause epileptic encephalopathy. *Nat Genet* 2015;47:393–399.
- 49 Gershon ES, Grennan K, Busnello J, Badner JA, Ovsiew F, Memon S, Alliey-Rodriguez N, Cooper J, Romanos B, Liu C: A rare mutation of CACNA1C in a patient with bipolar disorder, and decreased gene expression associated with a bipolar-associated common SNP of CACNA1C in brain. *Mol Psychiatry* 2014;19:890–894.
- 50 Sanders SJ, He X, Willsey AJ, Ercan-Sencicek AG, Samocha KE, Cicek AE, Murtha MT, Bal VH, Bishop SL, Dong S, Goldberg AP, Jinlu C, Keaney JF 3rd, Klei L, Mandell JD, Moreno-De-Luca D, Poultney CS, Robinson EB, Smith L, Solli-Nowlan T, Su MY, Teran NA, Walker MF, Werling DM, Beaudet AL, Cantor RM, Fombonne E, Geschwind DH, Grice DE, Lord C, et al: Insights into autism spectrum disorder genomic architecture and biology from 71 risk loci. *Neuron* 2015;87:1215–1233.