Single Case

Inframammary Dermatitis: A Case of Localized Late-Onset Darier's Disease

Dennis Linder^a Elena Marinello^b Pietro Maria Donisi^c
Roberto Salmaso^d Edoardo Zattra^b Anna Zampetti^e

^aSection of Biostatistics, University of Oslo, Oslo, Norway; ^bDermatology Unit, University of Padua, Padua, Italy; ^cPathology Department, 'SS. Giovanni e Paolo' Hospital, Venice, Italy; ^dPathology Unit, University of Padua, Padua, Italy; ^eRare Diseases and Periodic Fevers Research Centre, Department of Internal Medicine, Rome, Italy

Keywords

Adapalene · *ATP2A2* gene · Corticosteroids · Darier-White disease · Genodermatoses · Retinoids

Abstract

Darier's disease (DD) is an autosomal dominant inherited genodermatosis which is often under- or misdiagnosed. In the majority of cases, the disease manifests in adolescents or young adults with small brownish-yellow, warty, hyperkeratotic papules in multiple seborrheic areas of the body. Localized DD (LDD) is a clinical variant, first described by Kreibich in 1906; only a few cases are reported in the literature. We described the case of an aged woman presenting with LDD, and we review the literature on this subject.

© 2016 The Author(s)
Published by S. Karger AG, Basel

Case Report

A 71-year-old woman presented with several symmetrically distributed, small, very itchy, partly excoriated red papules surrounded by an erythematous halo in the infra-

mammary and presternal area. The lesions – approximately 4–8 mm in diameter – had allegedly appeared 5 years before and were localized in a well-demarcated area (fig. 1). Pressure did not provoke pain. Previous treatments with topical steroids, prescribed elsewhere on the basis of the clinical presentation, had proved ineffective.

A skin biopsy showed acanthosis in the epidermis, suprabasal dyskeratosis, and acantholysis, with clefts over the papillary dermis, covered by a single layer of epithelium. Dyskeratotic cells with irregular pyknotic nuclei surrounded by a bright eosinophilic halo were seen in the stratum granulosum, while the stratum corneum presented with so-called grains (i.e. multiple, compressed cells with elongated nuclei and many keratohyaline granules) (fig. 2).

Discussion

Darier's disease (DD), also called 'Darier-White disease' or 'dyskeratosis follicularis', is an autosomal dominant inherited genodermatosis with high penetrance (95%) and an incidence of 1:36,000–100,000 per year [1]. A mutation in the *ATP2A2* gene at locus 12q23–24, which codes for SERCA2, a calcium pump channel of the endoplasmic reticulum involved in adhesion and differentiation processes of epithelial cells, has been identified [2]. In the majority of cases, the disease manifests in adolescents or young adults with small brownish-yellow, warty, hyperkeratotic papules in the seborrheic areas of the body, such as the scalp, retroauricular and temporal areas, back of the neck, anterior trunk, and folds; the palms, soles, nails, and mucosae can also be involved. Lesions tend to coalesce into thicker, greasy plaques which, in folds, may macerate, become infected, and smell unpleasantly. In 80% of the cases, moderate-to-severe itch is reported [3].

Localized DD (LDD) is a clinical variant, first described by Kreibich in 1906 [4], usually presenting a zosteriform, linear, or segmental distribution and, in 10% of the cases, situated along the Blaschko lines [5, 6], because of a genetic mosaicism induced by a mutation occurring during embryogenesis [7]. Only a few nonsegmentally distributed and histologically confirmed cases of LDD have been described to date; they presented in the vulva [8, 9], on the mammary areolae [10], in photo-exposed areas only [11], and, in one case, ectopically – in the cervix [12]. LDD seems to have a late onset and can be associated with HIV and previous radiotherapy [13]. The medical history in our patient was irrelevant, the HIV serology negative. Histology proved conclusive for LDD, since all the required pathognomonic features were present, especially dyskeratotic cells.

The patient was encouraged to avoid triggering factors such as UV exposure, heat, and trauma and to daily apply 0.1% adapalene cream, since topical retinoids had proved effective in similar cases [3]. The treatment led to complete resolution of the lesions, as well as of the itch.

Statement of Ethics

Signed consent was obtained from the patient for publication.

Disclosure Statement

The authors declare no conflicts of interest.

References

- 1 Munro CS: The phenotype of Darier's disease: penetrance and expressivity in adults and children. *Br J Dermatol* 1992;127:126–130.
- 2 Savignac M, Edir A, Simon M, et al: Darier disease: a disease model of impaired calcium homeostasis in the skin. *Biochim Biophys Acta* 2011;1813:1111–1117.
- 3 Sehgal VN, Srivastava G: Darier's (Darier-White) disease/keratosis follicularis. *Int J Dermatol* 2005;44:184–192.
- 4 Kreibich K: Zum Wesen der Psorospermosis Darier. *Arch Dermatol Syphilol (Wien)* 1906;80:367.
- 5 Barfield RL, Barrett KR, Moon CM, David-Bajar K: Pruritic linear papules on a 75-year-old woman: a case of localized Darier-White disease. *Cutis* 2002;70:225–228.
- 6 O'Malley MP, Haake A, Goldsmith L, et al: Localized Darier disease. Implications for genetic studies. *Arch Dermatol* 1997;133:1134–1138.
- 7 Milobratović D, Vukićević J: Localized Darier's disease. *Acta Dermatovenereol Alp Pannonica Adriat* 2011;20:213–215.
- 8 Salopek TG, Krol A, Jimbow K: Case report of Darier disease localized to the vulva in a 5-year-old girl. *Pediatr Dermatol* 1993;10:146–148.
- 9 Barrett JF, Murray LA, MacDonald HN: Darier's disease localized to the vulva. Case Report. *Br J Obstet Gynaecol* 1989;96:997–999.
- 10 Fitzgerald DA, Lewis-Jones MS: Darier's disease presenting as isolated hyperkeratosis of the breasts. *Br J Dermatol* 1997;136:290.
- 11 Kimoto M, Akiyama M, Matsuo I: Darier's disease restricted to sun-exposed areas. *Clin Exp Dermatol* 2004;29:37–39.
- 12 Adam AE: Ectopic Darier's disease of the cervix: an extraordinary cause of an abnormal smear. *Cytopathology* 1996;7:414–421.
- 13 Emadi SN, Izadi M, Poursaleh Z, Akhavan-Moghaddam J, Barikbin B, Emadi SE: Darier disease associated with HIV infection: a case report. *HIV Clin Trials* 2011;12:48–53.

Fig. 1. Itchy red papules were noticeable in the inframammary region.

Fig. 2. Histology of the skin biopsy showed acanthosis in the epidermis, suprabasal dyskeratosis, and acantholysis, with clefts over the papillary dermis, covered by a single layer of epithelium. Dyskeratotic cells were seen in the stratum granulosum, while the stratum corneum presented with so-called grains.