***Kcne4* Deletion Sex-Dependently Alters Vascular Reactivity**

Geoffrey W. Abbott^a Thomas A. Jepps^b

^aBioelectricity Laboratory, Departments of Pharmacology and Physiology and Biophysics, School of Medicine, University of California, Irvine, Calif., USA; ^bIon Channels Group, Department of Biomedical Sciences, University of Copenhagen, Copenhagen, Denmark

Key Words

Kv7 channels · KCNE subunits · KCNE4 · KCNQ · Potassium channels · Vascular physiology · Smooth muscle

Abstract

Voltage-gated potassium (Kv) channels formed by Kv7 (KCNQ) α -subunits are recognized as crucial for vascular smooth muscle function, in addition to their established roles in the heart (Kv7.1) and the brain (Kv7.2–5). In vivo, Kv7 α -subunits are often regulated by KCNE subfamily ancillary (β) subunits. We investigated the effects of targeted germline *Kcne4* deletion on mesenteric artery reactivity in adult male and female mice. *Kcne4* deletion increased mesenteric artery contractility in response to α -adrenoceptor agonist methoxamine, and decreased responses to Kv7.2–7.5 channel activator ML213, in male but not female mice. In contrast, *Kcne4* deletion markedly decreased vasorelaxation in response to isoprenaline in both male and female mice. *Kcne4* expression was 2-fold lower in the female versus the male mouse mesenteric artery, and *Kcne4* deletion elicited only moderate changes of other *Kcne* transcripts, with no striking sex-specific differences. However, Kv7.4 protein expression in females was twice that in males, and was reduced in both sexes by *Kcne4* deletion. Our findings confirm a crucial role for KCNE4 in regulation of Kv7 channel activity to modulate

vascular tone, and provide the first known molecular mechanism for sex-specificity of this modulation that has important implications for vascular reactivity and may underlie sex-specific susceptibility to cardiovascular diseases.

© 2016 S. Karger AG, Basel

Introduction

KCNE4 is a member of the KCNE family (KCNE1–5) of single transmembrane-spanning β -subunits. Although these proteins do not form functional channels by themselves, they coassemble with pore-forming α -subunits of voltage-gated potassium (Kv) channels to regulate the trafficking, biophysical properties and pharmacology of the resultant heteromeric channel complexes [1]. The KCNE β -subunits can dramatically affect the activity of different ion channels; however, our understanding of their functional impact in many native systems is still limited. Recently, we showed that KCNE4 (also known as MiRP3) plays a fundamental role in regulating vascular tone [2]. In mesenteric artery myocytes from adult male

Dr. Geoffrey W. Abbott
A-360 Medical Surge II, Department of Pharmacology, School of Medicine
University of California, Irvine
Irvine, CA 92697-4625 (USA)
E-Mail abbottg@uci.edu

Dr. Thomas A. Jepps
Department of Biomedical Sciences, University of Copenhagen
The Panum Institute, Blegdamsvej 3
DK-2200 Copenhagen N (Denmark)
E-Mail tjepps@sund.ku.dk

rats, KCNE4 coassembled with the Kv7.4 and Kv7.5 α -subunits (encoded by KCNQ4 and KCNQ5, respectively), and knockdown of KCNE4 coincided with reduced Kv7.4 membrane expression and a more depolarized membrane potential [2].

Kv7.4 is a member of the Kv7 subfamily of Kv channel α -subunits, and has been widely reported to regulate vascular (and nonvascular) smooth muscle resting membrane potential [3–5]. The Kv7 family is comprised of 5 members (Kv7.1–5), of which Kv7.1, Kv7.4, and Kv7.5 are expressed in different rodent and human arteries [3, 5, 6]. In vascular smooth muscle, Kv7.4 and Kv7.5 channels predominate functionally, and recent evidence suggests that these isoforms coassemble to form heteromultimeric channels [2, 7, 8]. Pharmacological blockade of Kv7 channels produces vasoconstriction in several rodent and human blood vessels, and activation of Kv7 channels mediates relaxation in precontracted vessels [5, 6, 8–10]. Moreover, Kv7 channel blockade, or Kv7.4 channel knockdown, impairs various receptor-mediated vasorelaxations in different arteries. Thus, the vascular Kv7 channels are regarded as functional mediators of different receptor-activated vasodilatations [7, 11–15].

To date, these studies have used mainly male animal models, even though there are many sex-dependent differences in vascular physiology [16, 17]. Susceptibility to certain vascular diseases also differs between genders. Vascular diseases, such as hypertension, are typically more common in males and postmenopausal women than in premenopausal women [18–21]. Sex-dependent differences have been associated with different circulating androgen levels; however, we still do not understand the molecular mechanisms responsible for these differences in the vasculature [17, 18, 22].

Recently, using a germline *Kcne4*-deleted (*Kcne4*^{−/−}) mouse line, we showed cardiac KCNE4 expression to be positively regulated by 5 α -dihydrotestosterone (DHT), which corresponded with male mice displaying higher ventricular myocyte Kv current densities than female mice [23]. Ventricular Kv current density in young, sexually mature female mice was unaffected by *Kcne4* deletion and was similar to that of male *Kcne4*^{−/−} mice. This difference was associated with KCNE4 upregulating Kv1.5 and Kv4.2 currents specifically in male *Kcne4*^{+/+} mice due to *Kcne4* expression upregulation by DHT.

Given the newly identified role for KCNE4 in the vasculature, here we investigated the effects of *Kcne4* deletion on vascular function in mice, and found that *Kcne4* sex-specifically regulates mouse vascular reactivity.

Methods

Animals

The *Kcne4*^{−/−} mouse line was generated by replacement of the sole coding exon of *Kcne4* with the *neo* and *LacZ* genes in 129/Sv embryonic stem cells for embryonic injection followed by implantation into a C57 breeder. Progeny were genotyped by PCR and for final confirmation, Southern blot (performed by Lexicon, The Woodlands, Tex. and Texas A&M Institute for Genomic Medicine, College Station, Tex., USA; data not shown) [23]. Heterozygotes (*Kcne4*^{+/-}) were then backcrossed for at least 10 generations into the C57BL/6 strain. We genotyped the colony by conventional PCR of genomic DNA isolated from tail tips, using primers of the following sequences: *Kcne4* forward 5'-CAACGACAGCA-GTGAAGGC-3', *Kcne4* reverse 5'-GCAGAGCAAAAGCAAAA-CCC-3', Neo3a 5'-GCAGCGCATCGCCTTCTATC [23]. Mice were housed in a pathogen-free facility; colony maintenance, genotyping, euthanasia and tissue isolations were approved by the Animal Care and Use Committees, University of California, Irvine, and were conducted in strict accordance with recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health.

Myography

Studies were performed on 6- to 8-month-old sexually mature male and female mice, after euthanasia by CO₂ asphyxiation. The intestines were immediately removed, placed in ice-cold physiological saline solution (PSS) and the 2nd- and 3rd-order mesenteric arteries were dissected. The arteries were cleaned of adherent tissue and segments (approx. 2 mm) were mounted on 40- μ m stainless-steel wire in a myograph (Danish Myo Technology) for isometric tension recording. The myograph chambers were filled with PSS, maintained at 37°C and aerated with 95% O₂/5% CO₂. The vessels were allowed to equilibrate at 37°C before undergoing a passive-force normalization procedure.

Increasing concentrations (0.1–30 μ M) of the α -adrenoceptor agonist, methoxamine (Sigma), were applied. The vessels were then washed in fresh PSS before they were precontracted with 10 μ M methoxamine, and then increasing concentrations of the Kv7.2–7.5 channel activator, ML213 (0.1–10 μ M; Tocris), or the β -adrenoceptor agonist, isoprenaline (0.1–3 μ M; Sigma), were applied. The isoprenaline concentration response curve was then repeated in the presence of the Kv7 channel blocker, linopirdine (10 μ M; Sigma).

Real-Time Quantitative PCR (qPCR)

We quantified relative expression of the *Kcne1*–5 and *Kcnq1*–5 isoforms in the mesenteric arteries of male and female *Kcne4*^{+/+} and *Kcne4*^{−/−} mice by qPCR analysis, as we described previously [5]. Briefly, RNA was extracted with the RNeasy micro extraction kit, including a DNase treatment, according to the manufacturer's instructions (Qiagen). The Nanoscript 2 kit (PrimerDesign Ltd., Southampton, UK) was used to reverse-transcribe the RNA. The cDNA samples (at a concentration of 1.5 ng/ μ l) were used for quantitative analysis of *Kcne* transcript expression using Precision PLUS-iC SYBR mastermix (PrimerDesign Ltd.) in 20- μ l samples containing 5 μ l of cDNA and 300 nM primer. Experiments were run with the following cycling conditions: initial activation at 95°C for 10 min, followed by 40 cycles of 95°C for 15 s and 60°C for 1 min on a CFX96 real-time PCR detection system (Bio-Rad, UK),

Table 1. *Kcne* and *Kcnq* primers used for real-time qPCR experiments

Gene	Primer sequence, (+) sense/(-) antisense	GenBank accession No.	Amplicon, bp	Region spanned
<i>Kcne1</i>	(+) 5'-ACTCGCACGACCCTTTCA-3' (-) 5'-TTTCAATGACATAGCAAGCTCTG-3'	NM_012973	114	284–397
<i>Kcne2</i>	(+) 5'-TGTCATTTAAGTCCATTCCAATCAT-3' (-) 5'-TGAGAAAGAAGGTTGAAAGATTTGT-3'	NM_133603	116	769–884
<i>Kcne3</i>	(+) 5'-TTATGATGTCTGAGGATTGTCTTCT-3' (-) 5'-TGACCTAACTCTCTTACCAATTTCT-3'	NM_022235	114	449–562
<i>Kcne4</i>	(+) 5'-CCCTTGAGTCCCATGTGTCT-3' (-) 5'-GTAGCCCAGCATGATTCCAAT-3'	NM_212526	113	135–247
<i>Kcne5</i>	(+) 5'-GTCAACGGCGTCTGGAG-3' (-) 5'-CAGCAGCAAGCGGTTCAA-3'	NM_00110100	96	27–122
<i>Kcnq1</i>	(+) 5'-CTCGGAGTCACACGCTTCT-3' (-) 5'-GCTTGAACCTTCTTCTTTTACCAT-3'	NM_008434	75	1,301–1,375
<i>Kcnq2</i>	(+) 5'-CCCTCATTTGGTGTCTCGTTCT-3' (-) 5'-GGTCCGCGTTTCTCAAAG-3'	NM_010611	111	1,026–1,136
<i>Kcnq3</i>	(+) 5'-GAAGAGGGGCGAGGAGGA-3' (-) 5'-CCTGTACTTGGCGTTGTTCC-3'	NM_152923	93	226–318
<i>Kcnq4</i>	(+) 5'-GTGGTCTTTGGCTTGGAGTATAT-3' (-) 5'-CGATGACACAGAAGGGTTTCC-3'	NM_001081142	112	424–535
<i>Kcnq5</i>	(+) 5'-GTCAGATAAGAAGAGCCGAGAGA-3' (-) 5'-CGATGGACTGGACCTGTTTCT-3'	NM_001160139	107	2,164–2,270

and data was collected during each cycling phase. Melt-curve analysis completed the protocol. RNA samples that were reverse-transcribed without the nanoscript enzyme and no-template controls (NTCs) were run alongside all reactions to assess contamination. Quantification cycle (Cq) values were determined using Bio-Rad CFX96 Manager v3.0 software. The optimal reference genes were β -actin and cytochrome c-1 as determined by geNorm analysis (Biogazelle). The expression levels of the *Kcne* isoform transcripts were calculated relative to these reference genes in each artery to give a relative isoform expression profile using $2^{-\Delta Cq}$ [24]. All primer assays were obtained from PrimerDesign Ltd. and the sequences for the *Kcne1–5* and *Kcnq1–5* primers are given in table 1.

Western Blotting

We homogenized approximately 5-mm-long mouse mesenteric artery tissue samples (1 sample each from 4–5 different mice per sex per genotype) in 100 μ l of PBS (pH 7.5) containing protease inhibitor cocktail (Thermo Fisher, Waltham, Mass., USA) and 10% SDS (w/v) using a motorized Eppendorf homogenizer. Samples were then centrifuged for 10 min at 3 g at room temperature (to avoid solidification arising from the high SDS content). The supernatants were resuspended in LDS gel loading buffer (Thermo Fisher) containing 25 mM Tris(2-carboxyethyl)phosphine, heated for 10 min at 65°C, vortexed, centrifuged for 3 min at 5 g, and then separated by SDS-PAGE. Proteins were transferred to PVDF membranes and Western-blotted with 1/500 rabbit polyclonal anti-KCNQ4 (Santa Cruz Biotechnology, Dallas, Tex., USA) and 1/1,000 rabbit polyclonal anti-GAPDH (Abcam, Cambridge, UK) antibodies, with chemiluminescent detection via 1/5000 HRP-conjugated goat anti-rabbit IgG secondary antibodies (BioRad,

Hercules, Calif., USA). Band densities were quantified using ImageJ software (NIH, Bethesda, Mass., USA) and Kv7.4 band density values each normalized to same-lane GAPDH band density.

Statistical Analysis

Results are shown as means $\pm$ SEM, unless otherwise stated. $p < 0.05$ was taken as showing significant differences between means. Individual LogEC₅₀ values (fig. 1c, 2c) were calculated using a standard slope, equal to a Hill slope (or slope factor) of 1.0 for methoxamine and -1.0 for ML213. Differences between means were assessed for statistical significance with unpaired Student's *t* tests. A two-way ANOVA followed by a Bonferroni posttest was used to determine the significance of linopirdine on the isoprenaline-concentration effect curves and to compare the effect of isoprenaline between *Kcne4*^{+/+} and *Kcne4*^{-/-} male and female mice. Unpaired Student's *t* tests were used to compare changes in specific *KCNE* and *KCNQ* isoform transcript expression. ANOVA with the post hoc Tukey HSD test was used to compare Kv7.4 Western blot band densities.

Results

Application of sequentially increasing concentrations of the α -adrenoceptor agonist, methoxamine, caused a similar concentration-dependent constriction of mesenteric artery segments from both female ($n = 8$) and male ($n = 6$) *Kcne4*^{+/+} mice, with mean LogEC₅₀ values of -5.2 ± 0.05 M and -5.2 ± 0.05 M ($p = 0.31$, according to an un-

Fig. 1. Effect of methoxamine on male and female mesenteric arteries from *Kcne4*^{+/+} and *Kcne4*^{-/-} mice. Representative isometric tension recordings from female (**a**) and male (**b**) *Kcne4*^{+/+} (black lines) and *Kcne4*^{-/-} (red lines; color refers to the online version only) to sequentially increasing concentrations of methoxamine. **c** Mean concentration-effect curves to methoxamine. **d** Scatter plot of the EC₅₀ values to methoxamine. *n* = 6–8. ** *p* < 0.001, unpaired *t* test.

paired *t* test), respectively (fig. 1). In addition, the maximum force developed by the female and male artery segments did not differ significantly (4.3 ± 0.46 mN and 4.5 ± 0.53 mN, respectively; *p* = 0.75, according to an unpaired *t* test). In the females, the response of mesenteric artery segments from the *Kcne4*^{-/-} mice to methoxamine (LogEC₅₀ of -5.1 ± 0.01 M) was not significantly different from that of the *Kcne4*^{+/+} females (*p* = 0.53, according to an unpaired *t* test; fig. 1c, d). In contrast, segments of mesenteric artery from male *Kcne4*^{-/-} mice (*n* = 6) were significantly more sensitive to increasing concentrations of methoxamine (LogEC₅₀ of -5.6 ± 0.08 M; *p* = 0.006, according to an unpaired *t* test) compared to the *Kcne4*^{+/+} mesenteric arteries (fig. 1c, d); however, the maximum force developed at 30-μM methoxamine was not different in mesenteric artery segments between the male *Kcne4*^{+/+}

and *Kcne4*^{-/-} mice (4.6 ± 0.89 mN; *p* = 0.97, according to an unpaired *t* test).

Application of ML213 produced concentration-dependent relaxations that were not different in the mesenteric artery from the male and female *Kcne4*^{+/+} mouse (*p* = 0.24 by unpaired *t* test; fig. 2). In the females, there was no significant difference of vascular reactivity to ML213 in the *Kcne4*^{-/-} mice (*n* = 6) compared to the *Kcne4*^{+/+} mice (*n* = 5; fig. 2c, d). In contrast, *Kcne4* deletion in the mesenteric artery segments of male mice shifted the LogEC₅₀ for vascular response to ML213 from -6.01 ± 0.1 M (*Kcne4*^{+/+}, *n* = 5) to -5.7 ± 0.04 M (*Kcne4*^{-/-}, *n* = 6; *p* = 0.005 in unpaired *t* test; fig. 2c, d).

As previously reported, isoprenaline-induced relaxations are mediated, in part, by Kv7 channels [11]. Figure 3 shows that isoprenaline caused robust, concentration-

Fig. 2. Effect of ML213 on female and male mesenteric arteries from *Kcne4*^{+/+} and *Kcne4*^{-/-} mice precontracted with methoxamine. Representative isometric tension recordings from female (a) and male (b) *Kcne4*^{+/+} (black lines) and *Kcne4*^{-/-} (red lines; color refers to the online version only) to sequentially increasing concentrations of ML213 (0.1–10 μ M). c Mean concentration-effect curves to ML213. d Scatter plot of the EC₅₀ values to ML213. n = 5–6. ** p < 0.001, unpaired t test.

dependent relaxations in precontracted mesenteric artery segments from the *Kcne4*^{+/+} male and female mice that were not statistically different between the sexes. In both male (fig. 3a) and female (fig. 3b) *Kcne4*^{+/+} mouse mesenteric arteries, the isoprenaline relaxations were inhibited by linopirdine (10 μ M; n = 5–6).

We next determined the impact of *Kcne4* deletion on isoprenaline-mediated relaxations in mesenteric arteries from male and female mice. Isoprenaline was markedly less effective at relaxing the mesenteric arteries of *Kcne4*^{-/-} male and female mice (>2-fold less relaxation at 1 and 3 μ M isoprenaline compared to *Kcne4*^{+/+}; p < 0.05; n = 4–8; fig. 4).

We next employed qPCR to quantify the expression of *Kcne1*–5 and *Kcnq1*–5 isoform transcripts using mesenteric artery cDNA from male and female *Kcne4*^{+/+} and *Kcne4*^{-/-} mice (fig. 5). Expression of *Kcne4* was >2-fold

higher in male mouse mesenteric artery tissue compared to that of the female mice (n = 7; p = 0.0061 by unpaired t test). *Kcne1* was detectable in 5 of 7 female mesenteric artery samples but not detectable in any male mouse mesenteric arteries (n = 7; p = 0.016 by unpaired t test). *Kcne3* displayed the highest levels of expression, and, as also observed for *Kcne2* and *Kcne5*, this expression level was sex-independent (fig. 5a). As expected, *Kcne4* transcript was not detectable in *Kcne4*^{-/-} mesenteric arteries from either sex, confirming efficient germline deletion (fig. 5b, c). *Kcne4* deletion was associated with an approximately 50% increased expression of the already highly expressed *Kcne3* transcript in both male and female mice, which was significant in the male (n = 6; p = 0.006) but not the female mesenteric arteries (n = 6; p = 0.104; fig. 5b, c). *Kcne4* deletion was not associated with any other changes in other

Fig. 3. Effect of the Kv7 channel blocker, linopirdine (grey), on the vasorelaxant effect of isoprenaline in 6 male (a) and 5 female (b) *Kcne4*^{+/+} mesenteric arteries. * *p* < 0.05 and *** *p* < 0.001, two-way ANOVA followed by a Bonferroni post hoc test.

Fig. 4. Effect of isoprenaline on male and female mesenteric arteries from *Kcne4*^{+/+} and *Kcne4*^{-/-} mice. Mean concentration-effect curves to isoprenaline (*n* = 4–8). A two-way ANOVA followed by a Bonferroni post hoc test compared the effect of isoprenaline in *Kcne4*^{+/+} (black) with the *Kcne4*^{-/-} (red; color refers to the online version only) in both sexes. Significance is denoted by * for females and § for males.

Kcne isoform expression in male or female mouse mesenteric arteries (fig. 5b, c). We also analyzed expression of *Kcnq1*–5 transcripts in male and female *Kcne4*^{+/+} and *Kcne4*^{-/-} mice mesenteric arteries. There was no change in *Kcnq* transcript expression between *Kcne4*^{+/+} male and female mesenteric arteries (*n* = 6; fig. 6a), nor did we detect any change in *Kcnq* transcript expression in *Kcne4*^{+/+} mice compared to *Kcne4*^{-/-} in both male and female mice (*n* = 6; fig. 6b, c).

In contrast, *Kcne4* deletion reduced Kv7.4 protein expression. Kv7.4 protein expression in mouse mesenteric artery tissue preparations from female mice was double that observed in male mice (*p* < 0.05), and was reduced in both sexes by *Kcne4* deletion (fig. 7a, b), with the comparison for female *Kcne4*^{+/+} versus male or fe-

male *Kcne4*^{-/-} mice achieving statistical significance by ANOVA after the Tukey HSD test (*p* = 0 < 0.01; fig. 7b).

Discussion

This is the first study to identify sex-dependent differences in the function and expression levels of an ion channel ancillary subunit in the vasculature. Our findings show that *Kcne4* expression is higher, and functional effects of its deletion more striking, in the mesenteric arteries of male mice compared to female mice. Using a *Kcne4*^{-/-} mouse line, we found that targeted germline *Kcne4* deletion renders male but not female mouse mesenteric arteries more sensitive to α -adren-

Fig. 5. Real-time quantitative PCR analysis comparing *Kcne1–5* mRNA expression in male and female *Kcne4*^{+/+} mice (**a**) and *Kcne4*^{+/+} compared to *Kcne4*^{-/-} in male (**b**) and female (**c**) mice (n = 5–6). * p < 0.05, ** p < 0.01 and *** p < 0.001, unpaired t tests.

ergic stimulation and less responsive to the Kv7.2–7.5 activator, ML213. These data suggest that males are more sensitive to changes in *Kcne4* expression in the vasculature than females. However, responses to the β -adrenoceptor agonist, isoprenaline, were inhibited in both male and female *Kcne4*^{-/-} mice, suggesting that *Kcne4* is important in mediating the effects of this receptor in both sexes.

Kv7 channels, particularly Kv7.4 and Kv7.5, are important regulators of vascular tone [3, 6–15, 25, 26]. These channels have been shown extensively to be expressed and functionally important in several human and rodent arteries. We previously reported that KCNE4 is expressed in several arteries, and that targeted KCNE4 knockdown in male rat mesenteric arteries using a morpholino strategy depolarized the smooth muscle resting-membrane potential and reduced vasorelaxations in response to a Kv7.2–7.5 activator (S-1) [2]. We found that KCNE4 coassembles

with Kv7.4 and Kv7.5 in mesenteric myocytes, left-shifting the voltage-dependence of Kv7.4 channel activation and increasing Kv7.4 membrane expression [2]. Our findings here, using male *Kcne4*^{-/-} mice, confirm these previous findings and highlight the crucial regulatory role KCNE4 has on vascular Kv7 channel function in male mesenteric arteries. Importantly, this study uncovers a novel role for the KCNE4 subunit in mediating β -adrenoceptor responses in the mesenteric artery. Kv7 channels in the vasculature contribute to different Gs-coupled-receptor-mediated relaxations, yet the exact signaling mechanisms are still unclear [7, 11, 12, 14, 15]. Following Gs-coupled-receptor activation, a cAMP-dependent mechanism [11] and the $\beta\gamma$ -subunits [15] have both been shown to enhance Kv7.4/Kv7.5 channel activity. This study confirms that the isoprenaline effects in the mesenteric artery are sensitive to Kv7 blockade (linopirdine) and reveals an important role for KCNE4 in mediating these vasorelax-

Fig. 6. Real-time quantitative PCR analysis comparing *Kcne1–5* mRNA expression in male and female *Kcne4*^{+/+} mice (**a**) and *Kcne4*^{+/+} compared to *Kcne4*^{-/-} in male (**b**) and female (**c**) mice (n = 5–6).

ant effects in the mesenteric arteries of both males and females. Although Kv7 channel function was unaffected in the female *Kcne4*^{-/-} mice, it is interesting that the responses to isoprenaline were equally impaired in the male and female *Kcne4*^{-/-} mice. These data suggest the β -adrenoceptor intracellular signaling relies on KCNE4 to activate the vascular Kv7 channels. Interestingly, the KCNE4 protein contains a consensus cAMP phosphorylation site and a putative $\beta\gamma$ -binding site [27] in close proximity to one another (amino acids 61–68; accession EAW70810.1) on the C-terminus; however, understanding the mechanisms by which the KCNE4 subunit interacts with Kv7.4 to facilitate the receptor-mediated vasorelaxant effects requires future attention.

Recently, we found that the KCNE4 β -subunit is positively regulated by DHT in the mouse cardiac myocytes [23]. Male *Kcne4*^{+/+} mice were shown to have a higher Kv current density compared with their age-matched, pre-

menopausal female counterparts. The *Kcne4*^{-/-} mice (males and females) had the same Kv current density as the *Kcne4*^{+/+} females. *Kcne4* deletion did not alter ventricular repolarization in premenopausal female mice, but it delayed ventricular repolarization in postmenopausal female mice, which, unlike premenopausal female mice, showed ventricular *Kcne4* expression levels equivalent to those of adult male mice [23]. Interestingly, in this study, we found no sex-dependent differences in the *Kcne4*^{+/+} mouse mesenteric artery responses to methoxamine, isoprenaline or ML213, even though *Kcne4* mRNA expression in the male mouse mesenteric arteries was more than double that of the females. Although posttranscriptional modifications are possible, when taken together with the finding that the female *Kcne4*^{-/-} mice showed similar responses to ML213 as their *Kcne4*^{+/+} counterparts, these data suggest that female mice are less reliant on KCNE4 to regulate Kv7 channels in the vascu-

Fig. 7. Effects of sex and *Kcne4* deletion on mouse mesenteric artery Kv7.4 protein expression. **a** Exemplar Western blots showing Kv7.4 (upper) and GAPDH (lower) expression in mouse mesenteric arteries. **b** Mean Kv7.4 band densities, normalized to same-lane GAPDH band densities, for blots as in **a** ($n = 4$ –5 mice per group, as indicated in parentheses). * $p < 0.05$, ** $p < 0.01$, ANOVA with the Tukey HSD test.

lature, possibly relying on other *Kcne* isoforms. Aside from *Kcne4*, the only *Kcne* isoform to exhibit sex-dependent expression levels here was *Kcne1*. As reported previously [5, 28], *Kcne1* mRNA was not detected here in the mesenteric arteries of male mice; however, it was expressed in some of the female mesenteric arteries (fig. 5a). KCNE1 reportedly increases Kv7.4 [5, 29] and Kv7.5 [30] currents in vitro. Based on these in vitro findings, one might expect KCNE1 to increase the function of the vascular Kv7.4/Kv7.5 channel, which could explain why the vascular responses to methoxamine and ML213 were normal in the female *Kcne4*^{-/-} mice. The effect of KCNE1 on the Kv7.4/7.5 heteromeric channel that is likely to be found in the vascular smooth muscle is yet to be determined; however, given the findings presented in this paper, the role of KCNE1 in the vasculature may be worth further exploration. Furthermore, the qPCR data in this study identified *Kcne3* mRNA as predominant in both

male and female mesenteric arteries, and upregulated by approximately 50% in males in response to *Kcne4* deletion. As the *Kcne3* expression levels in *Kcne4*^{+/+} and *Kcne4*^{-/-} mice were not sex-dependent, we consider it unlikely that *Kcne3* underlies sex-specific differences in mouse mesenteric artery reactivity. However, we previously found that *Kcne3* was upregulated in the gastric epithelium in response to germline *Kcne2* deletion and that this resulted in mistrafficking of Kcnq1 to the basolateral side of parietal cells, where it could not perform its normal function of recycling K⁺ back into the stomach lumen [31]. In addition, KCNE3 suppresses currents produced by Kv7.4 [29]. It will, therefore, be interesting in the future to examine the possible functional effects of vascular *Kcne3* remodeling in response to *Kcne4* deletion.

Vascular diseases are, in general, more common in males than females, but the precise mechanisms responsible for these differences are incompletely understood. The effects of sex hormones in the cardiovascular system are thought to be one important determinant of these differences [17, 18, 22]. One factor that has been extensively researched in the vasculature is the male sex hormone, testosterone [22, 32]. Together with higher levels of circulating testosterone, males have a higher incidence of vascular disease; however, reduced testosterone levels in aging men are known to increase the risk of developing cardiovascular diseases [33–35]. Therefore, it is still unclear whether testosterone is a positive or negative factor in determining susceptibility to cardiovascular diseases.

The expression levels of KCNE transcripts and protein can be subject to regulatory influences, particularly by androgens such as estrogen and testosterone, in at least some tissues. As mentioned, *Kcne4* transcript is positively regulated by DHT in mouse ventricular myocardium to the extent that it dictates sex-specific functional differences in ventricular repolarization currents [23]. *Kcne2* is upregulated by estrogen via a genomic mechanism [36], and may act as the female equivalent of *Kcne4* with respect to aspects of ventricular function, as both *Kcne2* and *Kcne4* regulate Kv4.2 and Kv1.5 channels in mouse ventricular myocytes [23, 37]. *Kcne3* protein expression in rat colonic crypts is reportedly reduced by estrogen via a nongenomic action involving *Kcne3* serine 82 [38], a PKC phosphorylation site [39]. This activity may affect colonic Kv7.1-KCNE3 current properties and density during the estrus cycle [38]. In future work, it will be important to define the different roles and interaction partners of all the KCNE subunits in the regulation of smooth muscle contractility, and investigate the possible role of sex hormones in regulating the vascular expression of

both the KCNEs and their α -subunit partners, and the potentially dynamic effects of this on vascular reactivity. An important aspect of this will be to determine how the female mesenteric arteries compensate for loss of *Kcne4*; is it simply less important in females and so compensation is easier, for example? Our Kv7.4 Western blot results may provide some clues. These data suggest that Kv7.4 protein expression or stability in mouse mesenteric artery is reduced by *Kcne4* deletion, which is reminiscent of recent findings that transient KCNE4 knockdown in the rat mesenteric artery impairs Kv7.4 surface expression [2]. In addition, Kv7.4 protein was higher in mesenteric arteries isolated from female mice compared to those from males. This was despite the lack of a sex-dependent difference in *KCNQ4* transcript (which encodes Kv7.4) expression, an apparent discrepancy that could be explained by, for example, the posttranslational effects of hormones, as previously reported for KCNE3 in the rat colon [39], described above. These findings raise the interesting possibility that, in contrast to males, female mouse mesenteric arteries maintain enough Kv7.4 protein to ensure proper Kv7 channel function following the destabilizing effects of *Kcne4* deletion. Regardless, this study emphasizes that, in future experiments, both sexes should be considered when determining the vascular roles of KCNE subunits.

Previously, Zhou et al. [40] showed that testosterone deprivation reduced Kv currents in rat aortic myocytes, which was associated with decreased Kv1.5 expression. We previously found that castration DHT-dependently reduced ventricular myocyte *Kcne4* expression, and that lower *Kcne4* expression was associated with lower Kv1.5 current [23]. It is possible that *Kcne4* deletion could affect ion channels other than Kv7.4 in the mesenteric artery, including Kv1.5; however, in this study, we focused on its regulation of the Kv7 channels, as this interaction had been previously reported in the vasculature. It will be important in future experiments to determine other possible α -subunits with which KCNE4 might coassemble, to fully

understand the impact of DHT (or other androgen)-mediated changes in KCNE4 expression.

In summary, our findings highlight fundamental sex differences in the sensitivity of expression changes in KCNE4 in the vasculature, with important implications for vascular reactivity. These findings are in line with our previous studies that showed KCNE4 is regulated by DHT in mice, and that KCNE4 is an important regulator of Kv7.4 vascular function in male rats. We also provide novel evidence that KCNE4 is involved in the Kv7 channel recruitment following isoprenaline-induced activation of β -adrenoceptors. Given the important regulatory role of the KCNE family, not only with respect to Kv7 channels [41] but also many other Kv channel subtypes [42, 43], pacemaker channels [44] and even reportedly L-type Ca^{2+} channels [45], much more work needs to be done to elucidate the functional role and the implications of sex-dependent differences in the expression of the KCNE ancillary subunits in blood vessels.

Acknowledgements

We are extremely grateful to Carsten Thorndahl and James Lee Park from Danish Myo Technology for loaning a myograph to Dr. Abbott's laboratory, and to Drs. Steve Goldstein and Daniel Levy for the gift of the *Kcne4*^{-/-} mouse line. We thank Dr. Shawn Crump, Dara Nguyen and Thuyen Nguyen for technical assistance.

T.A.J. received funding from the People Programme (Marie Curie Actions) of the European Union's Seventh Framework Programme (FP7/2007–2013) under REA grant agreement No. 608765 and a grant from the Lundbeck Foundation. The Lundbeck Foundation also provided an additional travel grant for T.A.J. to travel to The University of California, Irvine, to carry out many of these experiments. G.W.A. was supported by the US National Institutes of Health (GM115189).

Disclosure Statement

The authors have no conflicts of interest to report pertaining to this study.

References

- Abbott GW, Goldstein SA: A superfamily of small potassium channel subunits: form and function of the MinK-related peptides (MiRPs). *Q Rev Biophys* 1998;31:357–398.
- Jepps TA, Carr G, Lundegaard PR, Olesen SP, Greenwood IA: Fundamental role for the KCNE4 ancillary subunit in Kv7.4 regulation of arterial tone. *J Physiol* 2015;593:5325–5340.
- Ng FL, Davis AJ, Jepps TA, Harhun MI, Yeung SY, Wan A, Reddy M, Melville D, Nardi A, Khong TK, Greenwood IA: Expression and function of the K⁺ channel KCNQ genes in human arteries. *Br J Pharmacol* 2011;162:42–53.
- Stott JB, Jepps TA, Greenwood IA: K(V)7 potassium channels: a new therapeutic target in smooth muscle disorders. *Drug Discov Today* 2014;19:413–424.

- 5 Jepps TA, Bentzen BH, Stott JB, Povstyan OV, Sivaloganathan K, Dalby-Brown W, Greenwood IA: Vasorelaxant effects of novel Kv7.4 channel enhancers ML213 and NS15370. *Br J Pharmacol* 2014;171:4413–4424.
- 6 Yeung SY, Pucovsky V, Moffatt JD, Saldanha L, Schwake M, Ohya S, Greenwood IA: Molecular expression and pharmacological identification of a role for Kv7 channels in murine vascular reactivity. *Br J Pharmacol* 2007;151:758–770.
- 7 Chadha PS, Jepps TA, Carr G, Stott JB, Zhu HL, Cole WC, Greenwood IA: Contribution of kv7.4/kv7.5 heteromers to intrinsic and calcitonin gene-related peptide-induced cerebral reactivity. *Arterioscler Thromb Vasc Biol* 2014;34:887–893.
- 8 Brueggemann LI, Mackie AR, Cribbs LL, Fredda J, Tripathi A, Majetschak M, Byron KL: Differential protein kinase C-dependent modulation of Kv7.4 and Kv7.5 subunits of vascular Kv7 channels. *J Biol Chem* 2014;289:2099–2111.
- 9 Zhong XZ, Harhun MI, Olesen SP, Ohya S, Moffatt JD, Cole WC, Greenwood IA: Participation of KCNQ (Kv7) potassium channels in myogenic control of cerebral arterial diameter. *J Physiol* 2010;588:3277–3293.
- 10 Jepps TA, Chadha PS, Davis AJ, Harhun MI, Cockerill GW, Olesen SP, Hansen RS, Greenwood IA: Downregulation of Kv7.4 channel activity in primary and secondary hypertension. *Circulation* 2011;124:602–611.
- 11 Chadha PS, Zunke F, Zhu HL, Davis AJ, Jepps TA, Olesen SP, Cole WC, Moffatt JD, Greenwood IA: Reduced KCNQ4-encoded voltage-dependent potassium channel activity underlies impaired β -adrenoceptor-mediated relaxation of renal arteries in hypertension. *Hypertension* 2012;59:877–884.
- 12 Khanamiri S, Soltysinska E, Jepps TA, Bentzen BH, Chadha PS, Schmitt N, Greenwood IA, Olesen SP: Contribution of Kv7 channels to basal coronary flow and active response to ischemia. *Hypertension* 2013;62:1090–1097.
- 13 Stott JB, Barrese V, Jepps TA, Leighton EV, Greenwood IA: Contribution of Kv7 channels to natriuretic peptide mediated vasodilation in normal and hypertensive rats. *Hypertension* 2015;65:676–682.
- 14 Stott JB, Povstyan OV, Carr G, Barrese V, Greenwood IA: G-protein $\beta\gamma$ subunits are positive regulators of Kv7.4 and native vascular Kv7 channel activity. *Proc Natl Acad Sci USA* 2015;112:6497–6502.
- 15 Mani BK, Robakowski C, Brueggemann LI, Cribbs LL, Tripathi A, Majetschak M, Byron KL: Kv7.5 potassium channel subunits are the primary targets for PKA-dependent enhancement of vascular smooth muscle Kv7 currents. *Mol Pharmacol* 2016;89:323–334.
- 16 Vassalle C, Simoncini T, Chedraui P, Pérez-López FR: Why sex matters: the biological mechanisms of cardiovascular disease. *Gynecol Endocrinol* 2012;28:746–751.
- 17 Gonzales RJ: Androgens and the cerebrovasculature: modulation of vascular function during normal and pathophysiological conditions. *Pflugers Arch* 2013;465:627–642.
- 18 Rossi P, Francès Y, Kingwell BA, Ahimastos AA: Gender differences in artery wall biomechanical properties throughout life. *J Hypertens* 2011;29:1023–1033.
- 19 Sandberg K, Ji H: Sex differences in primary hypertension. *Biol Sex Differ* 2012;3:7.
- 20 Doumas M, Papademetriou V, Faselis C, Kokkinos P: Gender differences in hypertension: myths and reality. *Curr Hypertens Rep* 2013;15:321–330.
- 21 Dasinger JH, Alexander BT: Gender differences in developmental programming of cardiovascular diseases. *Clin Sci (Lond)* 2016;130:337–348.
- 22 Lopes RA, Neves KB, Carneiro FS, Tostes RC: Testosterone and vascular function in aging. *Front Physiol* 2012;3:89.
- 23 Crump SM, Hu Z, Kant R, Levy DI, Goldstein SA, Abbott GW: Kcne4 deletion sex- and age-specifically impairs cardiac repolarization in mice. *FASEB J* 2016;30:360–369.
- 24 Livak KJ, Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 2001;25:402–408.
- 25 Mackie AR, Brueggemann LI, Henderson KK, Shiels AJ, Cribbs LL, Scrogin KE, Byron KL: Vascular KCNQ potassium channels as novel targets for the control of mesenteric artery constriction by vasopressin, based on studies in single cells, pressurized arteries, and in vivo measurements of mesenteric vascular resistance. *J Pharmacol Exp Ther* 2008;325:475–483.
- 26 Mills TA, Greenwood SL, Devlin G, Shweikh Y, Robinson M, Cowley E, Hayward CE, Cottrell EC, Tropea T, Brereton MF, Dalby-Brown W, Wareing M: Activation of KV7 channels stimulates vasodilatation of human placental chorionic plate arteries. *Placenta* 2015;36:638–644.
- 27 Yevenes GE, Moraga-Cid G, Guzmán L, Haeger S, Oliveira L, Olate J, Schmalzing G, Aguayo LG: Molecular determinants for G protein betagamma modulation of ionotropic glycine receptors. *J Biol Chem* 2006;281:39300–39307.
- 28 Moreno-Domínguez A, Ciudad P, Miguel-Velado E, López-López JR, Pérez-García MT: De novo expression of Kv6.3 contributes to changes in vascular smooth muscle cell excitability in a hypertensive mice strain. *J Physiol* 2009;587:625–640.
- 29 Strutz-Seeböhm N, Seeböhm G, Fedorenko O, Baltaev R, Engel J, Knirsch M, Lang F: Functional coassembly of KCNQ4 with KCNE-beta-subunits in *Xenopus* oocytes. *Cell Physiol Biochem* 2006;18:57–66.
- 30 Roura-Ferrer M, Etzebarria A, Solé L, Oliveras A, Comes N, Villarreal A, Felipe A: Functional implications of KCNE subunit expression for the Kv7.5 (KCNQ5) channel. *Cell Physiol Biochem* 2009;24:325–334.
- 31 Roepke TK, King EC, Purtell K, Kanda VA, Lerner DL, Abbott GW: Genetic dissection reveals unexpected influence of β subunits on KCNQ1 K⁺ channel polarized trafficking in vivo. *FASEB J* 2011;25:727–736.
- 32 Reckelhoff JF: Sex steroids, cardiovascular disease, and hypertension: unanswered questions and some speculations. *Hypertension* 2005;45:170–174.
- 33 Ruige JB, Mahmoud AM, De Bacquer D, Kaufman JM: Endogenous testosterone and cardiovascular disease in healthy men: a meta-analysis. *Heart* 2011;97:870–875.
- 34 Fukai S, Akishita M, Yamada S, Ogawa S, Yamaguchi K, Kozaki K, Toba K, Ouchi Y: Plasma sex hormone levels and mortality in disabled older men and women. *Geriatr Gerontol Int* 2011;11:196–203.
- 35 Hyde Z, Norman PE, Flicker L, Hankey GJ, Almeida OP, McCaul KA, Chubb SA, Yeap BB: Low free testosterone predicts mortality from cardiovascular disease but not other causes: the Health in Men Study. *J Clin Endocrinol Metab* 2012;97:179–189.
- 36 Kundu P, Ciobotaru A, Foroughi S, Toro L, Stefani E, Eghbali M: Hormonal regulation of cardiac KCNE2 gene expression. *Mol Cell Endocrinol* 2008;292:50–62.
- 37 Roepke TK, Kontogeorgis A, Ovanes C, Xu X, Young JB, Purtell K, Goldstein SA, Christini DJ, Peters NS, Akar FG, Gutstein DE, Lerner DJ, Abbott GW: Targeted deletion of *kcne2* impairs ventricular repolarization via disruption of I(K_{slow}) and I(to,f). *FASEB J* 2008;22:3648–3660.
- 38 Alzamora R, O'Mahony F, Bustos V, Rapetti-Mauss R, Urbach V, Cid LP, Sepulveda FV, Harvey BJ: Sexual dimorphism and oestrogen regulation of KCNE3 expression modulates the functional properties of KCNQ1 K(+) channels. *J Physiol* 2011;589:5091–5107.
- 39 Abbott GW, Butler MH, Goldstein SA: Phosphorylation and protonation of neighboring MiRP2 sites: function and pathophysiology of MiRP2-Kv3.4 potassium channels in periodic paralysis. *FASEB J* 2006;20:293–301.
- 40 Zhou P, Fu L, Pan Z, Ma D, Zhang Y, Qu F, Guo L, Cao J, Gao Q, Han Y: Testosterone deprivation by castration impairs expression of voltage-dependent potassium channels in rat aorta. *Eur J Pharmacol* 2008;593:87–91.
- 41 Abbott GW: Biology of the KCNQ1 potassium channel. *New J Sci* DOI: 10.1155/2014/237431.
- 42 Abbott GW: The KCNE2 K(+) channel regulatory subunit: ubiquitous influence, complex pathobiology. *Gene* 2015;569:162–172.
- 43 Abbott GW: KCNE1 and KCNE3: the yin and yang of voltage-gated K(+) channel regulation. *Gene* 2016;576:1–13.
- 44 Qu J, Kryukova Y, Potapova IA, Doronin SV, Larsen M, Krishnamurthy G, Cohen IS, Robinson RB: MiRP1 modulates HCN2 channel expression and gating in cardiac myocytes. *J Biol Chem* 2004;279:43497–43502.
- 45 Liu W, Deng J, Wang G, Zhang C, Luo X, Yan D, Su Q, Liu J: KCNE2 modulates cardiac L-type Ca channel. *J Mol Cell Cardiol* 2014;72:208–218.