Impact of Diet and/or Exercise Intervention on Infrapatellar Fat Pad Morphology: Secondary Analysis from the Intensive Diet and Exercise for Arthritis (IDEA) Trial

Aarón Leonardo Pogacnik Murillo^a Felix Eckstein^a Wolfgang Wirth^a
Daniel Beavers^b Richard F. Loeser^c Barbara J. Nicklas^c Shannon L. Mihalko^d
Gary D. Miller^d David J. Hunter^{f, g} Stephen P. Messier^d

^aInstitute of Anatomy, Paracelsus Medical University Salzburg and Nuremberg, Salzburg, Austria; ^bDepartment of Biostatistical Sciences and ^cSection on Gerontology and Geriatric Medicine, Wake Forest School of Medicine, and ^dDepartment of Health and Exercise Science, Wake Forest University, Winston-Salem, NC, and ^eDivision of Rheumatology, Allergy and Immunology and Thurston Arthritis Research Center, University of North Carolina School of Medicine, Chapel Hill, NC, USA; ^fRheumatology Department, Royal North Shore Hospital, and ^gInstitute of Bone and Joint Research, Kolling Institute, University of Sydney, Sydney, NSW, Australia

Keywords

Infrapatellar fat pad · Quantitative imaging · Weight loss · Knee osteoarthritis

Abstract

Objectives: The infrapatellar fat pad (IPFP) represents intra-articular adipose tissue that may contribute to intra-articular inflammation and pain by secretion of proinflammatory cytokines. Here we examined the impact of weight loss by diet and/or exercise interventions on the IPFP volume. **Methods:** Intensive Diet and Exercise for Arthritis (IDEA) was a single-blinded, single-center, 18-month, prospective, randomized controlled trial that enrolled 454 overweight and obese older adults with knee pain and radiographic osteoarthritis. Participants were randomized to 1 of 3 groups: exercise-only control (E), diet-induced weight loss (D), and diet-induced weight loss + exercise (D+E). In a subsample ($n = 106$; E: $n = 36$, D: $n = 35$, and D+E: $n = 35$), magnetic resonance images were acquired at baseline and at the 18-month follow-up, from which we analyzed IPFP volume, surface areas, and thickness in this secondary analysis. **Results:** The average weight loss amounted to 1.0% in the E group, 10.5% in the D

group, and 13.0% in the D+E group. A significant ($p < 0.01$) reduction in IPFP volume was observed in the E (2.1%), D (4.0%), and D+E (5.2%) groups. The IPFP volume loss in the D+E group was significantly greater than that in the E group ($p < 0.05$) when not adjusting for parallel comparisons. Across intervention groups, there were significant correlations between IPFP volume change, individual weight loss ($r = 0.40$), and change in total body fat mass (dual-energy X-ray absorptiometry; $r = 0.44$, $n = 88$) and in subcutaneous thigh fat area (computed tomography; $r = 0.32$, $n = 82$). **Conclusions:** As a potential link between obesity and knee osteoarthritis, the IPFP was sensitive to intervention by diet and/or exercise, and its reduction was correlated with changes in weight and body fat.

© 2017 S. Karger AG, Basel

These data were submitted as a diploma thesis entitled “Impact of Diet and/or Exercise Intervention on Infrapatellar Fat Pad Morphology – A Secondary Analysis from a Randomized Controlled Trial” by Aarón Leonardo Pogacnik Murillo in April 2015 to obtain the academic degree of Medical Doctor (Dr. med. univ.) at Paracelsus Medical University, Salzburg, Austria. This diploma thesis is publicly available at http://www.rheumatologie.at/pdf/Pogacnik_Aaron_Diplomarbeit.pdf.

Abbreviations used in this paper

CT	computed tomography
D	diet intervention group
D+E	diet-plus-exercise intervention group
DXA	dual-energy X-ray absorptiometry
E	exercise intervention group
IDEA	Intensive Diet and Exercise for Arthritis
IL	interleukin
IPFP	infrapatellar fat pad
MR	magnetic resonance
OA	osteoarthritis

Introduction

Knee osteoarthritis (OA) represents one of the leading causes of chronic disability among the elderly population [Guccione et al., 1994] and it is known to be associated with structural alterations in most articular and para-articular tissues [Poole, 2012]. Obesity is one of the most important risk factors for knee OA and a reduced quality of life [Losina et al., 2011], despite being mostly preventable. It has been recently suggested that the obesity-related risk of knee OA incidence and progression may be conveyed not only by biomechanical factors but also by endocrine and inflammatory mechanisms [Issa and Griffin, 2012; Berenbaum et al., 2013; Pogacnik Murillo, 2015]. Adipokines, a group of adipocyte-derived cytokines, are thought to mediate intra-articular inflammation and upregulation of cartilage matrix degradation [Issa and Griffin, 2012; Richter et al., 2015].

The infrapatellar fat pad (IPFP) represents intra-articular (i.e., intra-capsular, extra-synovial) adipose tissue [Hoffa, 1904; Haug, 2014], with its macroscopic anatomy having been previously examined in relation to knee function, pathology, and surgical approaches by Gallagher et al. [2005]. These authors [Gallagher et al., 2005] found the IPFP to be constant in shape, consisting of a central body with medial and lateral extensions. More recently, the IPFP also has been identified as a source of leptin and other proinflammatory cytokines [Gegout et al., 2008; Distel et al., 2009; Gandhi et al., 2011; Klein-Wieringa et al., 2011; Hui et al., 2012]. In knee OA, the IPFP was found to yield high expression levels of enzymes that are involved in fat metabolism [Gandhi et al., 2011], suggesting that it represents a link between obesity and knee OA. The Intensive Diet and Exercise for Arthritis (IDEA) trial [Messier et al., 2009] showed that an 18-month diet-induced weight loss plus exercise resulted in a sig-

nificant reduction in knee symptoms, improvement in knee function, and better physical health-related quality-of-life scores compared to exercise alone [Messier et al., 2013]. This improvement was associated with a significant reduction in the mechanistic primary outcomes, i.e., knee compressive forces and systemic inflammatory marker levels, specifically interleukin (IL)-6 [Messier et al., 2013].

Few studies thus far have quantitatively analyzed IPFP morphology. In a recent paper, Diepold et al. [2015] reported that men displayed a significantly greater ratio of IPFP volume/body weight than women, similar amounts of inter-muscular fat, and strikingly less subcutaneous fat than women. Studies examining the relationship between IPFP volume and knee pain or radiographic OA status have provided partially contradictory results [Chuckpaiwong et al., 2010; Han et al., 2014; Cai et al., 2015; Cowan et al., 2015; Eckstein et al., 2015; Pan et al., 2015; Steidle-Kloc et al., 2015; Teichtahl et al., 2015]. Although the IPFP has been reported to be preserved under starvation conditions [Ioan-Facsinay and Kloppenburg, 2013], there appears to be a dynamic window, as a longitudinal increase in IPFP volume was reported in mice with high-fat feeding and weight gain [Chang et al., 2011]. However, to date, no study has examined the relationship between IPFP volume and weight change in humans, and no interventional trial has studied the sensitivity of IPFP morphology to exercise and/or weight loss. Therefore, in this secondary analysis of the IDEA trial [Messier et al., 2013], we examined whether an 18-month diet-induced weight loss (with and without exercise) was associated with changes in IPFP morphology in patients with knee pain and mild-to-moderate OA. Further, we determined whether changes in IPFP morphology differed between groups with exercise intervention only (E), diet-induced weight loss intervention only (D), and a combined diet-induced weight loss + exercise intervention (D+E). Finally, we examined whether changes in IPFP volume correlated with changes in body weight and fat, independently of the intervention.

Materials and Methods

Study Population

IDEA was a single blind, single-center, 18-month, randomized, controlled trial [Messier et al., 2009, 2013], which was conducted from July 2006 to June 2011 at Wake Forest University and the Wake Forest School of Medicine, Winston-Salem, NC, USA. The study was approved by the Human Subjects Institutional Review Board of Wake Forest Health Sciences. Informed consent was obtained in writing from all participants (trial registration No. NCT00381290) [Messier et al., 2009, 2013].

Fig. 1. a Sagittal MR image showing the segmentation of the IPFP anterior surface (red, facing the patellar ligament) and the posterior surface (blue, facing the inferior patellar pole, the distal femur, and the proximal tibia). **b** Three-dimensional reconstruction of the IPFP (posterior view). The anterior surface area is colored in red and the posterior (femoral and tibial) surface area in blue. MR, magnetic resonance; IPFP, infrapatellar fat pad.

The IDEA trial included 454 ambulatory, community-dwelling persons aged ≥ 55 years with: (1) Kellgren-Lawrence grade [Kellgren and Lawrence, 1957] 2–3 (mild to moderate) radiographic tibiofemoral OA or tibiofemoral plus patellofemoral OA of one or both knees; (2) pain on most days, due to knee OA; (3) a BMI ranging from 27 to 41; and (4) a sedentary lifestyle, i.e., <30 min/week of formal exercise in the past 6 months.

Participants were recruited from the community over a 37-month period (November 2006 to December 2009) [Messier et al., 2009, 2013]. A stratified-block randomization method assigned all eligible persons to 1 of the following 3 intervention arms, stratified by BMI and gender: exercise-only control (E), diet-induced weight loss only (D), and diet-induced weight loss plus exercise (D+E) [Messier et al., 2009, 2013]. The E group was designated as the comparison (control) group because previous work indicated that aerobic walking or resistance training should be part of the standard of care for knee OA patients [Ettinger et al., 1997], and because the original study [Messier et al., 2013] was designed [Messier et al., 2009] to test whether diet alone, or diet in combination with exercise, leads to greater improvements than the standard of care (exercise) alone. Detailed descriptions of the diet-induced weight loss and exercise interventions, the trial design and rationale, and the primary outcomes have been reported elsewhere [Messier et al., 2009, 2013]. In brief, the diet intervention was initially based on partial meal replacements and one daily meal of 500–750 kcal. The exercise intervention consisted of aerobic walking (15 min), strength training (20 min), another aerobic phase (15 min), and a cool-down (10 min), and it was conducted 3 times weekly, for 1 h, over the entire 18-month observation period [Messier et al., 2009, 2013].

Due to budget restrictions, magnetic resonance (MR) images were obtained on a randomly selected subsample of 106 participants at baseline and at the 18-month follow-up [Hunter et al., 2015]. The sample size per group was: E, $n = 36$; D, $n = 35$; and D+E, $n = 35$.

Image Acquisition and Analysis of the IPFP

A 1.5-T (SIGNA HDx; General Electric Medical Systems, Milwaukee, WI, USA) scanner and an extremity coil were used to obtain MR images of the most affected knee [Messier et al., 2013; Hunter et al., 2015]. Segmentation of the IPFP was performed us-

ing a sagittal non-fat-suppressed T1-weighted spin echo sequence (time of repetition, 600 ms; time of echo, 11 ms; contiguous slices with a thickness of 4.5 mm; in-plane resolution, 0.625×0.625 mm; field of view, 16 cm; and image matrix, 256×256 pixels). All segmentations in this study were performed by the first author of this study (A.L.P.M.), who was trained using standardized test data sets [Steidle-Kloc et al., 2016]. Baseline and follow-up images were read as pairs, but the reader (A.L.P.M.) was blinded to the time of image acquisition and the type of intervention during segmentation. Brightness, intensity, contrast, and gray value limits were adjusted manually in each image to warrant optimal contrast between the IPFP and surrounding tissue, with the reader processing all slices (from medial to lateral) that clearly depicted the IPFP. By applying different labels, the reader manually traced the anterior border of the IPFP (the one facing the patellar ligament) and the posterior border (the one facing the knee joint) (Fig. 1a). Clefts (supra- and infrahoffatic recesses) [Roemer et al., 2016] located in the periphery of the IPFP, and the anterior intermeniscal ligament, were excluded from the segmentation. However, small alterations completely surrounded by adipose tissue were included in the volume segmentations. Based on these segmentations, the IPFP volume, the size of the anterior and posterior surface areas, and the mean and maximal depth from the anterior surface to the posterior surface were computed using custom image analysis software (Fig. 1b) [Diepold et al., 2015; Steidle-Kloc et al., 2016], with the intra- and interobserver reliability having been reported [Steidle-Kloc et al., 2016]. For further methodological details, please see the paper by Pogacnik Murillo [2015].

Analysis of Body and Thigh Composition

The BMI of all of the participants was calculated as mass (measured in kg on a standard calibrated scale) divided by body height squared (measured in m). The whole body fat mass was measured at baseline and at the 18-month follow-up using dual-energy X-ray absorptiometry (DXA) in a subsample of the participants ($n = 88$). DXA scans were obtained with a fan-beam scanner (Delphi ATM; Hologic, Waltham, MA, USA) using the manufacturer's recommendations for patient positioning, scan protocols, and scan analysis, with the reliability (coefficients of variation of repeated measurements) being reported as 1.2% for whole-body fat mass [Messier et al., 2009]. Further, baseline and 18-month follow-up

Table 1. Baseline characteristics of the E, D, and D+E intervention groups

Intervention group	All (<i>n</i> = 106)	E (<i>n</i> = 36)	D (<i>n</i> = 35)	D+E (<i>n</i> = 35)
Demographics				
BMI	33.9±3.8	33.7±3.7	34.3±4.1	33.7±3.7
Body weight, kg	93.5±14.0	91.6±13.0	96.9±15.7	91.9±13.0
Sex				
Female	77 (72.6)	27 (75.0)	24 (68.6)	26 (74.3)
Male	29 (27.4)	9 (25.0)	11 (31.4)	9 (25.7)
IPFP parameter				
Volume, cm ³	26.7±7.0	25.3±6.3	28.0±7.6	26.6±6.9
Anterior surface area, cm ²	24.6±5.6	24.1±5.5	25.3±6.2	24.4±5.3
Posterior surface area, cm ²	37.3±7.8	35.9±7.4	38.7±8.9	37.4±6.9
Mean thickness, mm	6.5±0.6	6.4±0.5	6.6±0.7	6.5±0.7
Maximal thickness, mm	15.0±1.6	14.9±1.5	15.3±1.6	14.9±1.6

Values are means ± SD or *n* (%). IPFP, infrapatellar fat pad; E, exercise intervention group; D, diet intervention group; D+E, diet-plus-exercise intervention group.

Table 2. Changes in BMI, body weight, and IPFP morphology in the E, D, and D+E intervention groups between baseline and the 18-month follow-up

Intervention group	E (<i>n</i> = 36)	D (<i>n</i> = 35)	D+E (<i>n</i> = 35)
Anthropometrics			
BMI	−0.26 (−0.74 to 0.22)	−3.45 (−4.46 to −2.43)	−4.31 (−5.43 to −3.20)
Body weight, kg	−0.89 (−2.16 to 0.39)	−10.18 (−13.16 to −7.21)	−11.93 (−14.86 to −9.01)
IPFP parameter			
Volume, mm ³	−704 (−1,217 to −190) ^a	−1,074 (−1,607 to −540) ^b	−1,462 (−1,994 to −930) ^b
Anterior surface area, mm ²	−20.7 (−73.5 to 32.1)	−25.4 (−78.5 to 27.6)	−71.4 (−120.0 to −17.8) ^a
Posterior surface area, mm ²	−90.3 (−156.7 to −24.0) ^a	−141.9 (−210.2 to −73.6) ^b	−182.0 (−250.5 to −113.5) ^b
Mean thickness, mm	0.00 (−0.09 to 0.10)	0.00 (−0.09 to 0.10)	−0.01 (−0.10 to 0.09)
Maximal thickness, mm	0.00 (−0.25 to 0.25)	−0.16 (−0.42 to 0.09)	−0.06 (−0.32 to 0.19)

The results for IPFP morphology were adjusted for baseline values of the outcome, baseline BMI, and sex. Values are presented as means (95% CI). IPFP, infrapatellar fat pad; E, exercise intervention group; D, diet intervention group; D+E, diet-plus-exercise intervention group. ^a *p* < 0.01, ^b *p* < 0.001.

computed tomography (CT) scans of the thigh were obtained in a subsample (*n* = 82), and measurements of subcutaneous thigh fat, intramuscular thigh fat, and total thigh fat were obtained [Messier et al., 2009].

Statistical Analysis

Baseline anthropometric measures and quantitative measures of the IPFP are reported as means and SD. Within- and between-group comparisons of longitudinal changes in IPFP morphology were evaluated using ANCOVA, adjusting for baseline values of the outcome, baseline BMI, and sex, but not adjusting for multiple between-group comparisons in this post hoc exploratory analysis. IPFP volume was considered the primary analytic focus of the current analysis, and IPFP anterior surface area, posterior surface area, mean and maximal depths were exploratory measures. Cor-

relations between IPFP change and BMI change, fat mass change (DXA) or thigh composition change (CT) were studied across the entire sample, independently of the randomized controlled trial design and the specific type of intervention, using Spearman correlations. The level of significance for all comparisons was set at 0.05. All analyses were performed using SAS v9.4 (SAS Institute, Cary, NC, USA).

Results

There were no significant differences in age, sex, BMI, or IPFP morphology measures between the 3 intervention groups at baseline (Table 1). There was a significant

Table 3. Pairwise comparisons between intervention groups for mean changes in IPFP morphology between baseline and the 18-month follow-up, adjusted for baseline values of the outcome, baseline BMI, and sex

Intervention group	E vs. D	E vs. D+E	D vs. D+E
<i>IPFP parameter</i>			
Volume, mm ³	370 (−328 to 1,068)	759 (68 to 1,450) ^a	389 (−307 to 1,085)
Anterior surface area, mm ²	4.7 (−65.9 to 75.4)	50.7 (−19.7 to 121.0)	45.9 (−25.1 to 117.0)
Posterior surface area, mm ²	51.6 (−38.6 to 141.7)	91.7 (2.3 to 181.0) ^a	40.1 (−49.8 to 130.1)
Mean thickness, mm	0.00 (−0.13 to 0.13)	0.01 (−0.11 to 0.14)	0.01 (−0.11 to 0.14)
Maximal thickness, mm	0.17 (−0.18 to 0.51)	0.07 (−0.27 to 0.40)	−0.10 (−0.44 to 0.24)

Values in parentheses are 95% CI. E, exercise intervention group; D, diet intervention group; D+E, diet-plus-exercise intervention group; IPFP, infrapatellar fat pad. ^a $p < 0.05$.

Table 4. Spearman correlations between IPFP volume changes and changes in body composition predictors

	<i>n</i>	Correlation	<i>p</i> value
<i>Body or thigh composition predictor</i>			
Weight change (kg)	106	0.40	<0.0001
BMI change	106	0.39	<0.0001
DXA total fat mass change (kg)	88	0.44	<0.0001
CT subcutaneous thigh fat change (cm ³)	82	0.32	0.0030
CT intermuscular thigh fat change (cm ³)	82	0.29	0.0074

Sample sizes (*n*) vary based on the predictor. DXA, dual X-ray absorptiometry; CT, computed tomography; IPFP, infrapatellar fat pad.

($p < 0.01$) reduction in IPFP volume in each of the 3 intervention groups over the 18-month observation period (Table 2; Fig. 2) (E: −2.1%, 95% CI −0.8 to −4.8%; D: −4.0%, 95% CI −1.9 to −5.7%; and D+E: −5.2%, 95% CI −3.5 to −7.5%). As a reference, the mean percent body weight loss from baseline amounted to −1.0% in the E group, −10.5% in the D group, and −13.0% in the D+E group. Changes in kilograms and their 95% CI are shown in Table 2. There was a significant ($p < 0.01$) within-group decrease in IPFP posterior surface area for each intervention group, and a significant ($p < 0.001$) within-group decrease in IPFP anterior surface area in the D+E group; the change in IPFP volume therefore appeared to be driven more by a change in surface areas than by a change in IPFP depth (Table 2). Between-group comparisons revealed a significantly greater reduction in IPFP volume and IPFP posterior surface area in the D+E group than in the E (control) group ($p < 0.05$); other between-group dif-

Fig. 2. Mean change in IPFP volume between baseline and the 18-month follow-up in the E, D, and D+E groups, adjusted for baseline values of the outcome, baseline BMI, and sex. Error bars represent 95% CI. IPFP, infrapatellar fat pad; E, exercise intervention group; D, diet intervention group; D+E, diet-plus-exercise intervention group.

ferences in the longitudinal changes in IPFP morphology did not reach statistical significance (Tables 2, 3).

Across the 3 intervention groups, there was a significant ($p < 0.01$) correlation between IPFP volume change and weight loss ($r = 0.40$), BMI change ($r = 0.39$), total body fat mass change ($r = 0.44$, $n = 88$; DXA), subcutaneous thigh fat change ($r = 0.32$, $n = 82$; CT), and intermuscular thigh fat change ($r = 0.29$, $n = 82$; CT) (Table 4). Across all intervention types, each percent of weight loss was related to a 0.27% reduction in IPFP volume.

Discussion

In this first study investigating the effect of diet-induced weight loss, with and without exercise, on IPFP morphology, we observed a significant reduction of IPFP size in all intervention groups. Further, the findings suggest that the combination of D+E was more effective in reducing IPFP volume than E alone and that, independently of the type of intervention, the reduction in IPFP volume was significantly correlated with loss of body weight and fat mass.

A limitation of the current analysis is that only a subsample of the IDEA cohort was examined [Hunter et al., 2015]; however, the subsample was reasonably representative of the larger sample and sufficient to demonstrate responsiveness of the IPFP to diet and exercise intervention. A strength of the current study is its randomized controlled design, which allowed analysis of the effects of diet-induced weight loss, with and without exercise, on IPFP volume. Further, randomization of the subsample to the different intervention groups appeared to be without bias, as anthropometric and IPFP morphometric baseline values were similar between the intervention groups. Another limitation of the current study is that a standard no-treatment reference group (with neither exercise nor diet intervention) was not available because the intent of the original study was to test whether diet, in combination with exercise or not, was more effective in reducing clinical symptoms than exercise, which is part of the standard of care for people with knee OA, and that no additional reference group with or without a known OA status in other joints, e.g., those of an upper limb, was available.

Interestingly, an exercise intervention alone resulted in a small, but statistically significant, within-group reduction in IPFP volume despite a minimal and statistically nonsignificant reduction in body weight. Diet in combination with exercise was more effective in reducing the IPFP volume despite reductions in body weight being similar in the D and D+E groups. Although the mechanism behind exercise-related effects on the IPFP is unclear, this could potentially be due to modifications in inflammation and edema within the IPFP. These findings indicate that exercise may be effective in modifying IPFP morphology, at least partly independently of weight loss. Further interventional studies should examine whether specific types of exercises are more effective than others in reducing the IPFP volume, whether reduction of the IPFP volume is associated with less inflammatory activity and release of adipokines, and whether modification of

the IPFP (including surgical reduction) is related to joint health.

Previous studies examining the relationship between IPFP size and knee OA have revealed partly contradictory results; Chuckpaiwong et al. [2010] found no differences in IPFP volume between patients with knee OA and healthy controls in a small (and likely underpowered) sample. A smaller IPFP size was found to be associated to cartilage defects compared to the control group, but only in women [Duran et al., 2015]. Furthermore, differences in IPFP volume between patients with cartilage defects and healthy controls disappeared after adjustment for age and BMI [Duran et al., 2015]. The IPFP volume in joints with patellofemoral OA was reported to be greater than in those without patellofemoral OA [Cowan et al., 2015]. A greater IPFP maximal sagittal cross-sectional area, however, appeared to be beneficial in terms of structural changes and knee pain, but these observations were limited to a single slice [Han et al., 2014; Pan et al., 2015; Teichtahl et al., 2015]. In this context, a strength of the present study is that the complete IPFP volume was studied (not only a single slice, a 2-dimensional representation of its tissue) [Han et al., 2014; Pan et al., 2015]. This type of volumetric analysis is reproducible within and between observers [Steidle-Kloc et al., 2016]. Also, volumetric analysis permitted us to examine the responsiveness of different measures of IPFP morphology, indicating that a reduction of the (posterior) surface area may be more sensitive to exercise and diet-induced weight loss than IPFP depth.

The IPFP is a local source of proinflammatory mediators, including a variety of cytokines and adipokines, such as leptin and adiponectin [Gegout et al., 2008; Distel et al., 2009; Gandhi et al., 2011; Klein-Wieringa et al., 2011; Hui et al., 2012]. The inflammatory activity of the IPFP affects synovial inflammation [Eymard et al., 2014], while high levels of inflammatory mediators are associated with cartilage degradation [Ding et al., 2008; Bao et al., 2010; Kang et al., 2010; Stannus et al., 2015]. The close spatial relation of the IPFP to the synovium-lined knee joint cavity suggests that it may more directly influence inflammatory mediator levels involved in the pathogenesis of knee OA than systemic factors [Clockaerts et al., 2010]. Findings that the IPFP of patients with knee OA display high expression levels of enzymes involved in fat metabolism led to the conclusion that the IPFP may be responsive to the overall nutritional state; a reduction of the IPFP volume may thus be associated with a reduction of the IPFP inflammatory activity and the secretion of inflammatory mediators into synovial fluid, and potentially a reduction in cytokines involved in cartilage degradation, inflamma-

tion, and joint pain. The role of the IPFP in inflammatory processes of the knee, however, is still controversial [Berenbaum et al., 2013]. However, conditioned adipose tissue medium of the IPFP of OA patients was found to block catabolic processes induced by IL-1 β in articular cartilage [Bastiaansen-Jenniskens et al., 2012]. Further, local production of inflammatory mediators in the knee was attributed not only to the IPFP but also to osteophytes and the synovium [Presle et al., 2006; Gegout et al., 2008]. No intra-articular inflammatory or other cytokine markers were examined in our current study; hence, follow-up studies, preferably with intra-synovial aspiration, will need to address whether a direct relationship exists between diet- and exercise-induced reduction in IPFP volume and inflammation. Messier et al. [2013] reported, among overweight and obese adults with knee OA, knee compressive forces to be lower after 18 months of diet intervention compared to exercise, concentrations of IL-6 to be lower with diet-and-exercise intervention compared to exercise, and knee pain levels to be lower (and knee function better) with diet-and-exercise intervention compared to both diet and exercise interventions. The D+E group also had better physical health-related quality of life scores than the E group. In extension to the original study [Messier et al., 2013], it was further shown that: (1) weight loss led to a reduction in bone mineral density in the femoral neck and hip, irrespectively of whether the diet was combined with an exercise intervention or not [Beavers et al., 2014]; (2) a high amount of subcutaneous fat had significant associations with knee joint forces similar to those of abdominal fat despite its much smaller volume, and thus this could be an important therapeutic target for patients with knee OA [Messier et al., 2014]; (3) intentional total body fat mass and abdominal fat volume reductions were independently associated with significant reductions in inflammatory serum markers [Beavers et al., 2015]; and (4) despite the potent effects of weight loss on symptoms and mechanistic outcomes, there was no statistically significant difference between the 3 active interventions on radiographic joint space width change, quantitative cartilage thickness loss, or semiquantitative measures of Hoffa-synovitis and joint effusion [Hunter et al., 2015]. However, whether the symptomatic improvement and increased physical health-related quality-of-life scores observed with the diet-and-exercise intervention [Messier et al., 2013] is mediated by longitudinal changes in IPFP volume, or whether the latter should rather be considered an “innocent bystander” in this process, is beyond the focus and design of this report and may be elucidated in further studies.

In summary, the IPFP, a potential link between obesity and knee OA, was sensitive to intervention by diet-induced weight loss and/or exercise in overweight or obese patients with symptomatic knee OA. The reduction in IPFP volume correlated with the individual level of weight and body fat change. Diet in combination with exercise appeared to be more effective in reducing IPFP volume than exercise alone.

Acknowledgment

Support for this study was provided by grants from the National Institutes of Health (R01397 AR052528-01, P30 AG21332, and M01-RR00211) and General Nutrition Centers, Inc. Support for the image analysis of the IPFP was received from the Paracelsus Medical University Forschungsförderungsfond (PMU FFF; R-14/0/061/WIT).

The statistical analysis and writing of this article was independent from and not contingent upon approval from the study sponsors. The funding sources were not involved in the study design or in the statistical analysis of the data.

We would like to thank Anja Ruhdorfer, MD, PhD, Eva Steidle, PhD, and Torben Dannhauer, PhD, at the Institute of Anatomy at Paracelsus Medical University for their support and critical discussion of IPFP segmentation in this study.

Author Contributions

- Substantial contribution to the study conception and design: all authors
- Substantial contribution to the acquisition of data (IPFP): A.L. Pogacnik Murillo, F. Eckstein, and W. Wirth
- Statistical analysis: D. Beavers
- Analysis and interpretation of data: all authors
- Writing of the first manuscript draft: A.L. Pogacnik Murillo and F. Eckstein
- Critical manuscript revision and approval of the final manuscript: all authors

Disclosure Statement

F. Eckstein is CEO of Chondrometrics GmbH, a company providing MR image analysis services to academic researchers and to industry. He has provided consulting services to MerckSerono, Abbvie, and Servier, has prepared educational sessions for Medtronic, and has received research support from Pfizer, Eli Lilly, Merck Serono, GlaxoSmithKline, Centocor R&D, Wyeth, Novartis, Stryker, Abbvie, Kolon, Synarc, Ampio, Orthotrophix, and Samumed.

W. Wirth has part time employment with Chondrometrics GmbH and is coowner of Chondrometrics GmbH.

The other authors declare no competing interests.

References

- Bao, J., W. Chen, J. Feng, P. Hu, Z. Shi, L. Wu (2010) Leptin plays a catabolic role on articular cartilage. *Mol Biol Rep* 37: 3265–3272.
- Bastiaansen-Jenniskens, Y.M., S. Clockaerts, C. Feijt, A.-M. Zuurmond, V. Stojanovic-Susulic, C. Bridts, L. de Clerck, J. DeGroot, J.A.N. Verhaar, M. Kloppenburg, G.J.V.M. van Osch (2012) Infrapatellar fat pad of patients with end-stage osteoarthritis inhibits catabolic mediators in cartilage. *Ann Rheum Dis* 71: 288–294.
- Beavers, D.P., K.M. Beavers, R.F. Loeser, N.R. Walton, M.F. Lyles, B.J. Nicklas, S.A. Shapses, J.J. Newman, S.P. Messier (2014) The independent and combined effects of intensive weight loss and exercise training on bone mineral density in overweight and obese older adults with osteoarthritis. *Osteoarthritis Cartilage* 22: 726–733.
- Beavers, K.M., D.P. Beavers, J.J. Newman, A.M. Anderson, R.F. Loeser, B.J. Nicklas, M.F. Lyles, G.D. Miller, S.L. Mihalko, S.P. Messier (2015) Effects of total and regional fat loss on plasma CRP and IL-6 in overweight and obese, older adults with knee osteoarthritis. *Osteoarthritis Cartilage* 23: 249–256.
- Berenbaum, F., F. Eymard, X. Houard (2013) Osteoarthritis, inflammation and obesity. *Curr Opin Rheumatol* 25: 114–118.
- Cai, J., J. Xu, K. Wang, S. Zheng, F. He, S. Huan, S. Xu, H. Zhang, L. Laslett, C. Ding (2015) Association between infrapatellar fat pad volume and knee structural changes in patients with knee osteoarthritis. *J Rheumatol* 42: 1878–1884.
- Chang, W., J. DeMoe, C. Kent, S. Kovats, P. Gartheiser, S. Doblas, R. Towner, T.M. Griffin (2011) Infrapatellar fat pad hypertrophy without inflammation in a diet-induced mouse model of obesity and osteoarthritis. *Osteoarthritis Cartilage* 19: 66.
- Chuckpaiwong, B., H.C. Charles, V.B. Kraus, F. Guilak, J.A. Nunley (2010) Age-associated increases in the size of the infrapatellar fat pad in knee osteoarthritis as measured by 3T MRI. *J Orthop Res* 28: 1149–1154.
- Clockaerts, S., Y.M. Bastiaansen-Jenniskens, J. Runhaar, G.J. Van Osch, J.F. Van Offel, J.A. Verhaar, L.S. De Clerck, J. Somville (2010) The infrapatellar fat pad should be considered as an active osteoarthritic joint tissue: a narrative review. *Osteoarthritis Cartilage* 18: 876–882.
- Cowan, S.M., H.F. Hart, S.J. Warden, K.M. Crossley (2015) Infrapatellar fat pad volume is greater in individuals with patellofemoral joint osteoarthritis and associated with pain. *Rheumatol Int* 35: 1439–1442.
- Diepold, J., A. Ruhdorfer, T. Dannhauer, W. Wirth, E. Steidle, F. Eckstein (2015) Sex-differences of the healthy infra-patellar (Hoffa) fat pad in relation to intermuscular and subcutaneous fat content – data from the Osteoarthritis Initiative. *Ann Anat* 200: 30–36.
- Ding, C., V. Parameswaran, F. Cicuttini, J. Burgess, G. Zhai, S. Quinn, G. Jones (2008) Association between leptin, body composition, sex and knee cartilage morphology in older adults: the Tasmanian Older Adult Cohort (TASOAC) study. *Ann Rheum Dis* 67: 1256–1261.
- Distel, E., T. Cadoudal, S. Durant, A. Poignard, X. Chevalier, C. Benelli (2009) The infrapatellar fat pad in knee osteoarthritis: an important source of interleukin-6 and its soluble receptor. *Arthritis Rheum* 60: 3374–3377.
- Duran, S., E. Aksahin, O. Kocadal, C.N. Aktekin, O. Hapa, Z.B. Gencturk (2015) Effects of body mass index, infrapatellar fat pad volume and age on patellar cartilage defect. *Acta Orthop Belg* 81: 41–46.
- Eckstein, F., J. Diepold, A. Ruhdorfer, T. Dannhauer, W. Wirth, A. Guermazi (2015) Infra-patellar fat pad morphology and MRI signal distribution in advanced radiographic knee OA – data from the OAI. *Osteoarthritis Cartilage* 23: A219.
- Ettinger, W.H., R. Burns, S.P. Messier, W. Applegate, W.J. Rejeski, T. Morgan, S. Shumaker, M.J. Berry, M. O'Toole, J. Monu, T. Craven (1997) A randomized trial comparing aerobic exercise and resistance exercise with a health education program in older adults with knee osteoarthritis: the Fitness Arthritis and Seniors Trial (FAST). *JAMA* 277: 25–31.
- Eymard, F., A. Pigenet, D. Citadelle, C.-H. Flouzat-Lachaniette, A. Poignard, C. Benelli, F. Berenbaum, X. Chevalier, X. Houard (2014) Induction of an inflammatory and prodegradative phenotype in autologous fibroblast-like synoviocytes by the infrapatellar fat pad from patients with knee osteoarthritis. *Arthritis Rheumatol* 66: 2165–2174.
- Gallagher, J., P. Tierney, P. Murray, M. O'Brien (2005) The infrapatellar fat pad: anatomy and clinical correlations. *Knee Surg Sport Traumatol Arthrosc* 13: 268–272.
- Gandhi, R., M. Takahashi, C. Virtanen, K. Syed, J.R. Davey, N.N. Mahomed (2011) Microarray analysis of the infrapatellar fat pad in knee osteoarthritis: relationship with joint inflammation. *J Rheumatol* 38: 1966–1972.
- Gegout, P.P., P.J. Francin, D. Mainard, N. Presle (2008) Adipokines in osteoarthritis: friends or foes of cartilage homeostasis? *Joint Bone Spine* 75: 669–671.
- Guccione, A.A., D.T. Felson, J.J. Anderson, J.M. Anthony, Y. Zhang, P.W. Wilson, M. Kelly-Hayes, P.A. Wolf, B.E. Kreger, W.B. Kannel (1994) The effects of specific medical conditions on the functional limitations of elders in the Framingham Study. *Am J Public Health* 84: 351–358.
- Han, W., S. Cai, Z. Liu, X. Jin, X. Wang, B. Antony, Y. Cao, D. Aitken, F. Cicuttini, G. Jones, C. Ding (2014) Infrapatellar fat pad in the knee: is local fat good or bad for knee osteoarthritis? *Arthritis Res Ther* 16: R145.
- Haug, S.E. (2014) Der Hoffasche Fettkörper: Ein Literature-Review – Klinik für Orthopädie und Sportorthopädie; thesis, Technische Universität München, Munich.
- Hoffa, A. (1904) The influence of the adipose tissue with regard to the pathology of the knee joint. *JAMA* 43: 795–796.
- Hui, W., G.J. Litherland, M.S. Elias, G.I. Kitson, T.E. Cawston, A.D. Rowan, D.A. Young (2012) Leptin produced by joint white adipose tissue induces cartilage degradation via upregulation and activation of matrix metalloproteinases. *Ann Rheum Dis* 71: 455–462.
- Hunter, D.J., D.P. Beavers, F. Eckstein, A. Guermazi, R.F. Loeser, B.J. Nicklas, S.L. Mihalko, G.D. Miller, M. Lyles, P. DeVita, C. Legault, J.J. Carr, J.D. Williamson, S.P. Messier (2015) The Intensive Diet and Exercise for Arthritis (IDEA) trial: 18-month radiographic and MRI outcomes. *Osteoarthritis Cartilage* 23: 1090–1098.
- Ioan-Facsinay, A., M. Kloppenburg (2013) An emerging player in knee osteoarthritis: the infrapatellar fat pad. *Arthritis Res Ther* 15: 225.
- Issa, R.I., T.M. Griffin (2012) Pathobiology of obesity and osteoarthritis: integrating biomechanics and inflammation. *Pathobiol Aging Age Relat Dis* 2: 17470.
- Kang, E.H., Y.J. Lee, T.K. Kim, C.B. Chang, J.-H. Chung, K. Shin, E.Y. Lee, E.B. Lee, Y.W. Song (2010) Adiponectin is a potential catabolic mediator in osteoarthritis cartilage. *Arthritis Res Ther* 12: 231.
- Kellgren, J.H., J.S. Lawrence (1957) Radiological assessment of osteo-arthritis. *Ann Rheum Dis* 16: 494–502.
- Klein-Wieringa, I.R., M. Kloppenburg, Y.M. Bastiaansen-Jenniskens, E. Yusuf, J.C. Kwekkeboom, H. El-Bannoudi, R.G.H.H. Nelissen, A. Zuurmond, V. Stojanovic-Susulic, G.J.V.M. Van Osch, R.E.M. Toes, A. Ioan-Facsinay (2011) The infrapatellar fat pad of patients with osteoarthritis has an inflammatory phenotype. *Ann Rheum Dis* 70: 851–857.
- Losina, E., R.P. Walensky, W.M. Reichmann, H.L. Holt, H. Gerlovin, D.H. Solomon, J.M. Jordan, D.J. Hunter, L.G. Suter, A.M. Weinstein, A.D. Paltiel, J.N. Katz (2011) Impact of obesity and knee osteoarthritis on morbidity and mortality in older Americans. *Ann Intern Med* 154: 217–226.
- Messier, S.P., D.P. Beavers, R.F. Loeser, J.J. Carr, S. Khajanchi, C. Legault, B.J. Nicklas, D.J. Hunter, P. Devita (2014) Knee joint loading in knee osteoarthritis: influence of abdominal and thigh fat. *Med Sci Sports Exerc* 46: 1677–1683.
- Messier, S.P., C. Legault, S. Mihalko, G.D. Miller, R.F. Loeser, P. Devita, M. Lyles, F. Eckstein, D.J. Hunter, J.D. Williamson, B.J. Nicklas (2009) The Intensive Diet and Exercise for Arthritis (IDEA) trial: design and rationale. *BMC Musculoskelet Disord* 10: 93.

- Messier, S.P., S.L. Mihalko, C. Legault, G.D. Miller, B.J. Nicklas, P. DeVita, D.P. Beavers, D.J. Hunter, M.F. Lyles, F. Eckstein, J.D. Williamson, J.J. Carr, A. Guermazi, R.F. Loeser (2013) Effects of intensive diet and exercise on knee joint loads, inflammation, and clinical outcomes among overweight and obese adults with knee osteoarthritis: the IDEA randomized clinical trial. *JAMA* 310: 1263–1273.
- Pan, F., W. Han, X. Wang, Z. Liu, X. Jin, B. Antony, F. Cicuttini, G. Jones, C. Ding (2015) A longitudinal study of the association between infrapatellar fat pad maximal area and changes in knee symptoms and structure in older adults. *Ann Rheum Dis* 74: 1818–1824.
- Pogacnik Murillo, A.L. (2015) Impact of Diet and/or Exercise Intervention on Infrapatellar Fat Pad Morphology: a Secondary Analysis from a Randomized Controlled Trial; thesis, Paracelsus Medical University, Salzburg.
- Poole, A.R. (2012) Osteoarthritis as a whole joint disease. *HSS J* 8: 4–6.
- Presle, N., P. Pottier, H. Dumond, C. Guillaume, F. Lapique, S. Pallu, D. Mainard, P. Netter, B. Terlain (2006) Differential distribution of adipokines between serum and synovial fluid in patients with osteoarthritis: contribution of joint tissues to their articular production. *Osteoarthritis Cartilage* 14: 690–695.
- Richter, M., T. Trzeciak, M. Owecki, A. Pucher, J. Kaczmarczyk (2015) The role of adipocytokines in the pathogenesis of knee joint osteoarthritis. *Int Orthop* 39: 1211–1217.
- Roemer, F.W., M. Jarraya, D.T. Felson, D. Hayashi, M.D. Crema, D. Loeuille, A. Guermazi (2016) Magnetic resonance imaging of Hoffa's fat pad and relevance for osteoarthritis research: a narrative review. *Osteoarthritis Cartilage* 24: 383–397.
- Stannus, O.P., Y. Cao, B. Antony, L. Blizzard, F. Cicuttini, G. Jones, C. Ding (2015) Cross-sectional and longitudinal associations between circulating leptin and knee cartilage thickness in older adults. *Ann Rheum Dis* 74: 82–88.
- Steidle-Kloc, E., J. Dörrenberg, W. Wirth, A. Ruhdorfer, F. Eckstein (2015) Knee pain is not related to alterations in the morphology or MRI signal of the infra-patellar fat pad (IPFP): a within-person and between-person analysis using data from the Osteoarthritis Initiative (OAI). *Osteoarthritis Cartilage* 23: 59–60.
- Steidle-Kloc, E., W. Wirth, A. Ruhdorfer, T. Dannhauer, F. Eckstein (2016) Intra- and inter-observer reliability of quantitative analysis of the infra-patellar fat pad and comparison between fat- and non-fat-suppressed imaging-data from the Osteoarthritis Initiative. *Ann Anat* 204: 29–35.
- Teichtahl, A.J., E. Wulidasari, S.R.E. Brady, Y. Wang, A.E. Wluka, C. Ding, G.G. Giles, F.M. Cicuttini (2015) A large infrapatellar fat pad protects against knee pain and lateral tibial cartilage volume loss. *Arthritis Res Ther* 17: 318.