Conservative versus Surgical Management of Iatrogenic Tracheal Rupture

Nikolaos Panagiotopoulos^a Davide Patrini^a Matthew Barnard^b
Efstratios Koletsis^{a, c} Dimitrios Dougenis^c David Lawrence^a

^aDepartment of Cardiothoracic Surgery, The Heart Hospital, University College London Hospital, and

^bDepartment of Anaesthesiology and Intensive Care, St Bartholomew's Hospital, London, UK;

^cDepartment of Cardiothoracic Surgery, School of Medicine, University of Patras, Patras, Greece

Keywords

Tracheal rupture · Iatrogenic surgical emphysema ·
Intubation · Pneumomediastinum

Abstract

Iatrogenic tracheal rupture (ITR) represents a life-threatening condition requiring prompt diagnosis, management, and treatment. The management of ITR is challenging, and treatment options depend on tear location, size, injury extent, and the patient's respiratory status. Although this complication has been extensively reported in published literature, the best evidence practice, for the management, requires clarification. In this review, the authors focused on the establishment of a differential diagnosis and the potential mechanism of the injury, the decision-making process, and the therapeutic approaches. It is suggested that for small lacerations or stable patients, conservative management could be considered sufficient, whereas invasive surgical therapy would be more appropriate in cases of large defects with significant air leak and patient instability.

© 2017 S. Karger AG, Basel

Introduction

Iatrogenic tracheal rupture (ITR) represents a life-threatening condition that requires prompt diagnosis, management, and treatment. Prokakis et al. [1] reported that prevalence of ITR in elective orotracheal intubations was 1/20,000–75,000 patients and increased to 15% in emergently performed procedures, and for percutaneous dilatative tracheotomies the estimated incidence is 0.2–0.7%. In the last decade, the estimated incidence of ITR after endotracheal intubation is 0.05–0.37% [1], with a higher incidence of approximately 0.5–1% reported in cases of double-lumen intubation [2]. The presence of other concomitant pathologies of patients undergoing surgical procedure for this, or those who are treated in intensive care units for other reasons, could lead to an early adverse outcome, ventilator failure, airway obstruction, or death from tension pneumothorax, or late development of airway stenosis and recurrent pulmonary infections [1]. Therefore, although prompt diagnosis is essential for the survival of the patients, the diagnostic delay is 25.7 ± 22.9 h (range, 3–72 h) [3]. The decision making

regarding the treatment of these patients is highly challenging. The principal necessity is to secure and maintain the airway and to allow adequate ventilation. If a repair of the injury is to be performed, minimal impact on the respiratory function and subsequent quality of life must be ensured.

Diagnosis

In non-thoracic surgery patients, tracheal injuries were diagnosed based on presenting symptoms of dyspnoea, subcutaneous emphysema, and haemoptysis in the early postoperative period. In thoracic surgery patients, they were diagnosed intraoperatively by direct visualization. In patients intubated in intensive care units, injuries presented with subcutaneous emphysema and/or pneumothorax. Diagnosis can be established with different investigations. Chest CT could confirm the presence of pneumothorax, pneumomediastinum, mediastinitis, or pneumonia; fiberoptic bronchoscopy will allow direct evaluation of the whole trachea; oesophagoscopy may exclude an oesophageal lesion [4].

Mechanisms of Injury

ITR commonly occurs after intubation or tracheotomy procedures. The most common mechanisms are intubation by inexperienced clinicians, dilatative percutaneous tracheotomy, and change of the position of the intubation tube without deflation or overinflation of the tube cuff [2, 4]. Relative overinflation of the tube cuff occurs if the tracheal tube is inflated just above the carina and moved back to the narrower upper portion of the trachea. Intubation with “double lumen” endotracheal tubes causes ITR more frequently, probably owing to their larger diameter and rigidity [5]. This mechanism of injury explains the incidence of trachea laceration after intraoperative repositioning of the patient’s head or body, resulting in displacement of the endotracheal tube. In females, the trachea’s narrower diameter, less resistant wall, and shorter length are predisposing factors to this injury [6]. Characteristically, they are longitudinal lacerations of the posterior tracheal wall [4, 6]. These tracheal tears are frequently found on the right side because of the proximity to the oesophagus on the left side, stabilizing the trachea wall. Occasionally, tracheal lacerations can extend into the main bronchi [6].

Treatment Decision

Deja et al. [4] described a comprehensive algorithm regarding the diagnosis and management of ITR. Conservative treatment is favoured for patients who do not require mechanical ventilation [7]. When mechanical ventilation is possible without any loss of tidal volume and the emphysema is only mild and stable during ventilation, conservative management is indicated. This choice is independent of the length of the lesion and includes lesions close to the carina [4, 6, 7]. When the injury includes the full thickness of the posterior membrane and mediastinal structures protrude into the tracheal lumen, surgical repair is indicated.

This algorithm is a more conservative approach compared to previous authors who suggest that nonsurgical therapy must only be considered in stable patients with small, uncomplicated tears of lengths between <2 and <4 cm [8, 9]. They advise that conservative treatment can be considered if previous pulmonary or mediastinal surgery has not been performed [9].

Therapeutic Approaches

In patients with spontaneous breathing at the time of diagnosis, medical treatment includes a broad-spectrum antibiotic therapy against the tracheobronchial flora, antiseptic anti-inflammatory aerosol therapy, antitussive agents, and chest tube insertion, if required [9].

In patients still requiring mechanical ventilation at the time of diagnosis due to acute respiratory failure, coma, or multi-organ failure, conservative management includes positioning the tracheal tube distal to the tracheal rupture, continuous measurement of the cuff pressure, and chest tube insertion, if required [4]. This strategy is effective in patients with lacerations in the upper region of the trachea. In cases of mechanically ventilated patients with lacerations close to the carina, which is the more common scenario, it is not easy to position the cuff distal to the tracheal rupture and to avoid dislocation of the tube tip into the mediastinum. Therefore, the tube must be sited close to the carina, under bronchoscopy with continuous cuff pressure monitoring, for a limited period [4]. In these patients, early tracheostomy allows spontaneous breathing with the aid of lower positive inspiratory pressure and continuous positive airway pressure ventilation.

Special care must be taken in patients with acute lung injury or acute respiratory distress syndrome, using pres-

sure-controlled low tidal volume ventilation, prone positioning, and implementation of early spontaneous ventilation, inhalation of nitric oxide, and, in some patients, extracorporeal membrane oxygenation or pumpless extracorporeal lung assist, postponing surgery to improve the clinical condition and gas exchange prior to the operation procedure [4].

Conservative management can be facilitated by tracheal stents, which avoid tracheal tube migration and permit early extubation [4]. Their main disadvantage is dilatation of the trachea with a potential increase in the diameter.

Welter et al. [10] reported a small series of 4 patients with rupture of the membranous wall of the trachea. They proposed complete intraluminal repair using a 70-cm, 2-0 Vicryl suture (UCLX needle; Ethicon, Germany). The suture is anchored to one end with a small absorbable polydioxanone clip (ABSOLOK AP100; Ethicon). A running suture starting at the peripheral end of the tear includes the whole posterior wall. At the proximal end of the suture, one or two polydioxanone clips are applied to secure the suture.

When surgery is indicated, it should be performed promptly to optimize success and avoid complication. The surgical approaches are dependent on which anatomical region of the tracheobronchial tree is affected [11]. For the cervical trachea, a collar incision or left cer-

vicotomy for the posterior wall is indicated. A collar incision plus partial sternotomy can be performed for proximal mediastinal tracheal injuries. With a transcervical-transtacheal approach, one can also perform a repair of the membranous wall of the trachea with subsequent closure of the tracheal incision. A right thoracotomy approach is mandatory in carina injuries, the proximal part of the left mainstem bronchus, and the right bronchial tree. Injuries of the left mainstem bronchus close to the lung hilum and left bronchial tree necessitate a left thoracotomy approach [4, 10].

Conclusions

The treatment decision for ITR depends on tear location, injury extent, and size. For small lacerations and stable patients, conservative management is considered sufficient, while surgical operation is mandatory in cases of large defects with significant air leak and patient instability.

Disclosure Statement

The authors declare no conflicts of interest.

References

- 1 Prokakis C, Koletsis E, Dedeilias P, et al: Airway trauma: a review on epidemiology, mechanisms of injury, diagnosis and treatment. *J Cardiothorac Surg* 2014;9:117.
- 2 Altinok T, Can A: Management of tracheobronchial injuries. *Eurasia J Med* 2014;46:209–215.
- 3 Gomez-Caro Andres A, Moradiellos Diez FJ, Ausin Herrero P, et al: Successful conservative management in iatrogenic tracheobronchial injury. *Ann Thorac Surg* 2005;79:1872–1878.
- 4 Deja M, Menk M, Heidenhain C, et al: Strategies for diagnosis and treatment of iatrogenic tracheal ruptures. *Minerva Anesthesiol* 2011;77:1155–1166.
- 5 Ovari A, Just T, Dommerich S, et al: Conservative management of post-intubation tracheal tears – report of three cases. *J Thoracic Dis* 2014;6:E85–E91.
- 6 Conti M, Pougeoise M, Wurtz A, et al: Management of post-intubation tracheobronchial ruptures. *Chest* 2006;130:412–418.
- 7 Schneider T, Storz K, Dienemann H, et al: Management of iatrogenic tracheobronchial injuries: a retrospective analysis of 29 cases. *Ann Thorac Surg* 2007;83:1960–1964.
- 8 Carbognani P, Bobbio A, Cattalani L, et al: Management of post-intubation membranous tracheal rupture. *Ann Thorac Surg* 2004;77:406–409.
- 9 Jougon J, Ballester M, Choukron E, et al: Conservative treatment for post-intubation tracheobronchial rupture. *Ann Thorac Surg* 2000;69:216–220.
- 10 Welter S, Krbek Th, Hajder R, et al: A new technique for complete intraluminal repair of iatrogenic posterior tracheal lacerations. *Interact Cardiovasc Thorac Surg* 2011;12:6–9.
- 11 Koletsis E, Prokakis Ch, Baltayiannis N, et al: Surgical decision making in tracheobronchial injuries on the basis of clinical evidences and the injury's anatomical setting: a retrospective analysis. *Injury* 2012;43:1437–1441.