Review

The Emerging Roles of CIB1 in Cancer

Xianwang Wang^{a,b,c} Xiaochun Peng^a Xueqing Zhang^b Hanyi Xu^a
Chengbiao Lu^c Lian Liu^a Jiaxing Song^d Yingjie Zhang^e^aThe Second Clinical Medical School of Yangtze University & Jingzhou Central Hospital, Jingzhou,^bKey Laboratory of Oncology, Center for Molecular Medicine, School of Medicine, Yangtze University,Jingzhou ^cKey Laboratory of Neuronal Oscillation and Disease, School of Medicine, Yangtze University,Jingzhou, ^dSchool of Medicine, Shenzhen University, Shenzhen, Guangdong, ^eCollege of Biology, Hunan

University, Changsha, Hunan, China

Key Words

CIB1 • Calcium • Integrin • Cancer • Tumor progression

Abstract

Calcium and integrin-binding protein 1 (CIB1) is an EF-hand calcium binding protein, which is involved in many cellular processes, including calcium signaling, cell survival and proliferation, cell migration, cell adhesion and apoptosis. A number of studies have found that CIB1 is ubiquitously expressed and is related to various human diseases, such as cancer, Alzheimer's disease (AD), cardiac hypertrophy and male infertility. The mechanism of CIB1 in human diseases is still not clear, although multiple functions of CIB1 are modulated by interacting with numerous interacting partners. As a calcium binding protein, the roles of CIB1 in calcium signaling by binding calcium or modulating some key modulators, such as calcineurin, integrin, inositol 1,4,5-trisphosphate receptor (IP₃R) and taste 1 receptor member 2 (TAS1R2). The tumor promoting mechanisms of CIB1 have been described in different aspects, including promoting tumor cell cycle and proliferation, inhibiting tumor cell apoptosis, and mediating tumor cell migration and angiogenesis. In addition, multiple functions of CIB1, such as neural development, taste or gustation functions, and virus infection are also elucidated. These recent advances have significantly expanded our understanding of the knowledge of CIB1 and highlighted the potential mechanisms of CIB1 in tumor progression.

Introduction

© 2017 The Author(s)
Published by S. Karger AG, Basel

Calcium and integrin-binding protein 1 (CIB1, also known as calmyrin and KIP) is a 22 kD protein, identified originally as a binding partner of platelet integrin cytoplasmic domain alpha-IIb/beta3(αIIbβ3) [1], and later found to regulate αIIbβ3 activation [2, 3] and down regulate inositol 1, 4,5-trisphosphate receptor (IP₃R)-dependent calcium (Ca²⁺) release [4]. As a calcium-binding protein, the structural properties of CIB1 are most closely related to calcineurin B (CnB) (57% sequence similarity to CIB1) and calmodulin (CaM)

X. Wang and X. Peng contributed equally to this work.

Xianwang Wang
and Yingjie ZhangKey Laboratory of Oncology, Center for Molecular Medicine, School of Medicine,
Yangtze University, China and College of Biology, Hunan University, Changsha, (China)
E-Mail 275379987@qq.com and 21392705@qq.com

(56% similarity to CIB1) [1, 5]. Similar to these Ca^{2+} related proteins, as shown in Fig. 1, the human CIB1 contains four helix-loop-helix EF hand motifs, two C-terminal EF hand III and EF hand IV are responsible for Ca^{2+} and magnesium (Mg^{2+}) binding [1, 6-10]. However, CIB1 was found to bind two Ca^{2+} ions in a sequential manner with dissociation constants near 1.9 and 0.54 μM for sites EF-III and EF-IV, respectively. In contrast, CIB1 bound only one Mg^{2+} ion strongly to EF-III with dissociation constants about 120 μM [11]. Of note, CIB1 is usually found in monomeric form and distributed in both the nucleus and cytoplasm [12], and its subcellular localization might be influenced by association with its interacting partners and/or by $\text{Ca}^{2+}/\text{Mg}^{2+}$ levels. Both Ca^{2+} and Mg^{2+} binding induce conformational changes which stabilize the higher structure of CIB1, leading to remarkable increases in the thermal stability of the proteins [11]. Yamniuk et al. identified that low affinity Ca^{2+} binding events that influence the structures of the N- and C-terminal extensions of CIB1 under high Ca^{2+} crystallization conditions [10]. Studies suggested that Mg^{2+} -CIB1 and Ca^{2+} -CIB1 exhibit a more ordered 4-EF-hand structure with smaller hydrodynamic radius than a highly flexible molten globule-like CIB1 (not bound to any metals)[6, 8]. A consensus sequence 5'-MGXXXS/T-3' at the CIB1 N terminus appears to be crucial for its myristoylation modification and plasma membrane localization [4, 7, 13, 14]. The myristoylation at the CIB1 N terminus might interfere with CIB1 cellular function [13, 15-17]. Stabler et al. showed that a myristoylated CIB1 that preferentially interacts with Presenilin 2 (PS2) [13]. The Ca^{2+} -myristoyl switch function of CIB1, and its ability to facilitate agonist-induced plasma membrane localization of SK1, a location where SK1 is known to trigger oncogenic signaling [15]. In addition, CIB1 myristoylation is also important in shuttling CnB to plasma membrane [16].

A number of studies have further indicated that CIB1 is ubiquitously expressed and is involved in various physiological and pathological processes, including apoptosis [18], cell survival and proliferation [19], cancer [20, 21], cardiovascular diseases [16], and Alzheimer's disease (AD) [22-24]. The precise mechanism between CIB1 and its various binding partners is not clear, although numerous associated proteins of CIB1 have been identified (Table. 1). In this review, we discussed the potential mechanism of CIB1 in Ca^{2+} signaling and paid more attention to tumor promoting effect of CIB1. Last but not least, some other functions of CIB1 were also involved in this article.

CIB1 and Ca^{2+} signaling

In general, intracellular Ca^{2+} signals result either from its release from the important intracellular stores (such as endoplasmic reticulum (ER) and endo-lysosome), or by activation of Ca^{2+} -conducting channels at the plasma membrane, including voltage-dependent Ca^{2+} channels (VDCCs), $\text{Na}^{+}/\text{Ca}^{2+}$ exchanger (NCX), plasma membrane calcium-transporting ATPases (PMCA), cyclic nucleotide-gated ion channels (CNGCs), α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor (AMPA) and N-methyl-D-aspartate

Fig. 1 The basic structure of human CIB1. The CIB1 contains four EF hand motifs, EF-I, EF-II, EF-III and EF-IV. The EF-III and IV of CIB1 are essential for $\text{Ca}^{2+}/\text{Mg}^{2+}$ binding. The N-terminus (1-6 aa) of CIB1 is response for its myristoylation modification.

Table 1 The important interacting partners and potential functions of CIB1.

Partners	Potential functions	References
α IIb β 3	platelet aggregation; cell migration; calcium signaling	[1, 2, 61]
FVIII	platelet aggregation	[24]
Rac3	cell migration	[66]
WASP	platelet aggregation; cell migration	[65]
FAK/PTK2	cell migration	[64, 70]
PAK1	cell migration; tumor growth	[67-69]
IP ₃ R	calcium signaling	[4, 8, 32]
TAS1R2	calcium signaling; taste or gustation	[4]
Plk2/Snk	cell cycle; cancer progression; DNA damage response; synaptic plasticity	[7, 39, 45]
Plk3/Fnk	cell cycle; cancer progression; DNA damage response; synaptic plasticity	[20, 39, 44, 46]
PKD2	tumorigenesis	[46]
SK1	cell proliferation; antiapoptosis; tumor growth	[15, 17, 49]
ASK1	antiapoptosis; calcium signaling;	[7, 57]
IFI6/G1P3	cell survival; antiapoptosis	[18]
UBR5/EDD	DNA damage response; tumorigenesis	[56]
DNA-PKcs/TRF2	DNA damage response	[53, 54, 56]
hTERT	telomere homeostasis; cancer cell proliferation	[53, 54]
SCG10/stathmin2	brain development	[72]
NBR1/BRCA1/ FEZ1	neural and brain development	[73]
PAX3	neurogenesis and myogenic differentiation	[75]
PS 1/2	apoptosis; AD	[13, 14, 22, 23]
EphrinA2	virus infection	[78]
Cn/CnB	calcium signaling; cardiac hypertrophy; valvular heart disease	[16, 25, 30, 79]

receptor (NMDAR) etc [4, 25-29]. Release of intracellular Ca^{2+} from ER store through IP₃R and/or ryanodine receptors (RyR) [25]. Endo-lysosomal Ca^{2+} stores, on the other hand, are mobilized by two-pore channels (TPCs) and transient receptor potential mucolipin-1 (TRP-ML1) [25-27].

The CIB1 belong to calcium-binding protein family, and serve important roles in Ca^{2+} signaling pathways by binding to Ca^{2+} and some key modulators. For example, CIB1 can interact with CnB and strongly enhances CnB membrane localization and activation [16, 30]. Calcineurin (Cn) was shown to associate with the L-type Ca^{2+} channel, which is the major mediator of Ca^{2+} influx from the sarcoplasmic reticulum by activating RyR₂ in cardiac myocytes [31]. The binding of CIB1 to α IIb β 3 is identified to inhibit phospholipase C (PLC)/IP₃ signaling [1, 2, 4]. Of note, CIB1 is also reported to inhibit the IP₃R- Ca^{2+} release channel [32]. Binding of CIB1 to the N-terminal region of IP₃R is reduced at low free $[\text{Ca}^{2+}]$ and could be inhibited by mutations to EF hand-III or EF hand-IV, suggesting that Ca^{2+} -binding is the trigger for the CIB1-IP₃R interaction [7]. CIB1 interact with the intracellular C-terminal domain of rat sweet taste 1 receptor member 2 (TAS1R2) also reported to inhibit IP₃R- Ca^{2+} release [4]. Until now, two recognized intracellular Ca^{2+} signaling pathways, integrin-IP₃R- Ca^{2+} signaling and CD38- Ca^{2+} signaling have been established. CD38 is a multifunctional ectoenzyme that catalyzes the synthesis and hydrolysis of cyclic ADP-ribose (cADPR) from nicotinamide adenine dinucleotide (NAD⁺) to ADP-ribose (ADPR) and nicotinic acid adenine dinucleotide phosphate (NAADP) [28, 33, 34]. Similar to integrin, a transmembrane CD38 also trigger intracellular Ca^{2+} signaling pathways [33]. In addition to IP₃, cADPR and NAADP are already identified as important intracellular Ca^{2+} messengers, which elicit intracellular Ca^{2+} flux from both ER and endolysosome [25, 33, 34]. CD38 act as a NAD⁺ dependent enzymes, which catalyzing the synthesis of cADPR and NAADP to mediate ER- Ca^{2+} and lysosome- Ca^{2+} signaling, respectively [28, 34]. In addition to modulate IP₃R- Ca^{2+} signaling, CIB1 is possibly also implicated in CD38 triggering Ca^{2+} signaling pathway. As summarized in Fig. 2, In IP₃-ER- Ca^{2+} signaling, CIB1 acts as a partner of integrin, which affects the activation of PLC. As we known, IP₃ is produced by the activation of PLC and it acts on IP₃R to release Ca^{2+} from the ER stores. Besides, CIB1 also interact with the C-terminal domain of TAS1R2 to inhibit IP₃R- Ca^{2+} release. In CD38 triggering Ca^{2+} signaling pathway, it is interesting that whether CIB1 may also serve as an accessory protein of CD38, which synthesizes the second messengers, cADPR and NAADP. The synthetic cADPR and NAADP target to RYR and TPC/TRP-ML1, resulted in Ca^{2+} release from ER and endolysosome stores, respectively [25, 28, 34]. Furthermore, whether CIB1 is

Fig. 2 The potential mechanism of CIB1 in intracellular Ca^{2+} signaling

Table. 2 The roles for CIB1 in tumor progression. Abbreviations: Acute myeloid leukemia (AML); colorectal cancer (CRC); hepatocellular Carcinoma (HCC); triple negative breast cancer (TNBC).

Tumor or cell types	Targets	Potential mechanism	References
MCF-7 cells	EDD, DNA-PKcs	DNA damage response; tumorigenesis	[56]
T47D breast cancer	FAK, PAK1	cancer cell migration	[69, 70]
H1299, MCF7 and HT1080 cells	DNA-PKcs, hTERT	telomere homeostasis; cancer cell proliferation	[53]
TMK-1 cells	G1P3	inhibit apoptosis	[18]
Melanoma or Lewis lung carcinoma cells	PAK1, ERK1/2	angiogenesis	[68, 71]
HeLa, MCF-7 and DU145 cells, lung cancer, AML, CRC	Ras, SK1	cell proliferation; antiapoptosis; tumor growth	[15, 17]
HCC and HepG2 cells	PAK1, ERK1/2 and ASK1	cancer cell proliferation	[42]
Breast cancer cells	PLK3	calcium signaling, cancer cell proliferation	[20, 44]
Breast, bladder, cervix and colorectal cancer	PKD2	tumor cell invasion, tumor growth and angiogenesis	[40]
Neuroblastoma and breast cancer cells	PI3K/AKT, MEK/ERK and PAK1	cancer cell proliferation	[43]
Kaposi's sarcoma	EphrinA2, ERK1/2	virus entry	[78]
Pancreatic cancer	DNA-PKcs, PLK3, Rac3, Pax3, PS2	cancer cell proliferation	[21]
TNBC cells	PI3K/AKT, MEK/ERK	cancer cell proliferation	[41]

also likely to affect the RYR and TPC/TRP-ML1 activity directly may be the result of further mechanism that have not yet been elucidated. Of interest, whether CIB1 and the modulators in these two signaling pathways synergistically modulating intracellular Ca^{2+} signaling are not clear. Whether CIB1 also affects some other Ca^{2+} channels on cell membrane (such as NMDAR, L-type Ca^{2+} channel and NCX) are still need further investigation. Several studies suggested that targeting Ca^{2+} signaling possible serve as a potential strategy in cancer therapy [35-38]. Thus, gaining better understanding of the relationship between CIB1 and these Ca^{2+} modulators remain an important area to investigate in the future study.

Fig. 3 The potential mechanism of CIB1 in tumorigenesis.

The potential roles of CIB1 in tumor progression

Recently, carcinogenesis of CIB1 is in the spotlight. Increasing evidence has suggested that CIB1 perform important roles in cell survival, cell proliferation, cell apoptosis and cell migration, thereby mediating tumor growth and angiogenesis [17, 20, 39-43]. CIB1 may serve an important role in tumorigenesis by regulating several kinase partners or oncogenic signaling (Table. 1, 2 and Fig. 3). The summary of these studies are described as below.

CIB1 and tumor cell survival and death signaling

Polo-like kinases (Plks), a family of serine/threonine kinases, have been shown to play key roles in regulation of cell cycle progression. So far, Plk1, Plk2, Plk3 and Plk4 proteins have been identified in the mammalian cells [20, 39, 44]. Plk2, also called serum-inducible protein kinase (Snk), is combined and inhibited by CIB1 [39, 45]. Plk3, also known as FGF-inducible kinase (Fnk), is coupled to the CIB1 in two-hybrid-analyses and coimmunoprecipitation assays [46]. The interaction between Plk3 and CIB1 might play a role in cancer progression. Later studies showed that CIB1 constitutively interacts with Plk3 and inhibits its activity in a Ca^{2+} -dependent manner, thereby contributes to the regulation of the cell cycle [20]. Protein kinase D2 (PKD2), a member of the PKD family of serine/threonine kinases, has been reported as a potential promoter for tumor growth and angiogenesis [40, 47]. Armacki et al. suggested that a novel splice variant of CIB1 (called CIB1a), which regulates tumor cell invasion, tumor growth and angiogenesis by mediating PKD-induced the secretion of vascular endothelial growth factor [40]. SK1 plays an important role in cell proliferation, protection from apoptosis, and oncogenic rat sarcoma (Ras)-mediated neoplastic transformation [15, 17, 48]. Study showed that CIB1 interacts with SK1 and promotes the translocation of SK1 from cytoplasm to plasma membrane, mediating activation of SK1 and in turn anti-apoptotic signaling through the nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B) pathway [15]. As we known, SK1 is a downstream mediator of oncogenic signaling by Ras since chemical inhibitors of SK or a dominant-negative SK1 attenuated Ras-induced neoplastic transformation [48, 49]. Besides, studies indicated that oncogenic Ras induced plasma membrane localization of SK1 [17, 50]. Thus, CIB1 serves as

an important effector of oncogenic signaling by Ras through the enhancement of subcellular localization of SK1 to the cell membrane [17]. Phosphatidyl inositol 3-kinase (PI3K) and extracellular regulated protein kinases (ERK) are involved in cellular functions such as cell growth, proliferation [51, 52]. CIB1 is also implicated in cell survival and proliferation by modulating PI3K/AKT and MEK/ERK oncogenic signaling [41, 42]. Leisner et al. have been identified that CIB1 depletion increased nuclear distribution of GAPDH by interrupting AKT and ERK signaling, leading to caspase-independent cell death and proliferation inhibition [43]. Besides, CIB1 can associate with the catalytic subunit of human telomerase reverse transcriptase (hTERT), stimulating telomerase activity and telomere lengthening [53, 54]. CIB1 is required for the association between telomeric repeat-binding factor 2 (TRF2) and DNA-protein kinase catalytic subunit (DNA-PKcs), the interplay of these three proteins may provide a mechanism for the recruitment of DNA-PKcs to telomeres [54]. The interaction between CIB1 and DNA-PKcs also affects DNA-PKcs related DNA damage [53, 55]. Moreover, being a binding partner of E3 ubiquitin-protein ligase UBR5 (EDD), CIB1 is a possible target for E3-mediated degradation and potentially involved in DNA damage signaling [56]. Study showed that CIB1 inhibits apoptosis signal-regulating kinase 1 (ASK1) activity both *in vitro* and *in vivo*, thereby negatively regulates stress-activated mitogen-activated protein kinase (MAPK) signaling pathways and apoptosis [57]. The mitochondria serve important roles in intrinsic apoptotic pathway [58, 59]. A mitochondrial localized G1P3 (also known as interferon alpha-inducible protein 6 (IFI6) or interferon-induced protein 6-16) has been shown to inhibit mitochondrial-mediated apoptosis in gastric cancer cell line TMK-1 cells [18]. This anti-apoptotic function is mediated by interacting with CIB1, thereby inhibit the depolarization of mitochondrial membrane potential, release of cytochrome c [18]. Although overexpression of CIB1 and/or PS2 promotes cell death *in vitro*, but no published data to confirm the role of CIB1 in cancer cell apoptosis [13, 18], further studies are needed to conclude the involvement in the anti-apoptotic activity. As summarized in Fig. 3, PI3K/AKT, Ras/Raf/MEK/ERK, Plks, PDK2, SK1 and G1P3 may be potential oncogenic signaling in CIB1-driven tumor progression. As an important effector of DNA damage signaling, CIB1 may effect tumor progression by regulating the function of DNA-PKcs, TRF2, EDD and TERT. Furthermore, inhibitions of tumor cells apoptosis by inhibiting ASK1/MAPK signaling are also involved in CIB1-induced carcinogenesis.

CIB1 and tumor cell migration and angiogenesis

Moreover, cell migration plays an essential role in tumor growth and metastasis. CIB1 is also involved in tumor metastasis by effecting tumor cell migration and spreading [7]. Until now, multiple studies have indicated that CIB1 serve important roles in regulating platelet integrin activity, cell adhesion and migration by interacting with at least six binding partners, including α IIb β 3, the coagulation factor VIII (FVIII), Wiskot-Aldrich syndrome protein (WASP), the small GTPase Rac3, P21-activated kinases 1 (PAK1) and focal adhesion kinase (FAK). As a major adhesion receptors, integrin control many normal cellular processes, including migration, growth, differentiation, and proliferation [60]. Studies has suggested that CIB1 is coupled to cytoplasmic domain of α IIb β 3, thereby blocks α IIb β 3 activation and in turn platelet aggregation [1, 2]. Further study indicated that CIB1 could bind to all integrin complexes and act as a broad regulator of integrin function [3]. CIB1 also interacts with the FVIII in 227-336 regions, suggesting that CIB1 pay important roles in coagulation [24]. Base on the relationship between CIB1 and α IIb β 3, CIB1 is also involved in cell migration [61]. WASP binding to CIB1 affects the affinity of α IIb β 3 for fibrinogen (also called factor I), a soluble plasma glycoprotein is converted by thrombin into fibrin during blood coagulation [62-64]. CIB1 can bind to N terminus of the WASP and recruit it to α IIb β 3, increasing α IIb β 3-mediated cell adhesion in stimulated platelets [65]. Tsuboi et al. advanced that inhibition of WASP-CIB1 binding by streptolysin O decreases α IIb β 3 affinity for fibrinogen [65]. It is reported that CIB1 interacts with the small GTPase Rac3 and promotes α IIb β 3-mediated adhesion and spreading of cells through the fibrinogen receptor, such as α IIb β 3 [66]. As a downstream of Rac3, PAK1 is another key factor in cell migration. Multiple studies have indicated that

CIB1 is essential for PAK1 activation both *in vitro* and *vivo*, which further suggested that CIB1-PAK1 interaction is involved in cell migration [67-69]. Naik et al. suggested that CIB1 positively regulates cell migration and is necessary for the recruitment of FAK (also known as protein tyrosine kinase 2, PTK2, a key regulator in focal complex formation) to the focal adhesions. Furthermore, CIB1-induced cell migration is dependent on MAPK signaling and its function is attenuated by PAK1 [69]. Overexpression of CIB1 resulted in cell migration on fibronectin and ERK1/2 MAPK activation [69]. Moreover, by using Chinese hamster ovary cells as model, study showed that CIB1 induces cell migration by binding of FAK and increasing its activity. Overexpression of dominant-negative FAK and CIB1 inhibited cell migration completely [70]. Zayed et al. also found that CIB1 play a critical role in facilitating tumor growth and tumor-induced angiogenesis [68, 71]. As shown in Fig. 4, CIB1 can interact with the N-terminus of α IIb β 3 subsequent to integrin activation. CIB1 also bind to WASP and recruit it to α IIb β 3, which further promoting cell adhesion and spreading. The binding of CIB1 to FAK will also enhance FAK activation [64], and thereby triggering Rac3 and PAK1 induced adhesion and spreading.

In conclusion, CIB1 is widely up-regulated across a diverse range of human cancers. As an emerging potential oncogene [17], CIB1 may affect tumor progression by modulating various processes, including Ca^{2+} signaling, apoptosis, cell cycle and proliferation, cell migration and spreading, telomerase activity and DNA damage. Taken together, there are several potential mechanisms of CIB1 in cancer, including (i) promoting tumor cell cycle and proliferation, (ii) inhibiting tumor cell apoptosis, and (iii) mediating tumor cell migration and angiogenesis. Thus, CIB1 might become a good potential target for cancer therapy. Confirming the interaction mechanism between CIB1 and its interacting partners might provide the potential strategies for preventing cancers.

CIB1 and other pathological or physiological process

CIB1 is a versatile protein by interacting with more than 35 binding partners so far [7]. Of interest, CIB1 is involved in embryogenesis, neural development, AD etc. SCG10 protein (stathmin2), a microtubule-destabilizing factor, which is implicated in neuronal growth during brain development by tied up to the C-terminal of CIB1 and inhibited SCG10-mediated neurite outgrowth [72]. Whitehouse et al. have indicated that CIB1 interacts with NBR1 (named as next to BRCA1) and fasciculation and elongation protein zeta-1 (FEZ1) by using both the yeast two-hybrid assay and coimmunoprecipitation studies [73]. They also identified the subcellular co-localization of CIB1, NBR1 and FEZ1 proteins by immunofluorescence analysis, and speculated that the interactions between these proteins possible an important event in brain development [73]. A member of the paired class homeodomain family of transcription factors, paired box 3 protein (PAX3) [74], has been demonstrated to implicated in brain development by specifically fastening with CIB1 [75]. It is well established that Presenilin 1/2 (PS1/2) are important ageing genes. CIB1 has been demonstrated to fasten with PS2 and increase cell death [13]. By site-directed mutagenesis, Zhu et al. suggested that the interaction between CIB1 and PS2 is regulated by Ca^{2+} binding sequences in EF-hands III and IV [14]. Continues study showed that these binding appear to be in Ca^{2+} independent manner [22]. Moreover, the interaction between CIB1 and Fnk/Snk suggested that CIB1 play vital roles in synaptic plasticity [39]. Study has also demonstrated that CIB1 is associated with hTERT [53], and presented in human forebrain and altered distribution in AD brain [23], further suggesting that CIB1 might play important roles in AD [13, 14, 22, 23]. As

Fig. 4 The roles of CIB1 in cell migration.

shown in Fig. 5, CIB1 control brain development by interact with a variety of partners, including NBR1, SCG10, FEZ1 and PAX3. CIB1 is involved in AD by modulating PS1/2, TERT and Fnk/Snk signaling. Moreover, CIB1 is also related to spermatogenesis. Down-regulation of CIB1 in the sperm of patients with oligoasthenozoospermia [76], reflected that CIB1 may be involved in male subfertility [19, 76]. CIB1 binds to TAS1R2 also suggested that CIB1 might be also effect taste or gustation functions [4]. CIB1 also serve as a potential effector molecule in virus infection, such as human herpes virus 8 and human immunodeficiency virus type-1 [77, 78]. It is demonstrated that CIB1 interacts with EphrinA2 (EphA2) and increase Kaposi's sarcoma-associated herpes virus infection in 293 cells [78]. CIB1 is high expressed in human and rat heart and necessary for cardiac hypertrophy [16, 30]. CIB1 and Cn expression was increased in atrial fibrillation atrial tissue, suggesting that CIB1 may be involved in the pathogenesis of valvular heart disease [79]. As we known, structure determines function. To get insight into the function of CIB1, the structure of CIB1 has been explored extensively by sequence analysis, nuclear magnetic resonance spectroscopy (NMR), circular dichroism or X-ray crystallography [6-8, 10, 80]. Thus, we believed that along with the further analysis of CIB1 structure, more and more CIB1 partners and the precise mechanisms for multiple function of CIB1 will be identified in the future.

Fig. 5 The potential roles of CIB1 in brain development and AD.

Conclusion

In summary, as a multifunctional protein, CIB1 serving not only as a Ca^{2+} modulating protein, most surprisingly, but also as an important potential tumor promoting factor. Here we discussed the plausible mechanism of CIB1 in intracellular Ca^{2+} signaling conduction and paid more attention to an important pro-tumorigenic role for CIB1. The review of the signaling mechanism for CIB1 in tumor progression will highlight the great potential target for cancer therapy.

Abbreviations

CIB1 (Calcium and integrin-binding protein 1); $\alpha\text{IIb}\beta 3$ (Integrin alpha-IIb/beta3); IP_3R (Inositol 1,4,5-trisphosphate receptor); RyR (Ryanodine receptors); CnB (Calcineurin B); CaM (Calmodulin); AD (Alzheimer's disease,); TPCs (Two-pore channels); TRP-ML1 (Transient receptor potential mucolipin-1); PLC (Phospholipase C); TAS1R2 (taste 1 receptor member 2); cADPR (Cyclic ADP-ribose); NAADP (Nicotinic acid adenine dinucleotide phosphate); Plks (Polo-like kinases); Snk (Serum-inducible protein kinase); Fnk (FGF-inducible kinase); PKD2 (Protein kinase D2); SK1 (Sphingosine kinase 1); ASK1 (Apoptosis signal-regulating kinase 1); IFI6 (Interferon alpha-inducible protein 6); hTERT (Human telomerase reverse transcriptase); DNA-PKcs (DNA-protein kinase catalytic subunit); EDD (E3 ubiquitin-protein ligase UBR5); FVIII (Coagulation Factor VIII); WASP (Wiskot-Aldrich syndrome protein); PAK1 (P21-activated kinases 1); FAK (Focal adhesion kinase); FEZ1 (Fasciculation and elongation protein zeta-1); PAX3 (Paired box 3 protein); PS1/2 (Presenilin); EphA2 (EphrinA2); HCC (Hepatocellular carcinoma).

Acknowledgements

This research is supported by the National Natural Science Foundation of China (31700736), Hubei Province Natural Science Foundation of China (2016CFB180), Hubei Province Health and Family Planning Scientific Research Project (WJ2016Y07), Hubei Province Scientific and Technological Research Project (Q20171306), Jingzhou Science and Technology Development Planning Project (JZKJ15063), the Program for Medical Talents of Hubei Province, the Fundamental Research Funds for the Central Universities (531107040909, 14700502044001), the Development Fund from School of Medicine in Yangtze University, the Yangtze Fund for Youth Teams of Science and Technology Innovation (2016CQT04) and the College Students Innovative Entrepreneurial Training Program in Yangtze University (2016133). We are grateful to Professor Hongwu Xing for assistance in editing of the manuscript.

Disclosure Statement

The authors do not have conflict of interest.

References

- 1 Naik UP, Patel PM, Parise LV: Identification of a novel calcium-binding protein that interacts with the integrin α IIb cytoplasmic domain. *J Biol Chem* 1997;272:4651-4654.
- 2 Yuan W, Leisner TM, McFadden AW, Wang Z, Larson MK, Clark S, Boudignon-Proudmon C, Lam SC, Parise LV: CIB1 is an endogenous inhibitor of agonist-induced integrin α IIb β 3 activation. *J Cell Biol* 2006;172:169-175.
- 3 Freeman TC, Jr., Black JL, Bray HG, Dagliyan O, Wu YI, Tripathy A, Dokholyan NV, Leisner TM, Parise LV: Identification of novel integrin binding partners for calcium and integrin binding protein 1 (CIB1): structural and thermodynamic basis of CIB1 promiscuity. *Biochemistry* 2013;52:7082-7090.
- 4 Hennigs JK, Burhenne N, Stahler F, Winnig M, Walter B, Meyerhof W, Schmale H: Sweet taste receptor interacting protein CIB1 is a general inhibitor of InsP3-dependent Ca^{2+} release *in vivo*. *J Neurochem* 2008;106:2249-2262.
- 5 Rusnak F, Mertz P: Calcineurin: form and function. *Physiol Rev* 2000;80:1483-1521.
- 6 Gentry HR, Singer AU, Betts L, Yang C, Ferrara JD, Sondek J, Parise LV: Structural and biochemical characterization of CIB1 delineates a new family of EF-hand-containing proteins. *J Biol Chem* 2005;280:8407-8415.
- 7 Leisner TM, Freeman TC, Black JL, Parise LV: CIB1: a small protein with big ambitions. *FASEB J* 2016;30:2640-2650.
- 8 Huang H, Ishida H, Yamniuk AP, Vogel HJ: Solution structures of Ca^{2+} -CIB1 and Mg^{2+} -CIB1 and their interactions with the platelet integrin α IIb cytoplasmic domain. *J Biol Chem* 2011;286:17181-17192.
- 9 Gifford JL, Walsh MP, Vogel HJ: Structures and metal-ion-binding properties of the Ca^{2+} -binding helix-loop-helix EF-hand motifs. *Biochem J* 2007;405:199-221.
- 10 Yamniuk AP, Anderson KL, Fraser ME, Vogel HJ: Auxiliary Ca^{2+} binding sites can influence the structure of CIB1. *Protein Sci* 2009;18:1128-1134.
- 11 Yamniuk AP, Nguyen LT, Hoang TT, Vogel HJ: Metal ion binding properties and conformational states of calcium- and integrin-binding protein. *Biochemistry* 2004;43:2558-2568.
- 12 Sobczak A, Blazejczyk M, Piszczek G, Zhao G, Kuznicki J, Wojda U: Calcium-binding calmyrin forms stable covalent dimers *in vitro*, but *in vivo* is found in monomeric form. *Acta Biochim Pol* 2005;52:469-476.
- 13 Stabler SM, Ostrowski LL, Janicki SM, Monteiro MJ: A myristoylated calcium-binding protein that preferentially interacts with the Alzheimer's disease presenilin 2 protein. *J Cell Biol* 1999;145:1277-1292.
- 14 Zhu J, Stabler SM, Ames JB, Baskakov I, Monteiro MJ: Calcium binding sequences in calmyrin regulates interaction with presenilin-2. *Exp Cell Res* 2004;300:440-454.
- 15 Jarman KE, Moretti PA, Zebol JR, Pitson SM: Translocation of sphingosine kinase 1 to the plasma membrane is mediated by calcium- and integrin-binding protein 1. *J Biol Chem* 2010;285:483-492.

- 16 Heineke J, Auger-Messier M, Correll RN, Xu J, Benard MJ, Yuan W, Drexler H, Parise LV, Molkentin JD: CIB1 is a regulator of pathological cardiac hypertrophy. *Nat Med* 2010;16:872-879.
- 17 Zhu W, Gliddon BL, Jarman KE, Moretti PAB, Tin T, Parise LV, Woodcock JM, Powell JA, Ruszkiewicz A, Pitman MR, Pitson SM: CIB1 contributes to oncogenic signalling by Ras via modulating the subcellular localisation of sphingosine kinase 1. *Oncogene* 2017;36:2619-2627.
- 18 Tahara E, Jr., Tahara H, Kanno M, Naka K, Takeda Y, Matsuzaki T, Yamazaki R, Ishihara H, Yasui W, Barrett JC, Ide T, Tahara E: G1P3, an interferon inducible gene 6-16, is expressed in gastric cancers and inhibits mitochondrial-mediated apoptosis in gastric cancer cell line TMK-1 cell. *Cancer Immunol Immunother* 2005;54:729-740.
- 19 Yuan W, Leisner TM, McFadden AW, Clark S, Hiller S, Maeda N, O'Brien DA, Parise LV: CIB1 is essential for mouse spermatogenesis. *Mol Cell Biol* 2006;26:8507-8514.
- 20 Naik MU, Pham NT, Beebe K, Dai W, Naik UP: Calcium-dependent inhibition of polo-like kinase 3 activity by CIB1 in breast cancer cells. *Int J Cancer* 2011;128:587-596.
- 21 Nagayoshi Y, Nakamura M, Matsuoka K, Ohtsuka T, Mori Y, Kono H, Aso T, Ideno N, Takahata S, Ryo A, Takeda H, Ito T, Oda Y, Endo Y, Sawasaki T, Tanaka M: Profiling of autoantibodies in sera of pancreatic cancer patients. *Ann Surg Oncol* 2014;21:S459-465.
- 22 Blazejczyk M, Wojda U, Sobczak A, Spilker C, Bernstein HG, Gundelfinger ED, Kreutz MR, Kuznicki J: Ca²⁺-independent binding and cellular expression profiles question a significant role of calmyrin in transduction of Ca²⁺-signals to Alzheimer's disease-related presenilin 2 in forebrain. *Biochim Biophys Acta* 2006;1762:66-72.
- 23 Bernstein HG, Blazejczyk M, Rudka T, Gundelfinger ED, Dobrowolny H, Bogerts B, Kreutz MR, Kuznicki J, Wojda U: The Alzheimer disease-related calcium-binding protein Calmyrin is present in human forebrain with an altered distribution in Alzheimer's as compared to normal ageing brains. *Neuropathol Appl Neurobiol* 2005;31:314-324.
- 24 Fang X, Chen C, Wang Q, Gu J, Chi C: The interaction of the calcium- and integrin-binding protein (CIBP) with the coagulation factor VIII. *Thromb Res* 2001;102:177-185.
- 25 Guse AH, Lee HC: NAADP: a universal Ca²⁺ trigger. *Sci Signal* 2008;1:re10.
- 26 Zhang F, Jin S, Yi F, Li PL: TRP-ML1 functions as a lysosomal NAADP-sensitive Ca²⁺ release channel in coronary arterial myocytes. *J Cell Mol Med* 2009;13:3174-3185.
- 27 Wang X, Song J, Wu Z, Fan B, Mode X: Dual Roles of CD38 in Autophagy. *Yangtze Medicine* 2017;01:8-19.
- 28 Gerasimenko JV, Charlesworth RM, Sherwood MW, Ferdek PE, Mikoshiba K, Parrington J, Petersen OH, Gerasimenko OV: Both RyRs and TPCs are required for NAADP-induced intracellular Ca²⁺(+) release. *Cell Calcium* 2015;58:237-245.
- 29 Zhu S, Xu T, Luo Y, Zhang Y, Xuan H, Ma Y, Pan D, Li D, Zhu H: Luteolin Enhances Sarcoplasmic Reticulum Ca²⁺-ATPase Activity through p38 MAPK Signaling thus Improving Rat Cardiac Function after Ischemia/Reperfusion. *Cell Physiol Biochem* 2017;41:999-1010.
- 30 Frost RJ, Olson EN: Separating the good and evil of cardiac growth by CIB1 and calcineurin. *Cell Metab* 2010;12:205-206.
- 31 Tandan S, Wang Y, Wang TT, Jiang N, Hall DD, Hell JW, Luo X, Rothermel BA, Hill JA: Physical and functional interaction between calcineurin and the cardiac L-type Ca²⁺ channel. *Circ Res* 2009;105:51-60.
- 32 White C, Yang J, Monteiro MJ, Foscett JK: CIB1, a ubiquitously expressed Ca²⁺-binding protein ligand of the InsP3 receptor Ca²⁺ release channel. *J Biol Chem* 2006;281:20825-20833.
- 33 Howard M, Grimaldi JC, Bazan JF, Lund FE, Santos-Argumedo L, Parkhouse RM, Walseth TF, Lee HC: Formation and hydrolysis of cyclic ADP-ribose catalyzed by lymphocyte antigen CD38. *Science* 1993;262:1056-1059.
- 34 Aarhus R, Graeff RM, Dickey DM, Walseth TF, Lee HC: ADP-ribosyl cyclase and CD38 catalyze the synthesis of a calcium-mobilizing metabolite from NADP. *J Biol Chem* 1995;270:30327-30333.
- 35 Cui C, Merritt R, Fu L, Pan Z: Targeting calcium signaling in cancer therapy. *Acta Pharm Sin B* 2017;7:3-17.
- 36 Zhang Y, Zhang T, Wu C, Xia Q, Xu D: ASIC1a mediates the drug resistance of human hepatocellular carcinoma via the Ca²⁺/PI3-kinase/AKT signaling pathway. *Lab Invest* 2017;97:53-69.
- 37 Davis FM, Azimi I, Faville RA, Peters AA, Jalink K, Putney JW, Jr., Goodhill GJ, Thompson EW, Roberts-Thomson SJ, Monteith GR: Induction of epithelial-mesenchymal transition (EMT) in breast cancer cells is calcium signal dependent. *Oncogene* 2014;33:2307-2316.

- 38 Azimi I, Roberts-Thomson SJ, Monteith GR: Calcium influx pathways in breast cancer: opportunities for pharmacological intervention. *Br J Pharmacol* 2014;171:945-960.
- 39 Kauselmann G, Weiler M, Wulff P, Jessberger S, Konietzko U, Scafidi J, Staubli U, Bereiter-Hahn J, Strebhardt K, Kuhl D: The polo-like protein kinases Fnk and Snk associate with a Ca(2+)- and integrin-binding protein and are regulated dynamically with synaptic plasticity. *EMBO J* 1999;18:5528-5539.
- 40 Armacki M, Joodi G, Nimmagadda SC, de Kimpe L, Pusapati GV, Vandoninck S, Van Lint J, Illing A, Seufferlein T: A novel splice variant of calcium and integrin-binding protein 1 mediates protein kinase D2-stimulated tumour growth by regulating angiogenesis. *Oncogene* 2014;33:1167-1180.
- 41 Black JL, Harrell JC, Leisner TM, Fellmeth MJ, George SD, Reinhold D, Baker NM, Jones CD, Der CJ, Perou CM, Parise LV: CIB1 depletion impairs cell survival and tumor growth in triple-negative breast cancer. *Breast Cancer Res Treat* 2015;152:337-346.
- 42 Tong JR, Zhou HC, He F, Gao Y, Yang XQ, Luo ZM, Zhang H, Zeng JY, Wang Y, Huang YH, Zhang JL, Sun LH, He GL: Proteomic identification of CIB1 as a potential diagnostic factor in hepatocellular carcinoma. *J Biosciences* 2011;36:659-668.
- 43 Leisner TM, Moran C, Holly SP, Parise LV: CIB1 prevents nuclear GAPDH accumulation and non-apoptotic tumor cell death via AKT and ERK signaling. *Oncogene* 2013;32:4017-4027.
- 44 Naik MU, Naik UP: Calcium- and integrin-binding protein 1 regulates microtubule organization and centrosome segregation through polo like kinase 3 during cell cycle progression. *Int J Biochem Cell Biol* 2011;43:120-129.
- 45 Ma S, Liu MA, Yuan YL, Erikson RL: The serum-inducible protein kinase Snk is a G1 phase polo-like kinase that is inhibited by the calcium- and integrin-binding protein CIB. *Mol Cancer Res* 2003;1:376-384.
- 46 Holtrich U, Wolf G, Yuan J, Bereiter-Hahn J, Karn T, Weiler M, Kauselmann G, Rehli M, Andreesen R, Kaufmann M, Kuhl D, Strebhardt K: Adhesion induced expression of the serine/threonine kinase Fnk in human macrophages. *Oncogene* 2000;19:4832-4839.
- 47 Hao Q, Wang L, Zhao ZJ, Tang H: Identification of protein kinase D2 as a pivotal regulator of endothelial cell proliferation, migration, and angiogenesis. *J Biol Chem* 2009;284:799-806.
- 48 Xia P, Gamble JR, Wang L, Pitson SM, Moretti PA, Wattenberg BW, D'Andrea RJ, Vadas MA: An oncogenic role of sphingosine kinase. *Curr Biol* 2000;10:1527-1530.
- 49 Pitson SM, Moretti PA, Zebol JR, Xia P, Gamble JR, Vadas MA, D'Andrea RJ, Wattenberg BW: Expression of a catalytically inactive sphingosine kinase mutant blocks agonist-induced sphingosine kinase activation. A dominant-negative sphingosine kinase. *J Biol Chem* 2000;275:33945-33950.
- 50 Gault CR, Ebelen ST, Neumann CA, Hannun YA, Obeid LM: Oncogenic K-Ras regulates bioactive sphingolipids in a sphingosine kinase 1-dependent manner. *J Biol Chem* 2012;287:31794-31803.
- 51 Castaing-Berthou A, Malet N, Radojkovic C, Cabou C, Gayral S, Martinez LO, Laffargue M: PI3Kbeta Plays a Key Role in Apolipoprotein A-I-Induced Endothelial Cell Proliferation Through Activation of the Ecto-F1-ATPase/P2Y1 Receptors. *Cell Physiol Biochem* 2017;42:579-593.
- 52 Wang X, Chen WR, Xing D: A pathway from JNK through decreased ERK and Akt activities for FOXO3a nuclear translocation in response to UV irradiation. *J Cell Physiol* 2012;227:1168-1178.
- 53 Lee GE, Yu EY, Cho CH, Lee J, Muller MT, Chung IK: DNA-protein kinase catalytic subunit-interacting protein KIP binds telomerase by interacting with human telomerase reverse transcriptase. *J Biol Chem* 2004;279:34750-34755.
- 54 Khadka P, Lee JH, Baek SH, Oh SY, Chung IK: DNA-PKcs-interacting protein KIP binding to TRF2 is required for the maintenance of functional telomeres. *Biochem J* 2014;463:19-30.
- 55 Cen WJ, Pan Y, Li WX, Yang SQ: Comparison of DNA-dependent protein kinase catalytic subunit expression in two lung adenocarcinoma cell lines with different radiosensitivity. *Nan Fang Yi Ke Da Xue Xue Bao* 2009;29:2241-2243.
- 56 Henderson MJ, Russell AJ, Hird S, Munoz M, Clancy JL, Lehrbach GM, Calanni ST, Jans DA, Sutherland RL, Watts CK: EDD, the human hyperplastic discs protein, has a role in progesterone receptor coactivation and potential involvement in DNA damage response. *J Biol Chem* 2002;277:26468-26478.
- 57 Yoon KW, Cho JH, Lee JK, Kang YH, Chae JS, Kim YM, Kim J, Kim EK, Kim SE, Baik JH, Naik UP, Cho SG, Choi EJ: CIB1 functions as a Ca(2+)-sensitive modulator of stress-induced signaling by targeting ASK1. *Proc Natl Acad Sci U S A* 2009;106:17389-17394.
- 58 Wang X, Xing D, Liu L, Chen WR: BimL directly neutralizes Bcl-xL to promote Bax activation during UV-induced apoptosis. *FEBS Lett* 2009;583:1873-1879.

- 59 Wang X, He X, Hu S, Sun A, Lu C: Involvement of Bim in Photofrin-mediated photodynamically induced apoptosis. *Cell Physiol Biochem* 2015;35:1527-1536.
- 60 Hynes RO: Integrins: bidirectional, allosteric signaling machines. *Cell* 2002;110:673-687.
- 61 Shock DD, Naik UP, Brittain JE, Alahari SK, Sondek J, Parise LV: Calcium-dependent properties of CIB binding to the integrin alphaIIb cytoplasmic domain and translocation to the platelet cytoskeleton. *Biochem J* 1999;342 Pt 3:729-735.
- 62 Litvinov RI, Farrell DH, Weisel JW, Bennett JS: The Platelet Integrin alphaIIbbeta3 Differentially Interacts with Fibrin Versus Fibrinogen. *J Biol Chem* 2016;291:7858-7867.
- 63 Duval C, Ariens RAS: Fibrinogen splice variation and cross-linking: Effects on fibrin structure/function and role of fibrinogen gamma' as thrombomodulin II. *Matrix Biol* 2017;60-61:8-15.
- 64 Naik MU, Naik TU, Summer R, Naik UP: Binding of CIB1 to the alphaIIb tail of alphaIIbbeta3 is required for FAK recruitment and activation in platelets. *PLoS One* 2017;12:e0176602.
- 65 Tsuboi S, Nonoyama S, Ochs HD: Wiskott-Aldrich syndrome protein is involved in alphaIIb beta3-mediated cell adhesion. *EMBO Rep* 2006;7:506-511.
- 66 Haataja L, Kaartinen V, Groffen J, Heisterkamp N: The small GTPase Rac3 interacts with the integrin-binding protein CIB and promotes integrin alpha(IIb)beta(3)-mediated adhesion and spreading. *J Biol Chem* 2002;277:8321-8328.
- 67 Leisner TM, Liu M, Jaffer ZM, Chernoff J, Parise LV: Essential role of CIB1 in regulating PAK1 activation and cell migration. *J Cell Biol* 2005;170:465-476.
- 68 Zayed MA, Yuan W, Chalothorn D, Faber JE, Parise LV: Tumor growth and angiogenesis is impaired in CIB1 knockout mice. *J Angiogenesis Res* 2010;2:17.
- 69 Naik MU, Naik UP: Contra-regulation of calcium- and integrin-binding protein 1-induced cell migration on fibronectin by PAK1 and MAP kinase signaling. *J Cell Biochem* 2011;112:3289-3299.
- 70 Naik MU, Naik UP: Calcium- and integrin-binding protein regulates focal adhesion kinase activity during platelet spreading on immobilized fibrinogen. *Blood* 2003;102:3629-3636.
- 71 Zayed MA, Yuan W, Leisner TM, Chalothorn D, McFadden AW, Schaller MD, Hartnett ME, Faber JE, Parise LV: CIB1 regulates endothelial cells and ischemia-induced pathological and adaptive angiogenesis. *Circ Res* 2007;101:1185-1193.
- 72 Sobczak A, Debowska K, Blazejczyk M, Kreutz MR, Kuznicki J, Wojda U: Calmyrin1 binds to SCG10 protein (stathmin2) to modulate neurite outgrowth. *Biochim Biophys Acta* 2011;1813:1025-1037.
- 73 Whitehouse C, Chambers J, Howe K, Cobourne M, Sharpe P, Solomon E: NBR1 interacts with fasciculation and elongation protein zeta-1 (FEZ1) and calcium and integrin binding protein (CIB) and shows developmentally restricted expression in the neural tube. *Eur J Biochem* 2002;269:538-545.
- 74 Wang S, Sun Z, Zhang X, Li Z, Wu M, Zhao W, Wang H, Chen T, Yan H, Zhu J: Wnt1 positively regulates CD36 expression via TCF4 and PPAR-gamma in macrophages. *Cell Physiol Biochem* 2015;35:1289-1302.
- 75 Hollenbach AD, McPherson CJ, Lagutina I, Grosveld G: The EF-hand calcium-binding protein calmyrin inhibits the transcriptional and DNA-binding activity of Pax3. *Biochim Biophys Acta* 2002;1574:321-328.
- 76 Sun W, Guan Q, Wen J, Zhang Q, Yang W, Zhang B, Cui W, Zou Z, Yu Y: Calcium- and integrin-binding protein-1 is down-regulated in the sperm of patients with oligoasthenozoospermia : CIB1 expression in patients with oligoasthenozoospermia. *J Assist Reprod Genet* 2014;31:541-547.
- 77 Godinho-Santos A, Hance AJ, Gonçalves J, Mammano F: CIB1 and CIB2 are HIV-1 helper factors involved in viral entry. *Sci Rep* 2016;6:30927.
- 78 Bandyopadhyay C, Valiya-Veetil M, Dutta D, Chakraborty S, Chandran B: CIB1 synergizes with EphrinA2 to regulate Kaposi's sarcoma-associated herpesvirus macropinocytic entry in human microvascular dermal endothelial cells. *PLoS Pathog* 2014;10:e1003941.
- 79 Zhao F, Zhang S, Chen L, Wu Y, Qin J, Shao Y, Wang X, Chen Y: Calcium- and integrin-binding protein-1 and calcineurin are upregulated in the right atrial myocardium of patients with atrial fibrillation. *Europace* 2012;14:1726-1733.
- 80 Weljie AM, Yamniuk AP, Yoshino H, Izumi Y, Vogel HJ: Protein conformational changes studied by diffusion NMR spectroscopy: application to helix-loop-helix calcium binding proteins. *Protein Sci* 2003;12:228-236.