An Overview on Renoprotective Effects of Thymoquinone

Hanieh Shaterzadeh-Yazdi^{a, b} Mohammad-Foad Noorbakhsh^c
Saeed Samarghandian^d Tahereh Farkhondeh^e

^aResearch Center of Pardis Hospital, Mashhad, Iran; ^bNastaran Center for Cancer Prevention, Mashhad, Iran;
^cDivision of Pharmacology and Toxicology, Department of Basic Sciences, School of Veterinary Medicine, Shiraz University, Shiraz, Iran; ^dDepartment of Basic Medical Sciences, Neyshabur University of Medical Sciences, Neyshabur, Iran; ^eCardiovascular Diseases Research Center, Birjand University of Medical Sciences, Birjand, Iran

Keywords

Kidney · Inflammation · Oxidative stress · Renoprotective effects · Thymoquinone

Abstract

Background: Kidneys as vital organs remove waste material from blood. Additionally, they may also have a role in the electrolyte balance, regulation of blood pressure, and red blood cell genesis. Kidney diseases may be caused by several factors such as ischemia/reperfusion injury, diabetes, and nephrotoxic agents. Oxidative stress and inflammation are involved in the pathogenesis and progression of kidney diseases. Traditionally, natural antioxidants are used for treatment of renal failure in various countries. **Summary:** People usually select natural antioxidants since they have an opinion that herbal medicine has not any important side effects. *Nigella sativa* is a flavoring herb that is widely used as a condiment and as a remedy for many disorders. Thymoquinone (TQ), the most important component of black seeds, mainly oil, is considered as an active agent responsible for a lot of the seed's useful effects. This review describes the protective roles and related mechanisms of TQ against renal disorders. The search terms, including TQ, antioxidant, renal

ischemia-reperfusion, diabetic nephropathy, and nephrotoxic agent were searched in scientific databases. TQ showed anti-inflammatory and antioxidant properties in animal and in vitro models of several renal diseases caused by inflammation and oxidative stress. **Key Messages:** Experimental studies have shown beneficial effects of TQ against renal diseases; however, well-designed clinical trials in humans are required to confirm these effects.

© 2018 S. Karger AG, Basel

Background

Nigella sativa, generally known as black seed or black cumin, is an annual flowering bush plant from the family Ranunculaceae which is growing in Southwest Asia, North Africa, and southern Europe [1, 2]. The main compounds that have been isolated from *N. sativa* seed include proteins, carbohydrates, fixed oils, essential oil, crude fiber, alkaloids, minerals, vitamins, ash, and moisture. The other components are tannins, resin, saponin, carotene, glucosides, and sterols [3]. One of the major components of the essential oil is thymoquinone (TQ) [4, 5]. Based on the use of *N. sativa* in traditional medicine

as a natural treatment for some diseases, researchers have investigated its protective effects against asthma [6], hypertension [7], diabetes [8], and inflammation [9]. In addition, TQ is known to have antioxidant [10], analgesic, antipyretic [11], antischistosomal [12], antifungal [13], antibacterial [14], anticancer [15], anticonvulsant [16], hepatoprotective [17], and neuroprotective activities [18]. It has been found that TQ has beneficial protective effects against renal diseases through anti-inflammatory, antioxidant, and antiapoptotic activities [19]. The present review aimed to review the recent studies from 2007 to 2017 that describe the protective effects of TQ in the management of kidney diseases.

Evidence Acquisition

Online literature resources were checked using different search engines such as Medline, PubMed, Scopus, and Google Scholar from 2000 to 2017 to identify articles, editorials, and reviews about renoprotective effects of TQ. In this review article, the words TQ, antioxidant, renal ischemia-reperfusion, diabetic nephropathy, and nephrotoxic agents were applied to search the literature from scientific databases.

Results

Antioxidant and Anti-Inflammatory Effects of TQ

Reactive oxygen species (ROS) have been described as an important cause of renal disorders [20–23]. Reducing ROS improves graft maintenance and removal of acute inflammation. The scavenging function of free radicals is a property of black seeds which can prevent lipid peroxidation (LPO). The studies showed that TQ, a major constituent of *N. sativa* seeds, has strong antioxidant, anti-inflammatory, and cytoprotective abilities [24–28]. In this regard, Khat tab and Nagi [29] studied the protective effect of TQ in the rats with hypertension induced by 4 weeks' administration of N (omega)-nitro-L-arginine methyl esters (L-NAME) (50 mg/kg/day p.o.). TQ treatment (0.5 and 1 mg/kg/day p.o.) decreased the increase in systolic blood pressure induced by L-NAME. Treatment of the rats with TQ brought increased creatinine (Cr) and glutathione (GSH) to normal levels. TQ prevented the production of superoxide radical in enzymatic and non-enzymatic systems. This study supported the effects of TQ in the rats against L-NAME-induced hypertension and kidney injury through its antioxidant activity [29]. The inhibitory effects of TQ (100, 200, 300, 400, 500, 750, 1,000 nM) on activation of interleukin-6 (IL-6) and re-

dox-sensitive transcription factor nuclear factor kappa B (NF- κ B) were investigated in vitro. An important reduction of advanced glycation end products induced high expression of IL-6 and NF- κ B in the human proximal tubular epithelial cells. This result showed antioxidative properties of TQ [30].

Ozer et al. [31] investigated the protective effects of TQ on survival, vascular reactivity, mesenteric artery blood flow, oxidative, and inflammatory responses in rats subjected to cecal ligation and puncture (CLP). Intraperitoneal injection (i.p.) of TQ (1 mg/kg/day) for 3 days partially reduced aortic dysfunction and improved mesenteric hypoperfusion induced by CLP. TQ also prevented the increase in the serum levels of alanine aminotransferase, aspartate aminotransferase, blood urea nitrogen (BUN), lactate dehydrogenase, Cr, and inflammatory cytokines (IL-1 β , IL-6 and TNF- α) in the CLP group. The reduction of the GSH levels in spleen, liver, and kidney and the elevation of the malondialdehyde (MDA) levels in kidney, liver, lung, and spleen induced by CLP were prevented by TQ. Additionally, TQ showed the histopathological protective effects on multiple organ injury induced by CLP [31]. Another study supported the ameliorative effects of TQ in renal disorder caused by rheumatoid arthritis in rats. In this study, TQ (2 mg/kg dissolved in distilled water, i.p.) and methotrexate (MTX) (0.5 mg/kg, i.p.) importantly decreased the total clinical score of inflammation and leukocyte numbers, and ameliorated serum Cr and blood urea [32]. Hosseini et al. [33] investigated the effects of TQ and renin-angiotensin system blockade on the renal expression of angiotensin II and renal tissue injury after unilateral ureteral obstruction (UUO) in rats. TQ (10 mg/kg, i.p.) 3 days before UUO and 2 weeks postoperatively importantly ameliorated apoptosis, TNF- α expression, and oxidative damage, but significantly reduced the upregulation of monocyte chemoattractant protein (MCP-1) and angiotensin II compared with the UUO group. This study showed the ability of TQ to improve UUO-induced renal tissue injury, which was comparable to the well-known renin-angiotensin system inhibitors such as losartan and captopril [33].

Ischemia/Reperfusion Injury

Renal ischemia-reperfusion (I/R) is known as a well-characterized model of acute renal injury that leads to both local and far organ damage [34–37]. Various studies have shown that TQ has protective effects against I/R injury to the several organs such as kidneys, lungs, and liver in different experimental models [38, 39]. The effect of

TQ on hepatic and renal changes induced by renal I/R in the rats was investigated. In this study, administration of TQ (10 mg/kg, p.o.) before I/R injury led to the reversal of increased MDA content and decreased activity of superoxide dismutase (SOD) and glutathione-S-transferase (GST) in renal and hepatic tissues. Additionally, it decreased the raised alanine aminotransferase and Cr serum levels and induced CYP3A1 mRNA expression by I/R. TQ also decreased spermidine/spermine N-1-acetyltransferase mRNA expression as a catabolic enzyme that participates in polyamine metabolism in the liver and kidney which was induced by I/R [40]. A study was done to investigate the effect of TQ on the changes in parameters of kidney function after the warm renal I/R injury (I/RI) in rats. Oral consumption of TQ (10 mg/kg/day) 4 days prior to the I/RI for 6 days improved the I/RI effects on the tubular renal functional and hemodynamic parameters as well as the expression of some renal damage markers, profibrotic and proinflammatory cytokines, which showed TQ has renoprotective effects on I/RI-induced renal disorders [39].

Diabetic Nephropathy

Diabetic nephropathy is the main reason of morbidity and death in diabetic people. To inhibit the expansion of this disease and to ameliorate progressed renal damage, effective remedies are needed [41–43]. Streptozotocin (STZ) is an antibiotic extensively applied in experimental models as an agent capable of inducing type 1 diabetes mellitus, which is known as insulin-dependent diabetes mellitus [44–46]. Studies show that *Nigella sativa* oil (NSO) and its main bioactive element TQ have, antihyperglycemic, antioxidant, and renoprotective properties in STZ-induced diabetes in the animal models [47, 48]. The study was done by Sayed [49], which showed that TQ reduced NF- κ B activation induced by angiotensin II in a dose-dependent manner, with the maximum inhibitory effect of 500 nM. In addition, preincubation of the proximal tubular epithelial cells with TQ caused the disappearance of the second peak of NF- κ B. The results demonstrated the therapeutic effect of TQ which can be administered to delay end-stage kidney disorders in diabetics. The study performed by Kanter [50] showed that the treatment with TQ (50 mg/kg body weight) by oral gavage in diabetic rats for 12 weeks decreased the capsular thickening, glomerular size, the tubular and glomerular basement membranes, the elevated content of mesangial matrix, and tubular dilatation and renal function as compared with the untreated diabetic rats. The findings showed that TQ treatment leads

to the renal structural and functional progress in rats with STZ-induced diabetes. The study was carried out to investigate the feasible beneficial effects of proanthocyanidin (PA), TQ, and their combination to ameliorate the diabetic nephropathy in rats. Treatment of rats with PA (250 mg/kg), TQ (50 mg/kg), or PA + TQ (250 + 50 mg/kg, respectively) orally for 12 weeks showed a decrease in body weight, in addition to reduced SOD and GSH contents. The increased levels of nitric oxide (NO), urea, Cr, IL-6, and MDA were significantly decreased as a result of the treatment. The results indicated that PA and TQ treatment could be introduced as therapeutic protective agents in diabetes through reducing oxidative stress and ameliorating diabetic nephropathy. Therefore, TQ and PA may be clinically effective for supporting the diabetic kidney against oxidative stress [51]. Another study was done to investigate the effects of TQ in the rats with STZ-induced diabetes and to define the predictive cost of epithelial and mesenchymal markers in TQ treatment in rats with diabetic nephropathy. Treatment with TQ (50 mg/kg, orally) importantly ameliorated renal structural changes in both renal tubules and glomeruli and also the immunohistochemical expression of MMP-17 and mesenchymal markers Fsp1, desmin, and epithelial marker ZO-1 disappearance in the glomeruli of STZ-induced diabetic rats. These findings showed that both epithelial and mesenchymal markers act as good predictors of early renal injury and indexes of TQ reactivity in STZ-induced diabetic nephropathy [52]. A study was designed by Al-Trad et al. [53] to assess the effect of NSO and TQ treatment on podocyte injury, albuminuria, and the complex systems adjusting the extracellular angiogenesis and matrix protein accumulation in rats with STZ-induced diabetic nephropathy. The results indicated that treatment with 2 mL/kg NSO or 50 mg/kg TQ by oral consumption for 10 weeks importantly ameliorated increased albuminuria and the kidney weight/body weight ratio in STZ-induced diabetic nephropathy. The real-time PCR displayed that the treatment with NSO and TQ inhibited diabetes-induced mRNA overexpression of transforming growth factor- β 1 (TGF- β 1), vascular endothelial growth factor-A (VEGF-A) and collagen IV as well as down-regulation of mRNA expression of the podocyte-specific marker (podocin) in the diabetic kidney. NSO and TQ treatment reduced albuminuria in the diabetic nephropathy experimental models through the preservation of the podocyte function with the suppression of increased extracellular matrix gene expression by preventing TGF- β 1 production and angiogenesis [53].

Nephrotoxic Agents

It is well defined that many drugs, such as the chemotherapeutic drug (cisplatin) and aminoglycoside antibiotics are able to cause nephrotoxicity [54–56].

Chemotherapy Drugs

Most chemotherapeutic drugs can lead to nephrotoxicity [57, 58]. Hence, recent strategies have been focused on preventing chemotherapy-induced nephrotoxicity. *N. sativa* or its isolated constituent TQ are introduced as the potential agent in fighting chemotherapy-induced nephrotoxicity [59].

Doxorubicin

The study carried out by Elsherbiny and El-Sherbiny [60] showed that oral consumption of TQ (50 mg/kg/day) for 3 weeks in rats that received doxorubicin (DOX) as a chemotherapeutic drug, reversed the increased levels of serum Cr, urea, and urinary albumin excretion, whereas animals that received DOX without TQ showed DOX-induced nephrotoxicity. Additionally, TQ ameliorated increased MDA and LPO, and also, it increased the decreased activities of SOD and GST in rats with DOX-induced nephrotoxicity. TQ also decreased the increased renal levels of IL-6, TNF- α , and NADPH oxidase 4, and increased the decrease in nuclear factor erythroid 2-related factor 2 (Nrf2) mRNA levels, interleukin-10 (IL-10) levels, and nuclear binding activity. Renal histopathological changes induced by DOX improved with TQ [60].

Cisplatin and Diesel Exhaust Particles

Ali et al. [61] studied the interaction between cisplatin (CP) nephrotoxicity and a single exposure to diesel exhaust particles and the concurrent treatment with TQ (20 mg/kg, orally) in rats. The results showed that several biochemical, physiological, and histopathological alterations contained decreased growth and Cr clearance and increased IL-6, plasma neutrophil gelatinase-associated lipocalin, C-reactive protein, urea and Cr concentrations, and urinary N-acetyl-b-D-glucosaminidase (NAG) activities induced by CP, and were significantly abrogated by TQ. Also, TQ significantly resolved potentiated indices of oxidative injuries in the kidney tissues, and induced renal tubular necrosis induced by CP and diesel exhaust particles [61].

Morphine

A study was designed by Jalili et al. [62] to investigate protective effects of TQ against morphine-induced renal toxicity in mice. The results showed that intraperitoneal

administration of TQ (4.5, 9, and 18 mg/kg) and TQ with morphine importantly increased kidney weight, number, and mean diameter of the glomeruli. Additionally, TQ led to a decrease in BUN, serum NO and serum Cr levels in the groups treated with it compared to the morphine group. This study explained that the renoprotective effects of TQ against damage due to morphine toxicity may be related to its antioxidant and antiapoptotic effects [62].

Gentamicin

Sayed-Ahmed and Nagi [63] showed that TQ supplementation (50 mg/L in drinking water) in rat caused a complete reversal of the gentamicin (GM)-induced increase in BUN, Cr, thiobarbituric acid-reactive substances and total nitrate/nitrite and reduction in GSH, glutathione peroxidase (GPx), catalase (CAT), and ATP to control values. Furthermore, histopathological examination of renal tissues corroborated the biochemical data, wherein TQ supplementation protected kidney against GM-induced degenerative changes. The results of this study proposed that TQ is useful in preventing from the development of GM-induced acute renal failure by a mechanism related somewhat to its ability to reduce oxidative stress and to maintain the activity of the antioxidant enzymes, as well as its ability to prevent the energy reduction in renal tissues [63]. In another study, Samarghandian et al. [64] indicated that the significant elevation in the levels of serum Cr, BUN, MDA, IL-6, IL-18, IL-1 β , and TNF- α and also the reduction of GSH, SOD, GPx, and IL-10 in the GM nephrotoxicity were improved by TQ (10, 20, 30 mg/kg), in a dose-dependent manner. These investigators proposed that TQ may improve acute renal failure by modulation of the oxidative stress and inflammatory responses.

Methotrexate

The study by El-Sheikh et al. [65] indicated that TQ treatment (10 mg/kg) concurrently with MTX in rats restored hepatorenal functions, as well as their normal histology. TQ also reversed oxidative and nitrosative stress, as well as inflammatory and apoptotic signs induced by MTX alone. This study proposed that TQ may be the useful adjuvant that gives hepatorenal protection to MTX toxicity through anti-inflammatory, antioxidant, antinitrosative, and antiapoptotic mechanisms.

Acetaminophen

Aycan et al. [66] studied the efficacy of TQ (10 mg/kg, i.p.) in acetaminophen (APAP)-induced renal toxicity in rat. The Cr and urea levels in the APAP + TQ treated

group were importantly lower than in the APAP-treated group. In contrast to the group treated with APAP, the MDA levels and serum NO activity were significantly lower in the TQ + APAP group. Histopathological study showed lower tissue damage scores in the APAP + TQ group compared with the APAP group.

Cyclosporine A

Farag et al. [38] investigated the effects of chronic cyclosporine A (CsA) treatment and acute renal I/R on the liver and kidney in rats treated with TQ. TQ treatment (10 mg/kg per day) decreased serum indicators back to control levels and improved CsA-induced renal and hepatic histopathological alterations. In renal and hepatic tissues, CsA and renal I/R induced considerable elevation of MDA levels with remarkable reduction in decreased GSH levels and SOD activities, as oxidative stress markers were neutralized by TQ therapy. This study showed that kidney and liver damage related to CsA or renal I/R can be significantly decreased by TQ.

Lead

Mabrouk and Ben Cheikh [67] investigated the possible beneficial effect of TQ on lead (Pb)-induced renal antioxidant defense system dysfunction in the rats. TQ treatment (5 mg/kg/day, p.o.) improved the decreased SOD and GSH level, CAT, GPx, and glutathione reductase activities in the kidney tissue induced by Pb [67].

Arsenic

A study was carried out on the protective role of TQ against renal injury induced by arsenic in rats. Renal degenerative changes induced by arsenic were decreased by TQ (10 mg/kg, intragastric gavage for 15 days). Also, TQ ameliorated the increased MDA levels and increased the reduced SOD, CAT, and GSH-Px enzyme activities [68].

Cadmium

Erboga et al. [69] showed that TQ (50 mg/kg body weight) once a day orally in rats reduced the nephrotoxicity of cadmium (Cd) (1 mg/kg body weight). TQ preserved the normal histological architecture of the renal tissue. Furthermore, the immunohistochemical analysis indicated that TQ importantly reduced the Cd-induced overexpression of NF- κ B in the kidney and also decreased apoptotic cells. TQ significantly repressed LPO and compensated dearth in the antioxidant defenses of the kidney. These results proposed that the nephroprotective potential of TQ in Cd toxicity may be related to its antioxidant

and antiapoptotic effects, which could be helpful for obtaining optimum results in the Cd-induced nephrotoxicity [69].

Mercuric Chloride

Fouda et al. [70] investigated protective effects of TQ in mercuric chloride (HgCl₂)-induced nephrotoxicity. Their results showed that the deterioration of antioxidant enzymes, increase in serum Cr and histological injury induced by HgCl₂ were markedly ameliorated by TQ therapy (10 mg/kg/day). Additionally, apoptosis and proliferative reactions were decreased. The maximal protective effect proposed by TQ was 48 and 72 h after administration of HgCl₂, when histological damage, renal cell apoptosis, and proliferative reactions reached their maximum. The findings showed the antioxidant effect of TQ.

Kidney Calculi

Kidney calculi are the term which is used to explain renal stones, which are formed in the urine that is made up of acid and mineral salts. They can block urine and cause renal infections, injury, and pain when they move around the body [71, 72]. A study was done to investigate the effects of TQ on ethylene glycol-induced renal calculi in rats. Intraperitoneal injection of TQ (5 mg/kg) caused fewer deposits. Also, TQ (5 mg/kg or 10 mg/kg, i.p.) importantly reduced the size and number of calcium oxalate sediments in the renal tubules [73]. Table 1 gives data on the protective effects of TQ on kidney diseases.

TQ Toxicity

Few studies reported TQ toxicity signs and symptoms in animal models. It was found that intraperitoneal injection of TQ caused peritonitis and abdominal muscle contractions and the oral administration of TQ caused dyspnea at different dose frequency exposure (acute, subacute, and subchronic) in rats and mice [74]. The maximum tolerated dose (MTD) of TQ, which is defined as the highest dose that is safe to administer to animal models in the absence of intolerable adverse effects, was determined in male and female Wistar rats [75]. The findings indicated that the MTD for intraperitoneal injection was 22.5 mg/kg in male rats and 15 mg/kg in females, whereas for oral administration it was 250 mg/kg in both male and female rats. Two rats died after administration of 500 mg/kg of TQ due to bowel obstruction. However, a possible mechanism of TQ toxicity was not fully understood [75].

Table 1. Protective effects of TQ on kidney diseases

Experimental model	Effect	Ref.
Rat	Decreased systolic blood pressure induced by L-NAME, and Cr and also elevated GSH levels	[29]
In vitro	Prevented from superoxide radical production in enzymatic and non-enzymatic systems Prevented AGEs-induced IL-6 and NF- κ B-activation in cultivated human pTECs	[30]
Rat	Decreased aortic dysfunction and improved mesenteric hypoperfusion induced by CLP, via decreasing the levels of inflammatory cytokines (IL-1, IL-6, TNF- α) and MDA and also increasing the GSH in kidney tissue	[31]
	Decreased the total clinical score of inflammation and leukocyte number and ameliorated serum Cr and blood urea in renal disorder caused by rheumatoid arthritis	[32]
	Prevented CsA-induced nephrotoxicity via decreasing the levels of MDA and increasing GSH level and SOD activity	[38]
	Prevented renal damage induced by unilateral ureteral obstruction via decreasing apoptosis, TNF- α expression, oxidative damage, upregulation MCP-1 and angiotensin	[33]
	Prevented renal damage induced by IR via decreasing profibrotic and proinflammatory cytokines	[39]
	Prevented renal damage induced by IR via decreasing MDA and SSAT mRNA expression as a catabolic enzyme that participates in polyamine metabolism	[40]
In vitro	Prevented renal tubular cells against tubular injury induced by AT II via decreasing NF- κ B activation	[49]
Rat	Decreased the capsular thickening, glomerular size, tubular and glomerular basement membranes, elevated content of mesangial matrix and tubular dilatation, and renal function as compared untreated diabetic rats	[50]
	Prevented diabetic kidney via decreasing IL-6, NO, and MDA and also increasing SOD and GSH in kidney tissue	[51]
	Prevented renal structural changes induced by STZ via modulating the expression of the MMP-17 and mesenchymal markers Fsp1, desmin, and epithelial marker ZO-1 in the glomeruli	[52]
	Prevented diabetic nephropathy via inhibiting mRNA overexpression of TGF- β 1, VEGF-A and collagen IV as well as downregulation of mRNA expression of the podocyte-specific marker (podocin) in the kidney	[53]
	Prevented DOX-induced nephrotoxicity via decreasing the levels of MDA, LPO, IL-6, TNF- α , and NOX-4 and also increasing the activities of SOD and GST, and Nrf2 mRNA l and IL-10 levels	[60]
	Prevented CP-induced nephrotoxicity via decreasing the plasma levels of IL-6, NGALa, CRP, and NAG activity	[61]
Mice	Prevented morphine-induced nephrotoxicity via decreasing the levels of blood urea nitrogen, NO, and serum levels of Cr	[62]
Rat	Prevented GM-induced nephrotoxicity via decreasing the serum levels of BUN, Cr, MDA, IL-6, IL-18, IL-1 β , TNF- α and total NOx and increasing the levels of GSH, GPx, CAT, IL-10, and ATP	[63, 64]
	Prevented MTX-induced nephrotoxicity via modulating nitrosative stress, as well as inflammatory and apoptotic responses	[65]
	Prevented APAP-induced nephrotoxicity via decreasing the serum levels of MDA and NO	[66]
	Prevented Pb-induced nephrotoxicity via increasing the levels of SOD, GSH, CAT, GPx, and GR	[67]
	Prevented Ar-induced nephrotoxicity via decreasing the levels of MDA and increasing the activities of SOD, CAT and GSH-Px enzyme	[68]
	Prevented Cd-induced nephrotoxicity via modulating NF- κ B in kidney	[69]
	Prevented HgCl ₂ -induced nephrotoxicity via increasing antioxidant enzymes and decreasing apoptosis and proliferative responses	[70]
	Prevented ethylene glycol-induced nephrotoxicity via reducing the size and number of calcium oxalate sediments in the renal tubules	[73]

L-NAME, N(omega)-nitro-L-arginine methyl esters; GSH, glutathione peroxidase; AGEs, advanced glycation end products; pTECs, proximal tubular epithelial cells; IL, interleukin; CLP, cecal ligation and puncture; ALT, alanine aminotransferase; AST, aspartate aminotransferase; BUN, blood urea nitrogen; LDH, lactate dehydrogenase; Cr, creatinine; MDA, malondialdehyde; TNF- α , tumor necrosis factor; MCP-1, monocyte chemoattractant protein; UUO, unilateral ureteral obstruction; SOD, superoxide dismutase; GST, glutathione-S-transferase; I/R, ischemia-reperfusion; SSAT, spermidine/spermine N-1-acetyltransferase; IRI, ischemia-reperfusion injury; AT II, angiotensin II; pTECs, proximal tubular epithelial cells; SOD, superoxide dismutase; NO, nitric oxide; STZ, streptozotocin; TGF- β 1, transforming growth factor- β 1, VEGF-A, vascular endothelial growth factor-A; VEGF-A, vascular endothelial growth factor-A; DOX, doxorubicin; LPO, lipid peroxidation; GST, glutathione-S-transferase; NOX-4, NADPH oxidase 4; Nrf2, nuclear factor erythroid 2-related factor 2; NGAL, neutrophil gelatinase-associated lipocalin; CRP, C-reactive protein; NAG, N-acetyl-b-D-glucosaminidase; GM, gentamicin; TBARS, thiobarbituric acid-reactive substances; NOx, nitrate/nitrite; GSH, glutathione peroxidase; GPx, glutathione peroxidase; CAT, catalase; ARF, acute renal failure; MTX, methotrexate; APAP, acetaminophen; CsA, cyclosporine A; Pb, lead; Cd, cadmium; NF- κ B, nuclear factor kappa B; HgCl₂, mercuric chloride.

Conclusions

Scientific interest in medicinal plants and their ingredients has increased because of high efficiency of herb-derived drugs, increasing interest in natural products. *N. sativa* seeds (Ranunculaceae family), have been used for thousands of years as a condiment and food preservative. TQ as the bioactive component of fugacious oil of black seed has indicated potent medicinal effects in traditional medicine. The studies provided clear documents that TQ has antioxidant effects via strengthening the oxidant scavenging system, and thus antitoxic properties. Also, it has shown potential anti-inflammatory effects [76, 77]. This review indicated medicinal or protective features of TQ against inflammation and oxidative stress in renal disorders. TQ showed anti-inflammatory and antioxidant properties in animal and in vitro models of several renal diseases caused by inflammation and oxidative stress. TQ indicated anti-inflammatory effects through modulation of inflammatory molecules IL-1, IL-6, IL-18, IL-10, and TNF- α , and NF- κ B activation. It also reduced

the total clinical score of inflammation and leukocyte number. TQ decreased oxidative stress markers such as MDA and increased antioxidant content, including GSH, CAT, GST, and SOD, which have an important role in kidney disorders. It also ameliorated serum Cr and blood urea in renal disorder caused by rheumatoid arthritis or morphine in rats. In ethylene glycol-induced renal calculi, TQ led to lower deposits and decreased the size and number of calcium oxalate sediments in the renal tubules. Additionally, TQ improved the histopathological changes of the kidney induced by CLP, nephrotoxic agents, and diabetic nephropathies. Although experimental studies have shown the beneficial effects of TQ against renal diseases, well-designed clinical trials in humans are required to confirm these effects.

Conflict of Interest Statement

The authors declare that there is no conflict of interest regarding the publication of this article. There was no financial support or sponsorship for this work.

References

- 1 El-Tahir KE, Bakeet DM: The black seed *Nigella sativa* Linnaeus – a mine for multi cures: a plea for urgent clinical evaluation of its volatile oil. *J Taibah Univ Sci* 2006;1:1–9.
- 2 Aljabre SH, Alakloby OM, Randhawa MA: Dermatological effects of *Nigella sativa*. *J Dermatol Dermatol Surg* 2015;19:92–98.
- 3 Randhawa MA, Al-Ghamdi MS: A review of the pharmacotherapeutic effects of *Nigella sativa*. *Pak J Med Res* 2002;41:77–83.
- 4 Venkatachallam SK, Pattekhani H, Divakar S, Kadimi US: Chemical composition of *Nigella sativa* L. seed extracts obtained by supercritical carbon dioxide. *J Food Sci Technol* 2010; 47:598–605.
- 5 Samarghandian S, Farkhondeh T, Samini F, Borji A: Protective effects of carvacrol against oxidative stress induced by chronic stress in rat's brain, liver, and kidney. *Biochem Res Int* 2016;2016:2645237.
- 6 Samarghandian S, Farkhondeh T, Samini F, Borji A: Protective effects of carvacrol against oxidative stress induced by chronic stress in rat's brain, liver, and kidney. *Biochem Res Int* 2016;2016:2645237.
- 7 Dehkordi FR, Kamkhah AF: Antihypertensive effect of *Nigella sativa* seed extract in patients with mild hypertension. *Fundam Clin Pharmacol* 2008;22:447–452.
- 8 Samarghandian S, Borji A: Anticarcinogenic effect of saffron (*Crocus sativus* L.) and its ingredients. *Pharmacognosy Res* 2014;6:99–107.
- 9 Chehl N, Chipitsyna G, Gong Q, Yeo CJ, Arafa HA: Anti-inflammatory effects of the *Nigella sativa* seed extract, thymoquinone, in pancreatic cancer cells. *HPB (Oxford)* 2009; 11:373–381.
- 10 Samarghandian S, Azimi-Nezhad M, Farkhondeh T: Crocin attenuate tumor necrosis factor- α (TNF- α) and interleukin-6 (IL-6) in streptozotocin-induced diabetic rat aorta. *Cytokine* 2016;88:20–28.
- 11 Al-Ghamdi M: The anti-inflammatory, analgesic and antipyretic activity of *Nigella sativa*. *J Ethnopharmacol* 2001;176:45–48.
- 12 Samarghandian S, Azimi-Nezhad M, Farkhondeh T, Samini F: Anti-oxidative effects of curcumin on immobilization-induced oxidative stress in rat brain, liver and kidney. *Biomed Pharmacother* 2017;87:223–229.
- 13 Islam S, Ahsan M, Hassan CM, Malek MA: Antifungal activities of the oils of *Nigella sativa* seeds. *Pak J Pharm Sci* 1989;2:25–28.
- 14 Morsi NM: Antimicrobial effect of crude extracts of *Nigella sativa* on multiple antibiotics-resistant bacteria. *Acta Microbiol Pol* 1999; 49:63–74.
- 15 Mahmoud SS, Torchilin VP: Hormetic/cytotoxic effects of *Nigella sativa* seed alcoholic and aqueous extracts on MCF-7 breast cancer cells alone or in combination with doxorubicin. *Cell Biochem Biophys* 2013;66:451–460.
- 16 Raza M, Alghasham AA, Alorainy MS, El-Hadiyiah TM: Potentiation of valproate-induced anticonvulsant response by *Nigella sativa* seed constituents: the role of GABA receptors. *Int J Health Sci* 2008;2:15.
- 17 Kanter M, Coskun O, Budancamanak M: Hepatoprotective effects of *Nigella sativa* L and *Urtica dioica* L on lipid peroxidation, antioxidant enzyme systems and liver enzymes in carbontetrachloride-treated rats. *World J Gastroenterol* 2005;11:6684–6688.
- 18 Khazdair MR: The protective effects of *Nigella sativa* and its constituents on induced neurotoxicity. *J Toxicol* 2015;2015:841823.
- 19 Al Mofleh IA, Alhaider AA, Mossa JS, Al-Sohaibani MO, Al-Yahya MA, Rafatullah S, Shaik SA: Gastroprotective effect of an aqueous suspension of black cumin *Nigella sativa* on necrotizing agents-induced gastric injury in experimental animals. *Saudi J Gastroenterol* 2008;14:128.
- 20 Nath KA, Norby SM: Reactive oxygen species and acute renal failure. *Am J Med* 2000;109: 665–678.
- 21 Vaziri ND, Ni Z, Oveisi F, Liang K, Pandian R: Enhanced nitric oxide inactivation and protein nitration by reactive oxygen species in renal insufficiency. *Hypertension* 2002;39:135–141.
- 22 Ha H, Hwang IA, Park JH, Lee HB: Role of reactive oxygen species in the pathogenesis of diabetic nephropathy. *Diabetes Res Clin Pract* 2008;82:S42–S45.
- 23 Sedeek M, Nasrallah R, Touyz RM, Hébert RL: NADPH oxidases, reactive oxygen species, and the kidney: friend and foe. *J Am Soc Nephrol* 2013;24:1512–1518.

- 24 Erkan N, Ayranci G, Ayranci E: Antioxidant activities of rosemary (*Rosmarinus officinalis* L.) extract, blackseed (*Nigella sativa* L.) essential oil, carnosic acid, rosmarinic acid and sesamol. *Food Chem* 2008;110:76–82.
- 25 Shenawy EL, Nahla S, Soliman MF, Reyad SI: The effect of antioxidant properties of aqueous garlic extract and *Nigella sativa* as antischistosomiasis agents in mice. *Rev Inst Med Trop Sao Paulo* 2008;50:29–36.
- 26 Mariod AA, Ibrahim RM, Ismail M, Ismail N: Antioxidant activity and phenolic content of phenolic rich fractions obtained from black cumin (*Nigella sativa*) seedcake. *Food Chem* 2009;116:306–312.
- 27 Chehl N, Chipitsyna G, Gong Q, Yeo CJ, Arafat HA: Anti-inflammatory effects of the *Nigella sativa* seed extract, thymoquinone, in pancreatic cancer cells. *HPB (Oxford)* 2009;11:373–381.
- 28 Umar S, Zargan J, Umar K, Ahmad S, Katiyar CK, Khan HA: Modulation of the oxidative stress and inflammatory cytokine response by thymoquinone in the collagen induced arthritis in Wistar rats. *Chem Biol Interact* 2012;197:40–46.
- 29 Khattab MM, Nagi MN: Thymoquinone supplementation attenuates hypertension and renal damage in nitric oxide deficient hypertensive rats. *Phytother Res* 2007;21:410–414.
- 30 Sayed AA, Morcos M: Thymoquinone decreases AGE-induced NF- κ B activation in proximal tubular epithelial cells. *Phytother Res* 2007;21:898–899.
- 31 Ozer EK, Goktas MT, Toker A, Pehlivan S, Bariskaner H, Ugurluoglu C, et al: Thymoquinone protects against the sepsis induced mortality, mesenteric hypoperfusion, aortic dysfunction and multiple organ damage in rats. *Pharmacol Rep* 2017;69:683–690.
- 32 Faisal R, Shinwari L, Jehangir T: Comparison of the therapeutic effects of thymoquinone and methotrexate on renal injury in pristane induced arthritis in rats. *J Coll Physicians Surg Pak* 2015;25:597–601.
- 33 Hosseinian S, Rad AK, Bideskan AE, Soukhtanloo M, Sadeghnia H, Shafei MN, et al: Thymoquinone ameliorates renal damage in unilateral ureteral obstruction in rats. *Pharmacol Rep* 2017;69:648–657.
- 34 Wever KE, Warlé MC, Wagener FA, van der Hoorn JW, Masereeuw R, van der Vliet JA, Rongen GA: Remote ischaemic preconditioning by brief hind limb ischaemia protects against renal ischaemia-reperfusion injury: the role of adenosine. *Nephrol Dial Transplant* 2011;26:3108–3117.
- 35 Grynberg K, Ozols E, Mulley W, Nikolic-Paterson DJ, Ma FY: Conditional deletion of Jnk signalling in proximal tubular epithelial cells ameliorates renal ischaemia-reperfusion injury. *Nephrology* 2015;20:18.
- 36 Zhang J, Zou YR, Zhong X, Deng HD, Pu L, Peng K, et al: Erythropoietin pretreatment ameliorates renal ischaemia-reperfusion injury by activating PI3K/Akt signalling. *Nephrology* 2015;20:266–272.
- 37 Huang Q, Wang Q, Zhang S, Jiang S, Zhao L, Yu L, et al: Increased hydrogen peroxide impairs angiotensin II contractions of afferent arterioles in mice after renal ischaemia-reperfusion injury. *Acta Physiol* 2016;218:136–145.
- 38 Farag MM, Ahmed GO, Shehata RR, Kazem AH: Thymoquinone improves the kidney and liver changes induced by chronic cyclosporine A treatment and acute renal ischaemia/reperfusion in rats. *J Pharm Pharmacol* 2015;67:731–739.
- 39 Hammad FT, Lubbad L: The effect of thymoquinone on the renal functions following ischemia-reperfusion injury in the rat. *Int J Physiol Pathophysiol Pharmacol* 2016;8:152.
- 40 Awad AS, Kamel R, Sherief MA: Effect of thymoquinone on hepatorenal dysfunction and alteration of CYP3A1 and spermidine/spermine N-1-acetyl-transferase gene expression induced by renal ischaemia-reperfusion in rats. *J Pharm Pharmacol* 2011;63:1037–1042.
- 41 Tervaert TW, Mooyaart AL, Amann K, Cohen AH, Cook HT, Drachenberg CB, et al: Pathologic classification of diabetic nephropathy. *J Am Soc Nephrol* 2010;21:556–563.
- 42 Navarro-González JF, Mora-Fernández C, De Fuentes MM, García-Pérez J: Inflammatory molecules and pathways in the pathogenesis of diabetic nephropathy. *Nat Rev Nephrol* 2011;7:327–340.
- 43 Fried LF, Emanuele N, Zhang JH, Brophy M, Conner TA, Duckworth W, et al: Combined angiotensin inhibition for the treatment of diabetic nephropathy. *N Engl J Med* 2013;369:1892–1903.
- 44 Zhu W, Chen M, Shou Q, Li Y, Hu F: Biological activities of Chinese propolis and Brazilian propolis on streptozotocin-induced type 1 diabetes mellitus in rats. *Evid Based Complement Alternat Med* 2011;2011:468529.
- 45 Ozawa Y, Kurihara T, Sasaki M, Ban N, Yuki K, Kubota S, et al: Neural degeneration in the retina of the streptozotocin-induced type 1 diabetes model. *Exp Diabetes Res* 2011;2011:108328.
- 46 Luippold G, Klein T, Mark M, Grempler R: Empagliflozin. A novel potent and selective SGLT-2 inhibitor, improves glycaemic control alone and in combination with insulin in streptozotocin-induced diabetic rats, a model of type 1 diabetes mellitus. *Diabetes Obes Metab* 2012;14:601–607.
- 47 Abdelmeguid NE, Fakhoury R, Kamal SM, Al Wafai RJ: Effects of *Nigella sativa* and thymoquinone on biochemical and subcellular changes in pancreatic β -cells of streptozotocin-induced diabetic rats. *J Diabetes* 2010;2:256–266.
- 48 Alimohammadi S, Hobbenaghi R, Javanbakht J, Kheradmand D, Mortezaee R, Tavakoli M, et al: Protective and antidiabetic effects of extract from *Nigella sativa* on blood glucose concentrations against streptozotocin (STZ)-induced diabetic in rats: an experimental study with histopathological evaluation. *Diagn Pathol* 2013;8:137.
- 49 Sayed AA: Thymoquinone protects renal tubular cells against tubular injury. *Cell Biochem Funct* 2008;26:374–380.
- 50 Kanter M: Protective effects of thymoquinone on streptozotocin-induced diabetic nephropathy. *J Mol Histol* 2009;40:107–115.
- 51 Sayed AA: Thymoquinone and proanthocyanidin attenuation of diabetic nephropathy in rats. *Eur Rev Med Pharmacol Sci* 2012;16:808–815.
- 52 Omran OM: Effects of thymoquinone on STZ-induced diabetic nephropathy: an immunohistochemical study. *Ultrastruct Pathol* 2014;38:26–33.
- 53 Al-Trad B, Al-Batayneh K, El-Metwally S, Al-hazimi A, Ginawi I, Alaraj M, et al: *Nigella sativa* oil and thymoquinone ameliorate albuminuria and renal extracellular matrix accumulation in the experimental diabetic rats. *Eur Rev Med Pharmacol Sci* 2016;20:2680–2688.
- 54 Lopez-Novoa JM, Quiros Y, Vicente L, Morales AI, Lopez-Hernandez FJ: New insights into the mechanism of aminoglycoside nephrotoxicity: an integrative point of view. *Kidney Int* 2011;79:33–45.
- 55 Shahbazi F, Dashti-Khavidaki S, Khalili H, Lessan-Pezeshki M: Potential renoprotective effects of silymarin against nephrotoxic drugs: a review of literature. *J Pharm Pharm Sci* 2012;15:112–123.
- 56 McWilliam SJ, Antoine DJ, Sabbisetti V, Turner MA, Farragher T, Bonventre JV, et al: Mechanism-based urinary biomarkers to identify the potential for aminoglycoside-induced nephrotoxicity in premature neonates: a proof-of-concept study. *PLoS One* 2012;7:e43809.
- 57 Perazella MA, Moeckel GW: Nephrotoxicity from chemotherapeutic agents: clinical manifestations, pathobiology, and prevention/therapy. *Semin Nephrol* 2010;30:570–581.
- 58 Maru S, Abe T, Shinohara N, Sazawa A, Maruyama S, Harabayashi T, et al: Influence of baseline renal function and dose reduction of nephrotoxic chemotherapeutic agents on the outcome of metastatic urothelial carcinoma: a retrospective study. *Int J Urol* 2012;19:110–116.
- 59 Cascella M, Palma G, Barbieri A, Bimonte S, Amruthraj NJ, Muzio MR, et al: Role of *Nigella sativa* and its constituent thymoquinone on chemotherapy-induced nephrotoxicity: evidences from experimental animal studies. *Nutrients* 2017;9:625.
- 60 Elsherbiny NM, El-Sherbiny M: Thymoquinone attenuates doxorubicin-induced nephrotoxicity in rats: role of Nrf2 and NOX4. *Chem Biol Interact* 2014;223:102–108.
- 61 Ali BH, Al Za'abi M, Shalaby A, Manoj P, Waly MI, Yasin J, et al: The effect of thymoquinone treatment on the combined renal and pulmonary toxicity of cisplatin and diesel exhaust particles. *Exp Biol Med* 2015;240:1698–1707.

- 62 Jalili C, Salahshoor MR, Hoseini M, Roshankhah S, Sohrabi M, Shabanizadeh A: Protective effect of thymoquinone against morphine injuries to kidneys of mice. *Iran J Kidney Dis* 2017;11:142–150.
- 63 Sayed-Ahmed MM, Nagi MN: Thymoquinone supplementation prevents the development of gentamicin-induced acute renal toxicity in rats. *Clin Exp Pharmacol Physiol* 2007;34:399–405.
- 64 Samarghandian S, Azimi-Nezhad M, Mehrad-Majd H, Mirhafez SR: Thymoquinone ameliorates acute renal failure in gentamicin-treated adult male rats. *Pharmacology* 2015;96:112–117.
- 65 El-Sheikh AA, Morsy MA, Abdalla AM, Hamouda AH, Alhaider IA: Mechanisms of thymoquinone hepatorenal protection in methotrexate-induced toxicity in rats. *Mediators Inflamm* 2015;2015:859383.
- 66 Aycan İÖ, Tokgöz O, Tüfek A, Alabalık U, Evliyaoglu O, Turgut H, et al: The use of thymoquinone in nephrotoxicity related to acetaminophen. *Int J Surg* 2015;13:33–37.
- 67 Mabrouk A, Cheikh HB: Thymoquinone ameliorates lead-induced suppression of the antioxidant system in rat kidneys. *Libyan J Med* 2016;11:310–318.
- 68 Sener U, Uygur R, Aktas C, Uygur E, Erboga M, Balkas G, et al: Protective effects of thymoquinone against apoptosis and oxidative stress by arsenic in rat kidney. *Ren Fail* 2016;38:117–123.
- 69 Erboga M, Kanter M, Aktas C, Sener U, Erboga ZF, Donmez YB, et al: Thymoquinone ameliorates cadmium-induced nephrotoxicity, apoptosis, and oxidative stress in rats is based on its anti-apoptotic and anti-oxidant properties. *Biol Trace Elem Res* 2016;170:165–172.
- 70 Fouda AM, Daba MH, Dahab GM, Sharaf el-Din OA: Thymoquinone ameliorates renal oxidative damage and proliferative response induced by mercuric chloride in rats. *Basic Clin Pharmacol Toxicol* 2008;103:109–118.
- 71 Wu P, Wang L, Wang K: Supine versus prone position in percutaneous nephrolithotomy for kidney calculi: a meta-analysis. *Int J Biochem Cell Biol* 2011;43:67–77.
- 72 Okhunov Z, Friedlander JI, George AK, Duty BD, Moreira DM, Srinivasan AK, et al: STONE nephrolithometry: novel surgical classification system for kidney calculi. *Urology* 2013;81:1154–1160.
- 73 Hadjzadeh MA, Mohammadian N, Rahmani Z, Behnam Rassouli F: Effect of thymoquinone on ethylene glycol-induced kidney calculi in rats. *Urol J* 2008;5:149–155.
- 74 Abukhader MM: The effect of route of administration in thymoquinone toxicity in male and female rats. *Indian J Pharm Sci* 2012;74:195–200.
- 75 Al-Ali A, Alkhawajah AA, Randhawa MA, Shaikh NA: Oral and intraperitoneal LD50 of thymoquinone, an active principle of *Nigella sativa*, in mice and rats. *J Ayub Med Coll Abbottabad* 2008;20:25–27.
- 76 Gali-Muhtasib H, Roessner A, Schneider-Stock R: Thymoquinone: a promising anticancer drug from natural sources. *Int J Biochem Cell Biol* 2006;38:1249–1253.
- 77 Ramadan MF: Nutritional value, functional properties and nutraceutical applications of black cumin (*Nigella sativa* L.): an overview. *J Food Sci Technol* 2007;42:1208–1218.