Challenging Lumen-Apposing Metal Stent Removal after Successful Drainage of a Pancreatic Pseudocyst

Athanasios D. Sioulas^a Konstantina Papadaki^a Dimitrios Schizas^b
Ilias Scotiniotis^a

^aDepartment of Gastroenterology, Hygeia Hospital, Athens, Greece; ^b1st Department of Surgery, National and Kapodistrian University of Athens, Laikon Hospital, Athens, Greece

Keywords

Complications · Pancreas · Pseudocyst drainage

Remoção desafiante de prótese metálica de aposição luminal após drenagem com sucesso de pseudoquisto pancreático

Palavras Chave

Complicações · Pâncreas · Drenagem de pseudoquisto

A 45-year-old man with a symptomatic 14-cm-sized pancreatic body pseudocyst following alcohol-induced acute pancreatitis 19 months earlier was treated with placement of a lumen-apposing metal stent. The stent (Hot Axios, Boston Scientific Corporation, Marlborough, MA, USA) had a length of 1 cm and a diameter of 10 mm and was inserted under endosonographic guidance. The

patient experienced immediate relief of symptoms and imaging studies confirmed the complete resolution of the pseudocyst. Endoscopy 11 weeks later revealed epithelial overgrowth of its gastric flange (Fig. 1), though the stent was visible within the gastrocystic fistula (Fig. 2). Stent removal by means of a rat-tooth forceps was performed, but necessitated significant maneuvering in order the gastric flange to be released. No procedure-related complications were encountered. Thus, the recommended 8-week period for lumen-apposing stent removal should not be significantly exceeded, to prevent a “buried stent” syndrome secondary to epithelial overgrowth at its gastric end [1–3].

Disclosure Statement

All authors confirm that there is no conflict regarding this publication.

Fig. 1. Epithelialized stent prior to removal.

Fig. 2. Stent dislocated into the gastrocystic fistula.

References

- 1 Bang JY, Hasan M, Navaneethan U, Hawes R, Varadarajulu S: Lumen-apposing metal stents (LAMS) for pancreatic fluid collection (PFC) drainage: may not be business as usual. *Gut* 2017;66:2054–2056.
- 2 Rodrigues-Pinto E, Grimm IS, Baron TH: Removal of buried gastroduodenal stents after drainage of pancreatic fluid collections: Silence of the LAMS (with video). *Gastrointest Endosc* 2016;83:853–854.
- 3 Patil R, Ona MA, Papafragkakis C, Anand S, Duddempudi S: Endoscopic ultrasound-guided placement of AXIOS stent for drainage of pancreatic fluid collections. *Ann Gastroenterol* 2016;29:168–173.