Mucosal Immune Response in Nasal-Associated Lymphoid Tissue upon Intranasal Administration by Adjuvants

Hiromi Takaki^{a, b} Shingo Ichimiya^b Misako Matsumoto^a Tsukasa Seya^a

^aDepartment of Vaccine Immunology, Graduate School of Medicine, Hokkaido University, Sapporo, Japan;

^bDepartment of Human Immunology, Research Institute for Frontier Medicine, Sapporo Medical University School of Medicine, Sapporo, Japan

Keywords

Immunoglobulin A · Influenza vaccine · Mucosal immunity · Nasal-associated lymphoid tissue

Abstract

The nasal administration of vaccines directed against diseases caused by upper respiratory tract infections of pathogens, such as the influenza virus, mimics the natural infection of pathogens and induces immunoglobulin A (IgA) production in the nasal cavity to effectively protect viral entry. Therefore, the development of a nasally administered vaccine is a research objective. Because the antigenicity of influenza split vaccines is low, nasal inoculation with the vaccine alone does not induce strong IgA production in the nasal cavity. However, the addition of adjuvants activates the innate immune response, enhancing antigen-specific IgA production and the T-cell response. Although the development of suitable adjuvants for nasal vaccinations is in progress, the mechanism by which adjuvants promote the immune response is still unclear. In this review, we discuss the mucosal immune response, especially in the nasal-associated lymphoid tissue, induced in response to the intranasal inoculation of an influenza vaccine and adjuvants in animal models.

© 2018 S. Karger AG, Basel

Introduction

Influenza, caused by the influenza virus, spreads throughout the world every year and mortality in the elderly and infants is high. Therefore, individuals are vaccinated against the virus to prevent the severe pathological conditions it causes. However, subcutaneous inoculation with an influenza vaccine does not prevent the viral infection itself [1]. The influenza virus is susceptible to antigen shift and antigen drift, and the prediction of epidemic strains is also difficult, so many problems must be solved to develop a vaccine that has the desired infection-preventing effect [2].

Nasally administered vaccines that mimic natural infections have been shown to be effective in preventing influenza virus infection [1]. After the intranasal administration of an influenza vaccine, immunoglobulin A (IgA) is secreted into the nasal mucosa and can prevent infection after viral exposure [3]. Furthermore, because IgA is cross-reactive, even if the vaccine strain and the epidemic strain differ, an infection-preventing effect can be induced [4]. However, because safe split vaccines have low antigenicity, it is necessary to administer them intranasally together with an adjuvant to induce IgA production. The selection of an appropriate adjuvant is essential

Table 1. Comparison of nasal and intestinal mucosal immunity

	Nasal mucosal immunity	Intestinal mucosal immunity	Ref.
Inductive sites	NALT	Peyer's patch and isolated lymphoid follicles	7, 12
Effective sites	Upper airways, lacrimal, nasal, and salivary glands	Small and large bowels	7, 12
T cells	Naïve T cell	Memory T cell	18, 19
B cells	CD62L, $\alpha 4\beta 1$, CCR7, CCR10	$\alpha 4\beta 7$, $\alpha 4\beta 1$, CCR9, CCR10	13–17
DCs	CD103 ⁺ CD11b ⁻ CD103 ⁺ CD11b ⁺ PDCA-1 ⁺ pDC CD11b ⁺ CD64 ⁺ F4/80 ⁻	CD103 ⁻ CD11b ⁻ CD103 ⁺ CD11b ⁻ CD8 α +CX3CR1 ⁻ CD103 ⁺ CD11b ⁺ CX3CR1 ^{low} CD103 ⁻ CD11b ⁺ CX3CR1 ^{int}	9, 20, 21
Macrophages	CD11b ⁺ CD64 ⁺ F4/80 ⁺	CD64 ⁺ CX3CR1 ^{hi}	9, 20, 21
Immune response	Th1/Th2	Th2 > Th1	

DCs, dendritic cells; NALT, nasal associated lymphoid tissue.

for the development of a safe and effective vaccine, but the mechanism of action of adjuvants in the nasal mucosa is unclear. New findings have recently been reported regarding the adjuvant effect in the nasal-associated lymphoid tissue (NALT), which is considered to regulate the mucosal immune response in the nasal mucosa and upper respiratory tract. In this paper, the mechanisms of antibody production induced by adjuvants after their nasal administration are discussed.

Nasal-Associated Lymphoid Tissue

NALT is one of the mucosal-associated lymphoid tissues, and is located on the nasal cavity side of the upper palate in rodents, including rats, hamsters, and mice [5]. In rodents, the NALT is considered to be equivalent to Waldeyer's ring and the adenoid, tubal, palatine, and lingual tonsils in humans [5]. In contrast to other mucosal lymph nodes (Peyer's patches), the NALT develops after birth. Although the development of Peyer's patches requires lymphotoxin (LT)-a, LT-b, and LT-b receptor, these molecules are essential for the development of the NALT. Id2^{-/-} mice have an impaired NALT structure [6, 7]. The NALT differs from the lymphoid tissues involved in other mucosal immunity in terms of its organizational development. Furthermore, NALT formation is suppressed in germ-free mice compared with wild-type mice, suggesting that the formation of the NALT is established by an interaction with the indigenous bacteria in the nasal cavity after birth [8].

The NALT, which consists of T cells, B cells, dendritic cells (DCs), macrophages, and microfold cells, is located in the nasal cavity and is the first immune tissue exposed to inhaled antigens and pathogens [5, 9, 10]. Inhaled antigens are transported to the NALT by microfold cells located in the epithelium overlying the NALT and are taken up by DCs, resulting in their presentation to T cells [7]. After T-cell activation, a cytokine environment suitable for IgA is considered to form at the inductive site of mucosal immunity, promoting IgA class switching and affinity maturation in the germinal centers formed in the NALT. Activated T cells and IgA-producing B cells reach the effector site to produce antigen-specific IgA in the nasal cavity to protect it against pathogen invasion [11]. Therefore, the NALT is considered to be the inductive site of the mucosal immune response in the upper respiratory tract.

Intranasal vaccination induces strong local and systemic immune responses in the respiratory and genitourinary tracts but induces only weak immune responses in the local intestinal tract (Table 1) [7, 12]. One of the reasons for directivity of mucosal immunity induction by intranasal route is that mucosal B cells induced at the inductive site have different characteristics. B cells induced by NALT express $\alpha 4\beta 1$ -integrin and CCR10, and therefore strongly migrate to the respiratory and genitourinary tracts highly expressing their ligands, VACM-1 and CCL28 [13–15]. Indeed, B cells co-cultured with lung DCs have been shown to more extensively migrate to the lung than to the intestinal tract [16]. On the other hand, B cells induced by intestinal inductive sites, Peyer's

Fig. 1. Ligands for pattern recognition receptors and signaling pathways. TLRs are present in cell membranes and in endosomes. Surface TLRs recognize components of the bacterial membrane and endosomal TLRs recognize the nucleic acid components of pathogens. TLRs activate the transcription factors NF- κ B, IRF3, and IRF7 via the adaptors MyD88 and TICAM1, leading to the production of type I IFNs and inflammatory cytokines. TLR4, which recognizes lipopolysaccharide, uses both MyD88 and TICAM1 as adaptors, whereas TLR3 uses only TICAM1 as an adaptor. Cytosolic RIG-I and MDA5 sense RNA and signal to activate NF- κ B and IRF3 via MAVS on the mitochondria. Cytosolic DNA

is converted to cGAMP by cyclic GMP-AMP synthase (cGAS) and is sensed by STING, resulting in the activation of NF- κ B, IRF3, and IRF7. Orange and blue colors indicate the molecules involved in the RNA and DNA recognition pathways, respectively. CDN, cyclic dinucleotides; cGAMP, cyclic GMP-AMP; cGAS, cGAMP synthase; IFN, interferon; IRF, interferon regulatory factor; MDA, melanoma differentiation-associated protein; MyD, myeloid differentiation primary response gene; RIG-I, retinoic acid-inducible gene I; STING, stimulator of interferon genes; TICAM1, toll-like receptor adaptor molecule 1; TLR, toll-like receptor.

patches and isolated lymphoid tissues, express CCR9 and $\alpha 4\beta 7$ -integrin, and therefore, preferentially migrate to intestinal tracts expressing their ligands, CCL25 and MAdCAM-1 [15, 17]. Although T cells of NALT show Th0-type dominant cytokine signature and are able to differentiate into Th1 or Th2 cells immediately after antigen inoculation by the nasal route [18], T cells in Peyer's patches show memory phenotype [19]. DCs and macrophages are different between NALT and intestinal tracts (Table 1) [9, 20, 21]. NALT has a low frequency of CD103⁺ DCs, and CD11b⁺CD64⁻F4/80⁻ DCs occupy the majority of DC subsets in NALT. By contrast, Peyer's patches have a high frequency of CD103⁺ DCs [9]. Both CD103⁺ DCs in NALT and intestinal tracts are able to incorporate antigens and migrate to draining lymph nodes. Oral vaccination induces a strong mucosal immune response in intestinal tracts but not in the upper airways. Antigens require modification for oral vaccination to be protected from digestion before reaching the intestinal tract, and oral vaccines tend to induce immune tolerance to anti-

gens [22]. Taken together, in several immunological respects nasal vaccines are superior to oral vaccines against pathogens that infect the respiratory tract.

Mucosal Adjuvants for IgA Production in the Upper Respiratory Tract

Cholera toxin (CT) and *Escherichia coli* heat-labile enterotoxin are major mucosal adjuvants. These toxins activate the immune response to antigens, and the intranasal inoculation of influenza hemagglutinin (HA) antigen with these toxins strongly induces antigen-specific IgA production in the upper respiratory tract [23–26]. CT triggers Th2-type cytokine production [27] and enterotoxin increases the Th1- and Th2-type immune responses, leading to the production of IgA in the mucosa [26]. Derivatives of these toxins, the B subunit of enterotoxin, non-toxic enterotoxin K63, and the B subunit of CT, have been developed to improve the safety of the adjuvants [26, 28].

Table 2. Representative nasally administered influenza vaccine adjuvant and immune responses

Adjuvant	Receptor	Immune responses	Ref.
Cholera toxin, CTB	Ganglioside	IgA induction Protection of viral challenge T cell response	23, 24, 26, 27
<i>Escherichia coli</i> heat-labile enterotoxin Enterotoxin B subunit	Ganglioside	IgA induction Protection of viral challenge	25, 26
PolyI:C	TLR3 RIG-I MDA5	IgA induction Protection of viral challenge T cell response	4
Monophosphoryl lipid A	TLR4	IgA induction T cell response	29
Flagellin	TLR5	IgA induction Protection of viral challenge T cell response	30, 31
Unmethylated CpG oligodeoxynucleotide	TLR9	IgA induction Protection of viral challenge T cell response	32, 33
Cyclic-di-nucleotides	STING	IgA induction Protection of viral challenge T cell response	46
Type I IFN	IFNAR	IgA induction Protection of viral challenge	34, 35

CTB, cholera toxin B subunit; IFN, interferon; IFNAR, interferon alpha receptor; MDA5, melanoma differentiation-associated gene 5; polyI:C, polyinosine-polycytidylic acid; RIG-I, retinoic acid-inducible gene-I; STING, stimulator of interferon genes; TLR, toll-like receptor.

Pathogen-associated molecular patterns (PAMPs) are pathogen molecules sensed by pattern recognition receptors, including the toll-like receptors (TLRs) and RIG-I-like receptors (RLRs), in host cells (Fig. 1). The activation of the innate immune response by PAMPs evokes a subsequent acquired immune response against invading pathogens. Recent studies of the innate immune system have shown that PAMPs function as adjuvants (Table 2). The intranasal inoculation of a split vaccine against the influenza virus together with PAMPs promotes mucosal IgA production. For example, monophosphoryl lipid A, a ligand of toll-like receptor 4 (TLR4), drives the Th1 response and IgA production [29]. Polyinosine-polycytidylic acid (polyI:C), a synthetic analog of double-stranded RNA, is recognized by endosomal TLR3 and cytosolic RLRs, which then activate the innate immune response, resulting in IgA production [4]. Flagellin, a component of bacteria, is recognized by TLR5, which enhances IgA production [30, 31]. Unmethylated CpG oligodeoxynucleotide, a synthetic analog of short single-stranded DNA and

a ligand of TLR9, activates DCs and B cells, resulting in the induction of IgA [32, 33]. These PAMPs induce the activation of immune cells and produce type I interferons (IFNs), which confer resistance to viruses in both myeloid and nonmyeloid cells. Therefore, type I IFNs are also used as intranasal mucosal adjuvants to promote the expression of HA antigen-specific IgA [34, 35]. The activation of DCs by IFNs might be one explanation for the mechanisms underlying the adjuvanticity of IFNs [36]. However, studies of IFN- α receptor knockout (KO) mice have shown that the adjuvanticity of TLR ligands does not necessarily depend on type I IFNs [37, 38].

After antigen stimulation, IgA class switching occurs in B cells via T-cell-dependent (TD) and T-cell-independent (TI) mechanisms [39, 40]. Transforming growth factor (TGF)- β signaling in B cells is necessary for both TD and TI pathways. TD IgA class switching requires binding of CD40 on B cells to CD40L on CD4⁺ T cells. The interaction between CD40 and CD40L leads to the activation of the transcription factor NF- κ B, resulting in the transcrip-

Fig. 2. PolyI:C enhances IgA production in the NALT via the TLR3-TICAM1 pathway. After the intranasal administration of an HA vaccine and polyI:C, CD103⁺ DCs incorporate polyI:C and are activated through the TLR3-TICAM1 pathway. CD103⁺ DCs express BAFF and APRIL to activate B cells. PolyI:C also induces Th2-type cytokines, including IL-4, IL-6, and IL-10, in a TLR3-dependent manner. Furthermore, polyI:C enhances the formation of germinal centers and the expression of the *Aicda* gene, coding

AID protein. Th2-type cytokines and TGF- β signaling promote IgA class switching in the NALT. The secretory IgA produced is secreted to the nasal mucosa through the polymeric immunoglobulin receptor expressed on epithelial cells in the NALT. AID, activation-induced cytidine deaminase; BAFF, B-cell-activating factor; DC, dendritic cell; HA, hemagglutinin; TGF- β , transforming growth factor β ; TLR, toll-like receptor.

tion of the activation-induced cytidine deaminase (*Aicda*) gene, which is an essential factor for class switching [39]. TI IgA class switching requires TLR signaling or the binding of B-cell-activating factor (BAFF) or proliferation-inducing ligand (APRIL) to the transmembrane activator and calcium modulator and cyclophilin ligand interactor (TACI) on B cells rather than the CD40-CD40L interaction required by the TD pathway [39, 40]. TLR signaling and TACI binding induce NF- κ B to transcribe the *Aicda* gene. The IgA induction mechanism in mucosal tissues has been predominantly studied in the intestinal tract, and the mechanism of IgA induction in the nasal mucosa is still largely unclear. However, to develop an efficient vaccine against a virus that enters the host cells in the upper respiratory tract, it is necessary to determine the molecular mechanisms by which the vaccine induces IgA production in the nasal mucosa. The molecular mechanisms by which adjuvants induce IgA in the upper respiratory tract mucosa are becoming clearer.

PolyI:C Induces IgA Production via the TLR3 Pathway

PolyI:C, a synthetic analog of double-stranded RNA, is recognized by endosomal TLR3 and the cytosolic RLRs, RIG-I and MDA5, which induce type I IFN production

via their adaptor molecules, TLR adaptor molecule 1 (TICAM1, also called TRIF) and mitochondrial antiviral signaling protein (MAVS, also called VISA, IPS-1, and Cardif), respectively (Fig. 1) [41]. PolyI:C robustly up-regulated the production of antigen-specific IgA in the nasal mucosa when mice were intranasally inoculated with a combination of an HA split vaccine and polyI:C, protecting against viral challenge not only with the same influenza viral strain but also with different strains [4]. The intranasal inoculation of polyI:C upregulates the production of Th2- and Th1-type cytokines in the NALT [4]. A recent study using KO mice clearly showed that the adjuvanticity of polyI:C during vaccination is dependent on the TLR3-TICAM1 pathway, but not on the RLR-MAVS pathway [37]. IgA and IgG production, the T-cell response, the formation of the germinal centers, and expression of the *Aicda* gene after the intranasal inoculation of a vaccine with polyI:C were significantly impaired in TLR3-KO and TICAM1-KO mice, but not in MAVS-KO mice. The intranasally inoculated polyI:C was incorporated into the NALT CD103⁺ DCs, which express TLR3, and colocalized with endosomal TLR3. BAFF expression was induced on NALT CD103⁺ DCs by polyI:C in a TLR3-TICAM1-dependent manner. Mice lacking CD103⁺ DCs showed reduced vaccine-specific Ig production and a reduced T-cell response. Together, these data indicate that

the intranasal inoculation of polyI:C exerts its adjuvant activity by activating the TLR3-TICAM1 pathway in the NALT CD103⁺ DCs and the expression of cytokines that skew IgA production in the NALT (Fig. 2) [37]. In terms of adverse effects, an adjuvant that activates only TLR3 is expected because the expression of TLR3 is restricted among the myeloid cells to CD103⁺ and CD8α⁺ DCs [42, 43]. A TLR3 adjuvant caused the regression of tumor growth in mice without a cytokine stream [44].

Cyclic-di Nucleotides Induce IgA Production in a STING-Dependent Manner

Cyclic di-nucleotides (CDNs), including cyclic (c) guanosine monophosphate (GMP)–adenosine monophosphate (AMP) (cGAMP), c-di-GMP, and c-di-AMP, are recognized by STING, which is located on the endoplasmic reticulum membrane. CDNs exert potent adjuvant activity on the vaccine effect, including Ig production, the T-cell response, and the formation of germinal centers, via the STING pathway [38, 45–47]. The adjuvant activity of c-di-GMP is more effective than that of cGAMP because it enhances antigen uptake by DCs and the selective activation of pinocytosis-efficient cells [38]. CD11c-specific STING-KO mice displayed impaired Ig production and a defective T-cell response after intranasal inoculation with c-di-GMP [45], so the adjuvant activity of c-di-GMP mainly depends on the activation of STING-expressing DCs. Although c-di-GMP also induces interleukin 33 and thymic stromal lymphopoietin TSLP in epithelial cells via the STING-independent pathway, the molecular mechanism of STING-independent cytokine production is unknown.

Conclusions

Although intranasal vaccination is less painful and simpler than previous vaccines using syringes and needles, these vaccines need strong adjuvants without side effects. Successful intranasal vaccination requires the development of appropriate adjuvants, but it is essential to clarify the mechanism of action of adjuvants as a preliminary step. Antigen-producing B cells induced by the mucosal vaccine accumulate in the effector site close to the inductive site, so it is very important to vaccinate the site where pathogens are likely to invade. Due to directivity of mucosal immunity induction by intranasal route, the development of a nasal administration vaccine is expected as a vaccine against pathogens causing upper respiratory tract infection such as the influenza virus, RS virus, *Streptococcus pneumoniae*, and *Haemophilus influenzae*. Since nasal vaccination also upregulates IgA production in genitourinary tracts, it may be one of the effective routes for the administration of vaccines against HIV and HPV.

Acknowledgments

This research was supported by the Japan Agency for Medical Research and Development (grant No. 17fk0108212h0302) and the Japan Society for the Promotion of Science (grant No. 16K19134).

Disclosure Statement

The authors declare no conflict of interest.

References

- Houser K, Subbarao K: Influenza vaccines: challenges and solutions. *Cell Host Microbe* 2015;17:295–300.
- Iwasaki A: Exploiting mucosal immunity for antiviral vaccines. *Annu Rev Immunol* 2016;34:575–608.
- Tamura S, Ainal A, Suzuki T, Kurata T, Hasegawa H: Intranasal inactivated influenza vaccines: a reasonable approach to improve the efficacy of influenza vaccine? *Jpn J Infect Dis* 2016;69:165–179.
- Ichinohe T, Watanabe I, Ito S, Fujii H, Moriyama M, Tamura S, Takahashi H, Sawa H, Chiba J, Kurata T, Sata T, Hasegawa H: Synthetic double-stranded RNA poly(I:C) combined with mucosal vaccine protects against influenza virus infection. *J Virol* 2005;79:2910–2919.
- Heritage PL, Underdown BJ, Arsenault AL, Snider DP, McDermott MR: Comparison of murine nasal-associated lymphoid tissue and Peyer's patches. *Am J Respir Crit Care Med* 1997;156:1256–1262.
- Krege J, Seth S, Hardtke S, Davalos-Misslitz AC, Förster R: Antigen-dependent rescue of nose-associated lymphoid tissue (NALT) development independent of LTβR and CXCR5 signaling. *Eur J Immunol* 2009;39:2765–2778.
- Kiyono H, Fukuyama S: NALT – versus Peyer's-patch-mediated mucosal immunity. *Nat Rev Immunol* 2004;4:699–710.
- Jain R, Waldvogel-Thurlow S, Darveau R, Douglas R: Differences in the paranasal sinuses between germ-free and pathogen-free mice. *Int Forum Allergy Rhinol* 2016;6:631–637.
- Lee H, Ruane D, Law K, Ho Y, Garg A, Rahman A, Esterházy D, Cheong C, Goljo E, Sikora AG, Mucida D, Chen BK, Govindraj S, Breton G, Mehandru S: Phenotype and function of nasal dendritic cells. *Mucosal Immunol* 2015;8:1083–1098.
- Mutoh M, Kimura S, Takahashi-Iwanaga H, Hisamoto M, Iwanaga T, Iida J: RANKL regulates differentiation of microfold cells in mouse nasopharynx-associated lymphoid tissue (NALT). *Cell Tissue Res* 2016;364:175–184.
- Wu HY, Nguyen HH, Russell MW: Nasal lymphoid tissue (NALT) as a mucosal immune inductive site. *Scand J Immunol* 1997;46:506–513.

- 12 Yanagita M, Hiroi T, Kitagaki N, Hamada S, Ito HO, Shimauchi H, Murakami S, Okada H, Kiyono H: Nasopharyngeal-associated lymphoreticular tissue (NALT) immunity: fimbriae-specific Th1 and Th2 cell-regulated IgA responses for the inhibition of bacterial attachment to epithelial cells and subsequent inflammatory cytokine production. *J Immunol* 1999;162:3559–3565.
- 13 Lazarus NH, Kunkel EJ, Johnston B, Wilson E, Youngman KR, Butcher EC: A common mucosal chemokine (mucosae-associated epithelial chemokine/CCL28) selectively attracts IgA plasmablasts. *J Immunol* 2003;170:3799–3805.
- 14 Kunkel EJ, Kim CH, Lazarus NH, Viera MA, Soler D, Bowman EP, Butcher EC: CCR10 expression is a common feature of circulating and mucosal epithelial tissue IgA Ab-secreting cells. *J Clin Invest* 2003;111:1001–1010.
- 15 Johansen FE, Baekkevold ES, Carlsen HS, Farstad IN, Soler D, Brandtzaeg P: Regional induction of adhesion molecules and chemokine receptors explains disparate homing of human B cells to systemic and mucosal effector sites: dispersion from tonsils. *Blood* 2005;106:593–600.
- 16 Ruane D, Chorny A, Lee H, Faith J, Pandey G, Shan M, Simchoni N, Rahman A, Garg A, Weinstein EG, Oropallo M, Gaylord M, Ungaro R, Cunningham-Rundles C, Alexandropoulos K, Mucida D, Merad M, Cerutti A, Mehndru S: Microbiota regulate the ability of lung dendritic cells to induce IgA class-switch recombination and generate protective gastrointestinal immune responses. *J Exp Med* 2016;213:53–73.
- 17 Kunkel EJ, Butcher EC: Plasma-cell homing. *Nat Rev Immunol* 2003;3:822–829.
- 18 Hiroi T, Iwatani K, Iijima H, Kodama S, Yanagita M, Kiyono H: Nasal immune system: distinctive Th0 and Th1/Th2 type environments in murine nasal-associated lymphoid tissues and nasal passage, respectively. *Eur J Immunol* 1998;28:3346–3353.
- 19 Chao CC, Sandor M, Dailey MO: Expression and regulation of adhesion molecules by gamma delta T cells from lymphoid tissues and intestinal epithelium. *Eur J Immunol* 1994;24:3180–3187.
- 20 Persson EK, Scott CL, Mowat AM, Agace W: Dendritic cell subsets in the intestinal lamina propria: ontogeny and function. *Eur J Immunol* 2013;43:3098–3107.
- 21 Cerovic V, Bain CC, Mowat AM, Milling SW: Intestinal macrophages and dendritic cells: what's the difference? *Trends Immunol* 2014;35:270–277.
- 22 Azizi A, Kumar A, Diaz-Mitoma F, Mestecky J: Enhancing oral vaccine potency by targeting intestinal M cells. *PLoS Pathog* 2010;6:e1001147.
- 23 Tamura S, Samegai Y, Kurata H, Nagamine T, Aizawa C, Kurata T: Protection against influenza virus infection by vaccine inoculated intranasally with cholera toxin B subunit. *Vaccine* 1988;6:409–413.
- 24 Tamura S, Yamanaka A, Shimohara M, Tomita T, Komase K, Tsuda Y, Suzuki Y, Nagamine T, Kawahara K, Danbara H: Synergistic action of cholera toxin B subunit (and *Escherichia coli* heat-labile toxin B subunit) and a trace amount of cholera whole toxin as an adjuvant for nasal influenza vaccine. *Vaccine* 1994;12:419–426.
- 25 Tamura S, Asanuma H, Tomita T, Komase K, Kawahara K, Danbara H, Hattori N, Watanabe K, Suzuki Y, Nagamine T: *Escherichia coli* heat-labile enterotoxin B subunits supplemented with a trace amount of the holotoxin as an adjuvant for nasal influenza vaccine. *Vaccine* 1994;12:1083–1089.
- 26 Pizza M, Giuliani MM, Fontana MR, Monaci E, Douce G, Dougan G, Mills KH, Rappuoli R, Del Giudice G: Mucosal vaccines: non toxic derivatives of LT and CT as mucosal adjuvants. *Vaccine* 2001;19:2534–2541.
- 27 Matsuo K, Iwasaki T, Asanuma H, Yoshikawa T, Chen Z, Tsujimoto H, Kurata T, Tamura SS: Cytokine mRNAs in the nasal-associated lymphoid tissue during influenza virus infection and nasal vaccination. *Vaccine* 2000;18:1344–1350.
- 28 Barchfeld GL, Hessler AL, Chen M, Pizza M, Rappuoli R, Van Nest GA: The adjuvants MF59 and LT-K63 enhance the mucosal and systemic immunogenicity of subunit influenza vaccine administered intranasally in mice. *Vaccine* 1999;17:695–704.
- 29 Baldridge JR, Yorgensen Y, Ward JR, Ulrich JT: Monophosphoryl lipid A enhances mucosal and systemic immunity to vaccine antigens following intranasal administration. *Vaccine* 2000;18:2416–2425.
- 30 Wang BZ, Xu R, Quan FS, Kang SM, Wang L, Compans RW: Intranasal immunization with influenza VLPs incorporating membrane-anchored flagellin induces strong heterosubtypic protection. *PLoS One* 2010;5:e13972.
- 31 Hong SH, Byun YH, Nguyen CT, Kim SY, Seong BL, Park S, Woo GJ, Yoon Y, Koh JT, Fujihashi K, Rhee JH, Lee SE: Intranasal administration of a flagellin-adjuvanted inactivated influenza vaccine enhances mucosal immune responses to protect mice against lethal infection. *Vaccine* 2012;30:466–474.
- 32 Joseph A, Louria-Hayon I, Plis-Finarov A, Zeira E, Zakay-Rones Z, Raz E, Hayashi T, Takabayashi K, Barenholz Y, Kedari E: Liposomal immunostimulatory DNA sequence (ISS-ODN): an efficient parenteral and mucosal adjuvant for influenza and hepatitis B vaccines. *Vaccine* 2002;20:3342–3354.
- 33 Moldoveanu Z, Love-Homan L, Huang WQ, Krieg AM: CpG DNA, a novel immune enhancer for systemic and mucosal immunization with influenza virus. *Vaccine* 1998;16:1216–1224.
- 34 Cao M, Sasaki O, Yamada A, Imanishi J: Enhancement of the protective effect of inactivated influenza virus vaccine by cytokines. *Vaccine* 1992;10:238–242.
- 35 Bracci L, Canini I, Puzelli S, Sestili P, Venditti M, Spada M, Donatelli I, Belardelli F, Proietti E: Type I IFN is a powerful mucosal adjuvant for a selective intranasal vaccination against influenza virus in mice and affects antigen capture at mucosal level. *Vaccine* 2005;23:2994–3004.
- 36 Tovey MG, Lallemand C, Meritet JF, Maury C: Adjuvant activity of interferon alpha: mechanism(s) of action. *Vaccine* 2006;24(suppl 2):–46–47.
- 37 Takaki H, Kure S, Oshiumi H, Sakoda Y, Suzuki T, Aina A, Hasegawa H, Matsumoto M, Seya T: Toll-like receptor 3 in nasal CD103⁺ dendritic cells is involved in immunoglobulin A production. *Mucosal Immunol* 2018;11:82–96.
- 38 Blaauboer SM, Gabrielle VD, Jin L: MPYS/STING-mediated TNF- α , not type I IFN, is essential for the mucosal adjuvant activity of (3'-5')-cyclic-di-guanosine-monophosphate in vivo. *J Immunol* 2014;192:492–502.
- 39 Cerutti A: The regulation of IgA class switching. *Nat Rev Immunol* 2008;8:421–434.
- 40 Bemarck M, Boysen P, Lycke NY: Induction of gut IgA production through T cell-dependent and T cell-independent pathways. *Ann NY Acad Sci* 2012;1247:97–116.
- 41 Kawai T, Akira S: Antiviral signaling through pattern recognition receptors. *J Biochem* 2007;141:137–145.
- 42 Edwards AD, Diebold SS, Slack EM, Tomizawa H, Hemmi H, Kaisho T, Akira S, Reis e Sousa C: Toll-like receptor expression in murine DC subsets: lack of TLR7 expression by CD8 α^+ DC correlates with unresponsiveness to imidazoquinolines. *Eur J Immunol* 2003;33:827–833.
- 43 Takaki H, Oshiumi H, Matsumoto M, Seya T: Dendritic cell subsets involved in type I IFN induction in mouse measles virus infection models. *Int J Biochem Cell Biol* 2014;53:329–333.
- 44 Matsumoto M, Tatematsu M, Nishikawa F, Azuma M, Ishii N, Morii-Sakai A, Shime H, Seya T: Defined TLR3-specific adjuvant that induces NK and CTL activation without significant cytokine production in vivo. *Nat Commun* 2015;6:6280.
- 45 Blaauboer SM, Mansouri S, Tucker HR, Wang HL, Gabrielle VD, Jin L: The mucosal adjuvant cyclic di-GMP enhances antigen uptake and selectively activates pinocytosis-efficient cells in vivo. *Elife* 2015;4.
- 46 Madhun AS, Haaheim LR, Nøstbakken JK, Ebensen T, Chichester J, Yusibov V, Guzman CA, Cox RJ: Intranasal c-di-GMP-adjuvanted plant-derived H5 influenza vaccine induces multifunctional Th1 CD4⁺ cells and strong mucosal and systemic antibody responses in mice. *Vaccine* 2011;29:4973–4982.
- 47 Takaki H, Takashima K, Oshiumi H, Aina A, Suzuki T, Hasegawa H, Matsumoto M, Seya T: cGAMP promotes germinal center formation and production of IgA in nasal-associated lymphoid tissue. *Med Sci (Basel)* 2017;5.